



# City Research Online

## City, University of London Institutional Repository

---

**Citation:** Pickard, S., Cluley, V., Danely, J., Laceulle, H., Leon-Salas, J., Vanhoutte, B. and Romero-Ortuno, R. (2019). New horizons in frailty: the contingent, the existential and the clinical. *Age and Ageing*, 48(4), pp. 466-471. doi: 10.1093/ageing/afz032

This is the accepted version of the paper.

This version of the publication may differ from the final published version.

---

**Permanent repository link:** <https://openaccess.city.ac.uk/id/eprint/22455/>

**Link to published version:** <http://dx.doi.org/10.1093/ageing/afz032>

**Copyright and reuse:** City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

---

City Research Online:

<http://openaccess.city.ac.uk/>

[publications@city.ac.uk](mailto:publications@city.ac.uk)

---



**New Horizons in Frailty: the contingent, the existential and the clinical**

Journal:	<i>Age and Ageing</i>
Manuscript ID	AA-19-0199
Manuscript Category:	New Horizons
Keywords:	frailty, operationalisation, lived experience, social science, first-person perspective
Keypoints:	The rapid implementation of the frailty paradigm in clinical settings may have unintended consequences, These include disconnecting geriatric medicine from its roots in a holistic approach to health and illness in old age, They also introduce a disjunction between clinical views and patient experience of frailty, Methods drawn from the social sciences and humanities may prove useful in addressing this, They can be introduced into a multidisciplinary toolkit for practitioners across a range of settings

## New Horizons in Frailty: the contingent, the existential and the clinical

### Abstract

The increasing operationalisation of frailty including in primary care brings with it a risk of oversimplifying diagnostic processes and inadvertently barring access to comprehensive geriatric assessment services. The emphasis on measurement tools, despite updated guidance to the contrary, can also undermine the importance of clinical judgement, meaning that non-contractual evidence-based opportunities for medical optimisation (e.g. exercise promotion, nutrition optimisation) as well as appropriate access to social supports may be missed.

Another significant consequence is to open up an ever-widening gap between clinical approaches and lived experience, including overlooking strengths and resources in problem-based approach and inadvertently alienating the very group it is trying to assist.

This article argues that the next horizon in frailty should not be to continue to refine its operationalisations but rather to reconnect with more holistic approaches to health and illness in old age, which are more in tune with the tradition of geriatric medicine. Building on recent trends to shift conceptualisation of health and illness in old age, this project would benefit from sociological and humanities-based approaches that foreground older people's lived experience using theoretically informed description focused on a first-person perspective. In reconfiguring what we think frailty is, such a shift will bring us closer to the views and experience of older people who live with frailty. We see a brighter horizon for frailty in that direction.

## Introduction

In the clinical literature, for the past decade frailty has been dominated by two approaches: the phenotype model, proposed by Linda Fried and colleagues, and the Frailty Index proposed by Kenneth Rockwood's group, with a vast and multiplying literature devoted to applying and reaffirming one or other of those two positions. Both of these are centred on signs and symptoms of the individual body. Following the 2017 GMS contract, all primary care practices have been incentivised to identify older patients with moderate or severe frailty, extending the use of these and similar 'appropriate' tools into primary care. One of the unintended consequences of the dominance of the operationalisation of frailty in either of these forms include clinical assumptions that frailty is a diagnosis, rather than a syndrome or state caused by other factors, which brings the danger of oversimplifying diagnostic processes and inadvertently barring access to comprehensive geriatric assessment services. The emphasis on measurement tools, despite updated guidance to the contrary [1], can also undermine the importance of clinical judgement, meaning that non-contractual evidence-based opportunities for medical optimisation (e.g. exercise promotion, nutrition optimisation) as well as appropriate access to social supports may be missed. Another significant consequence is to open up an ever-widening gap between clinical approaches and lived experience and inadvertently alienate the very group it is trying to assist.

In this article we argue that the next horizon in frailty should not be to continue to refine its operationalisations and have them compete for their rapid implementation into clinical practice, but rather to take a step back, look critically for unintended consequences, and reconnect with more holistic approaches to health and illness in old age, which are more in tune with the tradition of geriatric medicine. As well as building on recent trends [2] to shift

1  
2  
3 conceptualisation of health and illness in old age, in so doing it would benefit from  
4  
5 sociological and humanities-based approaches that foreground older people's lived  
6  
7 experience using theoretically informed description focused on a first-person perspective. In  
8  
9 reconfiguring what we think frailty is such a shift will also bring us closer to the views and  
10  
11 experience of older people who live with frailty. We see a brighter horizon for frailty in that  
12  
13 direction.  
14  
15

### 20 **The disjunction between clinical approaches and lived experience**

21  
22  
23 There is an emerging body of research, coming from the social sciences, humanities, social  
24  
25 work and nursing and focused on empirical work with older people themselves, that is highly  
26  
27 critical of clinical representations of frailty. This critique focuses, inter alia, on its failure to  
28  
29 take into account the role played in precipitating frailty by domains outside the biological,  
30  
31 including social disadvantage [3]; its overlooking of positive aspects of frail  
32  
33 embodiment/resilience [4]; its incorporation of psycho-social aspects in a dualistic framework  
34  
35 that posits the separation of mind and body [5]; its perpetuation of decline narratives; and  
36  
37 ultimately its potentially (albeit unintentionally) harmful impact on older people who are  
38  
39 labelled 'frail', which may in turn trigger frailty identity crises [6] precipitating a 'cycle of  
40  
41 decline' [7]. The importance of subjectivity is also highlighted in clinical literature: for  
42  
43 example, negative representations of ageing may underpin a decline in objective walking  
44  
45 speed [8]; in addition, the English Longitudinal Study of Ageing suggested that older people  
46  
47 who have a more positive attitude to ageing may be at reduced risk of becoming physically  
48  
49 frail or pre-frail [9].  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 In the wider social domain, the influence of frailty discourses on reconfiguring old age have  
4 been significant, separating the 'third' age (productive, healthy, associated with 'successful'  
5 ageing) from the fourth age (devalued, decrepit, senescent, 'geriatric'), into which latter  
6 category a frailty label may serve as a key gateway. The fourth age as a social imaginary [10]  
7 undermines any concept that old age may contain value and meaning. It stresses separation  
8 rather than continuity both from other (more 'successfully' ageing) older people and from  
9 experiences of vulnerability common through the life course including those precipitated by  
10 structural conditions configuring socio-economic insecurity and increased social inequality  
11 that undermine well-being and increase poor health as summed up in the term 'precarity'  
12 [11]. For all these reasons, older people themselves fear and resist the label, thus potentially  
13 alienating them from practitioners and 'frailty' services. This has resulted in calls in some  
14 quarters for the term to be dropped altogether [12] or for its use to be limited *because* of its  
15 significant limitations, calling instead for a broader strengths-based approach to older  
16 people's health and illness, 'defined by what they can, as much as what they cannot, do' [13]  
17 and thinking "diversabilities" rather than just "disabilities".  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40

41 However, a similar understanding is emerging in medicine. This is reflected in the emerging  
42 concept of "intrinsic capacity". Defined in the World Health Organisation's (WHO) World  
43 Report on Ageing and Health [14] as the combination of physical and cognitive attributes of  
44 an individual, its interplay with the individual's social environment shapes the functional  
45 ability that the WHO posits as the definition of healthy ageing [2]. This concept moves away  
46 from a disease-centred approach to one that emphasizes the positive attributes that ordinarily  
47 support 'health' in the sense of being able to be and do what one values [15]. Professional  
48 assessment within this framework would require a longitudinal observational approach rather  
49 than a single 'snapshot' assessment; intervention can then move from a preventive or reactive  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 focus to an emphasis on supporting the everyday capacity of the individual. This conceptual  
4 apparatus has been hailed as a potentially revolutionary transformation to viewing health and  
5 illness in old age [2]; as such it is well-placed to capture the complexity of frailty, both as a  
6 clinical state and as a lived experience, as well as to support interventions aimed at restoring  
7 functionality.  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17

### 18 **The expertise of geriatric medicine: clinical and social**

19  
20  
21 Whilst agreeing with the rationale behind the critiques of frailty as it is currently  
22 operationalised – particularly the disjunction between clinical approaches and lived  
23 experience that has emerged- we are not advocating that the term be dropped. Clearly frailty  
24 exists and signifies something ‘real’. Nor do we think clinical ‘frailty’ is necessarily and  
25 inherently too limited to be able to encompass lived experience, even though since its  
26 operationalisation over the past two decades it has been almost entirely focused on  
27 physiological or functional limitations. Indeed, our view is that frailty is broader than clinical  
28 symptoms and encompasses more than a medical syndrome alone and that geriatric medicine  
29 has the particular knowledge-base to overcome this disjunction. Historically, geriatric  
30 medicine has roots in the understanding that social and medical problems are inextricably  
31 intertwined in the lives of older patients and indeed that this linkage forms part of the unique  
32 nature of health and illness in old age, recognition of which is part of the unique expertise of  
33 geriatrics. As found in the works of pioneering geriatricians such as Amulree, Sheldon,  
34 Brocklehurst and others, going all the way back to Marjory Warren, the discipline was  
35 established to take a holistic approach, remedy as much as possible and focus on restoring  
36 function in such a way as to enable patients to return to their own homes and their lives [16].  
37 Revisiting this may inspire the reintegration of not just the social and medical, as advocated  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3 by the WHO's approach to healthy ageing, but also of a rapprochement between clinical  
4  
5 knowledge and lay experience of frailty.  
6  
7  
8  
9

10  
11 There is recognition of the importance of a more holistic approach within the field of frailty  
12  
13 itself, especially through CGA [17]. Firstly, running alongside the trend to operationalise  
14  
15 frailty in terms of individual and largely physiological deficits is the agreement that frailty is  
16  
17 a state of pre-existing vulnerability which, combined with a stressor of some kind (either  
18  
19 illness or an external crisis or both), can have significant negative consequences [18, 19].  
20  
21 Campbell and Buchner point out: 'the interaction of the individual with the environment is  
22  
23 central to this concept of frailty' [18]. Although the kinds of stressors discussed usually relate  
24  
25 to health – a minor illness or infection, starting new medication and so on [19] - this  
26  
27 conceptual framework is elastic enough theoretically to encompass events in the social world  
28  
29 that are contingent (such as falls, illness and other risks arising from systemic and structural  
30  
31 conditions) or existential (dilemmas that affect everyone as well as those that are specific to  
32  
33 old age, such as bereavement, grief, loneliness, loss of meaning or value) [20]. The  
34  
35 connection between social events and health is corroborated in neuroendocrinology [21] and  
36  
37 potentially traceable through biomarkers [12] conceptualised through the allostatic load [23].  
38  
39 This framework can also take into account the strengths and resources that co-exist and which  
40  
41 may require additional support at this time.  
42  
43  
44  
45  
46  
47  
48  
49  
50

51  
52 Secondly, rather than seeing 'frail elderly people as complex systems on the brink of failure'  
53  
54 [24] there is increasing recognition that frailty is dynamic, involving improvement as well as  
55  
56 continued decline [25] and that over time, indeed, the 'usual' pattern of frailty involves older  
57  
58 people moving between pre-frail, robust and frail positions [25]. Thus the 'brink' may in  
59  
60

1  
2  
3 many instances be one of renewed or improved health. Given that prefrail older people are far  
4 more likely to move to the robust position than are those already classified as frail  
5  
6 (approximately 25 per cent compared to 3 per cent respectively) [25], this suggests the  
7  
8 importance of interventions that are both sensitive and timely. However, currently in primary  
9  
10 care the obligation on GPs, following routine identification, is to intervene only in cases of  
11  
12 severe frailty, thus missing an opportunity to delay or reverse frailty earlier in its trajectory  
13  
14 [1]. Moreover, there is no consistent approach to the kinds of interventions that GPs should  
15  
16 utilise. A recent review of the evidence [26] suggest that a combination of strength exercises  
17  
18 and protein supplementation is best for a combination of ease and effectiveness in improving  
19  
20 frailty and suggests that a toolkit should be made available to GPs summarising the most  
21  
22 effective interventions. Missing from this mooted toolkit, however, is any reference to social  
23  
24 support dimensions.  
25  
26  
27  
28  
29  
30  
31  
32  
33

### 34 **Bridging the gap**

35  
36  
37 Sociological studies comprising interviews and thick observational data report that older  
38  
39 people talk about adapting to and indeed thriving in the ‘ordinary’ conditions of ageing,  
40  
41 involving ‘diminished everyday competence’ [27] which comprises a range of ‘problems’ all  
42  
43 the way from weaker sight and hearing to stiff joints and frailty. Part of the experience of  
44  
45 adaptation is the learning of new skills, ways of coping with reduced strength and the  
46  
47 challenges of the environment. Disruption of this everyday competence, for example through  
48  
49 the occurrence of contingent or existential events is what leads to felt frailty. A mixture of  
50  
51 appropriate interventions at this point can support the older person to return to a pre-frail or  
52  
53 robust state rather than fluctuate into further frailty. The experience of older people strongly  
54  
55 suggests that, although frailty is a severe diminishment of everyday competence, it is still part  
56  
57  
58  
59  
60

1  
2  
3 of that continuum, can be reversed or delayed, and is not, therefore, something separate. This  
4  
5 is what is recognised in the clinical arena in claims that the aim of frailty interventions is to  
6  
7 stabilise frailty and reduce the more general vulnerability it signals [19] (rather than, for  
8  
9 example, 'combat' frailty). Earlier versions of the frailty index directly reference John  
10  
11 Brocklehurst's 'balance beam model' which incorporates social and relational factors into its  
12  
13 understanding of frailty as a dynamic state [28]. The task then for geriatricians is to go from  
14  
15 acknowledging this in the abstract, viewing biology and the social as important but separate  
16  
17 in the 'frailty syndrome', or seeing its existence as lying beyond the domain of medicine, to  
18  
19 an appreciation of this 'damage' as something that involves not just cells, tissues and organs  
20  
21 (where it might have been present for a considerable length of time) but the lived self. The  
22  
23 clinical gaze intuitively grasps these principles, in the 'know it when I see it' approach but  
24  
25 this 'art' of medicine is in danger of being subsumed in the increasingly technical approaches  
26  
27 that characterise approaches to frailty detection and management, especially in busy settings.  
28  
29  
30  
31  
32  
33  
34  
35  
36

### 37 **The contribution of approaches from social sciences and the humanities to bridging the** 38 39 **gap** 40 41

42 In this aim, medicine can learn from sociological approaches to the lived experience of health  
43  
44 and illness. A well-established interpretive tradition critiques the reductionist nature of  
45  
46 biomedical readings of the body and disease, and stresses the importance of embodied,  
47  
48 biographical and social factors in the illness experience. Narrative-based methods can aid the  
49  
50 geriatrician in understanding how illness is experienced as a biographical event related to  
51  
52 meaning and identity [29, 30]. Phenomenological methods, influenced by the work of  
53  
54 Merleau-Ponty, can aid the geriatrician to understand health and illness from the perspective  
55  
56 not of medicine but of the older person [31]. It contrasts the objective anatomical body  
57  
58  
59  
60

1  
2  
3 (Körper) with that of the lived body (Leib), the biological body with the existential body, and  
4  
5 stresses the importance of the latter in understanding how a person experiences illness,  
6  
7 including in its fluctuations [32]. Phenomenology aims at ‘direct description of our  
8  
9 experience as it is’[33], although more sociologically informed accounts look not so much for  
10  
11 ‘essential’ structures of experience, as sought by philosophers, but are more sensitised to the  
12  
13 way experiences are structured through social forces, which can in turn increase vulnerability,  
14  
15 highlighting the role of age, gender, class and so on in shaping and mediating the trajectories  
16  
17 of frailty. Interpretive concepts and methods can explain how, for example, the ‘normality’ of  
18  
19 long-term conditions and other illness may be experienced more in the oldest patients than in  
20  
21 the young-old for whom sudden illness/impairment may feel like a biographical ‘rupture’,  
22  
23 with adverse consequences for coping and well-being [34, 35]. Both experience and onset of  
24  
25 frailty have been shown to be strongly dependent on life history, with accumulated  
26  
27 disadvantages leading to an earlier experience of frailty and poorer outcomes [36].  
28  
29 Interpretive methods can tease out if and how, under such circumstances, the social context in  
30  
31 which frailty arises might be more impactful on the subsequent frailty trajectory than the  
32  
33 health crisis which precipitated it. Similarly, it is known that gender is a factor in the  
34  
35 experience of frailty in that whilst more women than men are frail, they are also more likely  
36  
37 to fluctuate between health and frailty, although frailty is less of a ‘risk’ for them, at all  
38  
39 stages, than men in a paradox that is currently not well understood [2]. Exploring lived  
40  
41 experience through an interpretive lens may shed light on this paradox and help tailor  
42  
43 interventions accordingly. For example, different interventions are appropriate for contingent  
44  
45 and existential frailty factors respectively. The former are potentially modifiable by  
46  
47 multidisciplinary interventions to address the risk of falls, poor nutrition, ensure aids and  
48  
49 adaptations; what is needed for the latter, by contrast, is social, relational and spiritual  
50  
51 support [37]. These are promising new horizons for frailty.  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6 A phenomenological toolkit for use by both patients/carers and clinicians to increase mutual  
7  
8 understanding and communication was developed by the philosopher Havi Carel [38]. It  
9  
10 includes three elements: (i) shifting the focus away from the disease entity toward the  
11  
12 experience of illness; (ii) highlighting the contrasting perspectives of clinicians, patient,  
13  
14 carers in ways that help mutual understanding; (iii) recognising and identifying the ways in  
15  
16 which illness changes the experience of one's body, identity, relationships, interaction with  
17  
18 the environment, habits, routine, meanings, values and norms. Such a toolkit acknowledges  
19  
20 the importance of both the biological and social but also, and most importantly perhaps, the  
21  
22 interweaving of both as factors jointly underpinning an older person's movement along the  
23  
24 continuum of robust, pre-frail and frail. In so doing it opens up the chance to restore healthy  
25  
26 ageing by sensitive means in appropriate domains in (iii).  
27  
28  
29  
30  
31  
32  
33  
34

35 CGA is a highly sensitive vehicle for delivering such interventions and the introduction of a  
36  
37 CGA toolkit into primary care is thus a promising development. However, there is still a  
38  
39 danger that frailty is decoupled from CGA and thus it will be important to ensure that these  
40  
41 principles are not just applied in older people with frailty syndromes of any stage, or with  
42  
43 severe frailty [17], but that they inform the understanding of health, illness and normality in  
44  
45 all older patients, as well as the concept of, and practices around, frailty itself. A  
46  
47 reconfiguring of the understanding of and approach to frailty to include a broader approach  
48  
49 lies at the heart of a reconceptualisation of health and illness in old age, which in turn gives  
50  
51 geriatric medicine, as it is deployed by all branches of medicine [39] a key role in an ageing  
52  
53 world, not just in terms of intervening in frailty but in supporting health ageing in the wider  
54  
55 population.  
56  
57  
58  
59  
60

## Conclusion

Geriatric medicine as a specialty has the skills and expertise to bridge the gap between older people's lived experience of both old age and frailty and the current clinical approach to frailty by placing commitment to the principles on which it was founded at the very heart of its approach. It can do this by reinstating the importance of the social alongside the medical, appreciating the two as both distinguishable elements and one that are always, like yin and yang, already intertwined. Together and relatedly this will involve a recognition of resources as well as deficits present in the state of being frail and of the continuum of health, illness and frailty. We suggest that concepts and methods from the social sciences and humanities will be particularly helpful in this project. Invigorated by such principles, such a medical approach has the powerful potential to influence cultural views towards ageing in a more positive way, reducing the dread towards the fourth age and the 'othering' of (frail) older people that is detrimental to the well-being not just of the older people themselves but of a society that fears, dreads and denies ageing. This is important if frailty as a construct is to realise its full potential to ameliorate the vulnerabilities of later life, without itself inducing undesirable effects, and thus to engage most productively and sensitively with the people whom it is dedicated to help.

## References

1. NHS England, Updated guidance on supporting the routine frailty identification and frailty care through the GP contract, 2017/18, Long Term Conditions Unit, Medical Directorate

- 1  
2  
3 2. Cesari, M, Araujo de Carvalho, I, Amuthavalli Thiyagarajan, J et al, Evidence for the  
4 domains supporting the construct of intrinsic capacity, *J Gerontol, A, Biol Sci Med Sci*, 2018,  
5  
6 73(2): 1653-1660  
7
- 8  
9  
10 3. Stolz, E Mayerl, H Waxeneger, A et al Impact of socioeconomic position on frailty  
11 trajectories in 10 European countries: evidence from the Survey of Health, Ageing and  
12 Retirement in Europe (20014-13). *Journal of Epidemiology and Community Health*, 2017  
13  
14 71(1):73-80  
15  
16
- 17  
18 4. Nicholson, C, Meyer, J, Flatley, M et al. Living on the margin: understanding the  
19 experience of living and dying with frailty in old age. *Social Science and Medicine* 2012:  
20  
21 75(8): 1426-32  
22  
23
- 24  
25 5. Pickard, S Health, illness and frailty in old age: a phenomenological exploration, *Journal*  
26  
27  
28 of Ageing Studies, 2018 doi:10.1016/j.jaging.2018.10.002  
29
- 30  
31 6. Fillit, H and Butler, RN (2009) The frailty identity crisis, *Journal of the American*  
32  
33 Geriatrics Society, 57(2): 348-52  
34
- 35  
36 7. Warmoth, K, Lang, I.A, Phoenix, C et al ‘Thinking you’re old and frail’: a qualitative  
37 study of frailty in older adults, *Ageing and Society*, 2016, 36, 1483-1500  
38  
39
- 40  
41 8. Robertson, DA, Savva, G.M, King-Kallimaris, B.L, Kenny, R.A Negative perceptions of  
42 ageing and decline in walking speed: a self-fulfilling prophecy, *PloS ONE*, 2014, 10(4)  
43  
44 :e0123260.doi:10.1371/  
45
- 46  
47 9. Gale, C.R, Mötus, R, Deary, I.J, Cooper, C and Sayer, A.A Personality and risk of frailty:  
48 the English Longitudinal Study of Ageing, *Ann Behav Med*, 2017: 51(1): 128-136  
49  
50
- 51  
52 10. Higgs, P and Gilleard, C Rethinking old age: theorising the fourth age, 2015, London:  
53  
54 Palgrave Macmillan  
55  
56  
57  
58  
59  
60

- 1  
2  
3 11. Grenier, A, Lloyd, L and Phillipson, C Precarity in late life: rethinking dementia as  
4  
5 'frailed' old age, *Sociology of Health and Illness*, 2017, 39(2): 318-330  
6  
7
- 8  
9 12. Age UK and the British Geriatrics Society. Britain Thinks. Frailty: Language and  
10  
11 Perceptions, 2015.  
12
- 13 13. Nicholson, C, Gordon, A.L, Tinker, A Changing the way 'we' view and talk about frailty,  
14  
15 *Age Ageing*, 2017, 46(3): 349-51; p. 350  
16  
17
- 18 14 World Health Organization (2015) Summary. World report on ageing and health:  
19  
20 <https://www.who.int/ageing/events/world-report-2015-launch/en/>  
21  
22
- 23 15. Rahman, S, *Living with frailty: from assets and deficits to resilience*, 2019, London:  
24  
25 Routledge  
26  
27
- 28 16. Martin, M *Medical Knowledge and Medical Practice: geriatric medicine in the 1950s*,  
29  
30 *Social History of Medicine*, 1995 8: 443-61  
31  
32
- 33 17. Gordon, A.L, Masud, T, Gladman, J.R.F Now that we have a definition for physical  
34  
35 frailty, what shape should frailty medicine take? *Age Ageing*, 214 (1): 8-9  
36  
37
- 38 18. Campbell AJ and Buchner DM Unstable disability and the fluctuations of frailty. *Age*  
39  
40 *Ageing*, 1997 26: 315-8. 3  
41
- 42 19. Turner and Clegg Best practice guidelines for the management of frailty: a British  
43  
44 Geriatrics Society, Age UK and Royal College of General Practitioners report, *Age Ageing*,  
45  
46 2014; 43(6): 744–747; p. 745  
47  
48
- 49 20. Laceulle, H Virtuous ageing and existential vulnerability, *Journal of Aging Studies*, 2017,  
50  
51 43: 1-8  
52  
53
- 54 21. McEwen, B.S, Brain on stress: how the social environment gets under the skin, *PNAS*,  
55  
56 2012, 109(suppl.2) 17180-17185  
57  
58  
59  
60



- 1  
2  
3 22. Kamiya, Y, Timonen, V, Romero-Ortuno, R and Kenny, R.A Back to basics or into a  
4 brave new world? The potential and pitfalls of biomarkers in explaining the pathways  
5 between social engagement and health, in Hoque, N, McGehee, M and Bradshaw, BS (eds)  
6 Applied Demography and Public Health, pp. 313-336, Springer, Dordrecht, 2013.  
7  
8  
9  
10  
11  
12 23. Van Deurzen, I and Vanhoutte, B. Trajectories of allostatic load in later life: the role of  
13 sex, birth cohorts and risk accumulation. Research on Ageing, 2018 doi:  
14  
15 10.1177/0164027518813839.  
16  
17  
18  
19 24. Rockwood, K and Mitniski, A Frailty defined by deficit accumulation and geriatric  
20 medicine defined by frailty, Clin Geriatr Med, 2011, 27: 17-26; p. 23  
21  
22  
23 25. Kojima, G, Taniguchi, Y, Illife, S et al, Transitions between frailty states among  
24 community-dwelling older people: a systematic review and meta-analysis, Ageing Research  
25 Review, 2019, 50: 81-88  
26  
27  
28  
29  
30 26. Travers, J, Romero-Ortuno, R, Bailey, J and Cooney, M-T Delaying and reversing frailty:  
31 a systematic review of primary care interventions, British Journal of General Practice, 2018:  
32 DOI: <https://doi.org/10.3399/bjgp18X700241>  
33  
34  
35  
36  
37 27. Torres, S and Hammarström, G Speaking of 'limitations' while trying to disregard them:  
38 a qualitative study of how diminished everyday competence and aging can be regarded,  
39 Journal of Aging Studies, 2006, 20: 291-302  
40  
41  
42  
43  
44 28. Rockwood, K, Fox, R.A, Stolee, P, et al Frailty in elderly people: an evolving concept,  
45 Canadian Medical Association Journal, 1994, 150, 4, 489-495  
46  
47  
48  
49 29. Williams, S.J Medicine and the Body, 2003, London and Thousand Oaks, Calif: Sage  
50  
51  
52 30. Bury, M Chronic illness as biographical disruption, Sociology of Health and Illness, 1982,  
53 4(2): 167-1822  
54  
55  
56 31. Pickard, S and Rogers, A. Knowing as practice: self-care in the case of chronic multi-  
57 morbidities, Social Theory and Health, 2012, 10(2):101-120  
58  
59  
60

- 1  
2  
3 32. Leder, D *The absent body*, 1990, Chicago: University of Chicago Press  
4  
5 33. Merleau-Ponty, M *The phenomenology of perception*, 1964, London: Routledge, p. vi  
6  
7 34. Pound, P, Gompertz, P and Ebrahim, S, *Illness in the context of older age: the case of*  
8  
9 *stroke*, *Sociology of Health and Illness*, 1998, 20(4): 489-506  
10  
11 35. Faircloth, CA, Boylstein, C, Rittman, M et al *Sudden illness and biographical flow in*  
12  
13 *narratives of stroke recovery*, *Sociology of Health and Illness*, 2004, 26(20):242-261  
14  
15 36. Marshall, A, Nazroo, J, Tampubolon, G and Vanhoutte, B, *Cohort differences in the*  
16  
17 *levels and trajectories of frailty among older people in England*, *Journal of Epidemiology and*  
18  
19 *Community Health*, 2015, 69(4): 316-321  
20  
21 37. Huber, M, Knoffnerus, JA, Green, L et al, *How should we define health?* *BMJ*, 2011:  
22  
23 <https://doi.org/10.1136/bmj.d4163>  
24  
25 38. Carel, H *Phenomenology of Illness*, 2016, Oxford University Press  
26  
27 39. Coni, N, *The unlikely geriatricians*, *JRSoc Med*, 1996, 89: 587-589  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60