



City Research Online

City, University of London Institutional Repository

Citation: Anagnostopoulou, N., Alba, A., Gaete, J., Dima, D. ORCID: 0000-0002-2598-0952 and Kyriakopoulos, M. (2018). Prevalence of psychotic symptoms in children needing inpatient care: Relationship to diagnosis. *European Psychiatry*, 48, S521-S521.

This is the accepted version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: <http://openaccess.city.ac.uk/22499/>

Link to published version:

Copyright and reuse: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

City Research Online:

<http://openaccess.city.ac.uk/>

publications@city.ac.uk

Prevalence of psychotic symptoms in children needing inpatient care: relationship to diagnosis

Anagnostopoulou N, Alba A, Gaete J, Dima D, Kyriakopoulos M

Background: Psychotic symptoms, marking impaired reality testing, are considered a hallmark of psychotic disorders. However, several lines of research point towards these symptoms also being highly prevalent both in non-clinical populations and in other psychiatric disorders. There is also evidence to suggest that their prevalence is even higher in children and young people, reaching up to 17% in epidemiological samples. There is limited research on the prevalence of psychotic symptoms in young clinical populations. The present study aimed to assess the prevalence of psychotic symptoms in a 6 to 12-year-old clinical population of an inpatient setting. **Method:** Clinical records of all patients of a national UK children's unit from 2009 to 2017 were examined for the presence of psychotic symptoms and their relationship to diagnosis at discharge. **Results:** One hundred and eighty-five (185) children were included in the study. Psychotic symptoms were highly prevalent in this young clinical population, reaching 63.8%, regardless of diagnosis at discharge. Hallucinations (58.4%) were more prevalent than delusions (35.1%). As expected, all children with schizophrenia spectrum disorders experienced psychotic symptoms. **Conclusions:** The present study is the first to examine the prevalence of delusions and hallucinations across diagnoses in children needing inpatient care. Psychotic symptoms were very common in children without a diagnosis of schizophrenia spectrum disorder. Given the high frequency of psychotic symptoms in this severely affected clinical population, it is possible that they represent a marker of clinical severity.

Table 1. Psychotic Symptoms Percentages among Diagnoses.

DIAGNOSIS	TOTAL
EARLY-ONSET SCHIZOPHRENIA	100%
INTELLECTUAL DISABILITY	86.7%
ANXIETY	67.2%
AUTISM SPECTRUM DISORDER	65.7%

<i>DEPRESSION</i>	65.5%
<i>HYPERKINETIC DISORDER</i>	65.4%
<i>EATING DISORDER</i>	60%
<i>OBSESSIVE COMPULSIVE DISORDER</i>	53.6%