

City Research Online

City, University of London Institutional Repository

Citation: Frangou, S., Modabbernia, A., Doucet, G.E., Moser, D., Dima, D. ORCID: 0000-0002-2598-0952 and ENIGMA Lifespan Working Group, . (2019). Cortical Thickness and Subcortical Volume Trajectories Across the Lifespan: Data From 14,600 Healthy Individuals Aged 6-90 Years. Biological Psychiatry, 85(10), S35-S36. doi: 10.1016/j.biopsych.2019.03.099

This is the accepted version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: https://openaccess.city.ac.uk/id/eprint/22551/

Link to published version: http://dx.doi.org/10.1016/j.biopsych.2019.03.099

Copyright and reuse: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

City Research Online: publications@city.ac.uk http://openaccess.city.ac.uk/

Cortical Thickness and Subcortical Volume Trajectories Across the Lifespan: Data From 14,600 Healthy Individuals Aged 6-90 Years

Sophia Frangou¹, Amirhossein Modabbernia¹, Gaelle Doucet¹, Dominik Moser¹, and Danai Dima², ENIGMA Lifespan Working Group

¹Icahn School of Medicine at Mount Sinai ²City University London

Background: Age is a critical determinant of brain morphometry. We formed the Lifespan Working group of the Enhancing Neuroimaging Genetics through Meta-Analysis(ENIGMA) Consortium to perform a large-scale analysis of cortical thickness data across more than 79 samples and 14,600 healthy individuals aged 6-90 years.

Methods: We used fractional polynomial (FP) regression to characterize age-related trajectories in brain morphometry. Normalized growth centiles were constructed using the para-metric Lambda, Mu, and Sigma (LMS) method. Inter-individual variability was analyzed using meta-analysis and one-way analysis of variance.

Results: The volume of subcortical regions followed an inverted U-shaped trajectory while cortical thickness in almost all regions decreased during the first two to three decades of life and showed an attenuated or plateaued slope afterwards. Exceptions were the entorhinal, anterior cingulate and temporopolar cortices that followed attenuated age-related U-shaped trajectories. Age and its FP combinations explained up to 48% variance in morphometry. Sample, strength of the MRI scanner magnetic field and FreeSurfer version jointly accounted for a median of 17% of the variance.

Conclusions: Our results could shed light on the uncertainties about age-related developmental trajectories of cortical thickness. The estimated centile values will provide scientists and clinicians with more efficient tools to detect brain morphological deviations and their association with behavioral, cognitive and clinical outcomes

Supported By: R01MH116147

Keywords: Brain Development and Aging, Volumetric Neu-oimaging, ENIGMA Consortium, Healthy Subjects, Lifespan