

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



LSHTM Research Online

Morgan, Melissa C; Maina, Beth; Waiyego, Mary; Mutinda, Catherine; Aluvaala, Jalemba; Maina, Michuki; English, Mike; (2017) Oxygen saturation ranges for healthy newborns within 24 hours at 1800 m. Archives of disease in childhood. Fetal and neonatal edition, 102 (3). F266-F268. ISSN 1359-2998 DOI: <https://doi.org/10.1136/archdischild-2016-311813>

Downloaded from: <http://researchonline.lshtm.ac.uk/4653849/>

DOI: <https://doi.org/10.1136/archdischild-2016-311813>

Usage Guidelines:

Please refer to usage guidelines at <https://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: <http://creativecommons.org/licenses/by/2.5/>

<https://researchonline.lshtm.ac.uk>



OPEN ACCESS

Oxygen saturation ranges for healthy newborns within 24 hours at 1800 m

Melissa C Morgan,¹ Beth Maina,² Mary Waiyego,² Catherine Mutinda,² Jalemba Aluvaala,^{3,4} Michuki Maina,⁴ Mike English^{4,5}

¹Department of Paediatrics, University of California San Francisco, San Francisco, California, USA

²Pumwani Maternity Hospital, Nairobi, Kenya

³Department of Paediatrics and Child Health, University of Nairobi, Nairobi, Kenya

⁴KEMRI-Wellcome Trust Research Programme, Nairobi, Kenya

⁵Nuffield Department of Medicine & Paediatrics, University of Oxford, Oxford, UK

Correspondence to

Dr Melissa C Morgan, Department of Paediatrics, University of California San Francisco, 550 16th Street, P.O. Box 1224, San Francisco, CA 94158, USA; Melissa.Morgan@ucsf.edu

Received 11 August 2016

Revised 4 January 2017

Accepted 9 January 2017

Published Online First

2 February 2017

ABSTRACT

There are minimal data to define normal oxygen saturation (SpO₂) levels for infants within the first 24 hours of life and even fewer data generalisable to the 7% of the global population that resides at an altitude of >1500 m. The aim of this study was to establish the reference range for SpO₂ in healthy term and preterm neonates within 24 hours in Nairobi, Kenya, located at 1800 m. A random sample of clinically well infants had SpO₂ measured once in the first 24 hours. A total of 555 infants were enrolled. The 5th–95th percentile range for preductal and postductal SpO₂ was 89%–97% for the term and normal birthweight groups, and 90%–98% for the preterm and low birthweight (LBW) groups. This may suggest that 89% and 97% are reasonable SpO₂ bounds for well term, preterm and LBW infants within 24 hours at an altitude of 1800 m.

INTRODUCTION

There is a large burden of neonatal mortality in low-income countries (LIC). Hypoxaemia occurs in a substantial portion of hospitalised neonates, and is significantly associated with mortality. Pulse oximetry is a non-invasive method of measuring the oxygen saturation of haemoglobin, and has become a critical tool in determining need for oxygen in sick newborns. Routine use may also aid identification of infants with clinically unrecognised respiratory abnormalities in LIC, where discharge often occurs within the first 24 hours of life.

Use of pulse oximetry relies on knowledge of normal oxygen saturation (SpO₂) values. It is important to define normal SpO₂ levels for neonates of different gestational ages (GA) at different time points after birth. SpO₂ values prior to 24 hours are lower and more variable than those seen after 24 hours. Similarly, SpO₂ values may be lower at higher altitudes, such as in Kenya where 20% of the population resides above 1500 m. Thilo *et al*¹ found that mean SpO₂ was 92%–93% at 24–48 hours among well term neonates at 1610 m. Ravert *et al*² found mean SpO₂ of 95%–97% at ~1371 m and 94%–95% at ~2073 m among well term newborns during the first 72 hours. Three studies evaluated SpO₂ in healthy preterm infants. Ng *et al*³ (N=33) found mean SpO₂ of 97% (median GA 33 weeks, median age 14 days); Harigopal *et al*⁴ (N=43) found median SpO₂ of 95% (median GA 33 weeks, median age 14 days) and Richard *et al*⁵ (N=55) found median SpO₂ of 99% (mean GA 35, mean age 1 day). All of these studies were conducted at sea level.^{3–5} Specific gaps remain with sparse data for newborns within the first 24 hours or for well preterm infants, with no

What is already known on this topic?

- ▶ Proper use of pulse oximetry to guide oxygen therapy relies on knowledge of normal oxygen saturation (SpO₂) values for a given population.
- ▶ Based on available evidence, it appears that well term infants born at moderate altitude should have SpO₂ values of ≥92% after the first 24 hours.
- ▶ The optimal SpO₂ for well newborns within 24 hours of life and for well preterm infants born at moderate altitude is unknown.

What this study adds?

- ▶ This is the largest study of SpO₂ ranges for well preterm infants at moderate altitude.
- ▶ This study suggests that a SpO₂ range of 89%–97% is suitable for well term and preterm infants within 24 hours at 1800 m.
- ▶ This study provides important data about SpO₂ in the first 24 hours, which are useful as discharge postdelivery occurs within 24 hours in many low-income countries.

published reference ranges for preterm infants born at increased altitude. The aim of this study was to establish within 24 hours the reference interval for preductal and postductal SpO₂ in healthy term and preterm neonates at 1800 m.

METHODS

Participants and setting

This study was conducted at Pumwani Hospital in Nairobi, Kenya, located at 1800 m. The hospital provides care to ~22 000 women and their infants each year. A 120-bed nursery provides care for all infants requiring medical attention. Inclusion criteria included being born alive at Pumwani Hospital during the study period and appearing well as defined by (1) normal vital signs (heart rate 110–180 beats/min, respiratory rate 30–60 breaths/min, temperature 36.5°C–37.5°C), (2) absence of respiratory distress and (3) absence of other signs of illness (eg, poor suck, lethargy). Infants were excluded if they required admission for any reason other than observation or if they were transferred to another facility within 4 hours for anomalies or severe medical problems.



CrossMark

To cite: Morgan MC, Maina B, Waiyego M, *et al*. *Arch Dis Child Fetal Neonatal Ed* 2017;102:F266–F268.

Sampling approach

We aimed to enrol 800 well neonates, stratified to include 400 term (≥ 37 weeks) and 400 preterm (< 37 weeks) neonates. GA was based on last menstrual period (LMP), and Ballard examination was conducted when LMP was unknown or incongruent with appearance. We screened all preterm neonates and a random proportion of term neonates.

Procedures

Enrolled infants underwent testing once between 1 and 24 hours with the Lifebox oximeter, which was developed for low-resource settings by WHO and World Federation of Societies for Anaesthesiologists (Acare Technology, New Taipei City, Taiwan). The measurement was recorded when there was a good waveform for ≥ 15 s and SpO₂ was stable over that period.

Analysis

We determined mean, SD, median, IQR and 5th–95th percentile range of preductal and postductal SpO₂ measurements, stratifying by GA (term or preterm), birth weight (normal (NBW), ≥ 2.5 kg or low (LBW), < 2.5 kg) and postnatal age (0–6, 6–12, 12–18, 18–24 hours). Statistical analyses were conducted using Stata V.13 (StataCorp, College Station, Texas, USA).

Ethical aspects

Written informed consent was obtained from parents/guardians. Ethical approval was received from the University of California, San Francisco and the Kenya Medical Research Institute-Wellcome Trust Research Programme.

RESULTS

A total of 555 infants were enrolled between January and December 2015. The mean GA was 38 weeks (SD 2.4), mean birth weight was 2.94 kg (SD 0.53), 50% were male and 93% were delivered vaginally. Among preterm infants, median GA was 35 weeks (range 27–36).

Mean preductal SpO₂ was 93%–94% across all GA and birth-weight groups. Mean postductal SpO₂ was 93%–94% for all GA groups, 93% for birth weights ≥ 2.5 kg and 95% for < 2.5 kg. Median preductal and postductal SpO₂ values were similar (table 1). The 5th–95th percentile range for preductal and postductal SpO₂ was between 89% and 97% for the term and NBW groups, and between 90% and 98% for the preterm and LBW groups.

Table 1 Preductal and postductal SpO₂ by gestational age and birth weight

| | Gestational age | | Birth weight | |
|---------------------------------|-----------------|-----------------|-----------------------|-------------------|
| | Term (n=420) | Preterm (n=135) | ≥ 2.5 kg (n=456) | < 2.5 kg (n=99) |
| Preductal SpO ₂ , % | | | | |
| Mean, SD | 93 (2.8) | 94 (2.8) | 93 (2.8) | 94 (2.6) |
| Median, IQR | 94 (91–95) | 94 (92–96) | 94 (91–95) | 94 (92–96) |
| 5%–95% range | 89–97 | 90–98 | 89–97 | 90–98 |
| Postductal SpO ₂ , % | | | | |
| Mean, SD | 93 (2.5) | 94 (2.5) | 93 (2.6) | 95 (2.3) |
| Median, IQR | 94 (92–95) | 95 (93–96) | 94 (92–95) | 95 (93–96) |
| 5%–95% range | 89–97 | 90–98 | 89–97 | 90–98 |

SpO₂, normal oxygen saturation.

Table 2 Preductal and postductal SpO₂ by postnatal age

| | Postnatal age (hours of life) | | | |
|---------------------------------|-------------------------------|--------------------|---------------------|---------------------|
| | 0–6 hours (n=109) | 6–12 hours (n=145) | 12–18 hours (n=147) | 18–24 hours (n=137) |
| Preductal SpO ₂ , % | | | | |
| Mean, SD | 94 (2.9) | 93 (2.9) | 93 (2.5) | 93 (2.9) |
| Median, IQR | 95 (92–96) | 94 (91–95) | 93 (91–95) | 93 (91–95) |
| 5%–95% range | 90–98 | 89–98 | 89–97 | 88–98 |
| Postductal SpO ₂ , % | | | | |
| Mean, SD | 94 (2.7) | 93 (2.4) | 94 (2.4) | 93 (2.5) |
| Median, IQR | 95 (92–96) | 94 (92–95) | 94 (92–95) | 93 (92–95) |
| 5%–95% range | 90–98 | 90–97 | 90–97 | 88–97 |

SpO₂, normal oxygen saturation.

When stratified by postnatal age, results were similar (table 2).

Scatter plots of preductal and postductal SpO₂ by hours are shown in figures 1 and 2, respectively, and also suggest absence of change across time.

DISCUSSION AND CONCLUSION

Data are sparse to inform our understanding of oxygen saturation norms for newborns within the first 24 hours and for well preterm infants. We found that mean preductal and postductal SpO₂ was 93%–95% in all groups within 24 hours. This is similar to findings among well term infants in studies conducted by Thilo *et al*¹ and Ravert *et al*.² The mean and median SpO₂ for preterm infants were only slightly lower than those found by Ng *et al*,³ Harigopal *et al*⁴ and Richard⁵ at sea level despite the early postnatal age and increased altitude in our study. We found that the 5th–95th percentile ranges for preductal and postductal SpO₂ were 89%–97% for the term and NBW groups and 90%–98% for the preterm and LBW groups. Thilo *et al* reported a 95% CI of 89% to 97%,¹ which corresponds with our findings.

This study has limitations. Our findings for preterm infants are based on a small sample of 135 infants. It was difficult to recruit well preterm infants as many such infants were admitted for medical therapy, making them ineligible. Additional research about SpO₂ reference ranges in preterm infants born at a variety of altitudes is needed. If methods and timing for collecting




Figure 1 Preductal normal oxygen saturation (SpO₂) by hours of life.




Figure 2 Postductal normal oxygen saturation (SpO₂) by hours of life.

measurements were sufficiently similar, these data could make a considerable contribution to a pooled sample for meta-analysis.

In this study, GA was based on LMP with Ballard examination conducted when LMP was unknown or incongruent with appearance. LMP is subject to bias and Ballard may differ from ultrasound by 1 to 2 weeks. However, ultrasound is unavailable in many LIC facilities due to cost and need for skilled sonographers. The Lifebox oximeter is not motion-resistant and has not been validated in neonates. To ensure we obtained the most accurate measurement possible, we recorded measurements only when there was a good waveform for ≥ 15 s and SpO₂ was stable over that period.

Reports suggest that pulse oximetry use can decrease mortality in children with unrecognised hypoxaemia. In newborns, it is increasingly being used for pre-discharge screening to diagnose occult respiratory and cardiac disease. This study suggests that 89% and 97% may be reasonable SpO₂ bounds for well term, preterm and LBW infants within 24 hours at 1800 m, although we lack outcome data that would confirm infants in this study remained well. This study provides important data about SpO₂ within the first 24 hours, which are useful as discharge post-delivery is rapid and typically without skilled assessment by a clinician or nurse in many LIC settings.

Twitter Follow Melissa Morgan @melissacmorgan

Acknowledgements We would like to acknowledge the parents who consented for their infants to participate in this study. We would like to acknowledge Aineah Adiema, Alex Maina and Sarah Jumba for assisting with data collection at Pumwani Hospital, and Jacintah Mwachiro, Judy Nganga and Elizabeth Kyala at KEMRI-Wellcome Trust Research Programme for assisting with study management and logistics. In addition, we thank all the staff of the newborn unit at Pumwani Hospital for their cooperation during the study. This manuscript is published with the permission of the Director of KEMRI.

Contributors MCM initiated the collaborative study, designed data collection tools, monitored data collection for the study, wrote the statistical analysis plan, cleaned and analysed the data and drafted and revised the paper. She is guarantor. BM, MW, CM, JA and MM revised data collection tools, implemented the study in Kenya, monitored data collection and revised the paper. ME initiated the collaborative project, revised data collection tools, contributed to data analysis and revised the paper.

Funding MCM received support from the University of California San Francisco Resource Allocation Program Global Health Policy Award to conduct this research. Funds from The Wellcome Trust (#097170) awarded to Mike English as a Senior Fellowship together with additional funds from a Wellcome Trust core grant awarded to the KEMRI-Wellcome Trust Research Programme (#092654) also supported this work.

Competing interests None declared.

Ethics approval University of California San Francisco, KEMRI-Wellcome Trust Research Programme.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement Data are available on application from the KEMRI-Wellcome Trust Research Programme Data Governance Committee. Applications can be directed in the first instance to Mike English at MEnglish@kemri-wellcome.org.

Open Access This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See: <http://creativecommons.org/licenses/by/4.0/>

REFERENCES

- 1 Thilo EH, Park-Moore B, Berman ER, *et al.* Oxygen saturation by pulse oximetry in healthy infants at an altitude of 1610 m (5280 ft): what is normal? *Am J Dis Child.* 1991;145:1137–40.
- 2 Ravert P, Detwiler TL, Dickinson JK. Mean oxygen saturation in well neonates at altitudes between 4498 and 8150 feet. *Adv Neonatal Care.* 2011;11:412–7.
- 3 Ng A, Subhedar N, Primhak RA, *et al.* Arterial oxygen saturation profiles in healthy preterm infants. *Arch Dis Child Fetal Neonatal Ed.* 1998;79:F64–6.
- 4 Harigopal S, Satish HP, Taktak AFG, *et al.* Oxygen saturation profile in healthy preterm infants. *Arch Dis Child Fetal Neonatal Ed.* 2011;96:F339–42.
- 5 Richard D, Poets CF, Neale S, *et al.* Arterial oxygen saturation in preterm neonates without respiratory failure. *J Pediatr.* 1993;123:963–8.