

Exercise training in Patients with Chronic Obstructive Pulmonary Disease. Effectiveness ten years after participation.

Ângela Maria Pereira^{1,2,3}, Duarte Vieira¹, Jessica Reis¹, João Silveira¹

¹Physiotherapy, Escola Superior de Saúde Egas Moniz; ²Centro de investigação interdisciplinar Egas Moniz; ³Hospital Garcia de Orta, Almada, Portugal



Background

Chronic Obstructive Pulmonary Disease (COPD) is a preventable and treatable disease caused by exposure to tobacco smoke and poisonous gases. It is characterized by progressive obstruction of the air-flow, irreversible once lesions occur in the parenchyma. (1). Patients with COPD may respond to exercise training in different ways compared to healthy subjects, since the determinants of exercise limitation appear to be widely multi-factorial (2). Such factors may include gas exchange abnormalities, dynamic lung hyperinflation, insufficient energy supply to the peripheral and respiratory muscles, morphological alterations in leg and diaphragm muscle fibers and reduced functional metabolic capacities (3). The strength exercise can improve the respiratory muscle performance with potential positive effects on blood oxygenation, and consequently, muscular performance of patients with COPD (4). This improvements may affect positively this individuals health and quality of life at short and long term, leading to a lower health service recurrence (HSR) (5).

Purpose

To assess the effectiveness of attending to an exercise program, combined (aerobic and strength exercise) or aerobic training alone in patients with COPD compared with subjects not submitted to exercise programs.

Methods

Thirty men with moderate COPD, were randomly assigned for 2 groups: 15 patients (age, 66.5±6.2 years; FEV1, 55.8±9.9%) to a combined exercise training program (CG), and 15 (age, 65.4±3.6 years; FEV1, 59.1±9.1%) to an aerobic training program (AG), for 10 weeks, three times a week. Outcome variables included cardiopulmonary function (cardiopulmonary exercise test (CPET) and 6-min walk distance (6MWD), muscular strength (1-RM); and quality of life (HRQL) with SF-36 and QRSF. Ten years after the programs, both groups were compared with ten patients who weren't submitted to exercise programs, by evaluating health service recurrence and mortality for respiratory cause. The study was approved by the Ethics Committee of the Garcia de Orta Hospital and all participants gave their written informed consent prior to inclusion in the investigation.

Results

Means and standard deviations values of the modification rate, before and after training for both groups are displayed in Table 1.

Baseline data: No significant differences were observed, between groups, for cardiorespiratory variables, except for SatO_{2peak} (%) and 6-min walk test distance (m). For HRQL we just observed significant differences in vitality dimension in the MOS-SF 36 questionnaire.

After training: Both exercise groups increased (p<.05) functional capacity, O₂pulse peak, CPET time and power and 6MWD.

HRQL improved immediately after exercise, with greater benefits for the CG group (p<.05) in all variables in St. George Respiratory Questionnaire.

With the generic instrument, MOS SF-36, significant differences were observed in modification rates for physical function, role physical and role emotional (Graph 1).

Results

Ten years later, there were no differences between the exercise groups on mortality and HSR.


Between exercise groups and subjects who didn't integrate the program, there weren't differences on mortality, but there were significant differences on HSR (p<.05).

Table 1 Mean and standard deviation values for modification rate of functional cardiorespiratory parameters, maximal oxygen uptake (VO_{2peak}), carbon dioxide production (VCO_{2peak}), minute ventilation (VE_{peak}), oxygen pulse (Pulse O_{2peak}), ventilatory equivalents for oxygen uptake (VE/VO_{2peak}), oxygen saturation peak (SatO₂), heart rate at basal and peak stages, exercise duration and 6-min walk distance, in combined exercise training group (CG) and in aerobic training group (AG).

	% Δ		p*
	CG	AG	
VO _{2peak} (mL.kg ⁻¹ .min ⁻¹)	24±16	26±25	0.002
VCO _{2peak} (L.min ⁻¹)	10±5.3	14±6.4	0.121
VE _{peak} (L.min ⁻¹)	2.8±18.1	26.9±27.7	0.072
O ₂ pulso _{peak}	24±0.1	21±0.2	0.000
VE/VO _{2peak}	(-)17.3±10.5	6.1±14.3	0.006
SatO _{2peak} (%)	2.0±0.3	(-)1.0±3.1	0.053
HR _{peak} b.min ⁻¹	(-)3.8±4.9	3.7±13.8	0.048
HR _{reste} b.min ⁻¹	(-)4.9±2.1	(-)4.2±2.4	0.108
Exercise duration (min.)	28.6±8	3.0±17	0.000
6-min Walk (m)	12.7±4	6.8±3	0.003

* Differences between groups p<0.05

Graph 1: Modification rate in Medical Outcomes Study SF-36



Conclusions

Combined exercise training was more effective than aerobic exercise alone with greater improvement in muscular strength, functional capacity and HRQL after program, reducing morbidity by improving functional capacity through exercise. It is still unknown if improvements in functional capacity are maintained in the long-term and if this leads to increased physical activity levels as measured by a free-living activity monitor. Nevertheless participation in exercise programs seems to reduce HSR at long-term follow-up

References

1. Eliason G, Abdel-Halim S, Arvidsson B, Kadi F, Piehl-Aulin K. Physical performance and muscular characteristics in different stages of COPD. *Scand J Med Sci Sports* 2009;19:865-870
2. Gloeckl R, Marinov B, Pitta F. Practical recommendations for exercise training in patients with COPD. *European Respiratory Journal*. 2013;22(128):178-186.
3. Neves L, Chiappa A, da Silva V, Vieira P, Cipriano G, Arena R, et al. Comparative effects of inspiratory muscle training and resistance training on respiratory and skeletal muscle strength in COPD: Responses of a pulmonary rehabilitation program. *European Respiratory Journal*. 2014;44(Suppl 58).
4. Egan C, Deering BM, Blake C, Fullen BM, McCormack NM, Spruit MA, et al. Short term and long term effects of pulmonary rehabilitation on physical activity in COPD. *Respiratory Medicine*. 2012;106(12):1671-9.
5. Qureshi H, Sharafkhan A., Hanania NA Chronic obstructive pulmonary disease exacerbations: latest evidence and clinical implications. *Ther Adv Chronic Dis*. 2014; 5(5): 212-22