

DR. REBECCA GOSLING (Orcid ID : 0000-0002-0829-0060)

DR. MUNA ANJUM (Orcid ID : 0000-0001-5278-7587)

A sensitive method for the recovery of *Escherichia coli* serogroup O55 including Shiga toxin-producing variants for potential use in outbreaks.

Miranda Kirchner¹, Ellie Sayers^{1,2}, Shaun Cawthraw¹, Nicholas Duggett¹, Rebecca Gosling¹, Claire Jenkins³, Timothy J Dallman³, Doris Mueller-Doblies¹, Muna F Anjum^{1#}.

1 Department of Bacteriology, Animal and Plant Health Agency, Addlestone, Surrey, KT15 3NB.

2 University of East Anglia / Quadram Institute Bioscience, Norwich Research Park, NR4 7UA.

3 Public Health England, Colindale Avenue, London, NW9 5HT.

corresponding author: Muna.Anjum@apha.gsi.gov.uk

Running title: O55 *Escherichia coli* isolation method

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi:

10.1111/jam.14345

This article is protected by copyright. All rights reserved.

Abstract

Aim

Shiga toxin-producing *Escherichia coli* (STEC) cause bloody diarrhoea, kidney failure and occasionally death. However, identifying the source of infection caused by STEC other than serogroup O157 is hampered by availability of sensitive methods for detecting these pathogens. In this study we developed novel tools for detecting *E. coli* O55 that is potentially associated with human outbreaks.

Method and Results

Overall specificity of immuno-magnetic separation (IMS) beads coated with anti-O55 serum was good with exception of cross reactivity with *E. coli* O22 and O23, which was eliminated using an O55 specific PCR. Limit of detection for *E. coli* O55 using O55-IMS-beads in spiked cattle faeces was on average 50 CFU ml⁻¹ (range 1-90), and improved to <10 CFU ml⁻¹ using the O55 specific PCR, following IMS on samples enriched for 2h with *E. coli* O55. Application of these tools to test cattle faeces collected on-farm allowed the isolation of O55:H19, which through whole genome sequencing was compared to STEC O55:H7 human outbreak strains.

Conclusion

These tools provide a sensitive method which could be used to screen samples for STEC O55, whether environmental or human clinical.

Significance and Impact of the Study

Several human outbreaks reported in England were caused by STEC O55:H7. Tools developed here could assist in identification of the environmental source for these isolates, which has not yet been established.

Key words: Pathogen, Enterohaemorrhagic *Escherichia coli*, immuno-magnetic separation, rapid diagnostics, environmental

Introduction

Escherichia coli, although a constituent of human and animal gut microbiota can also cause intestinal and extra-intestinal infections following acquisition of virulence factors. Shiga toxin-producing *E. coli* (STEC) is a pathotype that has acquired a number of virulence factors including Shiga toxins. Shiga toxins, are encoded by the *stx1* (variants a, c, and d) and *stx2* (variants a, b, c, d, e, f and g) genes and are carried by bacteriophage (Kruger and Lucchesi, 2015). STEC are important human pathogens which are sometimes acquired through consumption of contaminated food-stuffs or close animal contacts (Jenkins et al., 2015, Adams et al., 2016, Ihekweazu et al., 2012, Aktan et al., 2007, Anjum et al., 2014, Carter et al., 2008). STEC can cause life threatening infections which can include haemolytic uremic syndrome (HUS) and haemorrhagic colitis. *E. coli* O157 is the most common STEC serogroup associated with human infection, although there are many more *E. coli* serogroups which can cause STEC infections, with serogroups O111, O145, O26, O103, O121 and O45 regarded as the most common. These are considered the major cause of non-O157 STEC infections (Bosilevac and Koohmaraie, 2012, Eichhorn et al., 2015).

In 2014-2015, an outbreak of STEC occurred in the county of Dorset, England caused by STEC serotype O55:H7 carrying *stx2* and *eae*, encoding an adhesin present in the locus of enterocyte effacement. In total, 31 cases of STEC O55 infection were reported mainly in children, 13 of which developed HUS (McFarland et al., 2017). The serotype O55:H7 had not previously been associated with STEC infections in England and the source of STEC O55:H7 isolates causing infections in Dorset could not be determined despite extensive investigations (Paz, 2016). Moreover, STEC O55 was reported in the county of Surrey, England, in November 2017, where it was confirmed in two people and suspected in a further five; STEC O55 from this outbreak was very closely related to the Dorset outbreak strain, belonging to the same 10 SNP single linkage cluster as the Dorset strain (Brock, 2017). Outside the UK, sporadic cases of human infection with STEC O55 have been reported in Germany, Italy and the Czech republic (Johnson et al., 2006). In October 2018, environmental STEC O55:H7 was again identified as the causative organism in two children from Leicestershire who developed HUS, which resulted in fatality, and again the environmental source was not identified (<https://www.bbc.co.uk/news/uk-england-leicestershire-45838401>; Public Health England, unpublished data).

Immuno-magnetic separation (IMS) can be used for the isolation of *E. coli* of interest, which may be present in low concentrations, in a sample contaminated with many other bacteria including other *E. coli* serotypes. Commercially-produced magnetic beads are available for O157 and the six most common non-O157 STEC serogroups (Noll et al., 2016), which can be used for their isolation from food, faeces, water and other environmental samples, where they may be present in low numbers (Guy et al., 2014, Islam et al., 2006, Kanki et al., 2014, Lejeune et al., 2001) (Dynabead product information, <https://www.thermofisher.com>). The IMS method for the detection of O157 is well

established and includes an IMS step followed by plating on selective agar, specifically cefixime-tellurite sorbitol MacConkey agar (CT-SMAC). There are no selective agars for non-O157 STEC making differentiation of non-O157 STEC from other *E. coli* difficult.

There is currently no method for the selective isolation of *E. coli* O55 from faecal (human or animal) and environmental samples, which may aid in increasing the sensitivity of detection and source attribution. Therefore the aim of this study was to design and evaluate a rapid and sensitive IMS based method for the selection of *E. coli* O55, which could be used to screen samples for this serogroup.

MATERIALS and METHODS

Bead preparation

Tosylactivated M-280 Dynabeads (ThermoFisher scientific) were coated with antiserum for the *E. coli* O55 lipopolysaccharide (BioRad) according to published methods (https://tools.thermofisher.com/content/sfs/manuals/dynabeads_m280tosylactivated_man.pdf). Washed beads were coated with O55 antiserum for 14h at 37°C in 0.04 mol l⁻¹ borate buffer pH9.5, 1.2 mol l⁻¹ ammonium sulphate. Following overnight coupling, beads were washed in Phosphate Buffered Saline (PBS)+0.5% bovine serum albumin (BSA), followed by two further washes in PBS+0.1%BSA.

Immuno-magnetic separation

IMS was performed using a manual method except for STEC isolation, which used an automated system. A 1ml sample (overnight broth culture or spiked faeces suspended in BPW) was mixed with 20µl of coated beads and incubated for 30min at room temperature. A magnet was used to separate the beads, which were then washed in triplicate with

1000µl PBS-0.05% Tween before being re-suspended in 100µl of PBS or water. Beads were plated onto CHROMagar ECC (BioConnections, UK), a chromogenic agar on which most *E. coli* grow as blue colonies. An automated system was used to perform IMS of STEC isolates. As described above, a sample was mixed with beads and IMS performed using the Bead-Retriever (Dynal). Following washing in PBS-0.05% Tween, beads were plated on to CHROMagar O157 (BioConnections, UK). The *E.coli* /VTEC program pre-installed on the Bead-Retriever (Dynal) was used for IMS.

Testing the specificity and sensitivity of O55 IMS beads

The specificity of O55 IMS beads was tested with a panel of 155 *E. coli* with different O-antigens. Serogroups not tested were: O31, O39, O47, O63, O64, O67, O68, O72, O84, O94, O122, O138, O139, O141, O146, O161, O171 and O172, as they were unavailable. Overnight cultures of 151 *E. coli* were diluted to 10^3 CFU ml⁻¹ in PBS and mixed in pools of 4 or 5 strains. Triplicate 1ml aliquots of each pool were subjected to IMS using O55-beads and the beads plated on LB agar. Four serogroups, O22, O23, O83 and O104, were tested singularly for cross-reactivity with the O55-beads. Serial dilutions of overnight cultures grown in LB-broth were prepared to achieve concentrations of 10^2 and 10^3 CFU ml⁻¹. These were subjected to manual IMS with O55-beads. Percentage recovery was determined following enumeration of pre- and post-IMS samples plated on LB agar.

Seven *stx*-negative strains of *E. coli* serogroup O55 (Table 1) were also tested to determine the recovery rate of cells following IMS as described above. For STEC O55, overnight cultures were prepared in LB broth and a 10-fold dilution series prepared down to 10^{-7} . IMS using O55-beads was performed on 1ml of dilutions 10^{-5} , 10^{-6} and 10^{-7} and beads plated onto CHROMagar O157 to determine cell counts.

Limit of detection in spiked faeces.

The *E. coli* O55 strain (B1362) was grown overnight at 37°C in LB broth and this was used to prepare dilutions to achieve a final concentration of between 10^7 and 10^1 CFU ml⁻¹. One gram of fresh cattle faeces (1-2 days old) was mixed with 9 ml BPW and to this 1ml of O55 cell suspension (at concentration between 10^7 and 10^1 CFU ml⁻¹) was added; a control sample (1g faeces mixed with 9 ml BPW) was spiked with PBS only. One ml aliquots were taken from the spiked and unspiked faeces prior to enrichment, and after 2h incubation at 37°C. IMS was performed with O55-dynabeads according to the method above. IMS beads were suspended in 100µl of PBS and half of this plated onto CHROMagar ECC. Plates were incubated for 14-16h at 37°C and blue colonies counted as presumptive *E. coli*. Latex agglutination was performed on 3 blue colonies from positive plates as described below. The method was considered able to detect *E. coli* O55 when at least 1/3 colonies selected for latex agglutination test were positive. A cell count was performed to confirm the inoculum size, using the Miles Misra method on CHROMagar ECC (Miles et al., 1938). The experiment was performed in triplicate.

Latex agglutination testing of colonies recovered following IMS.

Suspected isolates of *E. coli* O55 recovered following IMS were tested by latex agglutination. A colony from selective plates was mixed with 10µl of saline on a reaction card (<http://www.ssidiagnostica.com/da/Products/Antisera/E-coli-antisera/E-coli-latex-test>) before the addition of one drop of O55 ImmuLex latex agglutination sera (as above). The reagents were thoroughly mixed and following 10 min incubation at room temperature the results were read.

Real-time PCR detection of O55 E. coli and virulence factors

IMS-PCR was used to confirm that samples recovered following IMS were positive for *E. coli* O55. A real-time PCR (RT-PCR) developed previously (DebRoy et al., 2005) was used to detect O55 *wzy* gene. DNA was prepared by lysis of 50 µl of beads (post-IMS) using heating to 99°C for 15 min. Once boiled, beads were removed and the supernatant used as a source of DNA. RT-PCR reactions were prepared as follows: each 20 µl reaction consisted of 10 µl QuantiTect multiplex PCR master mix (Qiagen), 10µMol of each primer, 4 µMol of probe, water and 2 µl of DNA. PCR consisted of: 95°C for 15 min followed by 45 cycles of 95°C for 15 sec and 60°C for 60 sec. Fluorescence intensity was measured at the end of each amplification cycle and the cycle threshold (Ct) value was determined.

Isolation of O55 E.coli from naturally contaminated cattle faeces

Fifty cattle faecal samples collected from a previous on-farm investigation were screened for the presence of *E. coli* O55 (Treacy et al., 2019). One gram of faeces was added to 9ml BPW and grown for 2h at 37°C. Following incubation, a 0.5ml sample was boiled for 15min. PCR was performed to detect the O55 *wzy* gene and IMS performed on PCR positive samples using the O55-specific IMS beads. Up to 10 blue colonies from CHROMagar ECC were tested by agglutination for O55 antigen, and positive samples prepared for DNA sequencing. Fresh growth of *E. coli* O55 on CHROMagar ECC was re-suspended in 200µl of water, boiled for 15min and DNA extracted from the lysate using the MagMAX CORE nucleic acid purification kit (ThermoFisher Scientific) and a KingFisher duo prime.

Genomic DNA was sequenced using a MiSeq v2 kit with 2x150bp, 300 cycles (Illumina). A core genome SNP alignment was produced by Snippy (<https://github.com/tseemann/snippy>) using *E. coli* str. K-12 sub-strain MG1655 (NC_000913.3) as a reference. RAxML (<https://academic.oup.com/bioinformatics/article/30/9/1312/238053>) was used to build the phylogenetic tree, which was subsequently annotated using ITOL (<https://www.ncbi.nlm.nih.gov/pubmed/27095192>). Virulence gene carriage was assessed using APHA in-house SeqFinder tool and an in-house virulence gene database (Duggett et al., 2017).

RESULTS

Determining the sensitivity and specificity of O55-beads

The sensitivity of O55-beads for the recovery of *E. coli* O55 isolates was tested with a panel of seven Stx-negative *E. coli* O55 (Figure 1). For each isolate of Stx-negative *E. coli* O55, the recovery rate was determined at an initial inoculum of 10^2 CFU ml⁻¹. Recovery rates were on average between 30-90% for Stx-negative *E. coli* O55 with the lowest recovery rate seen for strain 394/10, indicating a natural variability in recovery between strains (Figure 1.). With the exception of strain 394/10 the recovery rate was >60%. The sensitivity of O55-beads was also tested with three strains of STEC O55. These strains included a STEC O55 isolate from the Dorset outbreak (H2) (Paz, 2016). For all 3 STEC tested, it was possible to recover *E. coli* O55 following IMS at a concentration of 10^2 CFU ml⁻¹. At this concentration, an average of 1.7 blue colonies (range 1-5 colonies/plate) grew on chromogenic agar, post-IMS.

The specificity of O55 IMS-beads was also tested with 155 *E. coli* of different non-O55 serogroups to test their cross reactivity with O55-beads. Pools of 4-5 different serogroups were tested at a bacterial concentration of 10^2 CFU ml⁻¹. For each pool, the number of colonies recovered from plating of O55-IMS elute was determined and the percentage bacterial recovery calculated. For the 31 pools tested, comprising 151 serogroups (as 4 serogroups were tested individually), the recovery of *E. coli* was low with only ~1.95% (range 0.06%- 5.29%; data not shown) recovered.

However, some strains showed higher recovery probably due to cross-reactivity between anti-O55 sera and these serogroups (DebRoy et al., 2005), and were tested individually; these included *E. coli* of serogroups O22, O23, O83 and O104. One isolate of each serogroup was tested at 10^2 and 10^3 CFU ml⁻¹, and the rate of recovery post-IMS determined (Figure 2A and 2B). Serogroups O83 and O104 gave low rates of recovery of *E. coli* post-IMS. At a 10^3 CFU ml⁻¹ inoculum, the average recovery of *E. coli* O83 was $3.4\% \pm 1.2$ and for O104 this was $6.6\% \pm 3.9$. Much higher rates of *E. coli* recovery were seen for serogroups O22 and O23 at 10^2 and 10^3 CFU ml⁻¹, which was between ~46-48% for serogroup O22 and ~19-33% for serogroup O23 (Figure 2).

Due to the cross-reactivity seen with these serogroups an O55 specific PCR was used to provide a method which improved the specificity of testing for *E. coli* O55 recovered via IMS. The results showed *E. coli* O22 and O23 isolates recovered through IMS were negative when using the O55 specific PCR, whilst O55 isolates were positive (data not shown).

Sensitivity of O55-beads for recovery of O55 *E. coli* from faecal enrichments

The sensitivity of O55-beads for the recovery of *E. coli* O55 from spiked faecal samples was determined. Experiments were performed with cattle faeces (1-2 days old), which were spiked with *E. coli* O55. For each spiking study, a comparison was made between IMS-culture and IMS-PCR, with and without a 2h enrichment. The limit of detection for IMS-culture were comparable for samples not enriched (Table 2), with an average detection limit of 52 CFU ml⁻¹, compared to 54.4 CFU ml⁻¹ following enrichment. For samples eluted by IMS, on which O55 PCR was performed (IMS-PCR), the limit of detection was on average 52 CFU ml⁻¹ prior to enrichment, which improved to 4.9 CFU ml⁻¹ following 2h enrichment. The average Ct values for RT-PCR increased as the concentration of spiked *E. coli* O55 decreased to between 22.82 (at 10⁶ CFU ml⁻¹) and 36.99 (at 10 CFU ml⁻¹), for enriched samples. Control samples, spiked with PBS, were all negative by IMS-PCR in pre and post enriched samples (data not shown). Also in the absence of an IMS step, PCR directly from the enrichment broths gave higher Ct values than IMS-PCR (data not shown). In contrast, using IMS-culture method an average of 98.6 presumptive *E. coli* (which were blue on CHROMagar ECC) (range 40-138) were recovered from control faecal samples pre-enrichment. Following 2h enrichment an average of 604 presumptive *E. coli* (range 368-1000) were recovered. Latex agglutination of presumptive *E. coli* from control samples (3 blue colonies/sample) found that none of those tested were serogroup O55, indicating cross-reaction with other O-antigens.

Analysis of naturally contaminated cattle faeces.

Using the IMS-PCR method, followed by IMS-culture, *E. coli* O55 were isolated from two of 50 cattle samples collected from a dairy farm, and whole genome sequence (WGS) analysis was performed on these *E. coli*. WGS data was analysed using a number of methods. Both isolates were confirmed as serotype O55:H19 using serotype finder (DTU

Centre of Genomic epidemiology; <http://www.genomicepidemiology.org/>). Analysis of the core genomes for both isolates determined that there were 192 SNP (single nucleotide polymorphism) differences between isolates. A phylogenetic tree was generated from the core genomes of the two cattle isolates of *E. coli* O55, twelve sequences of *E. coli* O55 available on NCBI BioProjects (<https://www.ncbi.nlm.nih.gov/bioproject/>), four of which were clinical outbreak strains, and five O157:H7 isolates, including two from cattle on this farm (Figure 3). The cattle O55:H19 isolates were distinct from all *E. coli* O55 and O157 used for comparison and context, with >40000 SNP differences in the core genome between the O55:H19 isolated during this study and all other isolates included in the tree. Analysis of virulence gene carriage indicated the two O55:H19 isolates lacked the virulence factors associated with STEC (such as *stx* and *eae*) and other *E. coli* pathogroups, but they carried haemolysin genes *hlyE* and *hlyC*, which are widely distributed on the chromosome of *E. coli* (Murase et al., 2012).

DISCUSSION

Currently there are no protocols for the selective enrichment of *E. coli* O55 from human or animal faecal specimens. Such protocols are required to assist in the rapid identification of sources for STEC O55, especially during outbreaks. IMS, also commonly used for isolation of STEC O157, provides a sensitive isolation method when combined with selective agar. In this study an assessment of the specificity of O55-IMS beads produced in-house for isolation of STEC O55 was made with a panel of *E. coli*. It identified a low level of non-specific binding with cross-reaction between O55-beads and *E. coli* of serogroups O23 and O22, which has been previously reported (DebRoy et al., 2005). It is not uncommon for non-specific binding of other *E. coli* serogroups to IMS beads; cross-reaction between beads for O111 and *E. coli* O103 has been reported (Bai et al., 2012).

An enrichment step is used for the isolation of STEC serotypes such as O157:H7 and O26, as they may be present in low numbers, and was trialled in this study to assist in the isolation of *E. coli* O55 from cattle faeces. However, the inclusion of an enrichment step resulted in high cell densities, which inhibited binding of the target *E. coli* to IMS beads (Hallewell et al., 2017). Conrad *et al.* (2014) found the identification of single colonies of the correct serogroup when identifying O26, O45, O103, O111, O121, and O145 on isolation plates post-IMS to be very difficult, due to high background contamination (Conrad et al., 2014); but selective agar media such as CT-SMAC for O157 increases the likelihood of detection. Currently there are no such selective agars for STEC O55 and most other non-O157 STEC serogroups, therefore, for our studies high background interfered with the ability to identify colonies of *E. coli* O55. Further work is required to identify an O55 specific metabolic marker, which could be exploited to develop a novel isolation agar.

It has been demonstrated in many studies that the use of PCR provides a robust and sensitive method to screen samples for STEC. The inclusion of a PCR step in our assay improved the specificity and sensitivity of *E. coli* O55 identification. Importantly, background levels of non-O55 *E. coli* did not affect the limit of detection for IMS-PCR. IMS-PCR on samples that had been enriched for 2 h was able to detect O55 spiked in faeces at concentration of <10 CFU ml⁻¹. The infectious dose of STEC O157:H7 is thought to be less than 100 CFU (Smith et al., 2014). The infectious dose for STEC O55 is unknown but if we assume a similar dose is required our IMS-PCR method is sensitive enough to detect below this level. However, the draw-backs of using PCR alone is the lack of an O55 isolate for further characterisation, as PCR does not demonstrate viability of the *E. coli*, and it does not directly link the virulence factors (*stx*) to the serogroup and isolate.

Nevertheless, as implemented in this study, the PCR element of an IMS-PCR procedure provided a robust screening tool to identify positive samples, while plating a portion of the IMS-eluted beads on agar plates provided colonies for further analysis.

The tools developed in this study were also used to screen cattle faeces collected from a farm linked to an outbreak of STEC O157, for the presence of *E. coli* O55. Cattle faeces were screened by IMS-PCR to determine if STEC O55 may simultaneously be present on farm. Two *E. coli* O55:H19 were successfully isolated from cattle faeces, which lacked virulence factors such as *stx2* and *eae* and were shown by WGS analysis to be distinct from the strain of *E. coli* O55:H7 that caused the Dorset outbreak (Schutz et al., 2017). It is unknown how common *E. coli* O55 is in cattle as researchers seldom examine animal faeces for the presence of commensal *E. coli* of a particular serogroup. In the UK, *E. coli* O55 (non-STEC) has previously been isolated from cattle faeces, however, this was during an investigation of antimicrobial resistance on a dairy farm and the *E. coli* O55 was selected because it had a particular resistance trait (Horton et al., 2016). Furthermore, analysis of clinical submissions from diseased animals in the UK, did not identify any STEC O55 isolates between 2005-2008 (Hutchinson et al., 2011). In that study, the predominant STEC serogroup in cattle was O26. However, STEC O55 have been detected in cattle samples in other countries including surveys of beef cattle (faeces) in Japan (Mekata et al., 2014), and in Ireland STEC O55 has been reported from beef carcasses (Monaghan et al., 2012).

Core genome analysis demonstrated the two O55:H19 were distinct from O157:H7 and O55:H7 outbreak isolates, which were more similar to each other. Phylogenetic analysis of STEC O157:H7 and STEC O55:H7 has indicated that they have a common progenitor, and O157:H7 evolved through acquisition of the Shiga toxin genes (Schutz et al., 2017). Our analysis suggested that other non-STEC O55 which are unrelated to the outbreak clone may be present in the cattle faeces. Notably, these results indicate that the O55 IMS-PCR method developed in this study is a sensitive and specific tool that can be used to identify the presence of *E. coli* O55 in faecal specimens. The isolate can be purified subsequently by IMS-culture and characterised using other molecular methods such as WGS.

Therefore the use of IMS O55-beads on enriched samples, followed by PCR is a method for selection of *E. coli* O55 from contaminated samples, especially those from cattle faeces where *E. coli* O55 may be present at low levels. In the future, the use of these tools to examine other samples such as environmental samples could be explored, to help support investigation of human outbreaks of infection with STEC O55. The low infectious dose characteristic of STEC makes the need for detecting low-level contamination a key component of any STEC assay. Due to the lack of selective agars for the isolation of non-O157 STEC, in future, consideration will be given to the evaluation of selective media for the differentiation and recovery of O55-STEC from culture. Optimisation of enrichment could also be considered as the presence of high levels of non-target *E. coli* can interfere with binding of target *E. coli*, which could be influenced by enrichment media, temperature and time. In conclusion the tools developed in this study, could be used in future for rapid analysis of human, animal or environmental samples during outbreak investigations.

ACKNOWLEDGEMENTS

The authors would like to thank the Department for Environment, Food & Rural Affairs (Defra) for funding this research and Defra, Scottish Government and Welsh Government for funding zoonoses outbreak investigation under APHA devolved surveillance contract B FZ2100. We would also like to acknowledge the contribution of Charlotte Featherstone (APHA).

CONFLICT OF INTEREST

None to declare.

References

- ADAMS, N. L., BYRNE, L., SMITH, G. A., ELSON, R., HARRIS, J. P., SALMON, R., SMITH, R., O'BRIEN, S. J., ADAK, G. K. & JENKINS, C. 2016. Shiga Toxin-Producing *Escherichia coli* O157, England and Wales, 1983-2012. *Emerg Infect Dis*, 22, 590-7.
- AKTAN, I., CARTER, B., WILKING, H., LA RAGIONE, R. M., WIELER, L., WOODWARD, M. J. & ANJUM, M. F. 2007. Influence of geographical origin, host animal and stx gene on the virulence characteristics of *Escherichia coli* O26 strains. *J Med Microbiol*, 56, 1431-9.
- ANJUM, M. F., JONES, E., MORRISON, V., TOZZOLI, R., MORABITO, S., TOTH, I., NAGY, B., SMITH, G., ASPAN, A., NIELSEN, E. M., FACH, P., HERRERA-LEON, S., WOODWARD, M. J. & RM, L. A. R. 2014. Use of virulence determinants and seropathotypes to distinguish high- and low-risk *Escherichia coli* O157 and non-O157 isolates from Europe. *Epidemiol Infect*, 142, 1019-28.
- BAI, J., PADDOCK, Z. D., SHI, X., LI, S., AN, B. & NAGARAJA, T. G. 2012. Applicability of a multiplex PCR to detect the seven major Shiga toxin-producing *Escherichia coli* based on genes that code for serogroup-specific O-antigens and major virulence factors in cattle feces. *Foodborne Pathog Dis*, 9, 541-8.
- BOSILEVAC, J. M. & KOOHMARAIE, M. 2012. Predicting the presence of non-O157 Shiga toxin-producing *Escherichia coli* in ground beef by using molecular tests for

Shiga toxins, intimin, and O serogroups. *Appl Environ Microbiol*, 78, 7152-5.

BROCK, A. 2017. Guildford woman critically ill in hospital after suffering organ failure from 'confirmed case of E.coli' infection. *Get Surrey*.

CARTER, B., WU, G., WOODWARD, M. J. & ANJUM, M. F. 2008. A process for analysis of microarray comparative genomics hybridisation studies for bacterial genomes. *BMC Genomics*, 9, 53.

CONRAD, C. C., STANFORD, K., MCALLISTER, T. A., THOMAS, J. & REUTER, T. 2014. Further development of sample preparation and detection methods for O157 and the top 6 non-O157 STEC serogroups in cattle feces. *J Microbiol Methods*, 105, 22-30.

DEBROY, C., FRATAMICO, P. M., ROBERTS, E., DAVIS, M. A. & LIU, Y. 2005. Development of PCR assays targeting genes in O-antigen gene clusters for detection and identification of Escherichia coli O45 and O55 serogroups. *Appl Environ Microbiol*, 71, 4919-24.

DUGGETT, N. A., SAYERS, E., ABUOUN, M., ELLIS, R. J., NUNEZ-GARCIA, J., RANDALL, L., HORTON, R., ROGERS, J., MARTELLI, F., SMITH, R. P., BRENA, C., WILLIAMSON, S., KIRCHNER, M., DAVIES, R., CROOK, D., EVANS, S., TEALE, C. & ANJUM, M. F. 2017. Occurrence and characterization of mcr-1-harbouring Escherichia coli isolated from pigs in Great Britain from 2013 to 2015. *J Antimicrob Chemother*, 72, 691-695.

EICHHORN, I., HEIDEMANN, K., SEMMLER, T., KINNEMANN, B., MELLMANN, A., HARMSSEN, D., ANJUM, M. F., SCHMIDT, H., FRUTH, A., VALENTIN-WEIGAND, P., HEESEMANN, J., SUERBAUM, S., KARCH, H. & WIELER, L. H. 2015. Highly Virulent Non-O157 Enterohemorrhagic Escherichia coli (EHEC) Serotypes Reflect Similar Phylogenetic Lineages, Providing New Insights into the Evolution of EHEC. *Appl Environ Microbiol*, 81, 7041-7.

GUY, R. A., TREMBLAY, D., BEAUSOLEIL, L., HAREL, J. & CHAMPAGNE, M. J. 2014. Quantification of E. coli O157 and STEC in feces of farm animals using direct multiplex real time PCR (qPCR) and a modified most probable number assay comprised of immunomagnetic bead separation and qPCR detection. *J Microbiol Methods*, 99, 44-53.

HALLEWELL, J., ALEXANDER, T., REUTER, T. & STANFORD, K. 2017. Limitations of Immunomagnetic Separation for Detection of the Top Seven Serogroups of Shiga Toxin-Producing Escherichia coli. *J Food Prot*, 80, 598-603.

HORTON, R. A., DUNCAN, D., RANDALL, L. P., CHAPPELL, S., BRUNTON, L. A., WARNER, R., COLDHAM, N. G. & TEALE, C. J. 2016. Longitudinal study of CTX-M ESBL-producing E. coli strains on a UK dairy farm. *Res Vet Sci*, 109, 107-113.

HUTCHINSON, J. P., CHENEY, T. E., SMITH, R. P., LYNCH, K. & PRITCHARD, G. C. 2011. Verocytotoxin-producing and attaching and effacing activity of Escherichia coli isolated from diseased farm livestock. *Vet Rec*, 168, 536.

IHEKWEAZU, C., CARROLL, K., ADAK, B., SMITH, G., PRITCHARD, G. C., GILLESPIE, I. A., VERLANDER, N. Q., HARVEY-VINCE, L., REACHER, M., EDEGHERE, O.,

SULTAN, B., COOPER, R., MORGAN, G., KINROSS, P. T., BOXALL, N. S., IVERSEN, A. & BICKLER, G. 2012. Large outbreak of verocytotoxin-producing *Escherichia coli* O157 infection in visitors to a petting farm in South East England, 2009. *Epidemiol Infect*, 140, 1400-13.

ISLAM, M. A., HEUVELINK, A. E., TALUKDER, K. A., ZWIETERING, M. H. & DE BOER, E. 2006. Evaluation of immunomagnetic separation and PCR for the detection of *Escherichia coli* O157 in animal feces and meats. *J Food Prot*, 69, 2865-9.

JENKINS, C., DALLMAN, T. J., LAUNDERS, N., WILLIS, C., BYRNE, L., JORGENSEN, F., EPPINGER, M., ADAK, G. K., AIRD, H., ELVISS, N., GRANT, K. A., MORGAN, D. & MCLAUCHLIN, J. 2015. Public Health Investigation of Two Outbreaks of Shiga Toxin-Producing *Escherichia coli* O157 Associated with Consumption of Watercress. *Appl Environ Microbiol*, 81, 3946-52.

JOHNSON, K. E., THORPE, C. M. & SEARS, C. L. 2006. The emerging clinical importance of non-O157 Shiga toxin-producing *Escherichia coli*. *Clin Infect Dis*, 43, 1587-95.

KANKI, M., SETO, K. & KUMEDA, Y. 2014. Simultaneous Immunomagnetic Separation Method for the Detection of *Escherichia coli* O26, O111, and O157 from Food Samples. *J Food Prot*, 77, 15-22.

KRUGER, A. & LUCCHESI, P. M. A. 2015. Shiga toxins and *stx* phages: highly diverse entities. *Microbiology*, 161, 451-462.

LEJEUNE, J. T., BESSER, T. E., RICE, D. H. & HANCOCK, D. D. 2001. Methods for the isolation of water-borne *Escherichia coli* O157. *Lett Appl Microbiol*, 32, 316-20.

MCFARLAND, N., BUNDLE, N., JENKINS, C., GODBOLE, G., MIKHAIL, A., DALLMAN, T., O'CONNOR, C., MCCARTHY, N., O'CONNELL, E., TREACY, J., DABKE, G., MAPSTONE, J., LANDY, Y., MOORE, J., PARTRIDGE, R., JORGENSEN, F., WILLIS, C., MOOK, P., RAWLINGS, C., ACORNLEY, R., FEATHERSTONE, C., GAYLE, S., EDGE, J., MCNAMARA, E., HAWKER, J. & BALASEGARAM, S. 2017. Recurrent seasonal outbreak of an emerging serotype of Shiga toxin-producing *Escherichia coli* (STEC O55:H7 Stx2a) in the south west of England, July 2014 to September 2015. *Euro Surveill*, 22.

MEKATA, H., IGUCHI, A., KAWANO, K., KIRINO, Y., KOBAYASHI, I. & MISAWA, N. 2014. Identification of O serotypes, genotypes, and virulotypes of Shiga toxin-producing *Escherichia coli* isolates, including non-O157 from beef cattle in Japan. *J Food Prot*, 77, 1269-74.

MILES, A. A., MISRA, S. S. & IRWIN, J. O. 1938. The estimation of the bactericidal power of the blood. *J Hyg (Lond)*, 38, 732-49.

MONAGHAN, A., BYRNE, B., FANNING, S., SWEENEY, T., MCDOWELL, D. & BOLTON, D. J. 2012. Serotypes and virulotypes of non-O157 shiga-toxin producing *Escherichia coli* (STEC) on bovine hides and carcasses. *Food Microbiol*, 32, 223-9.

MURASE, K., OOKA, T., IGUCHI, A., OGURA, Y., NAKAYAMA, K., ASADULGHANI, M., ISLAM, M. R., HIYOSHI, H., KODAMA, T., BEUTIN, L. & HAYASHI, T. 2012. Haemolysin E- and enterohaemolysin-derived haemolytic activity of O55/O157

strains and other *Escherichia coli* lineages. *Microbiology*, 158, 746-58.

NOLL, L. W., BAUMGARTNER, W. C., SHRIDHAR, P. B., CULL, C. A., DEWSBURY, D. M., SHI, X., CERNICCHIARO, N., RENTER, D. G. & NAGARAJA, T. G. 2016. Pooling of Immunomagnetic Separation Beads Does Not Affect Detection Sensitivity of Six Major Serogroups of Shiga Toxin-Producing *Escherichia coli* in Cattle Feces. *J Food Prot*, 79, 59-65.

PAZ, S. 2016. Dorset *E. coli*: Investigation fails to find source. Available: <http://www.bbc.co.uk/news/uk-england-dorset-38300159> [Accessed 25-07-17].

SCHUTZ, K., COWLEY, L. A., SHAABAN, S., CARROLL, A., MCNAMARA, E., GALLY, D. L., GODBOLE, G., JENKINS, C. & DALLMAN, T. J. 2017. Evolutionary Context of Non-Sorbitol-Fermenting Shiga Toxin-Producing *Escherichia coli* O55:H7. *Emerg Infect Dis*, 23, 1966-1973.

SMITH, J. L., FRATAMICO, P. M. & GUNTHER, N. W. T. 2014. Shiga toxin-producing *Escherichia coli*. *Adv Appl Microbiol*, 86, 145-97.

TREACY, J., JENKINS, C., PARANTHAMAN, K., JORGENSEN, F., MUELLER-DOBLIES, D., ANJUM, M., KAINDAMA, L., HARTMAN, H., KIRCHNER, M., CARSON, T. & KAR-PURKAYASTHA, I. 2019. Outbreak of Shiga toxin-producing *Escherichia coli* O157:H7 linked to raw drinking milk resolved by rapid application of advanced pathogen characterisation methods, England, August to October 2017. *Euro Surveill*, 24.

Tables and Figures

Table 1. Details of O55 strains used in this study. The origin and year of isolation are unknown for B2884 as this is a type strain from the Staten Serum Institut collection (<http://www.ssidiagnostica.com/da/Products/Bacterial-strains/E-coli>).

Table 2 Limit of detection for IMS-culture and IMS-PCR from spiked cattle faeces. The spiking concentration at which IMS-culture or IMS-PCR could detect O55. The results represent the average of triplicate experiments.

Figure 1. Average percentage recovery of O55 strains post-IMS with O55-beads. Error bars indicate the calculate standard error from mean.

Figure 2. Average percentage recovery of O22, O23, O83, O104 post-IMS with O55-beads at different initial cell concentrations (A. 10^2 CFU ml⁻¹ and B. 10^3 CFU ml⁻¹). Error bars indicate the calculated standard error from mean.

Figure 3. Tree generated using the SNP distance of core genomes from the isolates collected in this study (O55-S17 and O55-S20), E. coli O55 available in BioProjects (NCBI), and O157:H7 strains (44-APHA and 432299-PHE) collected during a recent outbreak investigation on this farm, and representatives of O157:H7 lineages described elsewhere (SRR4788235, SRR1981384, SRR4195765) (Schutz et al., 2017). The E. coli K-12 genome from MG1655 was used for reference.

Tables:

Table 1. Details of O55 strains used in this study. The origin and year of isolation are unknown for B2884 as this is a type strain from the Staten Serum Institut collection (<http://www.ssidiagnostica.com/da/Products/Bacterial-strains/E-coli>).

Strain ID	Year of isolation	Origin	Stx present	Serotype
B1362	2000	Bovine	Negative	O55:K+:-
B2884	Unknown	Unknown	Negative	O55:H-
165/13	2013	Bovine	Negative	O55:ND:ND
394/10	2010	Bovine	Negative	O55:ND:ND
328/11	2011	Bovine	Negative	O55:ND:ND
329/11	2011	Bovine	Negative	O55:ND:ND
384/10	2010	Bovine	Negative	O55:ND:ND
H1	2015	Human	Positive	O55:H51
H2	2016	Human	Positive	O55:H7
H3	2016	Human	Positive	O55:H12

ND= Not determined

Table 2 Limit of detection for IMS-culture and IMS-PCR from spiked cattle faeces. The spiking concentration at which IMS-culture or IMS-PCR could detect O55. The results represent the average of triplicate experiments.

Enrichment time	Limit of detection (CFU ml ⁻¹)	
	IMS-culture	IMS-PCR
0	52 (range 1-90)	52 (range 1-90)
2h	54.4 (range 7.2-90)	4.9 (range 1-7.2)

Figure 1. Average percentage recovery of O55 strains post-IMS with O55-beads. Error bars indicate the calculate standard error from mean.




Figure 2. Average percentage recovery of O22, O23, O83, O104 post-IMS with O55-beads at different initial cell concentrations (A. 10^2 CFU ml⁻¹ and B. 10^3 CFU ml⁻¹). Error bars indicate the calculated standard error from mean.




Figure 3. Tree generated using the SNP distance of core genomes from the isolates collected in this study (O55-S17 and O55-S20), *E. coli* O55 available in BioProjects (NCBI), and O157:H7 strains (44-APHA and 432299-PHE) collected during a recent outbreak investigation on this farm, and representatives of O157:H7 lineages described elsewhere (SRR4788235, SRR1981384, SRR4195765) (Schutz et al., 2017). The *E. coli* K-12 genome from MG1655 was used for reference.

