

MICROWAVE-WAVE SPECTROSCOPY OF 5-METHYL HYDANTOIN

HIROYUKI OZEKI, MINAMI AWATSU, *Department of Environmental Science, Toho University, Funabashi, Japan*; KAORI KOBAYASHI, *Department of Physics, University of Toyama, Toyama, Japan*.

Hydantoin (Imidazolidine-2,4-dione, $C_3H_4N_2O_2$) is a five-membered heterocyclic compound, and has been regarded as a direct precursor of glycine, the simplest amino acid. The molecule was detected in carbonaceous chondrites together with several kinds of amino acids.^a We have measured pure rotational spectrum of the molecule in its ground and vibrationally excited states, and have provided the molecular line frequency list in the millimeter-wave region for astronomical search.^b 5-methyl hydantoin is the simplest chiral molecule among hydantoins, because either of the hydrogen atoms bonded to the C5 position of the five-membered ring is asymmetrically substituted. The molecule becomes a direct precursor of alanine, the simplest chiral amino acid. We have tried to observe pure rotational spectrum of 5-methyl hydantoin in the millimeter-wave to sub-millimeter-wave region. Guided with quantum chemical calculation, several spectral lines can be assigned to b-type transitions.

^aA. Shimomiya, and R. Ogasawara, *Orig. Life Evol. Biosph.* 32, 165 (2002)

^bH. Ozeki, R. Miyahara, H. Ihara, S. Todaka, K. Kobayashi, and M. Ohishi, *Astron. Astrophys.* 600 A44 (2017).