



# Half-lives of PAHs and temporal microbiota changes in commonly used urban landscaping materials

Marja I. Roslund<sup>1</sup>, Mira Grönroos<sup>1</sup>, Anna-Lea Rantalainen<sup>1</sup>, Ari Jumpponen<sup>2</sup>, Martin Romantschuk<sup>1</sup>, Anirudra Parajuli<sup>1</sup>, Heikki Hyöty<sup>3</sup>, Olli Laitinen<sup>3</sup> and Aki Sinkkonen<sup>1</sup>

<sup>1</sup> Faculty of Biological and Environmental Sciences, Ecosystems and Environment Research Programme, University of Helsinki, Lahti, Finland

<sup>2</sup> Division of Biology, Kansas State University, Kansas, Manhattan, United States of America

<sup>3</sup> Faculty of Medicine and Life Sciences, University of Tampere, Tampere, Finland

## ABSTRACT

**Background.** Polycyclic aromatic hydrocarbons (PAHs) accumulate in urban soils, and PAH contamination can change soil microbial community composition. Environmental microbiota is associated with human commensal microbiota, immune system and health. Therefore, studies investigating the degradation of PAHs, and the consequences of soil pollution on microbial communities in urban landscaping materials, are crucial.

**Methods.** Four landscaping materials (organic matter 1, 2, 13 and 56%) were contaminated with PAHs commonly found at urban sites (phenanthrene, fluoranthene, pyrene, chrysene and benzo(b)fluoranthene) in PAH concentrations that reflect urban soils in Finland ( $2.4 \mu\text{g g}^{-1}$  soil dry weight). PAHs were analyzed initially and after 2, 4, 8 and 12 weeks by gas chromatography-mass spectrometry. Half-lives of PAHs were determined based on 12-weeks degradation. Bacterial communities were analyzed at 1 and 12 weeks after contamination using Illumina MiSeq 16S rRNA gene metabarcoding.

**Results.** Half-lives ranged from 1.5 to 4.4 weeks for PAHs with relatively low molecular weights (phenanthrene, fluoranthene and pyrene) in landscaping materials containing 1–2% organic matter. In contrast, in materials containing 13% and 56% organic matter, the half-lives ranged from 2.5 to 52 weeks. Shorter half-lives of phenanthrene and fluoranthene were thus associated with low organic matter content. The half-life of pyrene was inversely related to the relative abundance of Beta-, Delta- and Gammaproteobacteria, and diversity of Bacteroidetes and Betaproteobacteria. Compounds with higher molecular weights followed compound-specific patterns. Benzo(b)fluoranthene was resistant to degradation and half-life of chrysene was shorter when the relative abundance of Betaproteobacteria was high. Temporal microbiota changes involved increase in the relative abundance of Deltaproteobacteria and decrease in genera *Flavobacterium* and *Rhodanobacter*. Exposure to PAHs seems to adjust microbial community composition, particularly within class Beta- and Deltaproteobacteria.

**Conclusions.** In this study, PAH degradation depended on the organic matter content and bacterial community composition of landscaping materials. Contamination seems to alter bacterial community composition in landscaping materials depending on material type. This alteration includes changes in bacterial phyla associated with human health and immune system. This may open new possibilities for managing urban environments by careful selection of landscaping materials, to benefit health and wellbeing.

Submitted 12 October 2017  
Accepted 26 February 2018  
Published 19 March 2018

Corresponding authors

Marja I. Roslund,  
[marja.roslund@helsinki.fi](mailto:marja.roslund@helsinki.fi)  
Aki Sinkkonen,  
[aki.sinkkonen@helsinki.fi](mailto:aki.sinkkonen@helsinki.fi)

Academic editor  
Valery Forbes

Additional Information and  
Declarations can be found on  
page 20

DOI 10.7717/peerj.4508

 Copyright  
2018 Roslund et al.

Distributed under  
Creative Commons CC-BY 4.0

**OPEN ACCESS**

**Subjects** Biodiversity, Ecology, Ecosystem Science, Environmental Sciences, Soil Science

**Keywords** Polycyclic aromatic hydrocarbons, Urban pollution, Biodegradation, Microbial community composition, Human health, Health-associated bacteria

## INTRODUCTION

Polycyclic aromatic hydrocarbons (PAHs) have been classified as priority environmental pollutants by the United States Environmental Protection Agency (USEPA) and by the European Environment Agency (EEA). PAHs constitute a large class of fused aromatic ring compounds that are present in petroleum products and formed from the incomplete combustion of fossil fuels and biomass (*Boström et al., 2002*). Combustion of fossil fuels is the most important source of PAHs in urban environments, and PAH exposure in large cities can be very high (*Boström et al., 2002; Ravindra, Sokhi & Van Grieken, 2008; Hong et al., 2016*). Airborne PAHs often accumulate in soils (*Haritash & Kaushik, 2009*) and the concentrations in urban soils can be 10–100 times higher than in unpolluted rural soils (*Hiller et al., 2015*). Microbial degradation of PAHs can be significant in soil depending on soil microbial activity and other factors, such as soil properties, climate and bioavailability of PAHs (*Koivula et al., 2004; Haritash & Kaushik, 2009; Crampon et al., 2014*). Besides microbial activity, other pathways for PAHs in the environment include volatilization, photo-oxidation, chemical oxidation, sorption to soil particles and leaching (*Haritash & Kaushik, 2009*).

Many bacterial species degrade PAHs, but only a limited number of bacteria degrade PAHs with five or more aromatic rings, mainly because of their low bioavailability (*Johnsen, Wick & Harms, 2005*). However, PAHs with high molecular weight may be degraded in a series of processes by consortia of microbes, such as bacteria and fungi (*Johnsen, Wick & Harms, 2005*). Many members of the phylum Proteobacteria degrade PAHs, and they are more abundant in PAH polluted soils compared to non-contaminated soils (*Singleton et al., 2006; Lors et al., 2010; Martin et al., 2012; Parajuli et al., 2017*). Typically, PAH contamination changes the abundances within the soil microbial communities, specifically including phyla Proteobacteria, Actinobacteria and Bacteroidetes (*Lors et al., 2010; Martin et al., 2012; Peng et al., 2013; Mukherjee et al., 2014; Sawulski, Clipson & Doyle, 2014; Ren et al., 2015; Parajuli et al., 2017*). For example, *Mukherjee et al. (2014)* reported that proteobacteria, particularly Betaproteobacteria and Gammaproteobacteria, dominated in the zones with high PAH pollution and the abundance of Actinobacteria declined in these sites, whereas *Sawulski, Clipson & Doyle (2014)* reported that Actinobacteria dominate in fluoranthene polluted soil, but compared to unpolluted soil more Gammaproteobacteria and less Bacteroidetes were identified in PAH polluted soil.

PAH exposure causes a variety of health risks, ranging from carcinogenic and mutagenic changes to respiratory problems, hematological changes and neurological dysfunctions, and PAHs may induce immune-suppression in humans (*Kim et al., 2013; Kamal et al., 2015*). In addition to the direct health consequences, PAHs may also change the abundance and diversity of human health-associated bacteria (*Parajuli et al., 2017*). Since the highest soil PAH concentrations are usually found in urban centers where source intensity is highest

(*Heywood et al., 2006; Haugland, Ottesen & Volden, 2008; Hiller et al., 2015*), microbial communities in urban soils are likely affected by PAH pollution. Altered environmental microbial communities may lead to changes in human commensal microbiota, in turn affecting the human immune system (*Ege et al., 2012; Hanski et al., 2012; Heederik & Mutius, 2012; Ruokolainen et al., 2015*). Many factors affect the composition of human commensal microbiota, and skin microbiota may differ between rural and urban children and this microbial composition is associated with allergic sensitization (*Hanski et al., 2012; Lehtimäki et al., 2017*). Despite this, studies connecting PAH pollution, microbial community composition in urban soils and health-associated bacteria are lacking.

Urban areas are often characterized by a high degree of artificial surfaces, both paved surfaces (asphalt, buildings etc.) and non-sealed surfaces (man-made gardening and landscaping materials) (*Artmann, 2014; Morabito et al., 2016*). Urban soil often differs drastically in composition and function from natural soils (*Morel, Chenu & Lorenz, 2015*). For example, due to mechanical compaction, urban soils have a high bulk density and can be composed of coarse anthropogenic materials with reduced biological activity (*Morel, Chenu & Lorenz, 2015*). Since more than half of the global human population lives in urban areas (*United Nations, 2014*), it is surprising how little attention urban landscaping materials have gained in pollution-oriented studies. Furthermore, laboratory studies with PAH contamination have often been carried out with unrealistically high concentrations rarely found in urban soils (*Johnsen, Wick & Harms, 2005; Crampon et al., 2014*). Therefore, there is a need for studies to investigate how fast realistic urban concentrations of pollutants are degraded in different landscaping materials, and what changes pollutants cause in bacterial communities in those materials.

In this study, we analyzed the bacterial community composition and diversity changes in landscaping materials contaminated with five different PAHs in a 12-week laboratory experiment. We specifically focused on PAH concentrations similar to those common in polluted urban soil. We hypothesized that the half-life and degradation of PAHs is connected to soil properties such as organic matter content and the pH of the landscaping materials. In addition, we sought to find potential associations between bacterial community composition and PAH degradation, and to estimate how much PAH degradation depends on the number of aromatic rings. We also hypothesized that realistic urban concentrations of PAHs cause changes in the microbial community composition in commonly used landscaping materials, and that these changes differ between different landscaping materials. We particularly expected to observe changes in the bacterial community composition, diversity and relative abundance of Actinobacteria, Bacteroidetes and Proteobacteria. Finally, we discuss bacterial community composition changes of human health-associated bacteria.

The type of landscaping material is of great importance, both for the PAH degradation in an urban environment and for the exposure of urban inhabitants to environmental microbes important for the human immune system. The current study provides information about landscaping materials for urban planners to build sustainable cities suitable for human health and wellbeing.

## MATERIALS AND METHODS

### Selection of landscaping materials

Four different landscaping materials were included in the study: Coarse peat-sand (Kekkilä Group, Vantaa, Finland), fine peat-sand (Kekkilä Group, Vantaa, Finland), gardening compost (Biolan, Eura, Finland) and sandy gravel of glacial origin (Rudus, Lahti, Finland) (Table S1). The selected landscaping materials represent the whole spectrum of the most widely used materials in Finland, ranging from microbiologically rich compost material to mineral aggregate fraction with a low microbial activity.

### Physicochemical analyses of materials

The organic matter content of each material was determined by a standardized loss-on-ignition method (SFS 3008, 1990). Moisture content was measured by drying samples in an oven (+105 °C) to constant weight and the water-holding capacity was determined with the method of Priha & Smolander (1999) and pH according to the SFS-ISO 10390 (2005) standard. Bioavailable nutrients (nitrate  $\text{NO}_3^-$ , ammonium  $\text{NH}_4^+$  and phosphate  $\text{PO}_4^{3-}$ ) were extracted and filtered (Whatman® 5892, White ribbon, by Schleicher and Schuell) as described by Sinkkonen *et al.* (2013) using the QuikChem 8000 flow injection analysis system (LACHAT Instruments Inc., Loveland, CO, USA). Approximately 10 g of fresh landscaping material was dried and crushed to measure elements (Al, Cu, Fe, Mn, P, Pb) with Inductive Coupled Plasma Mass Spectrometry (ICP-MS, Elan 6000; Perkin Elmer Inc., Waltham, MA, USA), and to measure carbon and nitrogen with the LECO C/N/S-200-analyzer (Leco Corporation, Saint Joseph, MI, USA).

### PAH degradation experiment

There were five replicates (500 ml glass pots with lids and passive aeration, 100 g dw per pot) per landscaping material that were contaminated artificially with a total of  $2.4 \mu\text{g g}^{-1}$  dw of PAHs phenanthrene, fluoranthene, pyrene, chrysene and benzo[b]fluoranthene (Table S2). These five PAHs are often found from urban sites (Boström *et al.*, 2002; Saarnio *et al.*, 2008; Tarvainen *et al.*, 2013). Contamination level for phenanthrene, fluoranthene and pyrene was  $0.6 \mu\text{g g}^{-1}$ , and for chrysene and benzo[b]fluoranthene it was  $0.3 \mu\text{g g}^{-1}$ . Total PAH concentrations were similar to those in man-made urban soils in Finland (approximately  $0.3\text{--}3.6 \mu\text{g g}^{-1}$ , mean  $1.3 \pm 1.2$ ) (Tarvainen *et al.*, 2013). For PAH measurements, one non-contaminated control replicate was included for each landscaping material to measure background PAH levels, i.e., to ensure that materials were uncontaminated.

Before mixing PAHs, each landscaping material was homogenized, sieved ( $\phi$  0.5 mm), and rehydrated to 60% of its water-holding capacity (Table S1). An aliquot (2 g) of each material (five replicates) was spiked with a total concentration of  $\Sigma 5\text{PAHs} = 120 \mu\text{g g}^{-1}$  diluted in acetone. Non-contaminated replicates were spiked with acetone. For bacterial sequencing, five 2 g subsamples were randomly drawn (five sampling points from each non-contaminated control at the depth of 1–6 cm). The five subsamples were combined to generate a separate non-contaminated sample per each landscaping material that represented bacterial community composition in non-contaminated conditions. This

method has been shown to be able to detect the dominant members of the bacterial community in heterogeneous soil (Manter, Weir & Vivanco, 2010). The spiked materials were mixed and kept in a fume hood for 1 day to allow acetone to evaporate. Then,  $2 \times 2$  g of each spiked material were added to  $2 \times 98$  g (in dry equivalent) of the corresponding wet material and mixed carefully and divided in two to obtain the initial concentration (Table S3). Spiked materials were stored at  $16 \pm 1$  °C in glass flasks (500 ml) with an air hole ( $\varnothing$  3 mm) in the dark and the moisture content was kept at 60% of their water-holding capacity. The experimental temperature represents an average temperature in the southern part of Finland during summer (Finnish Meteorological Institute, 2017). PAH analyses were carried out immediately after spiking and after four, eight and 12 weeks. Initial concentrations for chrysene and benzo[b]fluoranthene were confirmed after two weeks. This was done because the extractability, especially for high molecular weight PAHs, may decrease after two weeks (Luo et al., 2012). DNA was analyzed for bacterial (16S) communities at weeks 1 and 12 using Illumina MiSeq metabarcoding (Table S4).

### PAH analyses

PAH concentrations were determined using hexane and acetone mixture (1:1 v/v) extraction and analyzed as described by Honkonen & Rantalainen (2013) using gas chromatography-mass spectrometry (Shimadzu GC-MS-QP5000) system equipped with an AOC-20i autoinjector and a 30-m ZB-5MS column (0.25 mm i.d., 0.25  $\mu$ m film thickness). The mass spectrometer interface temperature was set to 280 °C. The oven temperature program was set as follows: 80 °C for 1 min, 10 °C/min to 250 °C, 7 °C/min to 280 °C and 20 °C/min to final 320 °C for 5 min. We used a deuterated PAH mixture (naphthalene-*d*8, acenaphthene-*d*10, phenanthrene-*d*10, chrysene-*d*12 and perylene-*d*12) (Dr. Ehrenstorfer GmbH, Germany) as a standard and Anthracene-*d*10 (Dr. Ehrenstorfer GmbH, Germany) as a recovery standard.

### Kinetic modeling and estimation of half-life times

Kinetic analysis was used to determine the degradation of PAHs (Oleszczuk & Baran, 2003; Zahed et al., 2011). Half-life of PAHs was determined on the basis of the kinetic equation of the first order (Zahed et al., 2011).

$$C = C_0 e^{-kt} \quad (1)$$

where  $C$  is the PAH concentration ( $\text{ng g}^{-1}$ ) at time  $t$  (weeks)  $C_0$  is the initial concentration of PAH ( $\text{ng g}^{-1}$ ) and  $k$  is rate constant of the change in the PAH concentration ( $\text{week}^{-1}$ ). Half-life times ( $t_{1/2}$ ) were calculated by Eq. (2) (Oleszczuk & Baran, 2003; Zahed et al., 2011).

$$t_{1/2} = \frac{\ln 2}{k} \quad (2)$$

where  $k$  is the rate constant of Eq. (1). We also calculate the percent concentrations ( $C\%$ ) at each time point based on the decrease in the total concentration of PAHs using the following Eq. (3).

$$C\% = 100 - \frac{(C_i - C_r)}{C_i} * 100 \quad (3)$$

where  $C_i$  is the initial and  $C_r$  the residual PAH concentration.

### DNA extraction, PCR and Illumina MiSeq sequencing

Samples for MiSeq sequencing were prepared as in [Veach, Dodds & Jumpponen \(2015\)](#). Samples were stored in the deep freezer ( $<-70$  °C) before DNA extraction. Total DNA was extracted from 0.25–0.3 g of materials using a PowerSoil<sup>®</sup> DNA Isolation Kit (MoBio Laboratories, Inc., Carlsbad, CA, USA) according to the manufacturer standard protocol. Sterile water was used as negative control. DNA was checked with agarose gel (1.5%) electrophoresis. Total DNA concentration was measured with Quant-iT<sup>™</sup> PicoGreen<sup>®</sup> dsDNA reagent kit (Thermo Fisher Scientific, Waltham, MA, USA). The DNA concentration was adjusted to 0.35–0.4 ng/ $\mu$ l for each sample. DNA was analyzed for bacterial (16S) communities using a two-step PCR approach to avoid a 3'-end amplification bias resulting from the sample-specific DNA tags ([Berry et al., 2011](#)). The V4 region within the 16S ribosomal RNA (rRNA) gene was amplified by primary PCR (three replicates from each sample) using 505F and 806R primers ([Caporaso et al., 2012](#)). Primary PCR was carried out in a reaction mixture (reaction volume 50  $\mu$ l) consisting of 1  $\mu$ l each of 10 mM deoxynucleoside triphosphates (dNTPs; Thermo scientific, MA, USA) 5  $\mu$ l forward primer 505F (10  $\mu$ M; 5'-GTGCCAGCMGCCGCGGTAA-3') and 5  $\mu$ l reverse primer 806R (10  $\mu$ M; 5'-GGACTACHVGGGTWTCTAAT-3'), 0.5  $\mu$ l 2 U/ $\mu$ l Phusion Green Hot Start II High-Fidelity DNA polymerase (Thermo Fisher Scientific, Waltham, MA, USA), 10  $\mu$ l 5 $\times$  Green HF PCR buffer (F-537), 5  $\mu$ l template DNA and 23.5  $\mu$ l sterile water. The PCR reaction was run in a thermocycler (MJ Research, Waltham, MA, USA) as follows: initial denaturation at 98 °C for 5 min, followed by 25 cycles with denaturation at 94 °C for 1 min, annealing for 10 sec at 50 °C and extension for 1 min at 72 °C, and then a final extension at 72 °C for 10 min. A positive control (*Cupriavidus necator* JMP134, DSM 4058) was included in PRC to ensure that the PCR worked, and a negative control (sterile water) to detect possible contaminations. DNA was detected with agarose gel (1.5%) electrophoresis. The PCR products were purified using Agencourt AMPure XP solution (Beckman Coulter Inc., Brea, CA, USA) (1:1 ratio of bead solution to PCR volume) to minimize the carryover of primary PCR primers. Three replicates of the cleaned amplicons were pooled and diluted 1:5.

Cleaned and diluted primary PCR products were targeted in the secondary PCR (TagPCR). Reaction mixture to the TagPCR was equal as above except reverse primer included a 12 bp unique Multiplexing Identifier tag (MID-806R). The amplification program was the same as above except that there were only ten cycles with initial denaturation and final extension. TagPCR products were visualized on an agarose gel, purified with Agencourt AMPure and three replicates were pooled similarly to primary PCRs. DNA concentration for each sample was measured with PicoGreen and samples were pooled to equal amounts–150 ng. The sequencing was performed at the Kansas State University using Illumina MiSeq platform with a 2  $\times$  300 bp version 3 kit sequencing kit according to manufacturer's protocol. The GeneRead DNA Library I Core Kit (catalog

# 180432; Qiagen, Hilden, Germany) was used to ligate Illumina's TruSeq adapters to amplicons.

## Bioinformatics

Raw sequencing data was processed using Mothur (version 1.38.1, [Schloss et al., 2009](#)). The sequence processing protocol partly followed the pipeline suggested by [Schloss, Gevers & Westcott \(2011\)](#) and [Kozich et al. \(2013\)](#). The paired sequences contained in reverse and forward fastq files were aligned into a contig. Sequences were trimmed and screened to remove any that had mismatches with primer or DNA-tag sequences, ambiguous bases or homopolymers larger than 8 bp long.

Bacterial sequences were aligned against a SILVA reference (version 123, [Pruesse et al., 2007](#)), preclustered to minimize sequencing errors ([Huse et al., 2010](#)) and screened for chimeras with UCHIME ([Edgar et al., 2011](#)), which uses the abundant sequences as a reference. The chimeric sequences were removed and non-chimeric sequences were classified using the Mothur version of Bayesian classifier ([Wang et al., 2007](#)) with the RDP training set version 14 ([Cole et al., 2009](#)) with 80% bootstrap threshold. Sequences were removed from the analyses if classified to Chloroplast, Mitochondria, unknown, Archaea or Eukaryota.

A pairwise distance matrix for unique sequences was calculated and OTUs clustered at 97% sequence similarity using nearest neighbor (single linkage) joining. All bacterial sequence data was accessioned into the Sequence Read Archive (Sequence read Archive Accession [SRX3195289–SRX3195338](#)). Low abundance OTUs ( $\leq 2$ ) were removed. Sequence profiles revealed similar taxonomic classifications between negative controls and sandy gravel, which is typical with samples containing a low microbial biomass ([Salter et al., 2014](#)). Altogether 14 abundant OTUs detected in negative controls were removed from sequence data and sandy gravel was discarded from community composition analyses because of the questionable sequence reads for bacteria. Other samples were subsampled to the smallest sample sequence depth at 6374 sequences for community composition analyses. Good's coverage index (average  $\pm$  SD:  $0.93 \pm 0.04$ ) was used to determine OTU coverage adequacy for diversity and community composition analyzes. OTU richness (the Chao index) and diversity (the Shannon index) were calculated with summary.single command ([Table S5](#)).

## Statistics

**Chemical dissipation.** Multivariate analysis of variance (MANOVA) with Bonferroni correction in SPSS (v24) was used to compare half-life of PAHs (log transformed) in different landscaping materials. The linear regression models were used to estimate the relationship between half-lives and properties of the landscaping materials, and between half-lives and bacteria relative abundances and diversities.

**Bacterial community analysis.** PERMANOVA (R v3.4.0, [R Core Team, 2017](#), function *adonis* in *vegan* with Bray-Curtis metric) was used to compare temporal differences in the bacterial community composition between weeks 1 and 12. The non-metric multidimensional scaling (NMDS) in R was used to visualize the difference in bacterial

community composition (*meta* MDS function in the *vegan* package). Bray-Curtis distance matrices (R-package *vegan*) were used to compare each non-contaminated pot with the five contaminated pots within each landscaping material. In detail, Bray-Curtis distances were calculated between composite non-contaminated sample and contaminated samples. This resulted in five Bray-Curtis distances per landscaping material within each time point. Shannon diversity index for bacterial phylotypes was determined by using the function *diversity* in the *vegan* package. Temporal microbiota changes were determined for dominant phyla (Actinobacteria, Bacteroidetes and Proteobacteria) and classes (Alpha-, Beta-, Delta and Gammaproteobacteria), and to genera which abundance was more than 1%. Finally, the differences in the richness (the Chao index), diversity (the Shannon index) and relative abundances between weeks 1 and 12 were determined using the paired *t*-test in R.

## RESULTS

### Degradation of PAHs

Recovery of phenanthrene from freshly spiked sandy gravel ranged from 30% to 42%, indicating that phenanthrene evaporated from sandy gravel when acetone was permitted to evaporate in the fume hood. Other recoveries from freshly spiked materials were on average  $101 \pm 15\%$  for phenanthrene,  $105 \pm 12\%$  for fluoranthene,  $110 \pm 13\%$  for pyrene,  $91 \pm 17\%$  for chrysene and  $113 \pm 21\%$  for benzo[b]fluoranthene (Table S3). Recoveries for benzo[b]fluoranthene were very high at week 2 ( $171 \pm 37$ ) compared to the initial recoveries and they were still high after 12 weeks ( $114 \pm 15$ ). Consequently, half-life for benzo[b]fluoranthene cannot be reasonably calculated. Surrogate standard recoveries were, on average,  $77 \pm 17\%$  for all PAH analytes (Table S3).

For the three PAHs with molecular weight  $\leq 202$  g mol<sup>-1</sup> (phenanthrene, fluoranthene and pyrene), the half-lives were shorter in coarse mineral soils with organic content of 1–2% (sandy gravel and coarse peat-sand) than in soils with organic content of 13% (fine peat-sand) and 56% (gardening compost) (MANOVA:  $F \geq 92$ ,  $df = 3$ ,  $p < 0.001$ ; Table 1, Table S6). In addition, half-life of pyrene was 36 weeks shorter in gardening compost than in fine peat-sand ( $F = 101$ ,  $df = 3$ ,  $p < 0.001$ ; Table 1, Table S6). Organic matter content was directly related to the half-life of phenanthrene ( $F = 13$ ,  $R = 0.65$ ,  $r^2 = 0.43$ ,  $df = 18$ ,  $p < 0.01$ ) and fluoranthene ( $F = 21$ ,  $R = 0.74$ ,  $r^2 = 0.55$ ,  $df = 17$ ,  $p < 0.001$ ) (Table S7). Half-life of pyrene was inversely related to the relative abundance of Beta-, Delta- and Gammaproteobacteria, and directly to the abundance of Actinobacteria and Alphaproteobacteria (Table 2). In addition, diversity of Bacteroidetes and Betaproteobacteria was inversely related to the half-life of pyrene (Table 2).

The two high molecular weight PAHs ( $\geq 228$  g mol<sup>-1</sup>: Chrysene and benzo[b]fluoranthene) were more persistent than the three compounds with lower molecular weights (Fig. 1). Half-life of chrysene was more than 3 times longer in sandy gravel than in other landscaping materials (MANOVA:  $F = 18$ ,  $df = 3$ ,  $p < 0.01$ ; Table S6). Half-life of chrysene was inversely related to the relative abundance of Proteobacteria ( $F = 6$ ,  $df = 11$ ,  $R = 0.59$ ,  $r^2 = 0.35$ ,  $p < 0.05$ ), particularly Betaproteobacteria ( $F = 8$ ,  $df = 11$ ,  $R = 0.64$ ,  $r^2 = 0.41$ ,  $p < 0.02$ ) (Table S7).



**Table 1** Half-lives of PAHs were determined based on 12 week's degradation. Half-lives are shown in weeks (Mean  $\pm$  SD, *n*, number of replicates).

	Coarse peat-sand	Fine peat-sand	Gardening compost	Sandy gravel
Phenanthrene	1.5 $\pm$ 0.1 ( <i>n</i> = 5)	2.8 $\pm$ 0.2 ( <i>n</i> = 5)	2.5 $\pm$ 0.1 ( <i>n</i> = 5)	1.5 $\pm$ 0.1 ( <i>n</i> = 5)
Fluoranthene	2.4 $\pm$ 0.2 ( <i>n</i> = 4)	39 $\pm$ 3.1 ( <i>n</i> = 4)	36 $\pm$ 7.0 ( <i>n</i> = 5)	2.8 $\pm$ 1.0 ( <i>n</i> = 5)
Pyrene	4.4 $\pm$ 0.6 ( <i>n</i> = 4)	52 $\pm$ 26 ( <i>n</i> = 5)	16 $\pm$ 1.8 ( <i>n</i> = 5)	2.7 $\pm$ 0.6 ( <i>n</i> = 5)
Chrysene	22 $\pm$ 4.7 ( <i>n</i> = 4)	49 $\pm$ 6.5 ( <i>n</i> = 5)	56 $\pm$ 18 ( <i>n</i> = 4)	198 $\pm$ 102 ( <i>n</i> = 4)
Sum of PAHs	5.6 $\pm$ 0.3 ( <i>n</i> = 4)	17 $\pm$ 0.7 ( <i>n</i> = 4)	14 $\pm$ 1.1 ( <i>n</i> = 5)	6.4 $\pm$ 1.6 ( <i>n</i> = 5)

**Table 2** Regression between half-life of pyrene and relative abundances and diversities of bacterial phyla and class. Half-life of pyrene correlated inversely with relative abundance of Beta-, Delta- and Gammaproteobacteria, and directly with the relative abundance of Actinobacteria and Alphaproteobacteria. In addition, regression was found between lower half-life of pyrene and higher diversity of Bacteroidetes and Betaproteobacteria.

	<i>df</i>	<i>F</i>	<i>R</i>	<i>r</i> <sup>2</sup>	<i>t</i>	<i>p</i> -value
<b>Abundance</b>						
Actinobacteria	13	9	0.64	0.41	3.0	0.010
Alphaproteobacteria	13	12	0.69	0.48	3.5	0.004
Betaproteobacteria	13	7	0.60	0.36	-2.7	0.018
Deltaproteobacteria	13	10	0.67	0.44	-3.2	0.007
Gammaproteobacteria	13	6	0.57	0.32	-2.5	0.027
<b>Diversity</b>						
Bacteroidetes	13	18	0.76	0.58	-4.3	0.001
Betaproteobacteria	13	7	0.60	0.36	-2.7	0.018


**Notes.**

*r*<sup>2</sup>, Coefficient of Determination; *df*, Degree of Freedom; *R*, Correlation Coefficient; *t*, *t*-test statistic.

## Bacterial community compositions


Three of the four landscaping materials contained detectable levels of bacterial sequence. Based on Illumina sequencing data, the rarefied data had 11,557 OTUs and an average of 27,463 bacteria sequences for coarse peat-sand, 79,943 for fine peat-sand and 58,600 for gardening compost.

Proteobacteria OTUs were most abundant in contaminated fine (37–44%) and coarse peat-sand (38–62%) and gardening compost (33–40%) throughout the experiment (Fig. 2). Other abundant bacterial phyla (mean abundance of more than 1%) were Bacteroidetes (11–41%), Actinobacteria (4–26%) and Acidobacteria (1–11%) in three landscaping materials (Fig. 2). Within phyla Proteobacteria, Alphaproteobacteria were most abundant in fine peat-sand (44–58%) and gardening compost (30–37%) throughout the experiment. In coarse peat-sand, Alphaproteobacteria were more abundant at week 12 (34–43%), while Gammaproteobacteria dominated in the week 1 samples (29–53%). The most abundant bacterial class within phyla Bacteroidetes in all soil types at week 1 was Sphingobacteriia (49–63% in coarse peat-sand, 38–59% in fine peat-sand and 78–90% in gardening compost). At week 12, Sphingobacteriia were still the dominant class in coarse peat sand (74–91%). However, Flavobacteriia was the most abundant identified class within Bacteroidetes in the fine peat-sand (10–27%) and gardening compost (12–28%).




**Figure 1** Decrease of PAHs. %-decrease of PAHs in the studied landscaping materials (mean  $\pm$  SD): (A) Total PAH concentration. (B) Phenanthrene. (C) Fluoranthene. (D) Pyrene. (E) Chrysenes. (F) benzo[b]fluoranthene.

Full-size DOI: 10.7717/peerj.4508/fig-1



**Figure 2** Relative abundances % at phylum level. Proteobacteria OTUs were most abundant in coarse peat-sand, fine peat-sand and gardening compost throughout the experiment.

Full-size DOI: 10.7717/peerj.4508/fig-2



**Figure 3** Temporal changes in the bacterial community composition. Bacterial community composition differed between contaminated landscaping materials and between weeks 1 and 12: (A) Genus level. (B) Class level. (C) Phylum level. Timepoints shown are: Circle, week 1; Triangle, week 12.

Full-size [DOI: 10.7717/peerj.4508/fig-3](https://doi.org/10.7717/peerj.4508/fig-3)

At the higher taxonomic levels, the three most abundant classified genera in the contaminated coarse peat-sand were *Pseudomonas*, *Arthrobacter* and *Terrimonas*, and *Arthrobacter*, *Rhodanobacter* and *Nitrospira* in the fine peat-sand, throughout the experiment (Table S8). The three most abundant classified genera in the gardening compost were *Aequorivita*, *Devosia* and *Flavobacterium* (Table S8), but the most abundant bacteria were unclassified Bacteroidetes ( $13 \pm 2\%$ ), Chloroflexi ( $3 \pm 1\%$ ) and other bacteria ( $5 \pm 2\%$ ). From the six most abundant classified bacteria genera *Arthrobacter* and *Devosia* were abundant in all three landscaping materials (Table S8).

### Temporal changes in bacterial community composition in landscaping materials

To determine temporal changes in bacterial communities, we compared bacterial community compositions between weeks 1 and 12 in contaminated landscaping materials. Bacterial community compositions differed between all three tested landscaping materials and between weeks 1 and 12 from OTU to phylum level (PERMANOVA:  $p < 0.001$ , Fig. 3, Table S9). In addition, community composition of Actinobacteria, Bacteroidetes and Proteobacteria (including Proteobacteria classes) differed between landscaping materials and weeks (PERMANOVA:  $p < 0.001$ , Table S9).

**Diversity and richness changes.** OTU diversity and richness were lower in fine peat-sand at week 1 compared to week 12 (Paired  $t$ -test:  $p < 0.001$ , Table S9). The changes in fine peat-sand involved increased diversity of Actinobacteria, Bacteroidetes, Proteobacteria, Alpha- and Gammaproteobacteria from week 1 to week 12 (Paired  $t$ -test:  $p < 0.01$ , Table S9). There were no diversity or richness changes at the OTU level in coarse peat-sand and in gardening compost, but diversity of Bacteroidetes and Deltaproteobacteria in coarse peat-sand and diversity of Gammaproteobacteria in gardening compost decreased during the experiment (Paired  $t$ -test:  $p < 0.05$ , Table S9).

**Relative abundance changes.** Relative abundance of Deltaproteobacteria was increased from week 1 to week 12 in all of the three landscaping materials (Paired  $t$ -test:  $p < 0.02$ ,

Table S9), whereas relative abundance of Betaproteobacteria was decreased in coarse peat-sand and in gardening compost (Paired  $t$ -test:  $p < 0.005$ , Table S9). Relative abundance of Bacteroidetes, Proteobacteria and Gammaproteobacteria decreased in gardening compost ( $p < 0.01$ ), whereas in fine peat-sand the relative abundance of Actinobacteria was decreased ( $p < 0.02$ ) and the relative abundance of Gammaproteobacteria increased from week 1 to week 12 ( $p < 0.01$ ) (Paired  $t$ -test: Table S9).

At a higher taxonomic level, the relative abundance of *Rhodanobacter* decreased from week 1 to week 12 in all three landscaping materials (Table 3). In coarse peat-sand and in gardening compost, the relative abundance of *Aerimonas* and *Flavobacterium* decreased during the experiment ( $p < 0.05$  and  $p < 0.001$ , respectively, Table 3). In fine peat-sand these abundances increased, but the relative abundances of these two genera were very low in fine peat-sand both in week 1 and 12 (Table 3). The relative abundance of *Arthrobacter* decreased both in fine peat-sand and in gardening compost and the relative abundance of *Pseudolabrys* decreased both in fine and coarse peat-sands (Table 3). In addition, temporal changes in gardening compost involved an increase in the relative abundances of *Gemmatimonas*, *Opitutus*, *Pseudolabrys* and *Terrimonas*, and decrease of *Aequorivita*, *Methylophilus* and *Pedobacter* (Table 3). In fine peat-sand, temporal changes additionally involved increase in the relative abundance of *Nitrosospira* and *Pseudomonas* and decrease in the relative abundance of *Devosia*, whereas the relative abundance of *Devosia* increased in coarse peat-sand (Table 3). One unclassified genus within the phylum Bacteroidetes and another within the class Gammaproteobacteria were increased in coarse ( $p < 0.05$ ) and fine peat-sands ( $p < 0.01$ ), whereas unclassified genus within family Xanthomonadaceae was decreased ( $p < 0.01$ ) and unclassified genus within order Myxococcales was increased in coarse peat-sand ( $p < 0.01$ ) and in gardening compost ( $p < 0.05$ ) (Paired  $t$ -test: Table S9).


### Bacterial community composition distances between contaminated and non-contaminated materials

Bacterial community composition distances (Bray-Curtis) were analyzed between contaminated (five replicates) and non-contaminated (a non-replicated composite sample from five subsamples) samples within each landscaping material. This resulted in five distances per landscaping material within each time point. Average distances were thereafter compared between different landscaping materials, and between the two time-points (weeks 1 and 12) within each landscaping material. The reasoning was to determine potential alterations associated with contamination.

Long bacterial community composition distances were observed particularly in fine peat-sand at week one (Table S5). At this time point, the relative abundance of unclassified Betaproteobacteria genus was high and Deltaproteobacteria, *Pseudomonas*, *Flavobacterium* and *Pedobacter* abundances were low in contaminated fine peat-sand compared to non-contaminated material (Fig. 4). In fine peat-sand, bacterial community composition distance at OTU level was  $0.58 \pm 0.01$  (Mean  $\pm$  SD) at week 1, whereas at week 12 this same distance was  $0.39 \pm 0.04$ . At different taxonomic levels, average bacterial community composition distances in fine peat-sand varied between 0.14–0.77 at week 1 and 0.10–0.49 at week 12 (Table S5). In coarse peat-sand, distances varied between 0.09–0.48 and 0.11–0.52

**Table 3** Relative abundances differed between week 1 and 12. According to *t*-tests results, relative abundances of abundant genera (abundance over 1%) changed during the experiment.

Genus	Coarse peat-sand				Fine peat-sand				Gardening compost			
	Mean week 1	Mean week 12	<i>t</i>	<i>p</i> -value	Mean week 1	Mean week 12	<i>t</i>	<i>p</i> -value	Mean week 1	Mean week 12	<i>t</i>	<i>p</i> -value
<i>Acinetobacter</i>	40 ± 49	15 ± 24	0.8	0.464	1 ± 1	25 ± 23	-2.0	0.110	108 ± 26	87 ± 101	0.4	0.680
<i>Aequorivita</i>	3 ± 2	0.2 ± 0.4	2.5	0.065	3 ± 2	2.4 ± 2	0.2	0.884	252 ± 80	91 ± 9	4.5	0.011
<i>Arenimonas</i>	121 ± 19	66 ± 25	3.7	0.022	1 ± 1	5 ± 2	-3.3	0.031	24 ± 3	17 ± 4	6.4	0.003
<i>Arthrobacter</i>	185 ± 41	175 ± 85	0.2	0.843	667 ± 181	361 ± 74	4.2	0.014	146 ± 32	91 ± 21	4.6	0.010
<i>Devosia</i>	81 ± 13	166 ± 37	-4.0	0.017	112 ± 11	92 ± 15	4.7	0.009	173 ± 17	164 ± 17	0.6	0.605
<i>Flavobacterium</i>	137 ± 26	13 ± 8	7.8	0.001	13 ± 5	22 ± 5	-2.9	0.045	245 ± 31	59 ± 7	12.3	0.000
<i>Gaiella</i>	83 ± 21	78 ± 18	0.4	0.701	77 ± 21	66 ± 11	0.8	0.489	1 ± 1	2 ± 1	-1.4	0.242
<i>Gemmatimonas</i>	71 ± 28	84 ± 32	-0.6	0.582	43 ± 12	37 ± 6	1.5	0.198	33 ± 6	56 ± 5	-4.3	0.012
<i>Methylophilus</i>	117 ± 36	89 ± 22	1.2	0.308	1 ± 1	2 ± 2	-1.3	0.256	68 ± 9	45 ± 6	3.8	0.019
<i>Nitrospira</i>	8 ± 3	11 ± 2	-1.7	0.163	55 ± 19	136 ± 14	-7.0	0.002	1 ± 1	1 ± 1	0.4	0.704
<i>Opitutus</i>	133 ± 72	154 ± 43	-0.6	0.595	13 ± 5	24 ± 10	-2.3	0.081	58 ± 15	94 ± 15	-3.5	0.026
<i>Pedobacter</i>	23 ± 3	10 ± 8	2.6	0.059	79 ± 8	95 ± 14	-1.7	0.163	224 ± 39	53 ± 9	9.1	0.001
<i>Pseudolabrys</i>	70 ± 16	39 ± 6	3.9	0.017	101 ± 11	75 ± 10	2.9	0.045	4 ± 3	9 ± 3	-4.4	0.012
<i>Pseudomonas</i>	748 ± 560	260 ± 235	1.5	0.199	22 ± 11	63 ± 29	-4.4	0.012	115 ± 19	40 ± 18	7.5	0.002
<i>Rhodanobacter</i>	15 ± 5	4 ± 2	5.3	0.006	191 ± 21	139 ± 32	3.1	0.038	80 ± 14	17 ± 4	11.4	0.000
<i>Terrimonas</i>	96 ± 45	224 ± 115	-2.5	0.068	1 ± 1	1 ± 0	0.0	1.000	16 ± 5	48 ± 7	-7.7	0.002



**Figure 4** Relative abundances of bacterial phylotypes. Relative abundance of seven bacteria phylotypes at week one are shown here in non-contaminated (blue) and contaminated (red) landscaping materials: (A) Coarse peat-sand. (B) Fine peat-sand. (C) Gardening compost.

Full-size DOI: 10.7717/peerj.4508/fig-4

at weeks 1 and 12, and in gardening compost between 0.06–0.42 and 0.04–0.36, respectively (Table S5).

Short bacterial community composition distances were observed at phylum level in all three landscaping materials (Mean variation at week 1 and 12 0.04–0.14, Table S5), indicating that contamination may not affect bacterial community composition at the phylum level. In contrast, the longest community composition distances at week one in coarse and fine peat-sands were observed within class Deltaproteobacteria ( $0.48 \pm 0.10$  and  $0.77 \pm 0.04$ , respectively). In addition, Betaproteobacteria community composition distances were long at week 1 both in fine peat-sand ( $0.66 \pm 0.06$ ) and in gardening compost ( $0.42 \pm 0.03$ ) compared to week 12 ( $0.32 \pm 0.03$  and  $0.34 \pm 0.05$ , respectively). At week 12, the longest community composition distance was observed within class Betaproteobacteria in coarse peat-sand ( $0.52 \pm 0.16$ ), whereas in fine peat-sand longest community composition distances were observed within classes Gammaproteobacteria ( $0.49 \pm 0.10$ ) and Deltaproteobacteria ( $0.48 \pm 0.07$ ), and in gardening compost within class Alphaproteobacteria ( $0.36 \pm 0.02$ ) (Table S5).

## DISCUSSION

### Connections between PAH degradation, bacterial community composition and temporal microbiota changes

We aimed to find potential associations between bacterial community composition and PAH degradation. In this study, higher abundances of Beta-, Delta- and Gammaproteobacteria were associated with a shorter half-life of pyrene. In addition, higher diversity of Bacteroidetes and Betaproteobacteria was associated with a shorter half-life of pyrene. However, there might be reasons (see ‘Connections between half-lives and properties of landscaping materials’ and ‘Connections between half-lives and molecular weight’) other than bacterial degradation ability that affected the half-life of pyrene. This is evident, since half-life of pyrene is short also in sandy gravel with undetectable bacterial sequence reads, whereas chrysene was very persistent in sandy gravel and higher relative abundance of Betaproteobacteria was associated with shorter half-life of chrysene. Betaproteobacteria abundance was higher in coarse peat-sand and gardening compost at week 1 when PAH concentrations were higher compared to week 12. In addition, in coarse peat-sand, where half-life of chrysene was shortest, Betaproteobacteria community composition had longest distance between contaminated and non-contaminated material compared to other distances at week 12. Based on previous studies, Betaproteobacteria may be an important indicator for the bioremediation process of PAHs (Lors *et al.*, 2010) and the primary phenanthrene degraders in soil (Martin *et al.*, 2012). In addition, in soils with high pyrene concentration, Betaproteobacteria abundance may increase, whereas Deltaproteobacteria abundance may concurrently decrease (Ren *et al.*, 2015). In agreement with these findings, we have observed a higher relative abundance of the Betaproteobacteria and a lower relative abundance of the Deltaproteobacteria in the creosote (includes ~85% PAH) contaminated soils as compared to non-contaminated soils in our previous study (Parajuli *et al.*, 2017).

In this study, temporal microbiota changes concentrated on dominant phyla and classes that have previously been associated with soil pollution, and to genera which abundance was more than 1%. Synchronous temporal microbiota changes in all three landscaping materials involved increase in relative abundance of Deltaproteobacteria as PAH levels decreased during the experiment. In contrast, the relative abundance of Betaproteobacteria decreased with decreasing PAH concentrations in coarse peat-sand and gardening compost. *Rhodanobacter* (Gammaproteobacteria) was the sole genus in which relative abundance decreased during the experiment in all three landscaping materials. *Rhodanobacter* genus includes denitrifying acid-tolerant bacteria ([Green et al., 2012](#)) that has been associated with PAH degradation in some studies ([Kanaly, Harayama & Watanabe, 2002](#); [Song et al., 2016](#); [Li et al., 2017](#)). However, *Rhodanobacter* genus involvement in *in situ* degradation remains questionable, and it seems that the genome lacks functional genes associated with PAH degradation ([Li et al., 2017](#)). Another genus for which the relative abundance decreased considerably during the experiment was *Flavobacterium* (Bacteroidetes). This decrease was obvious in coarse peat-sand and gardening compost. In fine peat-sand, the overall bacterial diversity and richness, and the diversity of Actinobacteria, Bacteroidetes and Proteobacteria, particularly Alpha- and Gammaproteobacteria, were lower at week 1 compared to week 12, whereas the diversity of Bacteroidetes and Deltaproteobacteria in coarse peat-sand, and the diversity of Gammaproteobacteria in gardening compost, decreased during the experiment. Decreased diversities might reflect the effect of the PAH contamination; previous studies have also reported that PAH contamination decreases bacterial diversity ([Mukherjee et al., 2014](#); [Ren et al., 2015](#)).

We additionally hypothesized that PAH concentrations typical to urban soils adjust microbial communities in commonly used landscaping materials and these changes differ between landscaping materials. In support of this hypothesis, we detected long distances between contaminated and non-contaminated fine peat-sand after one-week of exposure to PAHs. At this time point, the longest distances were observed within classes Beta- and Deltaproteobacteria ( $0.66 \pm 0.06$  and  $0.77 \pm 0.04$ , respectively) in fine peat-sand. Interestingly, the bacterial community composition distances of Beta- and Deltaproteobacteria seems to be associated with PAH contamination in all landscaping materials included in our study; this observation is in line with earlier investigations showing that these two classes are typically related to PAH contamination in soil ([Ren et al., 2015](#); [Parajuli et al., 2017](#)). However, it must be taken into account that we used a composite non-contaminated sample that represents bacterial composition at a community-wide scale, and we used this composite sample to calculate Bray-Curtis distances between non-contaminated and contaminated landscaping materials. On the one hand, this method can reduce variability in heterogeneous samples and detect the dominant phylotypes of the bacterial community ([Manter, Weir & Vivanco, 2010](#)). On the other hand, this method can influence the estimates of phylotype richness and diversity, and locally dominant but spatially rare phylotypes may become rare in the final sample ([Manter, Weir & Vivanco, 2010](#)). In support of our study findings, the phylotypes that we observed to have long distances between non-contaminated and contaminated samples include the same phylotypes that were detected in previous studies with PAH contamination ([Singleton et al.,](#)

2006; Lors *et al.*, 2010; Martin *et al.*, 2012; Mukherjee *et al.*, 2014; Ren *et al.*, 2015; Parajuli *et al.*, 2017). Additionally, the phylotypes that we observed to have long distances between contaminated and non-contaminated materials were the same as those associated with pyrene or chrysene degradation, i.e., Beta-, Delta- and Gammaproteobacteria.

After twelve-week follow-up, PAHs concentrations were lower and bacterial communities had greater similarity between contaminated and non-contaminated fine peat-sand. It is currently uncertain whether community composition changes are permanent or temporary in real urban environments where PAH deposition is continuous. However, previous research has revealed that the abundance of Betaproteobacteria may be greater in urban parks than in forests under evergreen trees (Hui *et al.*, 2017). This might be the consequence of the PAH deposition in urban parks, since Betaproteobacteria are associated with the PAH contamination in this and other studies (Lors *et al.*, 2010; Martin *et al.*, 2012; Mukherjee *et al.*, 2014; Ren *et al.*, 2015; Parajuli *et al.*, 2017).

### Connections between half-lives and properties of landscaping materials

We also hypothesized that the half-life of PAHs correlates with substrate properties, such as organic matter content and pH of the landscaping material. Half-lives of phenanthrene, fluoranthene and pyrene were considerably lower in landscaping materials with low organic contents of 1–2% (1.5–4.4 weeks) (sandy gravel and coarse peat-sand) than in fine peat-sand with organic content of 13% (2.8–52 weeks) and gardening compost with organic content of 56% (2.5–36 weeks). However, the organic matter content was unrelated to the half-life of pyrene. High organic matter content can decrease the bioavailability, degradation and diffusion of PAHs by increasing the sorption of PAHs to the organic fractions (Ran *et al.*, 2007; Yang *et al.*, 2010; Luo *et al.*, 2012). Thus, one reason for the longer half-lives for phenanthrene, fluoranthene and pyrene in fine peat-sand and in gardening compost might be the sorption of PAHs to the organic matter, while physico-chemical characteristics of each PAH and bacterial degradation ability may also have influences. Chemical persistence and physical properties of PAHs affect the biodegradation rate since bacteria are known to degrade only water dissolved compounds, and water solubility decreases and solid/water distribution ratio increases when the complexity of PAHs increases (Johnsen, Wick & Harms, 2005).

Both gardening compost and fine peat-sand had a carbon-to-nitrogen ratio of 25:1, which may have negatively impacted the biodegradation effectiveness, since a ratio of 10:1 has been shown to be more optimal (Teng *et al.*, 2010). Optimal carbon-to-nitrogen ratio offers enough nutrients for microbial growth, which may enhance the biodegradation (Teng *et al.*, 2010). In addition, there were differences in the nutrient concentrations between landscaping materials, which might have influenced the biodegradation effectiveness, since bioavailable nutrients may stimulate PAH degrading bacteria (Chaudhary *et al.*, 2012). Bacterial community composition can also be affected by pH gradient (Mukherjee *et al.*, 2014). However, pH was not correlated with PAH half-lives in this study.



## Connections between half-lives and molecular weight

Our goal was to estimate how much PAH degradation depends on the number of aromatic rings. Generally, a great diversity of bacteria can degrade the low molecular weight PAHs, but relatively few bacterial genera degrade PAHs with more than four aromatic rings, such as benzo[b]fluoranthene (Juhász & Naidu, 2000; Folwell, McGenity & Whitby, 2016). Thus, high molecular weight PAHs are more resistant to biodegradation (Johnsen, Wick & Harms, 2005; Haritash & Kaushik, 2009). Our results corroborate with these findings: benzo[b]fluoranthene was resistant to biodegradation and chrysene was very persistent in sandy-gravel and its half-life was over 17-weeks longer in coarse peat-sand and over 19-weeks longer in gardening compost than the half-lives of lower molecular weight PAHs. In fine peat-sand, the half-lives of pyrene (52 weeks) and chrysene (49 weeks) were almost the same, and approximately 10-weeks longer than the half-life of fluoranthene and over 46-weeks longer than the half-life of phenanthrene. Interestingly, the half-life of pyrene was 20-weeks shorter in gardening compost than the half-life of fluoranthene. This might be the effect of PAH mixture, since the degradation of PAH mixtures may be more multifaceted than degradation of individual compounds (Johnsen, Wick & Harms, 2005). With PAH mixtures, degradation of phenanthrene, fluoranthene and pyrene can be inhibited (Hennessee & Li, 2016), but phenanthrene can also enhance the degradation of higher molecular weight PAHs (McLellan, Warshawsky & Shann, 2002).

In sandy gravel, commonly used in urban landscaping (Morel, Chenu & Lorenz, 2015), persistence of PAHs with molecular weights  $\geq 228 \text{ g mol}^{-1}$  was high, resulting in the accumulation of higher molecular weight PAHs in urban soils and greater risks to human health. Long half-life of chrysene (198 weeks) confirms the persistence of chrysene in sandy gravel, but a first-order kinetic model is more appropriate for low molecular weight PAHs during a 12-weeks study (Crampon *et al.*, 2014). Thus, extrapolating a half-life calculation based on 12 weeks might be inaccurate for chrysene in sandy gravel, but it is used in this study to reflect only on the persistence (Tansel *et al.*, 2011). It is known that lower molecular weight PAHs are more volatile and phenanthrene, fluoranthene and pyrene exist in the gas and particle phase, whereas chrysene and benzo[b]fluoranthene exist exclusively in particle phase (Ravindra, Sokhi & Van Grieken, 2008). Therefore, in this study it is possible that phenanthrene, fluoranthene and pyrene volatilize out of porous materials with low organic matter content (low sorption to organic particles) like sandy gravel and coarse peat-sand. However, we did not quantify volatilization. Yet our study does not exclude the possible degradation of low molecular weight PAHs in sandy gravel despite the undetectable bacterial sequence reads. Preventing the volatilization of PAHs is important for human health. For example, fluoranthene is an experimental carcinogen and occurs at high concentrations in urban air (Boström *et al.*, 2002).

## Potential associations between bacterial community changes and health-associated bacteria

The “altered environmental microbiome” hypothesis, as an explanation for how environmental pollution is associated with microbiota and health, was first introduced in our previous study (Parajuli *et al.*, 2017). Bacterial community changes, i.e., reduced

bacterial diversity and altered Bacteroidetes and Proteobacteria communities, caused by PAH pollution (current study and others—[Singleton et al., 2006](#); [Lors et al., 2010](#); [Martin et al., 2012](#); [Peng et al., 2013](#); [Mukherjee et al., 2014](#); [Sawulski, Clipson & Doyle, 2014](#); [Ren et al., 2015](#); [Parajuli et al., 2017](#)) include changes in bacteria potentially connected to immune-mediated diseases ([Abrahamsson et al., 2012](#); [Hanski et al., 2012](#); [Knip & Siljander, 2016](#); [Zhou & Zhi, 2016](#); [Santoru et al., 2017](#)). PAH pollution often explains abundances of Beta- and Gammaproteobacteria ([Martin et al., 2012](#); [Peng et al., 2013](#); [Mukherjee et al., 2014](#); [Ren et al., 2015](#)) and these two classes may have a potential role in the early priming of the immune system ([Ruokolainen et al., 2015](#)). Decreased abundance of genus *Flavobacterium* within phyla Bacteroidetes in the human gut have been associated with a risk of inflammatory bowel disease, which is a chronic inflammatory disease of the gastrointestinal tract, including ulcerative colitis and Crohn's disease ([Santoru et al., 2017](#)). In addition, decreased abundance of *Bacteroides* have been associated with a risk of inflammatory bowel disease ([Zhou & Zhi, 2016](#); [Santoru et al., 2017](#)) and IgE-associated eczema ([Abrahamsson et al., 2012](#)). In contrast, Bacteroidetes dominance and the greater abundance of the genus *Bacteroides* have been associated to type 1 diabetes risk ([De Goffau et al., 2013](#); [Knip & Siljander, 2016](#)).

Our findings suggest that diversity of bacteria, particularly Bacteroidetes and Proteobacteria, may decrease with increased PAH concentration. Reduced overall bacterial diversity in the human gut is associated with type 1 diabetes ([Knip & Siljander, 2016](#)), eczema ([Abrahamsson et al., 2012](#); [Ismail et al., 2012](#)) and asthma ([Abrahamsson et al., 2014](#)). In addition, low genetic diversity of Gammaproteobacteria on the skin is associated with an increased risk of atopy and decreased expression of IL-10 which is an anti-inflammatory cytokine ([Hanski et al., 2012](#)). Because PAH contamination can change bacterial communities in polluted soils, it is also possible that PAH contamination can also directly affect the human microbiota. Bacteria, especially Proteobacteria, in the human gut contribute to a variety of biological functions, i.e., immune modulation and metabolism ([Bradley & Pollard, 2017](#)), and reduced bacterial functional diversity in the gut microbiome increases the risk of type 1 diabetes ([Brown et al., 2011](#)). Connecting bacterial community composition changes in PAH-contaminated landscaping materials at the laboratory scale and changes in human gut microbiota in patients with immunological disorders is highly hypothetical and we encourage further studies. In addition, it is unclear to what extent immune-mediated diseases are the cause and to what extent they are the consequence of bacterial community composition changes. Further studies are needed to determine how bacterial functions are affected during PAH contamination and to examine the possible associations between urban PAH pollution and microbial community changes both in the living environment and in the human gut. Such studies are especially important in children, because the immune system develops in early childhood.

### **Recommendations to manage urban environments in the context of human health**

As urban populations grow, the quality of the urban environment will play an important role in public health ([WHO, 2015](#)). In industrialized countries, approximately 78% of the

human population lives in urban areas with little green space and limited exposure to natural biodiversity (*United Nations, 2014*). According to the biodiversity hypothesis, this reduced contact with natural, resident microbial biodiversity negatively affects the human commensal microbiota and immune system (*Hanski et al., 2012*). Evidence exist that microbial communities in urban environments differ from those in rural environments, and that urbanization reduces the transfer of diverse environmental microbiota indoors (*Pakarinen et al., 2008; Qian et al., 2017; Parajuli et al., 2018*).

Two of the greatest challenges of urban societies are ecologically sustainable management of urban spaces, particularly soils and green areas (*Morel, Chenu & Lorenz, 2015*), and the control of rapidly increasing prevalence of immune-mediated diseases (*Hanski et al., 2012*). As a solution, we propose research targeting the development of novel gardening and landscaping materials that both speed up the degradation of PAHs and support the development of the human immune system. It must also be considered that vegetation in urban parks affects the soil microbiota (*Hui et al., 2017*) and effectiveness of PAH degradation (*Haritash & Kaushik, 2009*), and sometimes landscaping materials are mixed with other soil materials when used in urban areas. Novel gardening and landscaping materials could be designed to harbor biodiverse non-culturable microbial communities of forest floor to fill the gap in immunomodulation of urban dwellers that face only a fraction of the environmental microbiota present in natural and many rural ecosystems.

## CONCLUSIONS

Bacterial community composition, organic matter content and molecular weight of PAHs affects the half-life of PAHs in landscaping materials. We found a relationship between higher relative abundance of Betaproteobacteria and shorter half-lives of pyrene and chrysene. In addition, higher relative abundances of Gamma- and Deltaproteobacteria were correlated with shorter half-life of pyrene. High organic matter content ( $\geq 13\%$  in this study) might decrease the degradation efficiency by increasing the sorption of PAHs to organic fractions, whereas coarse-grained landscaping materials with low organic matter content (1–2%) might increase the probability of volatilization of PAHs with a molecular weight  $\leq 202$ .

Temporal microbiota changes were particularly evident within class Deltaproteobacteria, and genera *Flavobacterium* and *Rhodanobacter*. Relatively low PAH contamination seems to adjust Proteobacteria community composition, particularly Beta- and Deltaproteobacteria communities. PAH contamination may also result in a decreased bacterial diversity. Bacterial community composition alterations include changes in bacteria phyla associated with human health and immune-mediated diseases.

## ACKNOWLEDGEMENTS

The authors thank AlmaLab personnel from University of Helsinki for their help with laboratory analyses and Kansas State University for their help in Illumina sequencing and bioinformatic analyses. In addition, the authors thank Rebecca Hochstein and anonymous reviewer who provided constructive comments.

## ADDITIONAL INFORMATION AND DECLARATIONS

### Funding

This article is a part of ADELE (Autoimmune Defense and Living Environment)—strategic research opening funded by the Tekes—Finnish Funding Agency for Technology and Innovation (decision number 40333/14). This study was funded in part by Onni and Hilja Tuovinen foundation and OLVI-foundation. All the funding were received during this study. There was no additional external funding received for this study. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

### Grant Disclosures

The following grant information was disclosed by the authors:

Tekes—Finnish Funding Agency for Technology and Innovation: 40333/14.  
Tuovinen foundation.

Onni and Hilja Tuovinen foundation.

### Competing Interests

The authors declare there are no competing interests.

### Author Contributions

- Marja I. Roslund conceived and designed the experiments, performed the experiments, analyzed the data, contributed reagents/materials/analysis tools, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.
- Mira Grönroos and Aki Sinkkonen conceived and designed the experiments, analyzed the data, authored or reviewed drafts of the paper, approved the final draft.
- Anna-Lea Rantalainen conceived and designed the experiments, performed the experiments, contributed reagents/materials/analysis tools, authored or reviewed drafts of the paper, approved the final draft.
- Ari Jumpponen, Martin Romantschuk, Heikki Hyöty and Olli Laitinen authored or reviewed drafts of the paper, approved the final draft.
- Anirudra Parajuli prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.

### Data Availability

The following information was supplied regarding data availability:

Sequence Read Archive Accession [SRX3195289–SRX3195338](#) (Study [SRP117895](#)).

### Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.4508#supplemental-information>.

## REFERENCES

- Abrahamsson TR, Jakobsson HE, Andersson AF, Björkstén B, Engstrand L, Jenmalm MC. 2012. Low diversity of the gut microbiota in infants with atopic eczema. *Journal of Allergy Clinical Immunology* 129:434–440 DOI 10.1016/j.jaci.2011.10.025.
- Abrahamsson TR, Jakobsson HE, Andersson AF, Björkstén B, Engstrand L, Jenmalm MC. 2014. Low gut microbiota diversity in early infancy precedes asthma at school age. *Clinical & Experimental Allergy* 44:842–850 DOI 10.1111/cea.12253.
- Artmann M. 2014. Assessment of soil sealing management responses, strategies, and targets toward ecologically sustainable urban land use management. *AMBIO* 43:530–541 DOI 10.1007/s13280-014-0511-1.
- Berry D, Ben Mahfoudh K, Wagner M, Loy A. 2011. Barcoded primers used in multiplex amplicon pyrosequencing bias amplification. *Applied and Environmental Microbiology* 77:7846–7849 DOI 10.1128/AEM.05220-11.
- Boström C, Gerde P, Hanberg A, Jernström B, Johansson C, Kyrklund T, Rannug A, Törnqvist M, Victorin K, Westerholm R. 2002. Cancer risk assessment, indicators, and guidelines for polycyclic aromatic hydrocarbons in the ambient air. *Environmental Health Perspectives* 110:451–481 DOI 10.1289/ehp.02110s3451.
- Bradley PH, Pollard KS. 2017. Proteobacteria explain significant functional variability in the human gut microbiome. *Microbiome* 5(1):36 DOI 10.1186/s40168-017-0244-z.
- Brown CT, Davis-Richardson AG, Giongo A, Gano KA, Crabb DB, Mukherjee N, Casella G, Drew JC, Ilonen J, Knip M, Hyöty H, Veijola R, Simell T, Neu J, Wasserfall CH, Schatz D, Atkinson MA, Triplett EW. 2011. Gut microbiome metagenomics analysis suggests a functional model for the development of autoimmunity for type 1 diabetes. *PLOS ONE* 6(10):e25792 DOI 10.1371/journal.pone.0025792.
- Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Huntley J, Fierer N, Owens SM, Betley J, Fraser L, Bauer M, Gormley N, Gilbert JA, Smith G, Knight R. 2012. Ultra-high-throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms. *The ISME Journal* 6:1621–1624 DOI 10.1038/ismej.2012.8.
- Chaudhary P, Singh S, Chaudhry S, Nain L. 2012. Impact of PAH on biological health parameters of soils of an Indian refinery and adjoining agricultural area—a case study. *Environmental Monitoring and Assessment* 184:1145–1156 DOI 10.1007/s10661-011-2029-3.
- Cole JR, Wang Q, Cardenas E, Fish J, Chai B, Farris RJ, Kulam-Syed-Mohideen AS, McGarrell DM, Marsh T, Garrity GM, Tiedje JM. 2009. The Ribosomal Database Project: improved alignments and new tools for rRNA analysis. *Nucleic Acids Research* 37:141–145 DOI 10.1093/nar/gkn879.
- Crampon M, Bureau F, Akpa-Vinceslas M, Bodilis J, Machour N, Le Derf F, Portet-Koltalo F. 2014. Correlations between PAH bioavailability, degrading bacteria, and soil characteristics during PAH biodegradation in five diffusely contaminated dissimilar soils. *Environmental Science & Pollution Research* 21:8133–8145 DOI 10.1007/s11356-014-2799-6.

- Edgar RC, Haas BJ, Clemente JC, Quince C, Knight R. 2011.** UCHIME improves sensitivity and speed of chimera detection. *Bioinformatics* 27:2194–2200 DOI 10.1093/bioinformatics/btr381.
- Ege MJ, Mayer M, Schwaiger K, Mattes J, Pershagen G, Van Hage M, Scheynius A, Bauer J, Von Mutius E. 2012.** Environmental bacteria and childhood asthma. *Allergy* 67:1565–1571 DOI 10.1111/all.12028.
- Finnish Meteorological Institute. 2017.** Kesäsään tilastoja. Statistic from years 1981–2010 (in Finnish). Available at <http://ilmatieteenlaitos.fi/kesatilastot> (accessed on 13 June 2017).
- Folwell B, McGenity TJ, Whitby C. 2016.** Biofilm and planktonic bacterial and fungal communities transforming high-molecular-weight polycyclic aromatic hydrocarbons. *Applied and Environmental Microbiology* 82:2288–2299 DOI 10.1128/AEM.03713-15.
- De Goffau MC, Luopajarvi K, Knip M, Ilonen J, Ruohtula T, Härkönen T, Orivuori L, Hakala S, Welling GW, Harmsen HJ, Vaarala O. 2013.** Fecal microbiota composition differs between children with  $\beta$ -cell autoimmunity and those without. *Diabetes* 62:1238–1244 DOI 10.2337/db12-0526.
- Green SJ, Prakash O, Jasrotia P, Overholt WA, Cardenas E, Hubbard D, Tiedje JM, Watson DB, Schadt CW, Brooks SC, Kostka JE. 2012.** Denitrifying bacteria from the genus *Rhodanobacter* dominate bacterial communities in the highly contaminated subsurface of a nuclear legacy waste site. *Applied and Environmental Microbiology* 78:1039–1047 DOI 10.1128/AEM.06435-11.
- Hanski I, Von Hertzen L, Fyhrquist N, Koskinen K, Torppa K, Laatikainen T, Karisola P, Auvinen P, Paulin L, Mäkela MJ, Vartiainen E, Kosunen LU, Alenius H, Haahtela T. 2012.** Environmental biodiversity, human microbiota, and allergy are interrelated. *Proceedings of the National Academy of Sciences of the United States of America* 109:8334–8339 DOI 10.1073/pnas.1205624109.
- Haritash AK, Kaushik CP. 2009.** Biodegradation aspects of Polycyclic Aromatic Hydrocarbons (PAHs): a review. *Journal of Hazardous Materials* 169:1–15 DOI 10.1016/j.jhazmat.2009.03.137.
- Haugland T, Ottesen RT, Volden T. 2008.** Lead and polycyclic aromatic hydrocarbons (PAHs) in surface soil from day care centres in the city of Bergen, Norway. *Environmental Pollution* 153:266–272 DOI 10.1016/j.envpol.2007.08.028.
- Heederik D, Mutius E. 2012.** Does diversity of environmental microbial exposure matter for the occurrence of allergy and asthma? *The Journal of Allergy and Clinical Immunology* 130(1):44–50 DOI 10.1016/j.jaci.2012.01.067.
- Hennessee CT, Li GX. 2016.** Effects of polycyclic aromatic hydrocarbon mixtures on degradation, gene expression, and metabolite production in four *Mycobacterium* species. *Applied and Environmental Microbiology* 82:3357–3369 DOI 10.1128/AEM.00100-16.

- Heywood E, Wright J, Wienburg CL, Black HI, Long SM, Osborn D, Spurgeon DJ. 2006.** Factors influencing the national distribution of polycyclic aromatic hydrocarbons and polychlorinated biphenyls in British soils. *Environmental Science & Technology* **40**:7629–7635 DOI [10.1021/es061296x](https://doi.org/10.1021/es061296x).
- Hiller E, Lachká L, Jurkovič L, Vozár J. 2015.** Polycyclic aromatic hydrocarbons in urban soils from kindergartens and playgrounds in Bratislava, the capital city of Slovakia. *Environmental Earth Sciences* **73**:7147–7156 DOI [10.1007/s12665-014-3894-1](https://doi.org/10.1007/s12665-014-3894-1).
- Hong W, Jia H, Ma W-L, Sinha RK, Moon H-B, Nakata H, Minh NH, Chi KH, Li W-L, Kannan K, Sverko E, Li Y-F. 2016.** Distribution, fate, inhalation exposure and lung cancer risk of atmospheric polycyclic aromatic hydrocarbons in some Asian countries. *Environmental Science & Technology* **50**:7163–7174 DOI [10.1021/acs.est.6b01090](https://doi.org/10.1021/acs.est.6b01090).
- Honkonen O, Rantalainen A-L. 2013.** Impact of urbanization on the concentrations and distribution of organic contaminants in boreal lake sediments. *Environmental Monitoring and Assessment* **185**:1437–1449 DOI [10.1007/s10661-012-2643-8](https://doi.org/10.1007/s10661-012-2643-8).
- Hui N, Jumpponen A, Francini G, Kotze JD, Liu X, Romantschuk M, Strömmer R, Setälä H. 2017.** Soil microbial communities are shaped by vegetation type and park age in cities under cold climate. *Environmental Microbiology* **19**:1281–1295 DOI [10.1111/1462-2920.13660](https://doi.org/10.1111/1462-2920.13660).
- Huse SM, Welch DM, Morrison HG, Sogin ML. 2010.** Ironing out the wrinkle in the rare biosphere through improved OTU clustering. *Environmental Microbiology* **12**:1889–1898 DOI [10.1111/j.1462-2920.2010.02193.x](https://doi.org/10.1111/j.1462-2920.2010.02193.x).
- Ismail IH, Oppedisano F, Joseph SJ, Boyle RJ, Licciardi PV, Robins-Browne RM, Tang MLK. 2012.** Reduced gut microbial diversity in early life is associated with later development of eczema but not atopy in high-risk infants. *Pediatric Allergy and Immunology* **23**:674–681 DOI [10.1111/j.1399-3038.2012.01328.x](https://doi.org/10.1111/j.1399-3038.2012.01328.x).
- Johnsen AR, Wick LY, Harms H. 2005.** Principles of microbial PAH-degradation in soil. *Environmental Pollution* **133**:71–84 DOI [10.1016/j.envpol.2004.04.015](https://doi.org/10.1016/j.envpol.2004.04.015).
- Juhasz AL, Naidu R. 2000.** Bioremediation of high molecular weight polycyclic aromatic hydrocarbons: A review of the microbial degradation of benzo[a]pyrene. *International Biodeterioration & Biodegradation* **45**:57–88 DOI [10.1016/S0964-8305\(00\)00052-4](https://doi.org/10.1016/S0964-8305(00)00052-4).
- Kamal A, Cincinelli A, Martellini T, Malik R. 2015.** A review of PAH exposure from the combustion of biomass fuel and their less surveyed effect on the blood parameters. *Environmental Science and Pollution Research* **22**:4076–4098 DOI [10.1007/s11356-014-3748-0](https://doi.org/10.1007/s11356-014-3748-0).
- Kanaly A, Harayama S, Watanabe K. 2002.** *Rhodanobacter* sp. strain BPC1 in a benzo[a]pyrene-mineralizing bacterial consortium. *Applied and Environmental Microbiology* **68**:5826–5833 DOI [10.1128/AEM.68.12.5826-5833.2002](https://doi.org/10.1128/AEM.68.12.5826-5833.2002).
- Kim K-H, Jahan SA, Kabir E, Brown RJC. 2013.** A review of airborne polycyclic aromatic hydrocarbons (PAHs) and their human health effects. *Environment International* **60**:71–80 DOI [10.1016/j.envint.2013.07.019](https://doi.org/10.1016/j.envint.2013.07.019).

- Knip M, Siljander H. 2016.** The role of the intestinal microbiota in type 1 diabetes mellitus. *Nature Reviews* **12**:154–167 DOI [10.1038/nrendo.2015.218](https://doi.org/10.1038/nrendo.2015.218).
- Koivula TT, Salkinoja-Salonen M, Peltola R, Romantschuk M. 2004.** Pyrene degradation in forest humus microcosms with or without pine and its mycorrhizal fungus. *Journal of Environmental Quality* **33**:45–53 DOI [10.2134/jeq2004.4500](https://doi.org/10.2134/jeq2004.4500).
- Kozich JJ, Westcott SL, Baxter NT, Highlander SK, Schloss PD. 2013.** Development of a dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence data on the MiSeq Illumina sequencing platform. *Applied and Environmental Microbiology* **79**:5112–5120 DOI [10.1128/AEM.01043-13](https://doi.org/10.1128/AEM.01043-13).
- Lehtimäki J, Karkman A, Laatikainen T, Paalanen L, Von Herzen L, Haahtela T, Hanski I, Ruokolainen L. 2017.** Patterns in the skin microbiota differ in children and teenagers between rural and urban environments. *Scientific Reports* **7**:45651 DOI [10.1038/srep45651](https://doi.org/10.1038/srep45651).
- Li J, Zhang D, Song M, Jiang L, Wang Y, Luo C, Zhang G. 2017.** Novel bacteria capable of degrading phenanthrene in activated sludge revealed by stable-isotope probing coupled with high-throughput sequencing. *Biodegradation* **28**:423–436 DOI [10.1007/s10532-017-9806-9](https://doi.org/10.1007/s10532-017-9806-9).
- Lors C, Ryngaert A, Périé F, Diels L, Damidot D. 2010.** Evolution of bacterial community during bioremediation of PAHs in a coal tar contaminated soil. *Chemosphere* **81**:1263–1271 DOI [10.1016/j.chemosphere.2010.09.021](https://doi.org/10.1016/j.chemosphere.2010.09.021).
- Luo L, Lin S, Huang H, Zhang S. 2012.** Relationships between aging of PAHs and soil properties. *Environmental Pollution* **170**:177–182 DOI [10.1016/j.envpol.2012.07.003](https://doi.org/10.1016/j.envpol.2012.07.003).
- Manter DK, Weir TL, Vivanco JM. 2010.** Negative effects of sample pooling on PCR-based estimates of soil microbial richness and community structure. *Applied and Environmental Microbiology* **76**:2086–2090 DOI [10.1128/AEM.03017-09](https://doi.org/10.1128/AEM.03017-09).
- Martin F, Torelli S, Le Palier D, Barbance A, Martin-Laurent F, Bru D, Geremia R, Blake G, Jouanneau Y. 2012.** Betaproteobacteria dominance and diversity shifts in the bacterial community of a PAH-contaminated soil exposed to phenanthrene. *Environmental Pollution* **162**:345–353 DOI [10.1016/j.envpol.2011.11.032](https://doi.org/10.1016/j.envpol.2011.11.032).
- McLellan SL, Warshawsky D, Shann JR. 2002.** The effect of polycyclic aromatic hydrocarbons on the degradation of benzo[a]pyrene by *Mycobacterium* sp. strain RJGII-135. *Environmental Toxicology and Chemistry* **21**:253–259 DOI [10.1002/etc.5620210205](https://doi.org/10.1002/etc.5620210205).
- Morabito M, Crisci A, Messeri A, Orlandini S, Raschi A, Maracchi G, Munafó M. 2016.** The impact of built-up surfaces on land surface temperatures in Italian urban areas. *Science of the Total Environment* **551–552**:317–326 DOI [10.1016/j.scitotenv.2016.02.029](https://doi.org/10.1016/j.scitotenv.2016.02.029).
- Morel J, Chenu C, Lorenz K. 2015.** Ecosystem services provided by soils of urban, industrial, traffic, mining, and military areas (SUITMAS). *Journal of Soils and Sediments* **15**:1659–1666 DOI [10.1007/s11368-014-0926-0](https://doi.org/10.1007/s11368-014-0926-0).
- Mukherjee S, Juottonen H, Siivonen P, Quesada CL, Tuomi P, Pulkkinen P, Yrjälä K. 2014.** Spatial patterns of microbial diversity and activity in an aged creosote-contaminated site. *The ISME Journal* **8**:2131–2142 DOI [10.1038/ismej.2014.151](https://doi.org/10.1038/ismej.2014.151).



- Oleszczuk P, Baran S. 2003. Degradation of individual polycyclic aromatic hydrocarbons (PAHs) in soil polluted with aircraft fuel. *Polish Journal of Environmental Studies* 12:431–437.
- Pakarinen J, Hyvärinen A, Salkinoja-Salonen M, Laitinen S, Nevalainen A, Mäkelä MJ, Haahtela T, Von Hertzen L. 2008. Predominance of Gram-positive bacteria in house dust in the low-allergy risk Russian Karelia. *Environmental Microbiology* 10:3317–3325 DOI 10.1111/j.1462-2920.2008.01723.x.
- Parajuli A, Grönroos M, Kauppi S, Płociniczak T, Roslund MI, Galitskaya P, Laitinen O, Hyöty H, Jumpponen A, Strömmer R, Romantschuk M, Hui N, Sinkkonen A. 2017. The abundance of health-associated bacteria is altered in PAH polluted soils—implications for health in urban areas? *PLOS ONE* 12(11):e0187852 DOI 10.1371/journal.pone.0187852.
- Parajuli A, Grönroos M, Siter N, Puhakka R, Vari HK, Roslund MI, Jumpponen A, Nurminen N, Laitinen OH, Hyöty H, Rajaniemi J, Sinkkonen A. 2018. Urbanization reduces transfer of diverse environmental microbiota indoors. *Frontiers in Microbiology* 9:84 DOI 10.3389/fmicb.2018.00084.
- Peng J, Zhang Y, Su J, Qiu Q, Jia Z, Zhu Y-G. 2013. Bacterial communities predominant in the degradation of 13 C 4-4, 5, 9, 10-pyrene during composting. *Bioresource Technology* 143:608–614 DOI 10.1016/j.biortech.2013.06.039.
- Priha O, Smolander A. 1999. Nitrogen transformations in soil under *Pinus sylvestris*, *Picea abies* and *Betula pendula* at two forest sites. *Soil Biology and Biochemistry* 31:965–977 DOI 10.1016/S0038-0717(99)00006-1.
- Pruesse E, Quast C, Knittel K, Fuchs BM, Ludwig W, Peplies J, Glöckner FO. 2007. SILVA a comprehensive online resource for quality checked and aligned ribosomal RNA sequence data compatible with ARB. *Nucleic Acids Research* 35:7188–7196 DOI 10.1093/nar/gkm864.
- Qian L, Xie J, Zhou D, Hu Y, Zhu L, Zhao R, Kang S, Jiang L. 2017. Impact of rural and urban environmental microbial exposure on intestinal microbiota in early infancy. *Nanoscience and Nanotechnology Letters* 9:761–769 DOI 10.1166/nnl.2017.2283.
- Ran Y, Sun K, Ma X, Wang G, Grathwohl P, Zeng EY. 2007. Effect of condensed organic matter on solvent extraction and aqueous leaching of polycyclic aromatic hydrocarbons in soils and sediments. *Environmental Pollution* 148:529–538 DOI 10.1016/j.envpol.2006.11.028.
- Ravindra K, Sokhi R, Van Grieken R. 2008. Atmospheric polycyclic aromatic hydrocarbons: source attribution, emission factors and regulation. *Atmospheric Environment* 42:2895–2921 DOI 10.1016/j.atmosenv.2007.12.010.
- R Core Team. 2017. R: a language and environment for statistical computing. Vienna: R Foundation for Statistical Computing. Available at <http://www.r-project.org>.
- Ren G, Ren W, Teng Y, Li Z. 2015. Evident bacterial community changes but only slight degradation when polluted with pyrene in a red soil. *Frontiers in Microbiology* 6:22 DOI 10.3389/fmicb.2015.00022.
- Ruokolainen L, Hertzen L, Fyhrquist N, Laatikainen T, Lehtomäki J, Auvinen P, Karvonen AM, Hyvärinen A, Tillmann V, Niemelä O, Knip M, Haahtela T,

- Pekkanen J, Hanski I. 2015.** Green areas around homes reduce atopic sensitization in children. *Allergy* **70**:195–202 DOI [10.1111/all.12545](https://doi.org/10.1111/all.12545).
- Saarnio K, Sillanpää M, Hillamo R, Sandell E, Pennanen AS, Salonen RO. 2008.** Polycyclic aromatic hydrocarbons in size-segregated particulate matter from six urban sites in Europe. *Atmospheric Environment* **42**:9087–9097 DOI [10.1016/j.atmosenv.2008.09.022](https://doi.org/10.1016/j.atmosenv.2008.09.022).
- Salter S, Cox MJ, Turek EM, Calus ST, Cookson WO, Moffatt MF, Turner P, Parkhill J, Loman N, Walker AW. 2014.** Reagent contamination can critically impact sequence-based microbiome analyses. *BMC Biology* **12**:87–98 DOI [10.1186/s12915-014-0087-z](https://doi.org/10.1186/s12915-014-0087-z).
- Santorù M, Piras C, Murgia A, Palmas V, Camboni T, Liggi S, Ibba I, Lai M, Orrù S, Loizedda A, Griffin J, Usai P, Caboni P, Atzori L, Manzin A. 2017.** Cross sectional evaluation of the gut-microbiome metabolome axis in an Italian cohort of IBD patients. *Scientific Reports* **7**:9523 DOI [10.1038/s41598-017-10034-5](https://doi.org/10.1038/s41598-017-10034-5).
- Sawulski P, Clipson N, Doyle E. 2014.** Effects of polycyclic aromatic hydrocarbons on microbial community structure and PAH ring hydroxylating dioxygenase gene abundance in soil. *Biodegradation* **25**:835–847 DOI [10.1007/s10532-014-9703-4](https://doi.org/10.1007/s10532-014-9703-4).
- Schloss PD, Gevers D, Westcott SL. 2011.** Reducing the effects of PCR amplification and sequencing Artifacts on 16 srRNA-based studies. *PLOS ONE* **6**:e27310 DOI [10.1371/journal.pone.0027310](https://doi.org/10.1371/journal.pone.0027310).
- Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, Lesniewski RA, Oakley BB, Parks DH, Robinson CJ, Sahl JW, Stres B, Thallinger GG, Van Horn DJ, Weber CF. 2009.** Introducing mothur: open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Applied and Environmental Microbiology* **75**:7537–7541 DOI [10.1128/AEM.01541-09](https://doi.org/10.1128/AEM.01541-09).
- SFS 3008. 1990.** *Determination of dry matter and loss on ignition for water, sludge and sediment*. Helsinki: Finnish Standards Association (SFS) (In Finnish).
- SFS-ISO 10390. 2005.** *Soil quality—determination of pH*. Helsinki: Finnish Standards Association (SFS) (In Finnish).
- Singleton DR, Sangaiah R, Gold A, Ball LM, Aitken MD. 2006.** Identification and quantification of uncultivated Proteobacteria associated with pyrene degradation in a bioreactor treating PAH-contaminated soil. *Environmental Microbiology* **8**:1736–1745 DOI [10.1111/j.1462-2920.2006.01112.x](https://doi.org/10.1111/j.1462-2920.2006.01112.x).
- Sinkkonen A, Kauppi S, Pukkila V, Nan H, Plociniczak T, Kontro M, Strömmer R, Romantschuk M. 2013.** Previous exposure advances the degradation of an anthropogenic s-triazine regardless of soil origin. *Journal of Soils and Sediments* **13**:1430–1438 DOI [10.1007/s11368-013-0742-y](https://doi.org/10.1007/s11368-013-0742-y).
- Song M, Jiang L, Zhang D, Luo C, Wang Y, Yu Z, Yin H, Zhang G. 2016.** Bacteria capable of degrading anthracene, phenanthrene, and fluoranthene as revealed by DNA based stable-isotope probing in a forest soil. *Journal of Hazardous Materials* **308**:50–57 DOI [10.1016/j.jhazmat.2016.01.009](https://doi.org/10.1016/j.jhazmat.2016.01.009).

- Tansel B, Fuentes C, Sanchez M, Predoi K, Acevedo M. 2011.** Persistence profile of polyaromatic hydrocarbons in shallow and deep Gulf waters and sediments: effect of water temperature and sediment–water partitioning characteristics. *Marine Pollution Bulletin* **62**:2659–2665 DOI [10.1016/j.marpolbul.2011.09.026](https://doi.org/10.1016/j.marpolbul.2011.09.026).
- Tarvainen T, Hatakka T, Salla A, Jarva J, Pitkäranta P, Anttila H, Maidell-Münster L. 2013.** Geochemical baselines in the Helsinki metropolitan area. *Geological Survey of Finland, Report of Investigation* **201**:1–99.
- Teng Y, Luo Y, Ping L, Zou D, Christie P. 2010.** Effects of soil amendment with different carbon sources and other factors on the bioremediation of an aged PAH-contaminated soil. *Biodegradation* **21**:167–178 DOI [10.1007/s10532-009-9291-x](https://doi.org/10.1007/s10532-009-9291-x).
- United Nations. 2014.** World Urbanization Prospects: The 2014 Revision of World Urbanization Prospects. Available at <https://esa.un.org/Unpd/Wup/>.
- Veach A, Dodds W, Jumpponen A. 2015.** Woody plant encroachment, and its removal, impact bacterial and fungal communities across stream and terrestrial habitats in a tallgrass prairie ecosystem. *FEMS Microbiology Ecology* **91**(10):fiv109 DOI [10.1093/femsec/fiv109](https://doi.org/10.1093/femsec/fiv109).
- Wang Q, Garrity GM, Tiedje JM, Cole JR. 2007.** Naive Bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. *Applied and Environmental Microbiology* **73**:5261–5267 DOI [10.1128/AEM.00062-07](https://doi.org/10.1128/AEM.00062-07).
- World Health Organization (WHO). 2015.** Connecting global priorities: biodiversity and human health. Geneva, World Health Organization and Secretariat of the Convention on Biological Diversity. Available at <http://www.who.int/globalchange/publications/biodiversity-human-health/en/>.
- Yang Y, Zhang N, Xue M, Tao S. 2010.** Impact of soil organic matter on the distribution of polycyclic aromatic hydrocarbons (PAHs) in soils. *Environmental Pollution* **158**:2170–2174 DOI [10.1016/j.envpol.2010.02.019](https://doi.org/10.1016/j.envpol.2010.02.019).
- Zahed MA, Aziz HA, Isa MH, Mohajeri L, Mohajeri S, Kutty SRM. 2011.** Kinetic modeling and half life study on bioremediation of crude oil dispersed by Corexit 9500. *Journal of Hazardous Materials* **185**:1027–1031 DOI [10.1016/j.jhazmat.2010.10.009](https://doi.org/10.1016/j.jhazmat.2010.10.009).
- Zhou Y, Zhi F. 2016.** Lower level of bacteroides in the gut microbiota is associated with inflammatory bowel disease: a meta-analysis. *BioMed Research International* **2016**:Article 5828959 DOI [10.1155/2016/5828959](https://doi.org/10.1155/2016/5828959).