

Accepted Manuscript

Title: Circulating pathogen-specific plasmablasts in female patients with upper genital tract infection

Authors: Nina V. Palkola, Sari H. Pakkanen, Oskari Heikinheimo, Jussi M. Kantele, Anu Kantele



PII: S0165-0378(17)30348-0
DOI: <https://doi.org/10.1016/j.jri.2018.02.005>
Reference: JRI 2534

To appear in: *Journal of Reproductive Immunology*

Received date: 13-10-2017
Revised date: 12-1-2018
Accepted date: 21-2-2018

Please cite this article as: Palkola, Nina V., Pakkanen, Sari H., Heikinheimo, Oskari, Kantele, Jussi M., Kantele, Anu, Circulating pathogen-specific plasmablasts in female patients with upper genital tract infection. *Journal of Reproductive Immunology* <https://doi.org/10.1016/j.jri.2018.02.005>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

CIRCULATING PATHOGEN-SPECIFIC PLASMABLASTS IN FEMALE PATIENTS WITH UPPER GENITAL TRACT INFECTION

Nina V. Palkola^{1,2}, Sari H. Pakkanen¹, Oskari Heikinheimo³, Jussi M. Kantele⁴, Anu Kantele^{2,5,6}

¹ Department of Bacteriology and Immunology, University of Helsinki, Helsinki, Finland

² Department of Medicine, University of Helsinki, Helsinki, Finland

³ Department of Obstetrics and Gynaecology, University of Helsinki and Helsinki University Hospital, Helsinki, Finland

⁴ Occupational Health and Environmental Medicine, Department of Public Health, University of Turku, Turku, Finland

⁵ Inflammation Center, Clinic of Infectious Diseases, University of Helsinki and Helsinki University Hospital, Helsinki, Finland

⁶ Unit of Infectious Diseases, Department of Medicine / Solna, Karolinska Institutet, Stockholm, Sweden

Words: Abstract 230, text 3000

Address correspondence to Anu Kantele, Professor, Inflammation Center, Clinic of Infectious Diseases, University of Helsinki and Helsinki University Hospital, POB 348, FIN-00029 HUS, Finland; FAX +358-9-471 75900; tel. +358-50-309 7640; email anu.kantele@hus.fi

List of abbreviations used in this paper:

FGT – female genital tract

ASC – pathogen-specific antibody-secreting cell

CLA – cutaneous lymphocyte antigen

HR – homing receptor

CCR – chemokine receptor

PLN – peripheral lymph node

ISC – immunoglobulin-secreting cell

HIGHLIGHTS

(Circulating pathogen-specific plasmablasts in female patients with upper genital tract infection)

- Pathogen-specific plasmablasts appear in the human circulation in upper FGT infection
- The FGT-related ASC response is dominated by IgA rather than IgG ASC.
- The expression of $\alpha_4\beta_7$, L-selectin, and CLA on ASC is similar in FGT and urinary tract infections, suggesting a unique homing receptor profile characteristic of the urogenital tract.
- In line with nasal-genital crosstalk, the HR profile resembles that after i.n. immunization

ABSTRACT

Mucosal antibodies constitute the first line of adaptive immune defence against invaders in the female genital tract (FGT), yet the sequence of events leading to their production is surprisingly poorly characterized. We explored the induction of pathogen-specific antibody-secreting cells (ASC) as a response to an acute infection in the upper FGT.

We recruited 12 patients undergoing surgery due to an upper FGT infection (7/12 blood culture positive, 5/12 negative) and six healthy controls. Pathogens were sampled during surgery and PBMC collected in the acute phase of the disease (day 7-10). We searched by ELISPOT circulating pathogen-specific ASC and explored their frequency, immunoglobulin isotype distribution, and expressions of homing receptors ($\alpha_4\beta_7$, L-selectin, and CLA).

All patients had circulating ASC specific to the infective bacteria; the geometric mean was 434 (95%CI 155-1234) ASC (IgA + IgG + IgM) / 10^6 PBMC. IgA ASC predominated in 7/12, IgG ASC in 3/12, and IgM ASC in 2/12 cases. Of all the pathogen-specific ASC, 60% expressed $\alpha_4\beta_7$, 67% L-selectin, and 9% CLA.

This study is the first to show induction of pathogen-specific ASC in the peripheral blood in bacterial infection in the human FGT. Our findings reveal that such FGT-originating pathogen-specific ASC are predominated by IgA ASC and exhibit a homing receptor profile resembling that of ASC in acute urinary tract infection. The data thus suggest a characteristic profile shared by the urogenital tract.

Keywords: Lymphocyte homing, adhesion molecule, homing receptor, pathogen-specific antibody-secreting cell, plasmablast, upper female genital tract, ELISPOT

ACCEPTED MANUSCRIPT

1. INTRODUCTION

The female genital tract (FGT) is uniquely challenged to provide protection against pathogens without compromising reproduction. The FGT comprises upper (Fallopian tubes, uterus, endocervix) and lower (ectocervix, vagina) parts, each harbouring a characteristic immune defence influenced by hormonal changes (Wira et al. 2015). Mucosal antibodies act as key players providing the first specific line of defence against a variety of pathogens (Woof and Mestecky, 2005). Nonetheless, the induction of specific antibody response in the human FGT is insufficiently characterized as yet.

An antigen encountered at mucosal sites activates its cognate naïve B lymphocytes which enter the lymphatics and return, via blood circulation, to mucosal sites (Brandtzaeg and Johansen, 2005, Sigmundsdottir and Butcher, 2008). Consistent with this migration, effector B cells, plasmablasts, can be found in the human circulation after mucosal antigen encounter: circulating antigen-specific antibody-secreting cells (ASC) have been reported after oral (Czercinsky et al. 1987; Kantele, 1990; Kantele et al. 1986), rectal (Kantele et al. 1998; Kutteh et al. 2001), or intranasal vaccinations (Quiding-Järbrink et al. 1997), and in various mucosal infections such as gastroenteritis (Kantele et al. 1988, 2008; Kantele JM et al. 1996a; Pakkanen et al. 2010), pyelonephritis and cystitis (Kantele et al. 1994, 2008), and tonsillitis, sinusitis, and pneumonia (Palkola et al. 2012, 2016). Thus far, circulating plasmablasts have not been explored in acute bacterial infection in the genital tract.

Homing of lymphocytes from circulation into tissues is a selective process where tissue-specific homing receptors (HR) and chemokine receptors (CCR) on the circulating cells recognize their respective ligands in the target tissue (Brandtzaeg and Johansen, 2005; Sigmundsdottir and Butcher, 2008). Tissue-specific HR have been identified: $\alpha_4\beta_7$ guides the cells to the intestine (Berlin et al. 1993), L-selectin to peripheral lymph nodes

(PLN) (Camerini et al. 1989), and cutaneous lymphocyte antigen (CLA) to cutaneous sites (Berg et al. 1991). Lymphocytes typically home back to sites where the antigen was initially encountered (Sigmundsdottir and Butcher, 2008). Accordingly, after intestinal antigen encounter, a high proportion of plasmablasts express $\alpha_4\beta_7$ and lower proportions L-selectin (Kantele et al. 1997; Kantele JM et al. 1996a; Quiding-Järbrink et al. 1997) or CLA (Kantele et al. 2003), while antigen encounter at other mucosal sites elicits a different homing profile (Kantele et al. 2008; Palkola et al. 2015, 2016; Quiding-Järbrink et al. 1997). Likewise, in mice, the set of receptors operating in the homing process differs between the various mucosal sites (e.g. nasal vs. intestinal) (Csencsits et al. 1999). There is, however, a lack of data on HR expressions associated with antigen encounter in the FGT.

We sought to identify circulating FGT-originating plasmablasts in patients with acute bacterial infection in the upper FGT. We explored ASC specific to bacteria isolated from the site of infection and characterized the response in terms of magnitude, isotype distribution, and HR expressions.

2. METHODS

2.1. Study design

We investigated circulating plasmablasts (pathogen-specific ASC and all immunoglobulin-secreting cells, ISC) and their homing potentials in patients with bacterial upper FGT infection and healthy controls (Fig. 1).

PBMC were analyzed for ASC and ISC by ELISPOT. Immunomagnetic cell sorting was combined with ELISPOT to determine their expression of HR. Samples for bacterial cultures were obtained during surgery and the bacteria applied as an antigen to detect pathogen-specific ASC.

The Ethics Committee of Helsinki University Hospital approved the study protocol (411/E5/02). Written informed consent was obtained from volunteers.

2.2. Study subjects

A total of 12 patients (aged 18 to 51) with acute bacterial upper FGT infection were recruited at Helsinki University Hospital (Table 1). Blood culture proved positive for 7/12 (invasive) and negative for 5/12 (non-invasive) patients. Six healthy volunteers (35 to 47 years) served as controls.

2.3. Sorting of PBMC by homing receptor expression

PBMC were sorted into HR⁺ and HR⁻ cell populations by immunomagnetic cell sorting as described earlier (Kantele et al. 1997; Kantele JM et al. 1996b). In brief, mAbs specific to L-selectin (Leu-8), $\alpha_4\beta_7$ (ACT-1), or CLA (HECA-452) were incubated with PBMC. After washings, magnetic Dynal M-450 beads (coated with sheep anti-mouse IgG; Oslo) were

added to separate HR⁺ from HR⁻ cells. The efficiency of the separation has been reported previously (Kantele et al. 1997).

2.4. Enumeration of ASC and ISC

Unsorted PBMC and subpopulations HR⁺ and HR⁻ were analyzed for pathogen-specific ASC and ISC by ELISPOT, as described earlier (Kantele, 1990). Briefly, for ASC, each patients' own bacterial isolate or panel of these isolates (healthy controls) was used to coat microtiter plate wells. The isolates were applied as suspensions of formalin-killed whole bacteria (8×10^6 bacteria / mL PBS; 50 μ L / well for three hours at 37°C or overnight at 20°C). For controls, coating suspensions of six patients were tested, each in their individual wells. The concentration of the coating antigen was thus the same for patients and controls. For ISC, human IgA, IgM, (Dako, Glostrup, Denmark), or IgG (Sigma, Immuno Chemicals, St. Louis, MO) -specific antisera were used for coating. After washings and blocking, aliquots of unsorted PBMC or HR⁺ and HR⁻ subpopulations were allowed to secrete antibodies in the wells. Next, alkaline phosphatase-conjugated anti-human IgA, IgG (Sigma) or IgM (Southern Biotech, Birmingham, AL) were added, followed by the substrate (5-bromo-4-chloro-3-indolyl phosphate, Sigma) in melted agarose; spots were enumerated under a light microscope.

2.5. Statistical analysis

The numbers of pathogen-specific ASC and all ISC were given as geometric means of ASC/ISC (IgA+IgG+IgM) / 10^6 PBMC with 95% confidence intervals (95%CI) as counted using bootstrapping in four groups: all patients, blood-culture positive and negative patients, and healthy controls.

HR expressions were determined as percentages of ASC:

$\% \text{HR}^+\text{ASC} = 100 \times (\text{number of ASC in HR}^+ \text{ population}) / (\text{total of ASC in HR}^+ \text{ and HR}^- \text{ populations})$

or ISC:

$\% \text{HR}^+\text{ISC} = 100 \times (\text{number of ISC in HR}^+ \text{ population}) / (\text{total of ISC in HR}^+ \text{ and HR}^- \text{ populations})$.

The proportion of ASC or ISC expressing the various HR were given as arithmetic means with SD. To obtain reliable statistics in the HR analyses, we only included those with ≥ 20 identified spots.

Independent-samples Mann-Whitney U test and related-samples Wilcoxon Signed Rank test were applied for comparisons (SPSS 24.01; SPSS Inc). $P < 0.05$ was considered significant.

3. RESULTS

3.1. Number of ASC and ISC

Pathogen-specific ASC (IgA+IgG+IgM) were found in all patients (12/12) (Table 1); the geometric mean was $434/10^6$ PBMC (95%CI 155–1234). None of the responses were <10 ASC, four were in the range 10–100 and 100–1000, three in 1000–10000, and one in 10000–100000 ASC / 10^6 PBMC (Fig. 2). The figures appeared not to correlate with blood culture findings: those with the lowest responses (20 and 28 ASC / 10^6 PBMC) and the one with the highest (11575 ASC) had positive blood cultures; the low-responders were both treated at an intensive care unit. The geometric mean of ASC (IgA+IgG+IgM) / 10^6 PBMC in blood culture positive (7/12) patients was 257 (95%CI 26–4448) and negative (5/12) 903 (68–9947) ($p=0.343$). The healthy volunteers had no ASC response specific to the pool of pathogens.

The geometric mean for all ISC (IgA+IgG+IgM) was higher in the patients than in healthy controls ($p<0.0001$) (Fig. 2). The frequency of ISC was similar in invasive and non-invasive cases ($p=0.876$) (Fig. 2).

3.2. Isotype distributions of ASC and ISC

The geometric mean for IgA ASC was 140 (95%CI 28–866) / 10^6 PBMC, for IgG ASC 103 (22–405), and for IgM ASC 50 (13–181) (Fig. 3A). IgA ASC were detected in all patients; one patient had no IgG and another no IgM ASC. IgA ASC predominated in 7/10 and IgG in 3/10 patients with the highest responses and IgM in the two low-responders (Fig. 3A). Circulating ISC were seen in all three isotypes for all subjects, implying that none of the volunteers had agammaglobulinemia (Fig. 3B and C). Figure 3 presents the geometric

means of IgA, IgG, and IgM ISC among patients (B) and healthy controls (C). IgG ISC predominated for 11/12 patients and all controls, and IgA ISC for one patient (Fig. 3B and C).

3.3. Expression of $\alpha_4\beta_7$, L-selectin, and CLA on pathogen-specific ASC and all ISC

When the data of all patients (invasive and non-invasive cases) were pooled, the proportion of ASC (IgA+IgG+IgM) \pm SD expressing $\alpha_4\beta_7$ was 60% \pm 27, L-selectin 67% \pm 22, and CLA 9.0% \pm 6.7 (Fig. 4). HR-expressions did not differ by isotype of ASC (data not shown).

Figure 5 shows HR expressions for invasive vs. non-invasive infections. L-selectin-expressing ASC appeared more frequent in invasive (77% \pm 14) than non-invasive (55% \pm 26) infections, but the difference was non-significant ($p=0.177$). As for $\alpha_4\beta_7$ and CLA, the proportions were similar in the two groups ($p=1.00$ for both) (Fig. 5).

HR expressions of patients' ISC resembled those of the same patients' ASC and those of healthy controls' ISC (Fig. 4). No difference was seen in HR expressions between the ISC of those with invasive and non-invasive infections; 57% \pm 25 vs. 60% \pm 8 for $\alpha_4\beta_7$ ($p=1.00$), 68% \pm 15 vs. 55% \pm 11 for L-selectin ($p=0.202$), and 18% \pm 9 and 10% \pm 3 for CLA ($p=0.149$).

4. DISCUSSION

4.1. Introduction

This is the first study to show that pathogen-specific plasmablasts emerge in the human circulation in response to acute bacterial upper FGT infection, a result according with previous findings for infections at other mucosal sites (Kantele et al. 1988, 1994; Palkola et al. 2012, 2016). Detecting such cells also in upper FGT infections encourages their use as tools for evaluating immune responses in the FGT.

4.2. Timing of sampling

As plasmablasts only circulate transiently before homing into their target tissues, sampling timing is critical. We utilized our previous experience of acute mucosal infections in the urinary (Kantele et al. 2008) and respiratory tracts (Palkola et al. 2012, 2016), and the gut (Kantele et al. 1988), and the kinetics of plasmablasts in the circulation as reported in gastroenteritis (Kantele, 2012) and after oral (Kantele, 1990, 1991; Kantele et al. 1986) and rectal vaccinations (Kantele et al. 1998). Antigen-specific plasmablasts enter the blood approximately on the third day after antigen encounter, peak in frequency on day seven, and disappear gradually by days 14–16. Prolonged antigenic stimulus may lengthen the detection period (Kantele, 1991, 2012). Accordingly, to catch the peak of the response, we only recruited patients with a history of symptoms lasting 7–10 days. The timing proved successful: pathogen-specific ASC were found in all patients.

4.3. Numbers of circulating plasmablasts (ASC and ISC)

The vigorous ASC response reveals the remarkable inductive capacity of the upper FGT, consistent with previous reports (Wira, 2015; Iwasaki, 2010). Indeed, the magnitude of the response exceeded that seen in infections of the upper (Palkola et al. 2016) and lower (Palkola et al. 2012) respiratory tracts, yet resembled that in infections at anatomically close sites i.e. gastroenteritis (Kantele et al. 1988) or upper urinary tract infection (UTI) (Kantele et al. 2008).

As for ISC, representing the total of all ASC subpopulations at the time of sampling, the numbers proved exceptionally high: they exceeded by far those of ISC in healthy volunteers, even with the increase caused by pathogen-specific ASC taken into account. The high ISC frequency accords with previous reports on gastroenteritis (Kantele et al. 1988), respiratory tract infections (Palkola et al. 2012, 2016), and upper but not lower UTI (Kantele et al. 1994). As explanation, we have suggested polyclonal stimulation associated with severe natural infections (Kantele et al. 1988; Palkola et al. 2012). According with our findings, human memory B lymphocytes have been shown to differentiate into plasma cells in response to polyclonal stimuli, such as microbial products or bystander T cell help. Specific antibody-secreting cells have been found in such settings even in the absence of their cognate antigen (Bernasconi et al. 2002).

4.4. Immunoglobulin isotype distribution of pathogen-specific plasmablast response

While IgA predominated in the pathogen-specific plasmablast responses, also IgG ASC were abundant. Indeed, pathogen-specific IgA and IgG ASC have been shown to accumulate in the genital tract as a response to mucosal immunization or infection in animals (Bekri et al. 2017; Cuburu et al. 2009; Johansson et al. 1998) and in salpingitis in

humans (Kutteh et al. 1990). Both isotypes can contribute to protection against sexually-transmitted pathogens (Bomsel et al. 2011; Li et al. 2011; Mascola 2000; Shin and Iwasaki 2013). The high IgA ASC numbers accord not only with the previously reported upstream increase of IgA levels in FGT secretions (Kozlowski 2002; Quesnel et al. 1997), and predominance of IgA ASC in mucosae of the upper FGT (Kutteh et al. 1990), but also the active IgA transport (pIgR) operating on the epithelium of upper FGT (Iwasaki, 2010). Moreover, IgA is the general mucosal isotype, predominating among the circulating pathogen-specific ASC in gastroenteritis (Kantele et al. 1988; Kantele JM et al. 1996a; Pakkanen et al. 2010) and upper UTI (Kantele et al. 2008), and after oral vaccination (Kantele 1990; Kantele et al. 1986, Pakkanen et al 2010). Like the high IgG ASC numbers in this study, high IgG antibody levels have been associated with FGT, and the mechanisms transporting IgG (nFcR) exist throughout the FGT (Li et al. 2011). Indeed, in human cervico-vaginal secretions, IgG levels even exceed those of IgA (Li et al 2011; Kozlowski et al. 2002).

4.5. Homing profiles of plasmablasts

Homing molecules directing lymphocytes into the human FGT have been poorly described. Studies with rodents report recruitment of IgA cells into the uterus by the CCR10-CCL28 interaction (Cha et al. 2011; Cuburu et al. 2009). As for T lymphocytes, a unique memory-T-cell population co-expressing $\alpha_4\beta_7$ and CLA has been detected in the endocervix of patients with *Chlamydia* infection (Kelly et al. 2009), and ligands such as E-selectin (ligand to CLA), ICAM-1, and VCAM-1 have been proved inducible on human endometrium (Tabibzadeh et al. 1994). Moreover, $\alpha_4\beta_1$ -integrin-expressing T cells have been found in infected murine FGT (Davila et al. 2014). For the present study of circulating plasmablasts in human FGT infection, we chose to investigate $\alpha_4\beta_7$, L-selectin, and CLA, since these

markers have been extensively explored in infections of other mucosal sites (Kantele et al. 2008, Kantele JM et al. 1996a; Palkola et al. 2015, 2016; Pakkanen et al. 2010).

4.6. General characteristics of HR profile

$\alpha_4\beta_7$ and L-selectin were both expressed on two thirds of the circulating pathogen-specific plasmablasts and CLA on only a small proportion. Such a HR profile implies that $\alpha_4\beta_7$ and L-selectin contribute, but CLA less, in the dissemination of B cells from the upper FGT. Although we could not explore plasmablasts in the FGT tissues, this HR profile may, at least partly, represent that required for homing to the FGT, as migrating effector lymphocytes favour the initial activation site (Sigmundsdottir and Butcher, 2008). It should be highlighted that FGT lacks organized structures equivalent to intestinal Payer's patches. Instead, draining lymph nodes (the common iliac, internal iliac, external iliac, and the inguinal femoral lymph nodes), may serve directly as activation sites (Iwasaki, 2010). The anatomically close rectal and urogenital mucosal sites at least partly share the same draining lymph nodes. It is thus of interest to compare the HR profile in FGT infection to that of infections in various human anatomical compartments, especially those of the intestinal and urinary tract.

4.6.1. Comparison with intestinal HR profile

The HR profile of circulating ASC in the upper FGT infection differed from the gut-seeking profile with respect to $\alpha_4\beta_7$: although substantial, the proportion of $\alpha_4\beta_7$ -expressing cells (60%) did not reach that of gut-seeking lymphocytes (90–100%) reported after oral immunization (Kantele et al. 1997; Quiding-Järbrink 1997) and in acute gastroenteritis (Kantele JM et al. 1996a; Palkola et al. 2008). The main ligand for $\alpha_4\beta_7$, mucosal cell adhesion molecule (MadCAM-1) is highly abundant in the gut lamina propria (Briskin et al.

1997), whereas in FGT, MadCAM-1 expression has only been reported in tissue culture models of chlamydia infection (Kelly et al. 2001), not in non-infection settings (Johansson et al. 1999). L-selectin, on the other hand, may figure more importantly in infections of the FGT (67%) than the intestine (40%; Kantele et al. 2008; Kantele JM et al. 1996a). L-selectin-expressing plasmablasts dominate a systemic profile (proportion of L-selectin high, $\alpha_4\beta_7$ moderate and CLA low) induced in PLN after parenteral immunization (Kantele et al. 1997; 2008; Quiding-Järbrink et al. 1997). The substantial proportion of L-selectin-expressing cells in FGT infections may reflect both the systemic part of the immune protection and a contribution of L-selectin in targeting effector B cells into the FGT. We are unaware of any studies exploring whether the ligand of L-selectin, peripheral lymph node addressin (PNAd), is abundant within the FGT.

4.6.2. Comparison with HR profile in respiratory tract

It is interesting to compare the HR profiles of the FGT and the respiratory tract, for a cross-talk has been reported between nasal and genital surfaces (Johansen et al. 2005; Kozlowski et al. 2002). Antigen-specific ASC and cell-mediated immune responses have been reported in murine/primate FGT after intranasal (Cha et al. 2011; Johansson 1998) and sublingual (Cuburu et al. 2009) immunizations. Accordingly, in line with our data, a high to moderate proportion of both L-selectin- and $\alpha_4\beta_7$ -expressing ASC has been shown after intranasal cholera vaccination in humans (Quiding-Järbrink et al. 1997). In contrast, in respiratory tract infections (sinusitis, tonsillitis, pneumonia) the proportion of L-selectin-expressing cells appears higher (79–82%) and that of $\alpha_4\beta_7$ -expressing cells lower (15–44%) than in upper FGT infections (67% and 60%, respectively) (Palkola et al. 2015, 2016). The slight differences between the HR profiles reported for tonsillitis and sinusitis and after intranasal immunization may reflect differences in response induction, with

intranasal immunization only stimulating a restricted area in the Waldeyer's ring (resulting in high $\alpha_4\beta_7$ proportions) and natural infections impacting more scattered induction sites.

4.6.3. Comparison with HR profile for UTI

A comparison with the urinary tract is of interest, as it may share draining lymph nodes with FGT. Indeed, the homing profile of pathogen-specific ASC in acute upper UTI ($\alpha_4\beta_7$ 61%, L-selectin 51%, CLA 13%) (Kantele et al. 2008) clearly resembles that found for FGT. The present HR profile (high to moderate proportion of $\alpha_4\beta_7$ and L-selectin, and low of CLA-expressing cells) might thus represent one shared by the urogenital area.

4.7. Limitations

Three limitations should be mentioned: small number of subjects, limited coverage of various homing markers, and the fact that we did not assess the possible effect of hormonal status. Future studies should cover other potential receptors and the impact of hormonal status on HR expression.

4.8. Conclusions

This study is the first to show that pathogen-specific plasmablasts appear in the circulation in response to bacterial infection in the upper FGT, the response predominated by IgA, followed by IgG and IgM plasmablasts. These circulating cells provide a tool for future investigations seeking to noninvasively explore immune responses elicited at this site. A homing profile resembling that reported previously for the upper urinary tract was revealed, suggesting a common urogenital homing from these sites.

Acknowledgements

We thank MSc Jukka Ollgren for expert advice on statistical analyses, and the personnel of HUSLAB for help in providing the bacterial strains.

Author Contributions

Conceived and designed the experiments: NVP, JMK, AK. Performed the experiments: NVP, OH, SHP. Analyzed the data: NVP, SHP, JMK, AK. Contributed reagents/materials/to collection of patient samples/analysis tools: OH, JMK, AK. Wrote the paper: NVP, OH, SHP, JMK, AK.

Conflicts of interest

All authors declare no conflicts of interest.

Funding

This work was supported by the Maud Kuistila Memorial Foundation supporting medical research of high standards in Finland (NVP). Funder played no role in the research.

REFERENCES

- Bekri S, Bourdely P, Luci C, Dereuddre-Bosquet N, Su B, Martinon F, Braud VM, Luque I, Mateo PL, Crespillo S, Conejero-Lara F, Moog C, Le Grand R, Anjuère F. Sublingual Priming with a HIV gp41-Based Subunit Vaccine Elicits Mucosal Antibodies and Persistent B Memory Responses in Non-Human Primates. *Front Immunol* 2017; 8:63. doi: 10.3389/fimmu.2017.00063. eCollection 2017. PMID: 28203239
- Berg EL, Yoshino T, Rott LS, Robinson MK, Warnock RA, Kishimoto TK, et al. The cutaneous lymphocyte antigen is a skin lymphocyte homing receptor for the vascular lectin endothelial cell-leukocyte adhesion molecule 1. *J Exp Med* 1991; 174: 1461–1466. doi: 10.1084/jem.174.6.1461.
- Berlin C, Berg EL, Briskin MJ, Andrew DP, Kilshaw PJ, Holzmann B, Weissman IL, Hamann A, Butcher EC. $\alpha 4\beta 7$ integrin mediates lymphocyte binding to the mucosal vascular addressin MAdCAM-1. *Cell* 1993; 74:185-95. PMID: 7687523
- Bernasconi NL, Traggiai E, Lanzavecchia A. Maintenance of serological memory by polyclonal activation of human memory B cells. *Science* 2002; 298:2199–202. doi: 10.1126/science.1076071.
- Brandtzaeg P, Johansen FE. Mucosal B cells: phenotypic characteristics, transcriptional regulation, and homing properties. *Immunol Rev* 2005; 206:32-63. PMID: 16048541
- Briskin M, Winsor-Hines D, Shyjan A, Cochran N, Bloom S, Wilson J, McEvoy LM, Butcher EC, Kassam N, Mackay CR, Newman W, Ringler DJ. Human mucosal addressin cell adhesion molecule-1 is preferentially expressed in intestinal tract and associated lymphoid tissue. *Am J Pathol* 1997; 151:97–110. PMID: 9212736
- Bomsel M, Tudor D, Drillet AS, Alfsen A, Ganor Y, Roger MG, Mouz N, Amacker M, Chalifour A, Diomedea L, Devillier G, Cong Z, Wei Q, Gao H, Qin C, Yang GB,

Zurbriggen R, Lopalco L, Fleury S. Immunization with HIV-1 gp41 subunit virosomes induces mucosal antibodies protecting nonhuman primates against vaginal SHIV challenges. *Immunity* 2011; 34:269-80. doi: 10.1016/j.immuni.2011.01.015. PMID: 21315623

Camerini D, James SP, Stamenkovic I, Seed B. Leu-8/TQ1 is the human equivalent of the Mel-14 lymph node homing receptor. *Nature* 1989; 342:78-82. PMID: 2509939

Cha HR, KO HJ, Kim ED, Chang SY, Seo SU, Cuburu N, Ryu S, Kim S, Kweon MN. Mucosa-associated epithelial chemokine/CCL28 expression in the uterus attracts CCR10+ IgA plasma cells following mucosal vaccination via estrogen control. *J Immunol* 2011; 187:3044-52. doi: 10.4049/jimmunol.1100402.

Csencsits KL, Jutila MA, Pascual DW. Nasal-associated lymphoid tissue: phenotypic and functional evidence for the primary role of peripheral node addressin in naive lymphocyte adhesion to high endothelial venules in a mucosal site. *J Immunol* 1999; 163:1382-9.

Cuburu N, Kweon MN, Hervouet C, Cha HR, Pang YY, Holmgren J, Stadler K, Schiller JT, Anjuère F, Czerkinsky C. Sublingual immunization with nonreplicating antigens induces antibody-forming cells and cytotoxic T cells in the female genital tract mucosa and protects against genital papillomavirus infection. *J Immunol* 2009; 183:7851-9. doi: 10.4049/jimmunol.0803740. PMID: 19933861.

Czerkinsky C, Prince SJ, Michalek SM, Jackson S, Russell MW, Moldoveanu Z, McGhee JR, Mestecky J. IgA antibody-producing cells in peripheral blood after antigen ingestion: evidence for a common mucosal immune system in humans. *Proc Natl Acad Sci U S A* 1987; 84:2449-53. PMID: 3470804. PMCID: PMC304669

- Davila SJ, Olive AJ, Starnbach MN. Integrin $\alpha 4\beta 1$ is necessary for CD4+ T cell-mediated protection against genital *Chlamydia trachomatis* infection. *J Immunol* 2014; 192:4284–93. doi: 10.4049/jimmunol.1303238.
- Iwasaki A. Antiviral immune responses in the genital tract: clues for vaccines. *Nat Rev Immunol* 2010; 10:699-711. doi: 10.1038/nri2836. Epub 2010 Sep 10.
- Johansen FE, Baekkevold ES, Carlsen HS, Farstad IN, Soler D, Brandtzaeg P. Regional induction of adhesion molecules and chemokine receptors explains disparate homing of human B cells to systemic and mucosal effector sites: dispersion from tonsils. *Blood* 2005; 106:593-600. Epub 2005 Apr 12. PMID: 15827133
- Johansson EL, Rask C, Fredriksson M, Eriksson K, Czerkinsky C, and Holmgren J. Antibodies and antibody-secreting cells in the female genital tract after vaginal or intranasal immunization with cholera toxin B subunit or conjugates. *Infect Immun* 1998; 66:514–20. PMID: 9453604
- Johansson EL, Rudin A, Wassén L, Holmgren J. Distribution of lymphocytes and adhesion molecules in human cervix and vagina. *Immunology* 1999; 96:272–7. doi: 10.1046/j.1365-2567.1999.00675.x. PMCID: PMC2326729.
- Kantele A. Antibody-secreting cells in the evaluation of the immunogenicity of an oral vaccine. *Vaccine*. 1990; 8:321-6. PMID: 2396471. doi: 10.1016/0264-410X(90)90088-4
- Kantele A. Immune response to prolonged intestinal exposure to antigen. *Scand J Immunol* 1991; 33: 225-9. doi: 10.1111/j.1365-3083.1991.tb03753.x. PMID: 2017658
- Kantele A. Persistence of diarrheal pathogens is associated with continued recruitment of plasmablasts in the circulation. *Clin Dev Immunol* 2012; 2012:279206. doi: 10.1155/2012/279206. Epub 2012 Jan 19. PMID: 22312405

- Kantele A, Arvilommi H, Jokinen I. Specific immunoglobulin-secreting human blood cells after peroral vaccination against *Salmonella typhi*. J Infect Dis 1986; 153:1126-31. PMID: 3701119
- Kantele AM, Takanen R, Arvilommi H. Immune response to acute diarrhea seen as circulating antibody-secreting cells. J Infect Dis 1988; 158:1011-6. doi: 10.1093/infdis/158.5.1011. PMID: 305391830
- Kantele A, Papunen R, Virtanen E, Möttönen T, Räsänen L, Ala-Kaila K, Mäkelä H, Arvilommi H. Antibody-secreting cells in acute urinary tract infection as indicators of local immune response. J Infect Dis 1994; 169:1023-1028. doi: 10.1093/infdis/169.5.1023
- Kantele A, Kantele JM, Savilahti E, Westerholm M, Arvilommi H, Lazarovits A, Butcher EC, Mäkelä PH. Homing potentials of circulating lymphocytes in humans depend on the site of activation: oral, but not parenteral, typhoid vaccination induces circulating antibody-secreting cells that all bear homing receptors directing them to the gut. J Immunol 1997; 158:574-9. PMID: 8992970
- Kantele A, Häkkinen M, Moldoveanu Z, Lu A, Savilahti E, Alvarez RD, Michalek S, Mestecky J. Differences in immune responses induced by oral and rectal immunizations with *Salmonella typhi* ty21a: evidence for compartmentalization within the common mucosal immune system in humans. Infect Immun 1998; 66:5630–5. PMID: 9510871
- Kantele A, Savilahti E, Tiimonen H, Iikkanen K, Autio S, Kantele JM. Cutaneous lymphocyte antigen expression on human effector B cells depends on the site and on the nature of antigen encounter. Eur J Immunol 2003; 33:3275-83. PMID: 14635035. DOI: 10.1002/eji.200324311

Kantele AM, Palkola, NV, Arvilommi, HS, Kantele, JM. Distinctive homing profile of pathogen-specific activated lymphocytes in human urinary tract infection. *Clin Immunol* 2008; 128:427-34. doi: 10.1016/j.clim.2008.05.003. Epub 2008 Jun 27. PMID: 18585960

Kantele JM, Arvilommi H, Kontiainen S, Salmi M, Jalkanen S, Savilahti E, Westerholm M, Kantele A. Mucosally activated circulating human B cells in diarrhea express homing receptors directing them back to the gut. *Gastroenterology* 1996a; 110:1061-7. PMID: 8612994

Kantele JM, Kantele A, Arvilommi H. Circulating immunoglobulin-secreting cells are heterogeneous in their expression of maturation markers and homing receptors. *Clin Exp Immunol* 1996b; 104:525-30. PMID: 9099939

Kelly KA, Natarajan S, Ruther P, Wisse A, Chang MH, Ault KA. Chlamydia trachomatis infection induces mucosal addressin cell adhesion molecule-1 and vascular cell adhesion molecule-1, providing an immunologic link between the fallopian tube and other mucosal tissues. *J Infect Dis* 2001; 184:885-91. Epub 2001 Aug 30. PMID: 11550128. doi: 10.1086/323341

Kelly KA, Wiley D, Wiesmeier E, Briskin M, Butch A, Darville T. The combination of the gastrointestinal integrin ($\alpha 4\beta 7$) and selectin ligand enhances T-Cell migration to the reproductive tract during infection with Chlamydia trachomatis. *Am J Reprod Immunol* 2009; 61(6):446-52. doi: 10.1111/j.1600-0897.2009.00705.x. Epub 2009 Apr 22. PMID: 19392980

Kozlowski PA, Williams SB, Lynch RM, Flanigan TP, Patterson RR, Cu-Uvin S, Neutra MR. Differential induction of mucosal and systemic antibody responses in women after nasal, rectal, or vaginal immunization: influence of the menstrual cycle. *J Immunol* 2002; 169:566-74. PMID: 12077289

- Kutteh WH, Blackwell RE, Gore H, Kutteh CC, Carr BR, Mestecky J. Secretory immune system of the female reproductive tract. II. Local immune system in normal and infected fallopian tube. *Fertil Steril* 1990; 54:51-5. PMID:2141578
- Kutteh WH, Kantele A, Moldoveanu Z, Crowley-Nowick PA, Mestecky J. Induction of specific immune responses in the genital tract of women after oral or rectal immunization and rectal boosting with *Salmonella typhi* Ty 21a vaccine. *J Reprod Immunol* 2001; 52:61-75. PMID: 11600178
- Li Z, Palaniyandi S, Zeng R, Tuo W, Roopenian DC, Zhu X. Transfer of IgG in the female genital tract by MHC class I-related neonatal Fc receptor (FcRn) confers protective immunity to vaginal infection. *Proc Natl Acad Sci U S A*. 2011; 108:4388-93. doi: 10.1073/pnas.1012861108.
- Mascola JR, Stiegler G, VanCott TC, Katinger H, Carpenter CB, Hanson CE, Beary H, Hayes D, Frankel SS, Birx DL, Lewis MG. Protection of macaques against vaginal transmission of a pathogenic HIV-1/SIV chimeric virus by passive infusion of neutralizing antibodies. *Nat Med* 2000; 6:207–10. doi: 10.1038/72318
- Palkola NV, Pakkanen SH, Kantele JM, Rossi N, Puohiniemi R, Kantele A. Pathogen-specific circulating plasmablasts in patients with pneumonia. *PLoS One* 2012; 7(3):e34334. doi: 10.1371/journal.pone.0034334. PMID: 22479603
- Palkola NV, Pakkanen SH, Kantele JM, Pakarinen L, Puohiniemi R, Kantele A Differences in homing potentials of *Streptococcus pneumoniae*-specific plasmablasts in pneumococcal pneumonia and after pneumococcal polysaccharide and pneumococcal conjugate vaccinations. *J Infect Dis* 2015; 212(8):1279-87. doi: 10.1093/infdis/jiv208. PMID: 25838267
- Palkola NV, Blomgren K, Pakkanen SH, Puohiniemi R, Kantele JM, Kantele A. Immune defense in upper airways: A single-cell study of pathogen-specific plasmablasts and

their migratory potentials in acute sinusitis and tonsillitis. PLoS One 2016; 11(4):e0154594. doi: 10.1371/journal.pone.0154594. eCollection 2016. PMID: 27128095

Pakkanen SH, Kantele JM, Moldoveanu Z, Hedges S, Häkkinen M, Mestecky J, Kantele A. Expression of homing receptors on IgA1 and IgA2 plasmablasts in blood reflects differential distribution of IgA1 and IgA2 in various body fluids. Clin Vaccine Immunol 2010; 17 :393-401

Quesnel A, Cu-Uvin S, Murphy D, Ashley RL, Flanigan T, Neutra MR. Comparative analysis of methods for collection and measurement of immunoglobulins in cervical and vaginal secretions of women. J Immunol Methods 1997; 202:153-61. doi: 10.1016/S0022-1759(97)00003-3

Quiding-Järbrink M, Nordström I, Granström G, Kilander A, Jertborn M, Butcher EC, Lazarovits AI, Holmgren J, Czerkinsky C. Differential expression of tissue-specific adhesion molecules on human circulating antibody-forming cells after systemic, enteric, and nasal immunizations. A molecular basis for the compartmentalization of effector B cell responses. J Clin Invest 1997; 99:1281-6. PMID: 9077537. doi: 10.1172/JCI119286

Shin H, Iwasaki A. Generating protective immunity against genital herpes. Trends Immunol 2013; 34:487-94. doi: 10.1016/j.it.2013.08.001. Epub 2013 Sep 3. PMID: 24012144

Sigmundsdottir H, Butcher EC. Environmental cues, dendritic cells and the programming of tissue-selective lymphocyte trafficking. Nature Immunol 2008; 9:981-7. PMID: 18711435

Tabibzadeh S, Kong QF, Babaknia A. Expression of adhesion molecules in human endometrial vasculature throughout the menstrual cycle. J Clin Endocrinol Metab 1994; 79:1024–32. PMID: 7962270

Wira CR, Rodriguez-Garcia M, Patel MV. The role of sex hormones in immune protection of the female reproductive tract. *Nat Rev Immunol* 2015; 15(4):217-30. doi:

10.1038/nri3819. Epub 2015 Mar 6. Review. PMID: 25743222

Woof JM and Mestecky J. Mucosal immunoglobulins. *Immunol Rev* 2005; 206 (1): 64–82.

10.1111/j.0105-2896.2005.00290.x

ACCEPTED MANUSCRIPT

FIGURE LEGENDS




Fig. 1. Flow diagram of the study (ASC – pathogen-specific antibody-secreting cell, ISC – immunoglobulin-secreting cell)




Fig. 2. Numbers of pathogen-specific ASC and all ISC

The numbers of (IgA + IgG + IgM) ASC and -ISCs /10⁶ PBMC in the peripheral blood are given individually for the 12 female patients with bacterial upper genital tract infection, and 6 healthy controls. The patients were examined 7–12 days after onset of symptoms. Open triangles indicate non-invasive and black ones invasive cases. Horizontal lines show geometric means. The numbers proved higher in patients than controls when tested with independent samples Mann-Whitney U test.




Fig. 3. Ig isotype distributions of pathogen-specific ASC and all ISC

Triangles represent the number of pathogen-specific IgA, IgG or IgM ASC (A) or ISC (/10⁶ PBMC) in the circulation of patients with upper genital tract infection (B) and healthy controls (C). Individual patients' results are joined with a line; open triangles indicate non-invasive and black ones invasive cases. Horizontal lines show geometric means (n=12 for patients and 6 for controls).




Fig. 4. Proportions of $\alpha_4\beta_7$ -, L-selectin-, and CLA-expressing cells among ASC and ISC

The bars indicate the means of the percentages of ASC or ISC expressing the given HR among the ASC and ISC of the patients and the ISC of the controls. The number of study

subjects from whom the data were pooled is indicated under each bar. The proportions of cells expressing $\alpha_4\beta_7$, L-selectin and CLA were similar in comparisons (Related-samples Wilcoxon Signed Rank test for ASC vs. ISC of patients; Independent-samples Mann-Whitney U test for ASC/ ISC of patients vs. ISC of controls).




Fig. 5. Expression of $\alpha_4\beta_7$, L-selectin, and CLA on pathogen-specific ASC in acute invasive and non-invasive female reproductive tract infection

The bars indicate arithmetic means with SD of percentages of cells expressing the given HR among ASC (IgA + IgG+ IgM). The number of patients for whom the data were pooled is indicated under each bar. The proportions of $\alpha_4\beta_7$ -, L-selectin-, and CLA-expressing cells appeared similar in the invasive and non-invasive groups (Independent-samples Mann-Whitney U test).

Table 1. Patient demographics, clinical data, number of pathogen-specific antibody-secreting cells (ASC), and proportions of Ig-isotypes.

Sample no.	Age	Infection	Pathogen	Day ^a	Fever °C	CRP mg/L	ASC ^b /10 ⁶ PBMC	IgA ASC %	IgG ASC %	IgM ASC %
Non-invasive										
1.1	23	Salpingitis	<i>Escherichia coli</i>	7	39	189	9 947	82	6	12
1.2	45	Salpingitis	<i>Staphylococcus aureus</i>	8	38.6	178	1 164	13	75	12
1.3	42	Salpingitis	<i>Bacteroides fragilis</i>	8	39	250	785	52	31	17
1.4	43	Salpingitis	<i>Escherichia coli</i>	12	>39	360	68	44	25	31
1.5	40	Salpingitis	<i>Escherichia coli</i>	7	NA ^c	NA	971	50	34	16
Invasive										
2.1	26	Chorioamnionitis	<i>Haemophilus influenzae</i>	8	38,7	135	2 757	6	74	20
2.2	25	Chorioamnionitis	<i>Streptococcus pyogenes</i>	7	38	110	65	61	39	0
2.3	42	Chorioamnionitis	<i>Streptococcus pyogenes</i>	8	NA	NA	468	81	18	1
2.4	38	Endometritis	<i>Streptococcus pyogenes</i>	9	>39	>200	135	19	78	3
2.5	51	Endometritis	<i>Escherichia coli</i>	9	NA	NA	11 575	86	7	7
2.6	23	Chorioamnionitis	<i>Eikenella corrodens</i>	7	NA	NA	28	14	21	65
2.7	18	Chorioamnionitis	<i>Streptococcus pyogenes</i>	9	>39	300	20	16	0	84

^a number of days between onset of symptoms and blood sample

^b (IgA + IgG + IgM) ASC

^c NA = data not available

ACCEPTED MANUSCRIPT