

# Accepted Manuscript

Catheter-Directed Thrombolysis vs. Pharmacomechanical Thrombectomy for Upper Extremity Deep Venous Thrombosis: Cost-Effectiveness Analysis

O. Mahmoud, P. Vikatmaa, J. Räsänen, E. Peltola, E. Sihvo, L. Vikatmaa, K. Lappalainen, M. Venermo



PII: S0890-5096(18)30229-2

DOI: [10.1016/j.avsg.2018.01.104](https://doi.org/10.1016/j.avsg.2018.01.104)

Reference: AVSG 3783

To appear in: *Annals of Vascular Surgery*

Received Date: 30 November 2017

Revised Date: 21 January 2018

Accepted Date: 24 January 2018

Please cite this article as: Mahmoud O, Vikatmaa P, Räsänen J, Peltola E, Sihvo E, Vikatmaa L, Lappalainen K, Venermo M, Catheter-Directed Thrombolysis vs. Pharmacomechanical Thrombectomy for Upper Extremity Deep Venous Thrombosis: Cost-Effectiveness Analysis, *Annals of Vascular Surgery* (2018), doi: 10.1016/j.avsg.2018.01.104.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1  
2 Catheter-Directed Thrombolysis vs. Pharmacomechanical Thrombectomy for  
3 Upper Extremity Deep Venous Thrombosis: Cost-Effectiveness Analysis

4  
5 Mahmoud O<sup>1,2</sup>, Vikatmaa P<sup>1</sup>, Räsänen J<sup>3</sup>, Peltola E<sup>4</sup>, Sihvo E<sup>5</sup>, Vikatmaa L<sup>6</sup>, Lappalainen K<sup>4</sup>, Venermo M<sup>1</sup>

6  
7 <sup>1</sup>Department of Vascular Surgery, Helsinki University Hospital and Institute of Clinical Medicine, Faculty  
8 of Medicine, University of Helsinki, Finland; <sup>2</sup>Department of Vascular Surgery, Assiut University Hospital,  
9 Faculty of Medicine, Assiut University, Egypt, <sup>3</sup> Department of General Thoracic and Esophageal Surgery,  
10 Heart and Lung Centre, University of Helsinki and Helsinki University Hospital, Helsinki, Finland,  
11 <sup>4</sup>Department of Radiology, Helsinki University Hospital and Institute of Clinical Medicine, Faculty of  
12 Medicine, University of Helsinki, Finland; <sup>5</sup> Department of Surgery, Central Finland Central Hospital,  
13 Jyväskylä, Finland and University of Helsinki, Finland; <sup>6</sup> Department of Anesthesiology, Intensive Care  
14 and Pain Medicine, Helsinki University Hospital and University of Helsinki, Finland

15  
16 Short title: Invasive treatment of UEDVT

17 Declaration of conflicting interests: None declared

18 Funding: This research did not receive any specific grant from funding agencies in the public,  
19 commercial, or not-for-profit sectors.

20 **Correspondence to:**

21 Maarit Venermo

22 Helsinki University Central Hospital, Department of Vascular Surgery

23 P.O. Box 440, FI-00029 HUS

24 Helsinki, Finland

25 E-mail: maarit.venermo@hus.fi

26 **ABSTRACT**

27

28 **Background and Aims:** We compared the immediate and one-year results as well as total hospital costs  
29 between catheter-directed thrombolysis (CDT) and pharmacomechanical thrombolysis (PMT) in the  
30 treatment of symptomatic upper extremity deep venous thrombosis (UEDVT).

31 **Material and Methods:** From 2006 to 2013, 55 patients with UEDVT were treated with either CDT or  
32 PMT at Helsinki University Hospital. Of them, 43 underwent thoracoscopic rib resection later in order to  
33 relieve phlebography-confirmed vein compression. This patient cohort was prospectively followed up  
34 with repeated phlebographies. CDT was performed to 24 patients and 19 had PMT with a Trellis™  
35 device. Clinical evaluation and vein patency assessment were performed with either phlebography or  
36 ultrasound one year after the thrombolysis. Primary outcomes were immediate technical success, one-  
37 year vein patency, and costs of the initial treatment.

38 **Results:** The immediate overall technical success rate, defined as recanalization of the occluded vein and  
39 removal of the fresh thrombus, was 91.7% in the CDT group, and 100% in the PMT group (n.s.). The  
40 median thrombolytic time was significantly longer in CDT patients than PMT patients (21.1 hours vs.  
41 0.33 hours,  $P < 0.00001$ ). There were no procedure-related complications. The one-year primary assisted  
42 patency rate was similar in both groups (91.7% and 94.7%, respectively). There were no recurrences of  
43 clinical DVT. The hospital costs for the acute period were significantly lower in the PMT group than the  
44 CDT group (medians 11,476 € and 5,975 € in the in the CDT and PMT group, respectively ( $P < 0.00001$ )).

45 **Conclusions:** The clinical results of the treatment of UEDVT with CDT or PMT were similar. However,  
46 PMT required shorter hospital stay and less intensive surveillance, leading to lower total costs.

47

48 **Key words:** Upper extremity deep venous thrombosis, thrombolysis, thrombectomy, catheter-directed  
49 thrombolysis, pharmacomechanical thrombectomy

50

**51 INTRODUCTION**

52

53 Upper extremity deep vein thrombosis (UEDVT) represents approximately 2–3% of all deep vein  
54 thromboses (1). Primary UEDVT includes idiopathic and effort-related thrombosis (Paget–Schroetter  
55 syndrome). Effort-related UEDVT may be related to abnormal anatomy, or it may be a consequence of  
56 strenuous activity (2). Secondary UEDVT is mostly related to central venous catheters, pacemaker  
57 devices, or malignancy (3).

58

59 The clinical manifestations of UEDVT include edema of the affected extremity in 80%, extremity pain in  
60 30%–50%, and erythema in approximately 15% of the patients (4). Approximately 5% of patients have  
61 no symptoms (5,6). The incidence of post-thrombotic syndrome (PTS) in the upper limb ranges from 7%  
62 to 46%, and PTS may result in significant morbidity, especially if it occurs in the dominant arm (7).  
63 Treatment with anticoagulant therapy alone is associated with delayed resolution of acute symptoms,  
64 reduced venous outflow, and increased incidence of residue thrombus, chronic venous obstruction,  
65 venous valvular incompetence, and subsequent venous hypertension (8). Systemic thrombolysis has  
66 been associated with major hemorrhagic problems. In order to reduce the thrombolytic therapy dose  
67 and the bleeding risk, American College of Chest Physicians (ACCP) guidelines encourage catheter-based  
68 thrombolysis over systemic infusion in treatment of UEDVT with severe symptoms; with thrombus  
69 extending most of the subclavian and the axillary vein, symptoms for <14 days, good functional status,  
70 life expectancy of > 1 year, and a low risk for bleeding (9).

71

72 CDT has also been associated with major systemic hemorrhage, and long infusion times (8). PMT with  
73 the Trellis™ device (Trellis™ Peripheral Infusion System, Covidien, Mansfield, MA, USA) intended to  
74 overcome these disadvantages by combining mechanical clot disruption and pharmacological

75 thrombolysis within an isolated zone. However, due to problems with the sterilization process and  
76 errors in marking of the balloons, Trellis was withdrawn from the market in xxxx. The current study was  
77 conducted prior to the withdrawal. To our knowledge, this is the only study comparing PMT and CDT for  
78 UEDVT.

79

80 We report the immediate and mid-term results as well as the total hospital costs between conventional  
81 CDT and PMT with the Trellis™ device in the treatment of symptomatic UEDVT.

82

ACCEPTED MANUSCRIPT

83 **MATERIAL AND METHODS**

84

85 From 2006 to 2013, 72 patients with UEDVT were seen at the Helsinki University Hospital. All patients  
86 were clinically assessed and duplex ultrasonography (DUS) was used as the primary diagnostic method.

87 The coagulation profile was assessed at the time of the first visit. Computed tomography (CT) and

88 magnetic resonance imaging (MRI) were used in patients with inconclusive duplex data or if a pulmonary

89 embolism was suspected. Of the 72 patients, 17 with minimal symptoms were treated conservatively

90 using low-molecular-weight heparin (LMWH), and 55 patients with more severe symptoms were treated

91 invasively, with either CDT or PMT using the Trellis™ device. After CDT/PMT, all patients underwent

92 completion phlebography with provocation tests in order to assess the technical success and

93 completeness of the thrombolysis/thrombectomy, as well as a possible vein compression. Technical

94 success was defined as successful recanalization of the occluded vein and removal of the fresh

95 thrombus. Completeness of the thrombolysis/thrombectomy was graded to three classes: “complete” if

96 phlebography showed no further clot, “partial” if thrombolysis was incomplete, but less than 50% of the

97 thrombus remained, and “failed” when more than 50% of the thrombus was present after the

98 intervention (11,12). To detect compression, phlebography was performed both at rest (full adduction)

99 and at provocation (arm 90 degrees abducted with external rotation “hand-on-head position”).

100

101 As a second stage procedure, forty three (78.2%) of 55 patients who underwent early clot removal

102 underwent a thoracoscopic first rib resection due to an external compression of the vein detected in

103 post-thrombolysis phlebography (13). They were included in a prospective rib resection surveillance

104 program, which was analysed retrospectively from hospital records. The surveillance protocol included

105 both clinical control and patency examinations with DUS and phlebography, as well as an assessment for

106 the need for PTA based on the presence of symptoms and stenosis >20% or occlusion.

107

108 Twelve (12%) of the 55 patients underwent local thrombolysis only with no rib resection and were  
109 excluded, because there was no systematic follow-up imaging available for these patients. The reasons  
110 being malignancy in 2 patients, lack of extrinsic compression in control phlebography in 5 patients, local  
111 foreign material (pacemaker) in 2 patients, and chronic occlusion of the subclavian vein in 3 patients (all  
112 three had thrombophilia and minimal or no symptoms after CDT) (Figure 1).

113 Twenty-four of the study patients underwent CDT and 19 had PMT. In the beginning of the study period,  
114 in 2006 until 2012, CDT was used routinely. Since 2012, the Trellis™ device was introduced in our  
115 institution and became more popular. Patient selection was thus partly time-dependent; and, towards  
116 the end of the study period, depended upon whether the radiologist performing the procedure was  
117 familiar with the PMT procedure or not. Phlebographic images obtained at the time of treatment and  
118 during the follow-up were carefully reviewed.

119 To assess mid-term results, one-year duplex scan reports as well as phlebography images and reports  
120 were evaluated. Duplex scan only was performed to 25 patients, phlebography only to 12 patients and  
121 both duplex and phlebography to 6 patients mostly due to inconclusive result of duplex. In the end the  
122 mid term patency assessment was based on duplex scan in 25 cases and phlebography in 18 patients.

123

124 The end points were immediate technical success and one-year vein patency. Successful PTA was  
125 defined as residual stenosis of 0-20%.

126

127 The detailed hospital costs for all patients during lysis admission time were collected from the hospital  
128 cost database (Financial administration services of the Department of Surgery, Helsinki University  
129 Hospital).

130

131

**132 Treatment options****133 A Catheter-directed thrombolysis**

134 Percutaneous access was achieved with ultrasound guidance primarily through the basilic vein and  
135 secondarily through the cephalic, brachial, or cubital veins, using a 4-French sheath. A 0.035"  
136 hydrophilic wire (Radiofocus guide wire M, Terumo Co., Japan) was passed through in the CDT group,  
137 and a diagnostic phlebography was performed to assess the lesion, its extension, and the presence of  
138 collaterals.

139

140 A single dose of alteplase (10 mg) was administered through a multi-hole catheter (täähän katetrin  
141 tiedot) into the occlusion, and infusion at a rate of 1 mg/hour was started. The patient was observed at  
142 the intermediate care unit. After approximately 24 hours of thrombolysis, a second phlebography was  
143 performed to assess the lytic success and the need to continue thrombolysis for an additional 24 hours.  
144 In the final phlebography, the need for balloon angioplasty was assessed and in case of a significant  
145 stenosis a 8-12mm balloon was used (Figure 2).

146

**147 B Pharmacomechanical thrombectomy**

148 The access technique and assessment of the lesion were similar to the CDT procedure. The Trellis™  
149 catheter was positioned over a 0.035" guide wire through an 8-Fr sheath, leaving the area of treatment  
150 between the two inflated balloons. Thereafter, 6–10 mg of alteplase was injected through the side holes  
151 of the device catheter and a rotational technique started to disrupt the thrombus. For the next 10–20  
152 minutes, alteplase was injected slowly to promote thrombolysis. Finally, the melted thrombosis was  
153 aspirated with a 50-ml syringe through the side hole and the vein was evaluated with a manual injection  
154 of contrast media. After the first session, if thrombolysis was not complete, the pharmacomechanical



155 lysis was continued using 6–10 mg of alteplase for another 10–20 minutes. The maximum amount of  
156 alteplase was 20 mg. Seventeen (89%) patients had a single-session lysis. In 16 patients, the lysis lasted  
157 for 20 minutes and for 3 patients, 10 minutes due to a short occlusion with a small thrombus. Two (11%)  
158 patients required 2 sessions lasting a total of 30–35 minutes. Completion phlebography, possible PTA  
159 and anticoagulation were similar to the CDT group. Patients treated with PMT had no need for a stay in  
160 an intermediate care unit.

161

162 Low molecular weight heparin was started with a dose 1mg/kg twice a day immediately when the  
163 diagnosis was made and continued during the CDT and PMT. After thrombolysis, the patients were kept  
164 on LMWH and warfarin until the INR reached 2–2.5, after which warfarin treatment was continued for  
165 3–6 months.

166

167 The surgical decompressions were later performed with a video-assisted thoracoscopic first rib resection  
168 (VTRR) technique. The procedure is described in detail elsewhere (13).

169

#### 170 **Statistical analysis**

171 SPSS 22.00 was used in the statistical analysis. Continuous variables are expressed as median values  
172 (range). The prevalence of risk factors is expressed as percentages. Comparisons between the groups  
173 were made using the Mann–Whitney U test (continuous variables) and chi-square test (dichotomic  
174 variables).

175 The study protocol has been accepted by the Institutional Review Board (HUS/214/2016). Because of  
176 the retrospective nature of this study, no informed consent was obtained from the study subjects.

177

178

**179 RESULTS**

180 The CDT group included 24 patients with a median age of 31 years, and the PMT group included 19  
181 patients with median age of 26. There were no significant differences in patient demographics between  
182 the groups. The most common symptoms were swelling, pain, and numbness of the affected extremity,  
183 with an equal prevalence in the groups (Table I). Duplex US was used as the first diagnostic examination  
184 in 41 patients (95%), while additional imaging was employed in 5 (12%): MRI in 2, and CT in 3 patients.  
185 The median time between symptom onset and intervention was 4.5 days (range 1–12 days) and 4 days  
186 (range 1–7 days) in CDT and PMT groups, respectively (n.s.). The median thrombosis length in treated  
187 patients with Trellis™ was 116 mm (range 30-225 mm), and in the CDT group the median lesion length  
188 was 160.5 mm (range 45-254 mm). The median time from the early clot removal to the rib resection was  
189 92 days (range, 10-458 days), 68.5 days (range, 15-458 days) in the CDT group and 120 days (range, 10-  
190 265 days) in the PMT group.

191

192

**193 Immediate technical success**

194 Immediate overall technical success was 92% and 100% in the CDT and PMT groups, respectively. In the  
195 PMT group, complete lysis was achieved in 17 (90%) and partial lysis in 2 (11%) patients. In the CDT  
196 group, the therapeutic response was complete in 19 (79%) and partial in 3 (13%) cases, while the  
197 treatment failed in 2 (8%). The residual lesion after thrombolysis and the change in the lesion's  
198 topography before rib resection are shown in Table II.

199

200 In 2 patients (8%), the treatment was started with CDT, but due to persistent thrombosis after two days  
201 of thrombolysis, PMT was successfully initiated to remove the residual thrombosis. Ten (42%) patients in  
202 the CDT group and 8 (42%) in the PMT group underwent balloon angioplasty due to moderate or

203 significant stenosis in the completion phlebography. The immediate phlebographic results are presented  
204 in Table II. There was no pulmonary embolism found in the CT scan in patients with a clinical suspicion of  
205 PE, or other major complications during the hospital stay after either of the procedures. The treatment  
206 parameters of the PMT and CDT patients are shown in Table III.

207

### 208 **One-year follow-up for vein patency**

209 After a median follow-up of 13 months (range 10–36), the vein patency was assessed either by  
210 phlebography (n=18) or duplex US (n=25). A good flow with no significant stenosis (<20%) was observed  
211 in 18 (75%) patients in the CDT group, and in 17 (90%) patients in the PMT group (n.s.). No significant  
212 difference in symptoms or technical success were seen at one year (Table IV). During the follow-up  
213 period, 11 (46%) patients in the CDT group and 10 (53%) patients in the PMT group (ns) underwent  
214 balloon angioplasty due to stenosis >20% or occlusion associated persistent symptoms (Table IV). No  
215 stents were used. The overall assisted primary patency at one year was 92% (n=22) in the CDT group and  
216 95% (n=18) in the PMT group (ns). No patients suffered a recurrence of clinical DVT during the follow-  
217 up.

218

### 219 **Total hospital costs**

220 The median total procedural cost of the hospital stay per patient was 6,986 (range 6,100–8,564) € in the  
221 CDT group and 4,499 (range 3,782–5,120) € in the PMT group,  $P < 0.001$ . The median total hospital cost  
222 was 11,476 (range 8,468–17,467) €/patient in the CDT group and 5,975 (range 4,763–7,395) €/patient in  
223 the PMT group ( $P < 0.001$ ). (Table III).

224

## 225 **DISCUSSION**

226

227 In acute UEDVT, both CDT and PMT are effective treatment methods and work more quickly than  
228 anticoagulation in the recanalization of the occluded vein (8,10). Studies on the results of PMT are  
229 scarce. We report a consecutive case series of 43 patients with symptomatic UEDVT who underwent  
230 invasive treatment with either CDT or PMT using a Trellis™ device and a thoracoscopic rib resection  
231 thereafter. We compared the safety, efficacy, one-year results, and total hospital costs of the two  
232 treatment methods. We found that PMT was associated with a significantly shorter treatment time, as  
233 well as lower total hospital costs than CDT, with similar safety, efficacy, and one-year results.  
234 Furthermore, the immediate phlebographic success was more often successful after PMT.

235  
236 Our results are comparable with those reported in previous publications comparing CDT and PMT,  
237 although the majority of the patients in these studies have had lower-extremity DVT (LEDVT) (10,14).  
238 Kim et al. compared CDT and PMT in the treatment of 23 UEDVT and 44 LEDVTs in 36 patients (14).  
239 Catheter-directed thrombolysis was performed in 40 cases and pharmacomechanical thrombectomy  
240 with an Angiojet rheolytic thrombectomy catheter in 27 cases. The mean duration of the treatment was  
241 significantly longer in CDT when compared to PMT—48 and 26 hours, respectively. In addition, the  
242 consumption of urokinase was significantly lower in PMT. The authors achieved complete clot lysis in  
243 73% using CDT and 82% with PMT. Lin et al., in turn, compared CDT and PMT with an Angiojet rheolytic  
244 thrombectomy system in 98 patients (10). They reported complete lysis of the thrombus in 75% of the  
245 patients after PMT versus 70% after CDT (n.s.) and partial lysis in 25% and 30% of the patients,  
246 respectively.

247  
248 The largest benefit of PMT in comparison to CDT is the need for minimal or no intermediate care unit  
249 treatment and a shorter hospital stay. None of our PMT patients needed to be admitted to an  
250 intermediate care unit, and they spent an average 3 days in the hospital. The CDT patients required an

251 average of 2 days of treatment in the intermediate care unit, which was the duration of thrombolysis;  
252 and the median length of the hospital stay was 6 days. Lin et al. reported somewhat longer treatment  
253 periods (10).

254

255 The delay between the onset of symptoms and treatment has an impact on the success of thrombus  
256 removal. If thrombolysis is performed within a few days, the primary success rate is close to 100%. After  
257 two weeks, the success rate decreases to 85%; and after 6 weeks, down to 50% (15,16). The ACCP  
258 guidelines recommend that local thrombolysis should be performed in patients with severe symptoms  
259 of recent onset (<14 days) if appropriate expertise and resources are available (9). In our study, the  
260 median time between symptom onset and intervention was approximately 4 days in both groups.  
261 Probably due to the relatively short delay, we had a high immediate technical success rate with an  
262 overall thrombus removal of 100% in the PMT group and 92% in the CDT group (n.s.).

263

264 In many studies, the major drawback of CDT therapy has been hemorrhagic complications, which have  
265 been related to prolonged treatment duration (17-23). Our CDT patients received a median of 21 hours'  
266 infusion of the thrombolytic agent, as opposed to 20 minutes in PMT patients. We did not observe any  
267 bleeding complications, probably due to the small sample size.

268

269 The aim of CDT and PMT is to open the occluded vein and achieve immediate relief of the symptoms.  
270 However, long-term patency of the treated vein is also important. If a significant stenosis persists in  
271 provocation phlebography, a risk of rethrombosis exists and our treatment of choice is to perform a  
272 thoracoscopic first rib resection, and a postoperative balloon angioplasty of the vein when appropriate  
273 (13,23). The focus of this paper was to compare two different treatment options in the acute phase.  
274 One-year vein patency and the need for PTA was equal between the groups. No stents were used in

275 these patients because even if data are sparse in the literature, stent fractures have been found to be  
276 common in this position (24).

277 We used the Trellis™ device to achieve PMT with no major difficulties or complications. Unfortunately,  
278 the device was later withdrawn from the market. However, other devices for pharmacomechanical  
279 thrombectomy are still available. The results with the Angiojet rheolytic (Possis Medical, Minneapolis,  
280 MN) thrombectomy device are comparable to ours (10,11). We have had good experiences with PMT,  
281 the main benefit being the savings in intermediate care, and are now looking for a suitable device for  
282 routine use.

283  
284 The main limitation of our study is the small number of patients. Furthermore, the length of follow-up  
285 was limited. The treatments were performed during different time periods: CDT was used in the  
286 beginning of the study period and the method then changed to PMT, which was mostly used in the  
287 latter period. CDT has been performed in our institution for years in both upper and lower extremities  
288 and the procedure was familiar to all interventional radiologists. Altogether 7 interventionalists was  
289 performing CDT in this material. However, as PMT was initiated during the study period, there might be  
290 some learning curve effect. However, when started, all PMTs were performed by 2 interventional  
291 radiologists made all except three Trellis PMTs in this study. However, Otherwise nothing else in the  
292 treatment protocol changed; patients underwent similar rib resection after the initial treatment, and the  
293 medication after thrombolysis/thrombectomy was the same. In the beginning of this study, a  
294 phlebographic protocol before and after rib resection was designed, and we chose to include only  
295 patients with a complete phlebographic work-up.

296

297 **CONCLUSION**

298

299 The immediate and one-year clinical results of the treatment of subclavian vein thrombosis with CDT  
300 and PMT are equal. However, the need for admission to an intermediate care unit, hospital stay, as well  
301 as multiple phlebographic sessions and prolonged thrombolysis, were significantly more infrequent in  
302 patients treated with PMT than with CDT, leading to significantly lower total costs.

303

304

ACCEPTED MANUSCRIPT

## 305 REFERENCES

306

- 307 1. Lindblad B, Bornmyr S, Kullendorff B, et al. Venous haemodynamics of the upper extremity after  
308 subclavian vein thrombosis. *Vasa* 1990; 19: 218–222.
- 309 2. Hughes ESR. Venous obstruction in the upper extremity (Paget-Schroetter's syndrome). *Intl*  
310 *Abstracts of Surg* 1949; 88: 89–127.
- 311 3. Joffe HV, Goldhaber SZ. Upper-extremity deep vein thrombosis. *Circulation* 2002; 106: 1874–  
312 1880.
- 313 4. Thompson RW. Comprehensive management of subclavian vein effort thrombosis. *Semin*  
314 *Intervent Radiol* 2012; 29: 44–51.
- 315 5. Marinella MA, Kathula SK, Markert RJ. Spectrum of upper-extremity deep venous thrombosis in  
316 a community teaching hospital. *Heart Lung* 2000; 29: 113–117.
- 317 6. Joffe HV, Kucher N, Tapson VF, et al. Upper-extremity deep vein thrombosis: a prospective  
318 registry of 592 patients. *Circulation* 2004; 110: 1605–1611.
- 319 7. Elman EE, Kahn SR. The post-thrombotic syndrome after upper extremity deep venous  
320 thrombosis in adults: a systematic review. *Thromb Res* 2006; 117: 609–614.
- 321 8. Persson LM, Arnhjort T, Lärfars G, et al. Hemodynamic and morphologic evaluation of sequelae  
322 of primary upper extremity deep venous thromboses treated with anticoagulation. *J Vasc Surg*  
323 2006; 43: 1230–1235.
- 324 9. Meissner MH, Gloviczki P, Comerota AJ, Dalsing MC, Eklof BG, Gillespie DL, et al. Early thrombus  
325 removal strategies for acute deep venous thrombosis: clinical practice guidelines of the Society  
326 for Vascular Surgery and the American Venous Forum. *J Vasc Surg* 2012; 55: 1449-1462



- 327 10. Lin PH, Zhou W, Dardik A, et al. Catheter-direct thrombolysis versus pharmacomechanical  
328 thrombectomy for treatment of symptomatic lower extremity deep venous thrombosis. *Am J*  
329 *Surg* 2006; 192: 782–788.
- 330 11. Watson LI, Armon MP. Thrombolysis for acute deep vein thrombosis. *Cochrane Database Syst*  
331 *Rev* 2004: CD002783.
- 332 12. Okrent D, Messersmith R, Buckman J. Transcatheter fibrinolytic therapy and angioplasty for left  
333 iliofemoral venous thrombosis. *J Vasc Interv Radiol* 1991; 2: 195–197.
- 334 13. Mahmoud O, Sihvo E, Räsänen J, et al. Treatment of the Paget-Schroetter syndrome with a three  
335 stage approach including thoracoscopic rib resection at the second stage. *J Vasc Surg* 2017, in  
336 press.
- 337 14. Kim HS, Patra A, Paxton PE, et al. Catheter-Directed Thrombolysis with Percutaneous Rheolytic  
338 Thrombectomy Versus Thrombolysis Alone in Upper and Lower Extremity Deep Vein  
339 Thrombosis. *Cardiovasc Intervent Radiol* 2006; 29: 1003–1007.
- 340 15. Illig KA, Doyle AJ. A comprehensive review of Paget-Schroetter syndrome. *J Vasc Surg* 2010; 5:  
341 1538–1547.
- 342 16. Doyle A, Wolford HY, Davies MG, et al. Management of effort thrombosis of the subclavian vein:  
343 today's treatment. *Ann Vasc Surg* 2007; 21: 723–729.
- 344 17. Ruiz-Bailén M, Brea-Salvago JF, de Hoyos EA, et al. Post-thrombolysis intracerebral hemorrhage:  
345 data from the Spanish Register ARIAM. *Crit Care Med* 2005; 33: 1829–1838.
- 346 18. Mewissen MW, Seabrook GR, Meissner MH, et al. Catheter-directed thrombolysis for lower  
347 extremity deep venous thrombosis: report of a national multicenter registry. *Radiology* 1999;  
348 211: 39–49.
- 349 19. Prins MH, Hutten BA, Koopman MM, et al. Long-term treatment of venous thromboembolic  
350 disease. *Thromb Haemost* 1999; 82: 892–898.

- 351 20. Schweizer J, Kirch W, Koch R, et al. Short- and long-term results after thrombolytic treatment of  
352 deep venous thrombosis. *J Am Coll Cardiol* 2000; 36: 1336–1343.
- 353 21. Semba CP, Dake MD. Iliofemoral deep venous thrombosis: aggressive therapy with catheter-  
354 directed thrombolysis. *Radiology* 1994; 191: 487–494.
- 355 22. Verhaeghe R, Stockx L, Lacroix H, et al. Catheter-directed lysis of iliofemoral vein thrombosis  
356 with use of rt-PA. *Eur Radiol* 1997; 7: 996–1001.
- 357 23. Ohtsuka T, Wolf RK and Dunsker SB. Port access first rib resection. *Surg Endosc* 1999; 13: 940–  
358 942.
- 359 24. Urschel HC Jr, Patel AN. Paget Schroetter syndrome therapy: failure of intravenous stents. *Ann*  
360 *Thorac Surg* 2003; 75: 1693–1696.
- 361

362

363 **FIGURE LEGENDS**

364 **Figure 1.** Patient flow.

365 **Figure 2. A** Phlebography showing thrombosis of the axillo-subclavian segment. **B** Post-thrombolysis  
366 control phlebography with patent veins and a partial success (less than 50% thrombus remaining). **C**  
367 Angioplasty after local thrombolysis using PMT. **D** Mid-term phlebography with open veins.

368

369

ACCEPTED MANUSCRIPT

**Table I.** Demographic data and initial symptoms before thrombolysis. There were no significant differences between the groups.

	PMT	CDT
Number of patients	19	24
Age (median, IQR)	26 (17-54)	31 (23-49)
Male: Female	9:10	12:12
Effort history* n (%)	13 (68%)	21 (88%)
Thrombophilia n (%)	3 (16%)	4 (17%)
Family history n (%)	4 (21%)	5 (21%)
RT UL n (%)	12 (63%)	17 (71%)
LT UL n (%)	7 (37%)	7 (29%)
Arm pain	17 (90%)	22 (92%)
Arm swelling	19 (100%)	24 (100%)
Arm numbness	11 (58%)	14 (58%)
Arm weakness	2 (11%)	2 (8%)
Neck swelling	1 (5%)	2 (8%)
Dilated neck veins	1 (5%)	7 (29%)
Positive provocation test	15 (79%)	20 (83%)
Pulmonary embolism	0 (0%)	1 (4%)
Paresthesia	4 (21%)	7 (29%)

PMT, pharmacomechanical thrombolysis; CDT, catheter-directed thrombolysis;

RT UL, right upper limb; LT UL, left upper limb.

\*Heavy upper limb exercise as a probable etiology.

Values are presented as No (%) unless otherwise indicated.

**Table II.** Phlebographic results.

<b>Degree of success of lysis (11,12)</b>	<b>PMT</b>	<b>CDT</b>	<b>p-value</b>
Complete lysis >99%	17 (90%)	19 (79%)	NS
Partial lysis (50%-99%)	2 (11%)	3 (13%)	NS
Unsatisfactory lysis <50% / change line of treatment	0 (0%)	2* (8%)	NS
<b>Degree of residual stenosis after thrombolysis</b>			
No lesion/stenosis <20%	13 (68%)	7 (29%)	0.010
Moderate stenosis 20%-49%	4 (21%)	5 (21%)	NS
Significant stenosis 50%/>50%	2 (11%)	10 (42%)	0.024
Occlusion	0 (0%)	2 (8%)	NS
<b>Pre-rib resection phlebographic findings **</b>			
No lesion/stenosis <20%	14 (74%)	8 (33%)	0.009
Moderate stenosis 20%-49%	2 (11%)	3 (13%)	NS
Significant stenosis 50%/>50%	3 (16%)	10 (42%)	NS
Occlusion	0 (0%)	3 (13%)	NS
<b>Re-thrombosis (pre rib resection)</b>	0 (0%)	0 (0%)	NS

NS, not significant; CDT, catheter directed thrombolysis; PMT, pharmacomechanical thrombectomy.

\*Changed to Trellis

\*\*This phlebography was done prior to thoracoscopic first rib resection (median time from thrombolysis to rib resection was 90.5 days, range 10–450 days).

#P-value=0.011, tested with chi-square for the combined numbers of significant and occlusions.

Values are reported as No. (%).

**Table III.** Treatment parameters, use of resources and costs in patients treated with PMT and/or CDT.

Treatment group	PMT	CDT	P value
Infusion time	0.33 h (0.17–0.58 h)	21.12 h (16.11–47.25 h)	< 0.00001
Total alteplase dose	6 mg (6–15 mg)	32 mg (20–55.3 mg)	< 0.00001
Intermediate care unit (h)	0	48 h (48–72)	< 0.00001
Number of phlebographies	1	2 (2–3)	< 0.00001
Angiography and/or Interventional suite costs	4499.00 € (3782€–5120€)	6985.50 € (8564€–6100€)	< 0.00001
Length of hospital admission	3 days (1–8 days)	6 days (3–15 days)	0.0061
Total hospital costs	5975.00 € (4763€–7395€)	11476.00 € (8468€–17467€)	< 0.00001

*CDT*, catheter-directed thrombolysis; *PMT*, pharmacomechanical thrombectomy.

The variables are expressed as medians (range).

**Table IV.** One-year vein patency and symptom status. No significant differences were seen.

	<b>PMT N = 19</b>	<b>CDT N = 24</b>	<b>p-value</b>
<b>Treatment method</b>			
Good flow no lesion/stenosis <20%	17 (90%)	18 (75%)	NS
Moderate stenosis (20%-49%)	1 (5%)	2 (8%)	NS
Significant stenosis (50%/ > 50%)	0 (0%)	1 (4%)	NS
Occlusion	1 (5%)	3 (13%)	NS
Any PTA during the FU	10 (53%)	11 (46%)	NS
Good results of PTA	8/10 (80%)	10/11 (91%)	NS
Recoil/Failed PTA	2/10 (20%)	1/11 (9%)	NS
Patency in the final phlebography	18 (95%)	22 (92%)	NS
<b>Symptoms assessment</b>			
No symptoms	15 (79%)	16 (67%)	NS
Pain during rest	0 (0%)	0 (0%)	NS
Pain during exercise	1 mild (5%)	3 mild (13%)	NS
Swelling	1 mild (5%)	2 mild (8%)	NS
Numbness	2 mild (11%)	3 mild (13%)	NS
Paresthesia	0 (0%)	0 (0%)	NS
Weakness	0 (0%)	0 (0%)	NS
Complete improvement	15 (79 %)	16 (67%)	NS
No improvement	0 (0%)	0 (0%)	NS


Overall improvement	19 (100%)	24 (100%)	NS
---------------------	-----------	-----------	----

---

NS, not significant; CDT, catheter directed thrombolysis; PMT, pharmacomechanical thrombectomy.


Values are reported as No. (%).






UEDVT = upper extremity deep venous thrombosis; CDT = catheter-directed thrombolysis; LMWH = low molecular weight heparin; PMT = pharmacomechanical thrombolysis


Figure 2.




(A)



(B)



(C)



(D)