

## Higher step count is associated with greater bone mass and strength in women but not in men

Sanna Tolonen<sup>1,2,3</sup>, Harri Sievänen<sup>4</sup>, Mirja Hirvensalo<sup>5</sup>, Marika Laaksonen<sup>1</sup>, Vera Mikkilä<sup>1,6</sup>, Kristiina Pälve<sup>6</sup>, Terho Lehtimäki<sup>2</sup>, Olli Raitakari<sup>7</sup> & Mika Kähönen<sup>3</sup>

Department of Food and Environmental Sciences, University of Helsinki, Finland<sup>1</sup>

Department of Clinical Chemistry, Fimlab Laboratories and Faculty of Medicine and Life Sciences, University of Tampere, Tampere, Finland<sup>2</sup>

Department of Clinical Physiology, Tampere University Hospital and Faculty of Medicine and Life Sciences, University of Tampere, Tampere, Finland<sup>3</sup>

The UKK Institute for Health Promotion Research, Tampere, Finland<sup>4</sup>

Faculty of Sport and Health Sciences, University of Jyväskylä<sup>5</sup>

Research Centre of Applied and Preventive Cardiovascular Medicine, University of Turku and Heart Center, Turku University Hospital, Turku, Finland<sup>6</sup>

Research Centre of Applied and Preventive Cardiovascular Medicine, University of Turku and Department of Clinical Physiology and Nuclear Medicine, Turku University Hospital, Turku, Finland<sup>7</sup>

All authors of this manuscript declare that they have no conflict of interest.

### Summary

In this cross-sectional study, peripheral bone traits were examined relative to total daily steps measured with pedometer. Higher number of steps was associated with greater bone values at the calcaneus and tibia in women, but not in men. In women, dose-dependent associations at the radius were congruent with the weight-bearing bones.

### Abstract

*Introduction* Habitual physical activity measured as daily steps may contribute to bone density and strength at the calcaneus and other weight-bearing bones.

*Methods* Subgroups of 705-837 women and 480-615 men aged 31-46 years from the Cardiovascular Risk in Young Finns Study participated in the present study. Participants were instructed to use pedometer for one week and the total daily steps, divided into tertiles, were evaluated relative to quantitative ultrasound-measured bone traits at the calcaneus and peripheral quantitative computed tomography-measured bone traits at the tibia and radius. Analysis of covariance was used to examine the between-group differences.

*Results* In women, significant dose-dependent between-group differences were found in the weight-bearing bones and in non-weight-bearing radius. The differences in broadband ultrasound attenuation and speed of sound at the calcaneus were 3.8% and 0.5% greater in women within the highest tertile of daily steps compared to the lowest tertile (p-values for trend  $\leq 0.04$ ). In tibia, women in the highest tertile ( $> 8765$  steps/day) had on average 1-5.4% greater bone cross-sectional area, bone mineral content (BMC), trabecular density and bone strength index at the distal site and 1.6-2.7% greater bone areas, BMC and strength strain index (SSI) at the shaft compared to women with less daily steps (p-values for trend  $\leq 0.02$ ). Similarly, in radius, BMC and BSI at the distal site, and bone cross-sectional areas, BMC and SSI at the shaft were 1.7-3.4% greater in women within the highest tertile of daily steps compared to their peers (p-values for trend  $\leq 0.04$ ). In men, the differences in calcaneal, tibial and radial bone traits were mainly non-significant between the tertiles of daily steps.

*Conclusion* Observed significant positive associations between daily steps and various bone traits at the calcaneus, tibia and radius in women suggest that habitual physical activity may benefit skeletal health in adulthood.

**Keywords** peripheral quantitative computed tomography (pQCT), quantitative ultrasound (QUS), pedometer, step counts, physical activity.

## Introduction

Bone structure adapts to regular exercise that includes greater loading than induced by habitual daily activities. This is particularly well documented in the lower and upper extremities of athletes participating in moderate-high impact sports that include e.g. vertical jumps, rapid turns and stops or repetitive impact loading [1-4]. Among general population, adults are recommended to exercise 3-5 times per week endurance sports like running and brisk walking to load their skeletal structure [5-6]. Also a combination of strength training for a few times a week and a sufficient number of jumps daily are recommended as a part of bone enhancing exercise.

Walking is the most common type of physical activity in humans. Indeed, daily steps have been suggested to increase bone mass and strength. In two studies on elderly Japanese women, daily walking steps were positively associated with the ultrasound parameters of calcaneus and negatively with bone resorption [7-8]. In another study on 50-80 year old women and men, pedometer determined steps were positively associated with areal bone mineral density (aBMD) at the hip especially in those over 65 years of age [9]. Also, in a Finnish exercise intervention trial on perimenopausal women, weekly endurance training including mainly brisk walking resulted in better maintenance of femoral neck aBMD [10]. Additionally, in a study comparing four different physical activity methods and two bone strength indices in young girls aged 8-13 years, pedometer steps were significantly associated with bone strength index (BSI) at the distal sites of femur and tibia [11]. In another study of young boys and girls aged 7-9 years, estimates of cortical bone and strength strain index (SSI) at the tibial mid-shaft were modestly higher in children who were in the active group according to daily steps and fitness tests compared to children classified to the inactive group [12]. So far the longest follow-up of daily steps was collected in a study of Japanese women and men aged 65-84 years during a period of five years [13]. These researchers concluded that it would be beneficial to

take at least 7000-8000 steps per day and noted also that habitual physical activity performed at higher intensity than 3 metabolic equivalents (MET) was associated with calcaneal bone health.

In the present study, we examined if daily steps measured with pedometer were associated with various bone traits determined at calcaneus, tibia and radius. To the best of our knowledge, there are no previous studies that have investigated this association in a large sample of adults aged 31-46 years old.

## **Methods**

### *Participants*

Participants in the present study, aged 31-46 years old, were participants of the Cardiovascular Risk in Young Finns study which is an ongoing follow-up study of six age-cohorts born in 1962, 1965, 1968, 1971, 1974 and 1977 (Table 1.). At baseline in 1980, the study subjects were randomly chosen from the Finnish national register in the areas of five university cities Helsinki, Kuopio, Oulu, Tampere and Turku (n=3596 participated, 83% of those invited). Later follow-up studies have been performed in 1983, 1986, 1989, 2001, 2007 and 2012 with the participation rates of 60-80% of the baseline cohort. Those who were lost in the follow-up have been described elsewhere [14-15]. The study participants gave written informed consent and the study protocol was reviewed and approved by the ethics committees of the participating universities.

### *Assessment of steps*

In 2007, the pedometer study was conducted among 1876 individuals. Daily steps were gathered with the pedometer (Walking style One, HJ-152R-E, Omron Healthcare Co., LTD. Kyoto, Japan). Pedometers were instructed to be attached on the waistband or belt in the same position for 7 consecutive days and to be removed only while bathing or swimming [16]. Participants were asked to continue their habitual physical activity during the one week monitoring period. At the end of the day, the number of total steps was recorded in pedometer logs. Participants who had at least 4 monitoring days with at least 8 hours pedometer wear time were included in all the analyses (n=1853, 98.8% of those who completed the pedometer study). Sickness or injury status, exceptional step count recorded as an atypical day, or problems with pedometer use, were imputed by the mean of other days. Pedometer accuracy for measuring total steps over 1000 meters varied between 0.1-1.3 % in different walking situations [16] and the Spearman's rank correlation coefficient with steps measured by the accelerometer (GT1M, ActiGraph, Pensacola, FL, USA) was 0.966 (p<0.001) [17].

### *Bone measurements*

In 2008, all those who had participated the Cardiovascular Risk in Young Finns baseline study, excluding those who had died (n=77), moved abroad or forbid any further invitations, were invited to the peripheral quantitative computed tomography (pQCT) and quantitative ultrasound (QUS) bone measurements

(n=3386, 94.2% of the baseline cohort). Altogether, 1884 individuals (55.6% of those who were invited) from Turku, Helsinki, Tampere, Kuopio and Oulu attended the pQCT measurements between February and December 2008 [18]. The corresponding number of the QUS measurements was 1953 (57.7% of the invited).

Both distal and diaphyseal sites of two functionally different bones, non-weight-bearing radius and weight-bearing tibia, were measured with pQCT (XCT 2000R, Stratec Medizintechnik GmbH, Pforzheim, Germany). Pregnant women were excluded from the measurements. Radius was measured on the non-writing hand and tibia on the left leg if no contraindication emerged, e.g. previous fracture of the given site. Distal sites were 4 % for radius and 5 % for tibia relative to the distal cortical endplate. The diaphyseal sites were 30 % for both bones. Approximately 98% of the pQCT measurements were successfully scanned and analyzed. The main reasons for the failed pQCT investigation were missing scan images, inability to fit leg into the pQCT gantry, inaccurate positioning of the forearm or leg, or movement artefacts during scanning. For the distal sites, the contour and peel modes 2 were used according to the previously described procedures [19]. For the shafts, a density threshold of 710 mg/cm<sup>3</sup> was applied to separate the cortical bone from other tissues. The pQCT measurements yielded the following bone traits: bone mineral content (BMC, mg), trabecular and cortical bone mineral densities (mg/cm<sup>3</sup>), and total and cortical bone cross-sectional areas (mm<sup>2</sup>). Additionally, three bone strength indices were calculated: SSI (mm<sup>3</sup>), BSI (g<sup>2</sup>/cm<sup>4</sup>) and cortical strength index (CSI). The conventional SSI was calculated as a density-weighted polar section modulus. It predicts the torsional bone strength and was determined for the diaphysis only. For the distal sites, BSI reflecting the compressive bone strength was calculated as a product of squared total bone mineral density and total cross-sectional area (total density<sup>2</sup> x total area) [1]. The value of CSI corresponded to the ratio of cortical bone area and total bone area (cortical area/total area).

Additionally, the QUS technique (Sahara Clinical Bone Sonometer, Hologic Inc, Waltham, MA, USA) was used to measure the speed of sound (SOS, m/s) and broadband ultrasound attenuation (BUA, dB/MHz) at the calcaneus mainly from the left heel. Totally, ~23 % of the QUS measurements were unsuccessful since there were problems in the propagation of ultrasound. The precision of the pQCT and QUS methods were evaluated in a subgroup (n=39) with in vivo coefficients of variation (%), which varied between 0.3 % and 4.8 % [18].

### *Clinical measurements and lifestyle*

In 2007, body weight, height and body mass index (kg/m<sup>2</sup>) were determined with standard methods [20]. On the same year, energy and nutrient intakes were assessed with a 131-item food frequency questionnaire developed by the Finnish National Institute for Health and Welfare [21]. The quantities of nutrients were based on the Finnish Food Composition database, Fineli® [22]. Serum 25-hydroxy vitamin D (later calcidiol) levels (in nmol/l) of the study subjects were determined using radioimmunoassay (DiaSorin, Stillwater, Minnesota). Information on alcohol consumption (drinks during the previous week), smoking habits (daily or less frequently) and oral corticosteroid medication for at least one month (yes/no) were collected with self-administered questionnaires in 2007-2008. Maximal oxygen consumption (VO<sub>2</sub>max) and work rate (WRmax) were obtained from the exercise tests performed on electronically braked cycle ergometers during the years 2007-2009 in Tampere and Turku (Lode Corival 906900, Lode BV, Groningen, Netherlands). During the tests, 12-lead electrocardiography was recorded (Corina ECG amplifier and CardioSoft acquisition software ver. 4.2, GE Medical Systems, Freiburg, Germany) and breath-by-breath

measurements performed with ventilator gas analyzers (V-max 29C, SensorMedics, Yorba Linda, CA, USA and Jaeger Oxycon Pro, VIASYS Healthcare GmbH, Hoechberg, Germany) .  $VO_2$ max was determined as the highest oxygen uptake during the last 30-second averaged interval and  $WR_{max}$  was calculated by adding the work during the last full minute interval to the fraction of the work performed over the final, interrupted interval. These tests and measurements have been described earlier in more detail [23].

Physical activity index in 2007 was calculated based on a self-report questionnaire which included items on intensity of physical activity, frequency of vigorous physical activity, hours spent on vigorous physical activity, average duration of a physical activity session and participation in organized exercise.

### *Statistical analyses*

For the statistical dose-dependence association analyses, participants were divided into tertiles by their mean daily steps. The following values were used as the cut-off values: < 6317 and > 8765 in women (min 1145, max 22879) and < 5514 and > 7962 in men (min 1548, max 19351). After the group classifications, associations with the characteristic variables were analyzed with the analysis of variance (ANOVA) or Brown-Forsythe-test for continuous and Chi square-test for categorical variables. In the main analyses, pQCT and QUS bone parameters were compared between the daily step tertiles with the analysis of covariance using age, height, weight, serum calcidiol level and physical activity index as covariates. We also tested the linear trends with polynomial contrast because of an assumption of the dose-responsive effect of daily steps on bone traits. Analyses were done separately for women and men because of different endocrine profiles. Statistical analyses were carried out with IBM SPSS Statistics 24. P-values  $\leq 0.05$  were considered statistically significant.

## **Results**

The background characteristics of the study groups divided into tertiles by mean daily steps are shown in Table 1. In women, differences were seen in body height, weight and body mass index (BMI) (p-values  $\leq 0.02$ ). Body weight and BMI were the lowest among those women who had taken more than 8765 daily steps. Women in the highest tertile of steps had also higher mean serum calcidiol level, physical activity score and maximal work rate value compared to less active women (p-values  $\leq 0.002$ ). On the contrary, they smoked less than the other women (p-value 0.02). In men, similar differences as in women were seen in body weight, BMI, physical activity score and maximal work rate value (Table 1, p-values  $\leq 0.01$ ). Mean serum calcidiol level was, instead, the highest among men in the middle tertile of daily steps (p-value  $< 0.001$ ). In addition, the  $VO_2$ max value reflecting physical endurance capacity was the highest among men in the third tertile of daily steps (p-value 0.001). On average, women had taken more daily steps than men during the monitoring week (7827 vs. 7063 steps/day, p-value  $< 0.001$ ).

In women, the differences in calcaneal BUA and SOS values were 3.8% and 0.5% greater in women in the highest tertile of daily steps compared to the lowest tertile (Table 2, p-values for trend  $\leq 0.04$ ). Similarly in distal tibia, women with over 8765 daily steps had 1.1% larger total cross-sectional area, 3.2% higher BMC, 3.7% higher trabecular density and 5.4% greater BSI than women in the lowest tertile of steps (Table 3, p-values for trend  $\leq 0.02$ ). In tibial shaft, bone area was 1.7%, cortical bone area 1.6%, BMC 1.7% and SSI 2.7%

greater among women in the highest tertile of daily steps compared to the other women (Table 3, p-values for trend  $\leq 0.02$ ). In radius, women in the highest tertile of daily steps had 2.2% higher BMC and 3.4% greater BSI at the distal site compared to women in the lowest tertile (Table 4, p-values for trend  $\leq 0.04$ ). In radial shaft, total cross-sectional area was 2.3%, cortical area 1.7%, BMC 1.7% and SSI 2.4% greater among women in the highest tertile of daily steps than in the lowest or middle tertile of steps (Table 4, p-values for trend  $\leq 0.03$ ).

In men, the differences in calcaneal and radial bone traits were non-significant between the tertiles of daily steps (Table 2 and 4). In distal tibia, total cross-sectional area and BMC were the highest in men within the middle tertile of steps (Table 3, p-values for trend  $\leq 0.03$ ). In tibial shaft, bone area and SSI were, however, the greatest in men with least daily steps (Table 3, p-values for trend  $\leq 0.04$ ). The differences between the highest and the lowest values ranged from 0.8% to 2%.

## Discussion

In the present study, women aged 31-46 years whose mean daily steps exceeded 8765 had larger total and cortical bone cross-sectional areas, higher bone mineral mass and density, and greater proximal BSI and diaphyseal SSI at the weight-bearing tibia compared to their peers. Also, the calcaneal ultrasound parameters were higher among these more walking women. Similarly, in the non-weight bearing radius, greater number of steps was associated with larger bone areas, higher bone mineral mass and greater strength indices in women. In men, on the contrary, the results were non-significant at the calcaneus and radius. In tibia, some statistically significant but conflicting associations of daily steps with bone areas, BMC and SSI were found in men. These results may reflect different physical activity habits between women and men in the present population. On an average, women had taken more daily steps than men during the monitoring week (difference 764 steps/day). Thus, women may have also exercised more frequently so-called Nordic walking which was a popular leisure time activity at that time in Finland. In Nordic walking, sticks are used along with walking which makes it a good exercise for the arms and upper body too. This may explain the higher values in women's radii with the highest number of daily steps compared to the other less walking women. In men, the associations between daily steps and studied bone traits were mainly non-significant why they should be interpreted cautiously. It may be, for instance, that men in the lowest tertile of daily steps took their steps at higher intensity which then contributed to slightly higher total area and SSI at the tibial shaft compared to the other men with more steps. This explanation cannot be, however, verified with pedometer which was used here to assess usual ambulatory activity.

Although, the loading effect of habitual daily steps on calcaneal, tibial and radial bone traits seemed to be relatively small in women, it may have some clinical relevance. In a study where walking was the primary activity, four hours per week of walking was associated with 41% lower hip fracture risk in postmenopausal women compared to peers with less than one hour of walking per week [24]. Also the walking pace seemed to influence the risk of hip fractures among these women, so that those who walked briskly were in the lowest fracture risk. However, in the meta-analysis of eight randomized controlled trials including postmenopausal women, no effect of walking interventions on lumbar spine BMD was found while the effect on femoral neck was modest [25]. The researchers concluded that the effect of walking on bone density as a sole exercise may be too small that it would reduce the fracture risk in these older women.

Previously, the positive association between daily steps and bone traits has been found in older women and men [7-9, 13] and in children [11-12]. However, in a study of young students, no correlation was found between daily steps and ultrasound bone parameters [26]. Scheffler and others discussed that the reason behind this was too a homogenous population and that students may have been more health-conscious than the average population. It is also possible that the associations between different bone traits and habitual physical activity measured as total steps may vary with age as the total amount of physical activity may decrease or increase in different phases of life. The mean number of steps in the aforementioned studies varied from 6896 to 15495 steps per day which is almost the same range as recommended for relatively healthy, younger adults [27].

We also found that daily steps were inversely associated with body weight in both genders and positively with serum calcidiol levels in women. Participants in the highest tertile of daily steps had on average 6.8-7.4 % lower body weight and also 4.6-5.7% lower BMI than their study peers in the lowest tertile. In women, mean serum calcidiol level was on average 9.3% higher within the highest tertile of daily steps compared to the lowest tertile. In addition, the proportion of participants who smoked daily was the lowest in the highest tertile of daily steps in both genders. Participants with more daily steps had also higher physical activity score in general and better maximal work output in the cycle ergometer exercise test than their less walking counterparts. These health-related differences between the daily step groups may have also influenced the studied bone traits. For instance, vitamin D has many known effects on the physiological pathways relevant to bone mineral homeostasis and it is also a common supplement along with calcium in patients with osteoporosis [28]. Also the differences in mean body weight and general physical activity may have confounded the results to some extent. In the earlier study from the Young Finns study Cohort, overweight in childhood was associated with greater bone size later in adulthood [29]. Also obesity in adulthood seemed to decrease the risk of low trabecular bone density at the distal sites of radius and tibia in the Young Finns cohort [18].

When considering the strengths of the present study, the observations were drawn from a large cohort including both women and men aged 31-46 years. To our knowledge, no similar studies in this age group have been done previously. In addition, three different peripheral bones, calcaneus, tibia and radius were studied relative to objectively measured step counts which make our congruent results more reliable. Additionally, many health-related factors known to affect bone mass were examined in the present population and considered when interpreting the results.

There are some limitations in relation to the studied associations of step counts and peripheral bone parameters. The present study was a cross-sectional in design and therefore we do not know the outcomes between accumulated lifetime steps and different bone traits. However, prospective data on physical activity estimated with a self-report questionnaire from the present population was earlier shown to predict the level of physical activity in adulthood since childhood [30]. This same physical activity index was also used in our previous pQCT study where frequent habitual physical activity at the age of 9-18-years was associated with higher tibial bone size and strength indices in adulthood [31]. One limitation is also that steps were collected for a quite short period of time and can therefore be confounded by varying weather or participants' health conditions. As earlier reported, the main reasons for non-participation and interruption of pedometer use were lost or broken pedometer (n=75), illness (n=30) and other reasons like untypical day (n=22) [16]. In addition, subjects who participated the pedometer sub-study were more likely to have a higher socioeconomic position than all those who participated in the examinations of follow-up 2007. This may have resulted in higher number of reported steps compared to the general Finnish

population. Pedometer itself has also some shortcomings since it cannot measure accurately the intensity of training or other types of exercises than locomotion e.g. cycling, swimming, weight training, yoga, skiing and similar exercises. These exercises can, however, be useful in promoting both general and bone health through developing the muscle strength. Despite these limitations, the objectively measured step data provided a relevant estimate of habitual physical activity in this population.

## Conclusion

In conclusion, women aged 31-46 years who had taken more than 8765 daily steps had on average 1-5.4 % greater bone cross-sectional areas, bone mineral content, trabecular density, and strength indices at the weight-bearing tibia compared to their study peers. Also the calcaneal ultrasound parameters were 0.5-3.8 % greater among more walking women. In non-weight bearing radius, the results were similar as in the weight-bearing bones. In men, the associations of daily steps and peripheral bone traits were mainly non-significant. These results suggest that habitual daily steps may preserve bone mass and strength in women, but not in men. Obviously, more studies are needed to confirm these results.

## Acknowledgements

The Young Finns Study has been financially supported by the Academy of Finland: grants 286284, 134309 (Eye), 126925, 121584, 124282, 129378 (Salve), 117787 (Gendi) (ST), and 41071 (Skidi); the Social Insurance Institution of Finland; Competitive State Research Financing of the Expert Responsibility area of Kuopio, Tampere and Turku University Hospitals (grant X51001); Juho Vainio Foundation (ST); Paavo Nurmi Foundation; Finnish Foundation for Cardiovascular Research ; Finnish Cultural Foundation (ST); Tampere Tuberculosis Foundation; Emil Aaltonen Foundation; Yrjö Jahnesson Foundation; Signe and Ane Gyllenberg Foundation and Diabetes Research Foundation of Finnish Diabetes Association.

## References

1. Kontulainen S, Sievänen H, Kannus P, Pasanen M, Vuori I. Effect of long-term impact-loading on mass, size, and estimated strength of humerus and radius of female racquet-sports players: a peripheral quantitative computed tomography study between young and old starters and controls. *J Bone Miner Res* 2003; 18:352-359.
2. Nikander R, Sievänen H, Uusi-Rasi K, Heinonen A, Kannus P. Loading modalities and bone structures at nonweight-bearing upper extremity and weight-bearing lower extremity: a pQCT study of adult female athletes. *Bone* 2006; 39:886-894.

3. Nikander R, Kannus P, Dastidar P, Hannula M, Harrison L, Cervinka T, Narra NG, Aktour R, Arola T, Eskola H, Soimakallio S, Heinonen A, Hyttinen J, Sievänen H. Targeted exercises against hip fragility. *Osteoporos Int* 2009; 20:1321-1328.
4. Nikander R, Kannus P, Rantalainen T, Uusi-Rasi K, Heinonen A, Sievänen H. Cross-sectional geometry of weight-bearing tibia in female athletes subjected to different exercise loadings. *Osteoporos Int* 2010; 21:1687-1694.
5. Kohrt WM, Bloomfield SA, Little KD, Nelson ME, Yingling VR; American College of Sports Medicine. American College of Sports Medicine Position Stand: physical activity and bone health. *Med Sci Sports Exerc* 2004; 36:1985-1996.
6. Nikander R, Lepola V, Karinkanta S, Sievänen H. Exercising briskly 3 to 4 hours a week: Does it diminish hip fractures by one third? (only abstract in English) *Suomen Lääkärilehti* 22/2008 vsk 63: 2033-2040.
7. Kitagawa J, Omasu F, Nakahara Y. Effect of daily walking steps on ultrasound parameters of the calcaneus in elderly Japanese women. *Osteoporos Int* 2003; 14:219-224.
8. Kitagawa J, Nakahara Y. Associations of daily walking steps with calcaneal ultrasound parameters and a bone resorption marker in elderly Japanese women. *J Physiol Anthropol* 2008; 27: 295-300.
9. Foley S, Quinn S, Jones G. Pedometer determined ambulatory activity and bone mass: a population-based longitudinal study in older adults. *Osteoporos Int* 2010; 21:1809-1816.
10. Heinonen A, Oja P, Sievänen H, Pasanen M, Vuori I. Effect of two training regimens on bone mineral density in healthy perimenopausal women: a randomized controlled trial. *J Bone Miner Res* 1998; 13:483-490.
11. Farr JN, Lee VR, Blew RM, Lohman TG, Going SB. Quantifying bone-relevant activity and its relation to bone strength in girls. *Med Sci Sports Exerc* 2011; 43:476-483.
12. Duckham RL, Rantalainen T, Ducher G, Hill B, Telford RD, Telford RM, Daly RM. Effects of habitual physical activity and fitness on tibial cortical bone mass, structure and mass distribution in pre-pubertal boys and girls: The Look Study. *Calcif Tissue Int* 2016; 99: 56-65.
13. Shephard RJ, Park H, Park S, Aoyagi Y. Objective Longitudinal Measures of Physical Activity and Bone Health in Older Japanese: the Nakanojo Study. *J Am Geriatr Soc* 2017; 65:800-807.
14. Juonala M, Viikari JS, Hutri-Kähönen N, Pietikäinen M, Jokinen E, Taittonen L, Marniemi J, Rönnekaa T, Raitakari OT. The 21-year follow-up of the Cardiovascular Risk in Young Finns Study: risk factor levels, secular trends and east-west difference. *J Intern Med* 2004; 255:457-468.
15. Raitakari OT, Juonala M, Rönnekaa T, Keltikangas-Järvinen L, Räsänen L, Pietikäinen M, Hutri-Kähönen N, Taittonen L, Jokinen E, Marniemi J, Jula A, Telama R, Kähönen M, Lehtimäki T, Åkerblom HK, Viikari JS. Cohort profile: the Cardiovascular Risk in Young Finns Study. *Int J Epidemiol* 2008; 37:1220-1226.
16. Hirvensalo M, Telama R, Schmidt MD, Tammelin TH, Xiaolin Yang, Magnussen CG, Viikari JS, Raitakari OT. Daily steps among Finnish adults: variation by age, sex, and socioeconomic position. *Scand J Public Health* 2011; 39:669-677.
17. Mansikkaniemi K, Juonala M, Taimela S, Hirvensalo M, Telama R, Huupponen R, Saarikoski L, Hurme M, Mallat Z, Benessiano J, Jula A, Taittonen L, Marniemi J, Kähönen M, Lehtimäki T, Rönnekaa T, Viikari J, Raitakari OT. Cross-sectional associations between physical activity and selected coronary heart disease risk factors in young adults. The Cardiovascular Risk in Young Finns Study. *Ann Med* 2012; 44:733-744.
18. Laaksonen MML, Sievänen H, Tolonen S, Mikkilä V, Räsänen L, Viikari J, Lehtimäki T, Kähönen M, Raitakari OT, The Cardiovascular Risk in Young Finns Study Group. Determinants of bone strength

- and fracture incidence in adult Finns: Cardiovascular Risk in Young Finns Study (the GENDI pQCT study). *Arch Osteoporos* 2010; 5: 119-130.
19. Sievänen H, Koskue V, Rauhio A, Kannus P, Heinonen A, Vuori I. Peripheral quantitative computed tomography in human long bones: evaluation of in vitro and in vivo precision. *J Bone Miner Res* 1998; 13:871-882.
  20. Raiko JR, Viikari JS, Ilmanen A, Hutri-Kähönen N, Taittonen L, Jokinen E, Pietikäinen M, Jula A, Loo BM, Marniemi J, Lehtimäki T, Kähönen M, Rönnemaa T, Raitakari OT, Juonala M. Follow-ups of the Cardiovascular Risk in Young Finns Study in 2001 and 2007: levels and 6-year changes in risk factors. *J Intern Med* 2010; 267:370-384.
  21. Paalanen L, Männistö S, Virtanen MJ, Knekt P, Räsänen L, Montonen J, Pietinen P. Validity of a food frequency questionnaire varied by age and body mass index. *J Clin Epidemiol* 2006; 59: 994-1001.
  22. National Institute for Health and Welfare, Nutrition Unit. Fineli. Finnish food composition database. Release, 10; 2009. <http://www.fineli.fi>. Helsinki.
  23. Hulkkonen J, Aatola H, Pälve K, Lehtimäki T, Hutri-Kähönen N, Viikari JS, Raitakari OT, Kähönen M. Determinants of exercise peak arterial blood pressure, circulatory power, and exercise cardiac power in a population based sample of Finnish male and female aged 30 to 47 years: the Cardiovascular Risk in Young Finns Study. *BMC Cardiovasc Disord*. 2014; 14:35.
  24. Feskanich D, Willett W, Colditz G. Walking and leisure-time activity and risk of hip fracture in postmenopausal women. *JAMA* 2002; 288:2300-2306.
  25. Martyn-St James M, Carroll S. Meta-analysis of walking for preservation of bone mineral density in postmenopausal women. *Bone* 2008; 43:521-531.
  26. Scheffler C, Gniosdorj B, Staub K, Rühli F. Skeletal robustness and bone strength as measured by anthropometry and ultrasonography as a function of physical activity in young adults. *Am J Hum Biol* 2014; 26:215-220.
  27. Tudor-Locke CE, Myers AM. Methodological considerations for researchers and practitioners using pedometers to measure physical (ambulatory) activity. *Res Q Exerc Sport* 2001; 72:1-12.
  28. Bhattoa HP, Konstantynowicz J, Laszcz N, Wojcik M, Pludowski P. Vitamin D: Musculoskeletal health. *Rev Endocr Metab Disord* 2016. doi:10.1007/s11154-016-9404-x.
  29. Uusi-Rasi K, Laaksonen M, Mikkilä V, Tolonen S, Raitakari OT, Viikari J, Lehtimäki T, Kähönen M, Sievänen H. Overweight in childhood and bone density and size in adulthood. *Osteoporos Int* 2012; 23:1453-1461.
  30. Telama R, Yang X, Viikari J, Välimäki I, Wanne O, Raitakari O. Physical activity from childhood to adulthood: a 21-year tracking study. *Am J Prev Med* 2005; 28:267-273.
  31. Tolonen S, Sievänen H, Mikkilä V, Telama R, Oikonen M, Laaksonen M, Viikari J, Kähönen M, Raitakari OT. Adolescence physical activity is associated with higher tibial pQCT bone values in adulthood after 28-years of follow-up – the Cardiovascular Risk in Young Finns Study. *Bone* 2015; 75:77-83.

**Table 1.** Subject characteristics as mean values and standard deviations (SD) or frequencies (% , n) in the groups of daily steps.

	Women			P-value*	Men			P-value*
	N=295-353	N=304-355	N=299-354		N=206-263	N=221-265	N=221-263	
	< 6317 steps/day	6317-8765 steps/day	> 8765 steps/day		< 5514 steps/day	5514-7962 steps/day	> 7962 steps/day	
Age, years	37.4 (4.9)	37.9 (5.1)	38.0 (5.0)	0.21	37.5 (5.1)	37.6 (5.2)	37.8 (4.9)	0.74
Height, cm	166 (5.9)	166.4 (6.2)	165.1 (5.9)	0.02	180.9 (6.7)	179.6 (6.9)	178.7 (6.5)	0.001
Weight, kg	71.9 (15.9)	69.5 (13.9)	67.3 (11.8)	<0.001	89.0 (17.3)	85.8 (14.4)	82.9 (14.3)	<0.001
Body mass index, kg/m <sup>2</sup>	26.1 (5.5)	25.1 (5.0)	24.7 (4.1)	0.001	27.1 (4.6)	26.5 (3.9)	25.9 (3.9)	0.003
Energy intake, kcal/day	2144 (701)	2157 (689)	2177 (615)	0.82	2683 (900)	2635 (864)	2763 (812)	0.27
Protein intake, g/day	94 (32.8)	93 (32.9)	94 (28.5)	0.92	116 (39.2)	116 (39.8)	117 (32.6)	0.91
Protein intake, E%	17.7 (2.4)	17.4 (2.3)	17.4 (2.3)	0.21	17.5 (2.6)	17.7 (2.5)	17.2 (2.2)	0.06
Dietary calcium intake, mg/day	1333 (573)	1361 (545)	1391 (525)	0.41	1454 (631)	1574 (664)	1575 (600)	0.06
Serum calcidiol, nmol/l	57.5 (19.3)	61.3 (21.7)	63.4 (19.6)	<0.001	52.3 (17.2)	59.4 (16.4)	58.4 (16.2)	<0.001
Alcohol consumption, drinks/day	0.6 (0.7)	0.5 (0.7)	0.5 (0.7)	0.84	1.5 (2.0)	1.4 (1.5)	1.2 (2.1)	0.27
Current daily smokers, %	18.8 % (66 women)	12.8 % (45)	11.9 % (42)	0.02	22.1 % (58 men)	24.2 % (64)	16 % (42)	0.05
Oral corticosteroid medication ≥ 1 month, %	6.4 % (19 women)	6.9 % (21)	8.4 % (25)	0.64	3.9 % (8 men)	1.4 % (3)	2.7 % (6)	0.26
Physical activity index <sup>a</sup>	8.3 (1.6)	9.0 (1.6)	9.2 (1.8)	<0.001	8.1 (1.8)	8.9 (1.8)	9.2 (2.0)	<0.001
VO <sub>2</sub> max, ml/kg/min <sup>b</sup>	25.8 (6.9)	26.9 (6.7)	27.3 (7)	0.35	32.3 (7.3)	35.4 (7.6)	37.3 (8.4)	0.001
WRmax, W <sup>b</sup>	154 (26.8)	163 (26.3)	170 (30.2)	0.002	246 (45.6)	260 (39.1)	269 (52.1)	0.01

\* Tested with ANOVA, Brown-Forsythe-test or Chi square-test.

<sup>a</sup>Physical activity scores ranged from 5 to 15, where 5 is indicating low activity. <sup>b</sup>VO<sub>2</sub>max, maximal oxygen consumption; WRmax, maximal work rate. VO<sub>2</sub>max and WRmax were measured with maximal cycle ergometer exercise test in a subpopulation of Young Finns (women, N (total)=237-243 and men, N (total)=211).

**Table 2.** Unadjusted mean values (SD) of quantitative ultrasound (QUS) bone parameters separately for women and men in the groups of daily steps.

	Women			ANCOVA <sup>b</sup>		Men			ANCOVA <sup>b</sup>	
	N=223	N=231	N=251	P-value	P-value	N=151	N=168	N=161	P-value	P-value
QUS parameters of calcaneus <sup>a</sup>	< 6317 steps/day	6317-8765 steps/day	> 8765 steps/day		for trend	< 5514 steps/day	5514-7962 steps/day	> 7962 steps/day		for trend
BUA, dB/MHz	77.6 (14.3)	78.8 (15.6)	80.7 (16.6)	0.10	0.04	80.1 (15.7)	83.6 (16.3)	82.6 (16.8)	0.40	0.40
SOS, m/s	1555 (28.7)	1557 (27.9)	1563 (30.2)	0.02	0.01	1555 (27.6)	1561 (28.8)	1562 (29.7)	0.57	0.32

<sup>a</sup>BUA, broadband ultrasound attenuation; SOS, speed of sound. <sup>b</sup>Covariates used in the model: age, height, weight, serum calcidiol level and physical activity index.

**Table 3.** Unadjusted mean values (SD) of tibial peripheral quantitative computed tomography (pQCT) bone parameters separately for women and men in the groups of daily steps.

	Women			ANCOVA <sup>b</sup>		Men			ANCOVA <sup>b</sup>	
	N=267	N=285	N=285	P-value	P-value for trend	N=195	N=213	N=207	P-value	P-value for trend
pQCT bone traits <sup>a</sup>	< 6317 steps/day	6317-8765 steps/day	> 8765 steps/day			< 5514 steps/day	5514-7962 steps/day	> 7962 steps/day		
<i>Distal tibia (5% from the cortical endplate)</i>										
Total area, mm <sup>2</sup>	809.8 (101.3)	815 (118)	818.7 (103.8)	0.04	0.02	993.4 (136.7)	997.8 (128.7)	990.1 (147.6)	0.08	0.03
BMC, mg	511.1 (69.4)	521.1 (71.7)	527.8 (70.6)	<0.001	<0.001	698.8 (103.4)	712.8 (104.4)	708.4 (102.2)	0.04	0.01
Trabecular density, mg/cm <sup>3</sup>	224.2 (28.4)	229.6 (31.1)	232.7 (30.7)	0.001	<0.001	252.3 (33.3)	257.3 (33.4)	256.9 (32.1)	0.75	0.51
BSI, g <sup>2</sup> /cm <sup>4</sup>	0.819 (0.189)	0.848 (0.196)	0.866 (0.208)	0.002	0.001	1.25 (0.29)	1.30 (0.32)	1.29 (0.30)	0.24	0.10
<i>Tibial shaft (30%)</i>										
Total area, mm <sup>2</sup>	341 (44.6)	338.9 (42.1)	344.9 (41.9)	0.001	0.01	438.8 (55.9)	433.9 (49.2)	434.8 (54.1)	0.04	0.04
Cortical area, mm <sup>2</sup>	245.3 (30.7)	245.9 (30)	249.2 (30.3)	0.02	0.01	321.3 (42.1)	320 (40.3)	319.9 (43)	0.19	0.13
Cortical density, mg/cm <sup>3</sup>	1168 (21.6)	1169 (17.5)	1170 (19.3)	0.90	0.65	1144 (21.9)	1147 (22.5)	1148 (22.2)	0.62	0.54
BMC, mg	572.8 (69.2)	574.7 (68.7)	582.5 (68.3)	0.02	0.02	734.2 (89.9)	733.2 (88.9)	733.7 (94.2)	0.17	0.09
CSI	0.72 (0.05)	0.73 (0.05)	0.72 (0.05)	0.36	0.93	0.73 (0.05)	0.74 (0.05)	0.74 (0.05)	0.77	0.68
SSI, mm <sup>3</sup>	1296 (241.9)	1287 (234.3)	1322 (225.3)	0.002	0.01	1842 (325.6)	1818 (301.7)	1830 (329.9)	0.02	0.02

<sup>a</sup>BMC, bone mineral content; BSI, bone strength index; CSI, cortical strength index; SSI, stress-strain index. <sup>b</sup>Covariates used in the model: age, height, weight, serum calcidiol level and physical activity index.

**Table 4.** Unadjusted mean values (SD) of radial peripheral quantitative computed tomography (pQCT) bone parameters separately for women and men in the groups of daily steps.

	Women			ANCOVA <sup>b</sup>		Men			ANCOVA <sup>b</sup>	
	N=267	N=284	N=281	P-value	P-value for trend	N=193	N=213	N=205	P-value	P-value for trend
pQCT bone traits <sup>a</sup>	< 6317 steps/day	6317-8765 steps/day	> 8765 steps/day			< 5514 steps/day	5514-7962 steps/day	> 7962 steps/day		
<i>Distal radius (4% from the cortical endplate)</i>										
Total area, mm <sup>2</sup>	310.9 (42.2)	312.2 (47.5)	314.7 (46.7)	0.25	0.12	418.7 (63.2)	418 (66.1)	420.9 (67.9)	0.26	0.11
BMC, mg	195.2 (28.9)	197.9 (28.4)	199.6 (30.6)	0.04	0.01	305.2 (43.4)	303.3 (46.2)	305.3 (46.7)	0.27	0.17
Trabecular density, mg/cm <sup>3</sup>	205.8 (27.8)	206.8 (28.3)	208.5 (30.2)	0.48	0.22	246.8 (31)	249.5 (33.1)	247.2 (31.5)	0.84	0.96
BSI, g <sup>2</sup> /cm <sup>4</sup>	0.314 (0.085)	0.321 (0.080)	0.325 (0.090)	0.13	0.04	0.565 (0.125)	0.561 (0.138)	0.565 (0.137)	0.58	0.42
<i>Radial shaft (30%)</i>										
Total area, mm <sup>2</sup>	92.8 (13.2)	92.6 (14.2)	94.8 (13.8)	0.008	0.01	140.1 (19.7)	138.7 (19.4)	138.1 (20.1)	0.65	0.42
Cortical area, mm <sup>2</sup>	74.4 (8.4)	74.7 (8.4)	75.7 (8.2)	0.01	0.006	109.7 (12.7)	108.7 (13.2)	108.3 (13.4)	0.73	0.56
Cortical density, mg/cm <sup>3</sup>	1211 (19.5)	1213 (21)	1210 (20)	0.16	0.19	1178 (23.2)	1182 (23)	1183 (20)	0.19	0.17
BMC, mg	180.1 (20.2)	181.2 (20.2)	183.2 (19.6)	0.03	0.01	258.3 (29.6)	256.8 (30.3)	256.2 (31.5)	0.68	0.42
CSI	0.81 (0.06)	0.81 (0.06)	0.81 (0.06)	0.12	0.50	0.79 (0.05)	0.79 (0.05)	0.79 (0.05)	0.77	0.48
SSI, mm <sup>3</sup>	203.8 (38.4)	203.9 (42.1)	208.8 (39)	0.03	0.03	350.3 (67.6)	345.7 (63.8)	344.2 (68.5)	0.60	0.38

<sup>a</sup>BMC, bone mineral content; BSI, bone strength index; CSI, cortical strength index; SSI, stress-strain index. <sup>b</sup>Covariates used in the model: age, height, weight, serum calcidiol level and physical activity index.