A method for automatic creation of a vegetation map using high-resolution aerial photographs of unmanned aerial vehicles



Masatoshi DENDA • Yuichi KAYABA


River restoration team, Water environment group, Public Works Research Institutes Ministry of Infrastructures, Transport and Tourism, Japan government

http://www.pwri.go.jp/team/rrt/index.html, denda@pwri.go.jp




Introduction: Woodland Overgrowth

- Wood overgrowth in river channels degrades the discharge capacity and biodiversity; thus, it is an important river management issue in Japan.
- Many researchers and river managers have pointed out the degradation of riverbeds and reduce of flood disturbance due to gravel pits.
- However, there have been few studies on using ecological modeling to quantify the mechanism. Therefore, our aim was to conduct quantitative research on the mechanism and discuss operation methods to address overgrowth.




Purpose: developments of automatically monitoring method

■Mechanism of Woodland overgrowth




■River Restoration method : channel excavation




First step goal:

Development of automatic creation of a vegetation map using high-resolution aerial photographs of unmanned aerial vehicles .

Method: Overview of the Chikuma river






slope of the river bed: 1/200

Average depth: 0.75m

Introduction: Background River ecosystem change in middle reach of the Chikuma river

In the past river space in middle reach of the Chikuma river has good habitat condition such as sand and gravel riverbed spread along various fluvial condition. And pools, riffles, and ecotones formed along the river, the condition supplied to the good habitat for the aquatic biota.




And, degradations of sand and gravel river bed and ecotone damaged the habitat such as little tern.

^{*}Committee on planning of conservation and restoration on sand and gravel river land in middle reach of the Chikuma river

Typical pattern on bio-diversity degradation in the Chikuma river





Gravel pit

Degradation of disturbances due to floods

Degradation of riverbed

Decrease of habitat of characteristic spices on river land

Growth land spread of Invaded species



wormwood(Arte misia capillaris Thunb)

little tern




Over growth of wood land due to locust




Growth of bur-cucumber

Overview of Restoration projects in the Chikuma river




Results of Restoration projects in the Chikuma river



Flow of vegetation map creation and Replaceable part by machine learning







Overview of Object Based Classification

Object-oriented image classification involves identification of image objects, or segments, that are spatially contiguous pixels of similar texture, color, and tone (Green and Congalton, 2012)


One pixel has enough data for recognize boundary of vegetation community



Recognition by Similar pixels sounding area











Decipher of Vegetation community Using decision tree




Vegetation community map using decision tree




- We success to reconstruct vegetation map
- The precision of the automatically created vegetation map is approximately 65%, and 35% of the errors
- Due to extractions by OBC are too high resolution ?



A Result of automatically creation vegetation map using object-based classification



Human creation vegetation map



Automatically creation vegetation map

LegendWoodland areaWater areaGravel bed

Automatically creation vegetation map reconstructed the spatial distributing character of sand-bar landscape

