

The internal Cdc20 binding site in BubR1 facilitates both spindle assembly checkpoint signaling and silencing.

Tiziana Lischetti¹, Gang Zhang¹, Garry G. Sedgwick¹, Victor M. Bolanos-Garcia² and Jakob Nilsson^{1*}

¹ The Novo Nordisk Foundation Center for Protein Research, University of Copenhagen, Blegdamsvej 3b, 2200 Copenhagen, Denmark

² Department of Biological and Medical Sciences; Oxford Brookes University; Gypsy Lane, Headington; Oxford, OX3 0BP, England

* For correspondence:

Email: jakob.nilsson@cpr.ku.dk

Phone: +45 21328025

Fax: +45 35325001

Running Title: Cdc20 kinetochore interactions

ABSTRACT

Improperly attached kinetochores activate the spindle assembly checkpoint (SAC) and by an unknown mechanism catalyze the binding of two checkpoint proteins, Mad2 and BubR1, to Cdc20 forming the mitotic checkpoint complex (MCC). Here, to address the functional role of Cdc20 kinetochore localization in the SAC, we delineate the molecular details of its interaction with kinetochores. We find that BubR1 recruits the bulk of Cdc20 to kinetochores through its internal Cdc20 binding domain (IC20BD). We show that preventing Cdc20 kinetochore localization by removal of the IC20BD has a limited effect on the SAC because the IC20BD is also required for efficient SAC silencing. Indeed, the IC20BD can disrupt the MCC providing a mechanism for its role in SAC silencing. We thus uncover an unexpected dual function of the second Cdc20 binding site in BubR1 in promoting both efficient SAC signaling and SAC silencing.

INTRODUCTION

Accurate chromosome segregation during mitosis is ensured by the spindle assembly checkpoint (SAC), which is a conserved mechanism requiring the Aurora B, Mad1, Mad2, Mps1, Bub1, BubR1 (Mad3 in yeast) and Bub3 proteins^{1,2}. The proper attachment of microtubules to kinetochores is monitored by the SAC and a single incorrectly attached kinetochore is able to activate the checkpoint^{3,4}. The SAC inhibits chromosome segregation and mitotic exit by blocking the degradation of Securin and Cyclin B1, respectively. Both of these proteins are targeted for degradation by an E3 ubiquitin ligase, the Anaphase Promoting Complex/Cyclosome (APC/C) in complex with its co-activator Cdc20⁵. APC/C activity is inhibited by the cooperative binding of three checkpoint proteins, Mad2, BubR1 and Bub3, to Cdc20, which results in the formation of the mitotic checkpoint complex (MCC)⁶⁻¹¹. Mad2 can bind stably to Cdc20 independently of BubR1 while BubR1 requires Mad2 for its stable association with Cdc20¹²⁻¹⁵. Bub3 does not directly bind to Cdc20 but is required for kinetochore recruitment of BubR1¹⁶⁻¹⁸. Mad2 binds to a short conserved sequence in the N-terminus of Cdc20 while BubR1 interacts with the WD40 domain of Cdc20 and additional contacts between Mad2 and BubR1 stabilize the entire MCC^{19,20}. The protein p31 binds to Mad2 and removes it from the MCC leading to sub stoichiometric levels of Mad2 in the MCC²¹. An N-terminal KEN box motif in BubR1 is critical for stable binding to Cdc20 and binds to conserved residues on the top side of the WD40 domain^{17,19,22-25}. BubR1 in vertebrates also contains an internal Cdc20 binding domain (referred to here as IC20BD) that binds the Cdc20 WD40 domain in a Mad2 independent manner^{6,26,27}. IC20BD spans residues 490-560 in human

BubR1 and binding to Cdc20 depends on a conserved stretch of six amino acids within this otherwise poorly conserved part of BubR1²⁶. The exact function of the IC20BD is not clear but it appears largely dispensable for SAC signaling^{14,17,28}.

Cdc20 inhibitory complexes might be assembled at the kinetochore as the removal of outer kinetochore proteins inactivates the checkpoint²⁹. In line with this, a small fraction of all checkpoint proteins as well as Cdc20 accumulate at unattached kinetochores and turns over rapidly here³⁰⁻³². Indeed, elegant biochemical experiments have shown that kinetochores can stimulate complex formation between Mad2 and Cdc20¹³. Understanding how the kinetochore stimulates the binding of checkpoint proteins to Cdc20 is a key unresolved question but requires a detailed understanding of how Cdc20 interacts with kinetochores, which is currently missing.

Here, we show that the IC20BD of BubR1 is the major kinetochore receptor for Cdc20 and that this domain has dual functions during SAC - both in signaling and silencing. We propose that this dual function could couple BubR1 localization to MCC production or disassembly.

RESULTS

Localization of Cdc20 to kinetochores requires BubR1

To understand the mechanism of Cdc20 kinetochore localization, we first aimed at identifying the kinetochore receptor during prometaphase. A Cdc20 monoclonal antibody stained kinetochores and this staining was absent in cells depleted of Cdc20 by RNAi showing that it is specific (Supplementary Figure 1A). Epitope mapping by peptide array identified an epitope in the N-terminus of Cdc20 and the antibody could immunoprecipitate MCC components (Supplementary Figure 1B-C). We analyzed Cdc20 localization as HeLa cells progressed through an unperturbed mitosis and observed that the levels were highest in prometaphase at unattached kinetochores and decreased in metaphase and anaphase (Supplementary Figure 1D-E). Thus, Cdc20 shows the same localization pattern as canonical SAC components as expected and argued that Cdc20 was very likely recruited to kinetochores through interactions with SAC components.

To determine which protein(s) Cdc20 interacts with at the kinetochore, we focused on Mad2 and BubR1, as these were obvious candidates. We performed RNAi depletion of BubR1 and Mad2 (Figure 1A-C and Supplementary Figure 1F for depletion efficiency). The depletion of BubR1 resulted in a $\approx 65\%$ decrease in Cdc20 kinetochore levels while Mad2 depletion resulted in an $\approx 35\%$ increase (Figure 1B-C).

We used a live cell approach to further investigate the role of Mad2 and BubR1 in Cdc20 kinetochore localization. To this end, we generated a stable HeLa cell line expressing Venus-Cdc20 (for expression levels see

Supplementary Figure 1G), which has previously been shown to complement the Cdc20 RNAi phenotype^{12,33}. This cell line and all other cell lines described here are stable isogenic inducible cell lines made using a HeLa FRT/TRex cell line. When cells expressing Venus-Cdc20 entered mitosis, Cdc20 accumulated on kinetochores and this signal decreased upon alignment of chromosomes (Figure 1D, Supplementary Movie 1). In an unperturbed mitosis Cdc20 kinetochore localization depended on BubR1 but not Mad2 (Figure 1D, Supplementary Movie 2-3). Quantification of the kinetochore intensity of Venus-Cdc20 from the movies revealed similar results as obtained with endogenous Cdc20 (Figure 1E). For this analysis we only analyzed cells that progressed rapidly through mitosis ensuring that the RNAi against BubR1 and Mad2 had worked (see also Supplementary Figure 1H for phenotypic effect of Mad2 and BubR1 RNAi). Similar qualitative results were obtained in cells treated with nocodazole (Supplementary Figure 1I).

Our results reveal a role for BubR1 in the recruitment of Cdc20 to kinetochores, similar to that reported in mice and flies^{27,34}. BubR1 appears to be a major receptor for Cdc20 at kinetochores but additional receptor(s) likely exist as $\approx 35\%$ Cdc20 remains after very efficient BubR1 depletion. In agreement with this, we have found that depletion of KNL1 results in a more efficient removal of Cdc20 from kinetochores, which is not due to more efficient BubR1 removal from kinetochores (Supplementary Figure 1I). Given that BubR1 played a clear role in recruiting Cdc20 to kinetochores, we here focus our efforts on understanding the interaction between these two proteins at kinetochores.

Cdc20 KEN box binding is needed for kinetochore localization

The interaction between Cdc20 and BubR1 within the MCC strongly depends on binding of the N-terminal KEN box of BubR1 to residues of the WD40 domain of Cdc20^{19,25}. To determine if this interaction was still required for kinetochore localization of Cdc20, we mutated amino acids 377-380 of Cdc20 to alanine (mutant referred to as Cdc20 4A), as the available Cdc20-BubR1 structures predict this should disrupt KEN-box binding of Cdc20 (Figure 2A)^{19,25}.

We imaged live cells expressing Venus Cdc20 4A by spinning disk confocal microscopy and observed weaker kinetochore localization of Cdc20 4A compared to wild type Cdc20 (Figure 2B-C). In contrast, when we mutated the Mad2 binding site of Cdc20 by introducing the R132A mutation there was an increase in Cdc20 kinetochore levels (Figure 2B-C). These differences were not due to differences in the overall expression levels of the different Cdc20 proteins (Figure 2D). The kinetochore localization pattern of the Cdc20 mutants was also confirmed by staining the stable cell lines for exogenous Cdc20 (Figure 2E) and this is in agreement with the results from our RNAi depletion of Mad2 and BubR1 (Figure 1).

To ensure that the Cdc20 4A was indeed a functional protein and its inability to locate to kinetochores was not due to misfolding, we further analyzed this mutant. First, we determined the ability of Cdc20 4A to co-immunoprecipitate known interactors from mitotic cells when endogenous Cdc20 was depleted (Figure 2 F-G). Cdc20 4A showed a clear reduction in BubR1-Bub3 binding but maintained binding to Mad2 as predicted. In agreement with the fission yeast MCC structure¹⁹, which revealed that BubR1 and p31 compete for the

same binding site on Mad2, more p31^{comet} was co-purifying with Cdc20 4A. Indeed Mad2 dissociated much more readily from Cdc20 4A when the SAC was silenced (Supplementary Figure 2A). As predicted the Cdc20 R132A mutant had reduced Mad2 binding but still maintained some binding to BubR1-Bub3 although at reduced levels.

We next analyzed mitotic progression in stable HeLa cells expressing Venus tagged Cdc20, Cdc20 R132A and Cdc20 4A that were depleted of endogenous Cdc20. The depletion of Cdc20 delayed cells in mitosis and this could be rescued by expressing Venus-Cdc20 (Figure 2H). The expression of Cdc20 R132A and Cdc20 4A allowed mitotic progression supporting that the mutant proteins were functional and properly folded and this was further confirmed by *in vitro* APC/C ubiquitination assays (Supplementary Figure 2B). Furthermore, both Cdc20 R132A and Cdc20 4A progressed faster through mitosis (Figure 2H) in agreement with the biochemical analysis showing weaker binding to MCC components. As the depletion of Cdc20 by RNAi in HeLa cells does not give a robust mitotic arrest, we also analyzed the ability of Cdc20 4A to support mitotic progression in mouse embryonic fibroblasts (MEFs). We obtained MEFs from the Malumbres lab where exon 2 of Cdc20 is flanked by loxP sites and can be efficiently removed by Cre recombinase³⁵. The MEFs were transfected with the different Venus-Cdc20 constructs (human and mouse Cdc20 are almost identical) and removal of endogenous Cdc20 was induced by addition of 4-hydroxytamoxifen (4-OHT) 24 hours prior to time-lapse imaging. Without the introduction of exogenous Cdc20 there was a strong mitotic delay indicative of efficient Cdc20 removal (Figure 2I, Supplementary Figure 2C). This was fully suppressed by reintroducing the

different forms of Cdc20, showing that Cdc20 4A and Cdc20 R132A are functional (Figure 2I, Supplementary Figure 2C). When we challenged the MEFs with taxol to activate the SAC, a prolonged mitotic arrest was observed when we reintroduced Cdc20 but not Cdc20 R132A and Cdc20 4A confirming that these mutants are checkpoint defective (Figure 2I).

The analysis of Cdc20 4A reveals that this is a functional protein able to interact with Mad2 but defective in stable binding to BubR1-Bub3 and therefore unable to support SAC signaling. The failure of Cdc20 4A to efficiently locate to kinetochores further supports the role of BubR1 as a kinetochore receptor for Cdc20.

The IC20BD of BubR1 recruits Cdc20 to kinetochores

Since Cdc20 4A is inefficiently recruited to kinetochores it suggested to us that the N-terminal KEN box of BubR1 recruits Cdc20 to kinetochores. To test this we generated a panel of stable cell lines expressing inducible Venus tagged siRNA resistant BubR1 constructs (Figure 3A). We then depleted endogenous BubR1 and induced the expression of exogenous BubR1 and treated cells with nocodazole prior to fixation and staining. We stained with a BubR1 antibody recognizing an epitope present in all the BubR1 constructs assayed and only analyzed cells that had a similar level of BubR1 at kinetochores as control treated cells. The kinetochore localization of Cdc20 could be restored to normal levels by expressing exogenous BubR1 (Figure 3B-E). Surprisingly, a BubR1 mutant in which the first KEN box was mutated to AAA (BubR1 KEN/AAA) also efficiently recruited Cdc20 to kinetochores (Figure 3B-E). BubR1 KEN/AAA was clearly defective in forming the MCC as

predicted (see Figure 4D). When we compared the ability of BubR1 1-483 and BubR1 1-715 in recruiting Cdc20 to kinetochores there was a striking difference. BubR1 1-715 clearly recruited Cdc20 while BubR1 1-483 did not (Figure 3B-E). The major difference between these two BubR1 constructs is that BubR1 1-715 contains the IC20BD. In agreement with this, when we deleted the IC20BD of BubR1 (BubR1 Δ 490-560), the BubR1-dependent kinetochore localization of Cdc20 was lost (Figure 3B-E) indicating that the IC20BD and not the N-terminal KEN box of BubR1 recruits Cdc20 to the kinetochores. Removal of six conserved residues of the IC20BD (BubR1 Δ 530-535) was enough to prevent Cdc20 kinetochore localization. Both BubR1 Δ 490-560 and BubR1 Δ 530-35 appeared to be folded properly as they migrated as wild type BubR1 on a size exclusion column with no signs of larger aggregates (Supplementary Figure 3A). Indeed a short peptide encompassing this region of BubR1, but not a mutant peptide, could bind purified Cdc20 arguing that the loss of Cdc20 kinetochore localization upon deletion of the IC20BD is due to loss of a Cdc20 binding site and not misfolding of BubR1 (Supplementary Figure 3B). In agreement with this a fragment of BubR1 encompassing the IC20BD could recruit Cdc20 to kinetochores (Supplementary Figure 3C). Upon overexpression (approximately 300-fold endogenous levels) of Venus tagged BubR1 proteins encompassing the IC20BD in HEK293 cells, we could detect Cdc20 binding confirming previous observations (Supplementary Figure 3 D-E)^{26,27}. However at endogenous levels, IC20BD-containing BubR1 fragments do not co-purify detectable levels of Cdc20 (data not shown). This could either be due to a weak interaction that is not maintained during our purification conditions or

that only a small fraction of BubR1 is interacting with Cdc20 through its IC20BD.

KEN-box binding residues of Cdc20 binds the IC20BD

Initially, we had hypothesized that the defect in kinetochore localization of Cdc20 4A was due to its inability to bind the N-terminal KEN box of BubR1. However, given our observation that BubR1 KEN/AAA still recruited Cdc20 to kinetochores this result was inconsistent with this model. We reasoned that possibly the same residues of Cdc20 binding to the N-terminal KEN-box of BubR1 are also required for binding to IC20BD even though the IC20BD does not contain a KEN-box motif.

To test this hypothesis, we expressed Cdc20 and Cdc20 4A in HEK293 cells and purified the proteins using a strep-tag. We then compared the ability of the Cdc20 proteins to bind a recombinant FLAG tagged fragment of BubR1 (amino acids 516-715) that encompasses the conserved residues of the IC20BD that we found necessary for Cdc20 kinetochore recruitment. The BubR1 fragment was first bound to beads using its FLAG tag, next purified Cdc20 and Cdc20 4A were titrated in and following incubation, the beads were washed. Proteins bound to the beads were analyzed by both quantitative western blotting and by coomassie staining (Figure 3F-G and Supplementary Figure 3F). From these experiments, it was clear that Cdc20 bound BubR1 516-715 while this binding was strongly reduced in Cdc20 4A explaining its reduced kinetochore localization.

Cdc20 kinetochore localization facilitates SAC signaling

As the IC20BD of BubR1 was required for kinetochore localization of Cdc20, we anticipated that it could contribute to SAC signaling although previous work has shown that the N-terminal half of BubR1 is largely sufficient for a functional SAC^{14,17,27}.

To analyze the function of IC20BD, we compared mitotic progression in stable cell lines depleted of endogenous BubR1 and complemented with either Venus-BubR1 or Venus-BubR1 Δ 490-560 at close to endogenous levels (Figure 4A-B, Supplementary Figure 4A). We only analyzed cells with comparable levels of BubR1 as judged from the intensity of the Venus signal. In an unperturbed mitosis, we did not observe any effect on mitotic progression when we removed residues 490-560 (Figure 4A-B). We then challenged the cells with either taxol or nocodazole and monitored the length of mitotic arrest (Figure 4C). In agreement with previous reports, taxol-arrested cells had very few or no Mad2 positive kinetochores while all kinetochores in nocodazole arrested cells were Mad2 positive (Supplementary Figure 4B). In nocodazole-arrested cells, the median time of arrest was only slightly reduced in BubR1 Δ 490-560 complemented cells (BubR1 $t=405$ min., BubR1 Δ 490-560 $t=370$ min.) but this was not statistically significant. In taxol-arrested cells, the length of mitotic arrest was significantly lower in BubR1 Δ 490-560 (BubR1 $t=430$ min., BubR1 Δ 490-560 $t=325$ min., $p\leq 0.01$) but MCC formation was not detectably affected, likely reflecting that the biochemical assay is less sensitive in detecting small differences in MCC composition (Figure 4D). A similar result was obtained with BubR1 Δ 530-535 in taxol (Supplementary Figure 4C). Since the effect that we saw could be due either to the impaired binding of Cdc20 to the IC20BD of BubR1 or to Cdc20

impaired localization to the kinetochore, we prevented BubR1 kinetochore localization by mutating the Bub3 binding site in BubR1 (BubR1 E412K/E413K) and BubR1 Δ 490-560 (referred to as BubR1 Δ Bub3 and BubR1 Δ Bub3/ Δ 490-560) which strongly reduced SAC strength (Figure 4E). In this situation, we exclude the effect of BubR1 kinetochore localization and thereby also of Cdc20, so we can assess the importance of Cdc20 binding alone. The duration of a taxol induced mitotic arrest was the same in BubR1 Δ Bub3 and BubR1 Δ Bub3/ Δ 490-560 complemented cells suggesting that it is not the binding of Cdc20 to IC20BD but the kinetochore localization of Cdc20 that contributes to SAC signaling in taxol (Figure 4E). Our results therefore suggest that Cdc20 kinetochore localization by the IC20BD contributes to SAC signaling under conditions where Mad2 kinetochore levels are low likely by facilitating Mad2-Cdc20 complex formation. In agreement with this hypothesis, depletion of p31^{comet}, which stabilizes the Mad2-Cdc20 complex, resulted in similar timing of BubR1 and BubR1 Δ 490-560 in taxol treated cells (Figure 4F).

The IC20BD is required for SAC silencing

Our analysis of the IC20BD reveals that it makes a small contribution to SAC signaling despite being critical for Cdc20 kinetochore localization. This was puzzling as BubR1 kinetochore localization was critical for an efficient SAC (Figure 4E and¹⁷). However, we observed that in an unperturbed mitosis BubR1 Δ Bub3/ Δ 490-560 complemented cells spent longer time in mitosis than BubR1 Δ Bub3 (BubR1 Δ Bub3/ Δ 490-560 $t=67.5$ min, BubR1 Δ Bub3 $t=50$ min, Mann-Whitney test $p\leq 0.0001$) (Figure 5A). Thus, under conditions where

the SAC is weakened and MCC production lower, the IC20BD seemed important for cells to exit mitosis and thus had the opposite effect than observed in the taxol and nocodazole challenge experiments. This pointed to a role of the IC20BD in SAC silencing.

To directly test this, we challenged BubR1 and BubR1 Δ 490-560 complemented cells with nocodazole and then treated them with reversine to silence the SAC and monitored time of exit by live cell microscopy (Figure 5B-C). This revealed a clear role for the IC20BD in mediating efficient exit in that BubR1 exited with a median time of 36.7 min while BubR1 Δ 490-560 took 50.9 min.

Given that both the BubR1 N-terminus and the IC20BD required the KEN box binding residues of Cdc20 for interaction, a possibility was that the IC20BD could facilitate SAC silencing through competition. To test this, we asked whether recombinant BubR1 516-715 could dissociate BubR1 from the MCC. Using a GFP affinity resin, we purified Venus-Cdc20 from nocodazole-arrested cells, which co-purified Mad2 and BubR1 (Figure 5D). As previously discussed for Venus-BubR1, at close to endogenous levels of proteins we cannot detect binding to the IC20BD, arguing that what we purify with Venus-Cdc20 is a mix of checkpoint complexes (Mad2-Cdc20-BubR1-Bub3, Cdc20-BubR1-Bub3 and Mad2-Cdc20). Next, we added increasing concentrations of recombinant BubR1 516-715 to our affinity purified Venus-Cdc20. Following a one hour incubation at room temperature, the beads were washed and the amount of MCC components remaining bound to Cdc20 analyzed by quantitative western blotting (Figure 5 D-E). With increasing concentrations of BubR1 516-715, we observed a gradual reduction of endogenous BubR1

bound to Cdc20 while the Mad2 levels were only slightly affected. In agreement with this when we overexpressed BubR1 fragments containing the IC20BD, but lacking their Bub3 binding site, the SAC was weakened in taxol arrested cells. Importantly effective weakening of the SAC depended on the conserved residues of the IC20BD (Figure 5F).

Our results show that the IC20BD of BubR1 is required for efficient SAC silencing and a possible mechanism is by competing with the Mad3 homology region of BubR1 for binding to Cdc20.

DISCUSSION

Here, we have dissected the major mechanism of Cdc20 kinetochore recruitment and show that this requires the IC20BD of BubR1 and KEN box binding residues of Cdc20. Preventing Cdc20 kinetochore localization by removing the IC20BD has a limited effect on the SAC but as the IC20BD is also required for SAC silencing these activities of the IC20BD are antagonizing each other potentially to make the checkpoint dynamic and responsive to BubR1 localization.

Our finding that BubR1 is the major kinetochore receptor for Cdc20 in human cells, and that this in turn requires the internal Cdc20 binding site of BubR1, is in agreement with a number of previous observations. Firstly, a clear role for BubR1 but not Mad2 in recruiting Cdc20 to kinetochores has been observed in flies and in mice^{27,34} and fits with the temporal order of recruitment of BubR1 and Cdc20³⁶. Secondly, removal of the Mad2 binding site from Cdc20 does not prevent its kinetochore localization in PtK₂ and LLC-PK1 cells similar to what we observe in human cells with Cdc20 R132A^{30,31}. As the binding between Cdc20 and IC20BD does not require Mad2 binding to Cdc20^{6,26}, our data explains why Mad2 is not needed for bulk Cdc20 recruitment. Elegant FRAP studies of Cdc20 in PtK₂ cells revealed two equal sized populations of Cdc20 at kinetochores, one with a fast turnover and one with a slower turnover³⁰. The Cdc20 population with a slower turnover was dependent on the N-terminus of Cdc20 and an active SAC. The same study also detected two populations of BubR1 with similar slow and fast turnover as Cdc20. These data together with the data presented here suggests that Cdc20 is binding the IC20BD of both a slow and a fast pool of BubR1 at kinetochores and that the

slow BubR1 pool depends on an active SAC (Figure 6). In flies there appears to be only a fast pool of Cdc20 which might reflect a difference in how BubR1 interacts with kinetochores^{34,37}.

Collectively our data does not support that Cdc20 exists in MCC-like complexes at kinetochores as both preventing Mad2 binding or mutating the N-terminal KEN box of BubR1 did not affect Cdc20 localization. This suggests that the full assembly of the MCC occurs in the cytoplasm either through maturation of a partly assembled MCC or binding of soluble BubR1 to Mad2-Cdc20 (Figure 6).

Although we find that BubR1 recruits the bulk of Cdc20 to kinetochores our work also suggests that at least one other binding partner must exist at kinetochores as even after efficient BubR1 depletion $\approx 35\%$ Cdc20 remains. The stronger effect of KNL1 RNAi could point to a role of Bub1 as an additional binding partner for Cdc20 at kinetochores and indeed human Bub1 has been shown to bind Cdc20 (Kang et al., 2007). Further experiments are needed to address the role of Bub1 in BubR1 and Cdc20 kinetochore localization.

The exact function of the IC20BD of BubR1 has not been clear and several studies have shown that it is not required for a functional SAC^{14,17}. In the study from the Cleveland lab it was however noted that cells complemented with full length BubR1 maintained a nocodazole induced arrest for longer time than cells complemented with BubR1 1-477 potentially revealing a role of the C-terminal half of BubR1 in prolonged arrest¹⁴. Although the IC20BD can inhibit Cdc20 activity *in vitro*⁶, our work and that of others indicate that *in vivo*

the interaction of Cdc20 with this region of BubR1 is either of low affinity or restricted to binding a small proportion of Cdc20.

Our comparison of BubR1 and BubR1 deleted of its IC20BD only revealed a small contribution of this domain to SAC signaling specifically in taxol-arrested cells. We favor that the function of the IC20BD is to weakly bind and hereby concentrate Cdc20 at kinetochores bringing it in proximity of the Mad1-Mad2 complex to facilitate Mad2-Cdc20 complex formation. The reason we favor this is that we only see a SAC defect when Mad2 signaling from kinetochores is low and this defect is suppressed by p31^{comet} removal. Furthermore the defect must relate to Cdc20 kinetochore localization as preventing BubR1 kinetochore localization abolishes the effect of removing the IC20BD. However we cannot exclude that the IC20BD plays a more active role in the SAC.

We also find a requirement of the IC20BD in SAC silencing and the true effect on the SAC upon removal of the IC20BD is likely masked by this function. We roughly estimate the median times to be 50 min for BubR1 and 30 min for BubR1 Δ 490-560 in an unperturbed mitosis without the effect of the IC20BD on SAC silencing. This is as severe an effect on the SAC as mutating the Bub3 binding site in BubR1. Our data would support a model in which the IC20BD contributes to SAC signaling by recruiting Cdc20 to kinetochores to facilitate interaction with Mad2 and in SAC silencing by destabilizing the BubR1-Cdc20 interaction within the MCC through competition (Figure 6). As these two activities of the IC20BD are counteracting each other, it depends on the experimental conditions what activity is the most dominant. This would explain why we only see a clear effect on the SAC in taxol arrested cells

where the role of the IC20BD in recruiting Cdc20 to kinetochores becomes very critical.

BubR1 is recruited to improperly attached kinetochores and thus its function in recruiting Cdc20 to the kinetochore to facilitate interaction with Mad2 and at the same time recruiting PP2A³⁸⁻⁴⁰ to stabilize kinetochore-microtubule interactions shows the remarkable ability of BubR1 to integrate important kinetochore activities through short interaction motifs. Once proper kinetochore-microtubule interactions are established BubR1 leaves the kinetochore and the IC20BD then acts to destabilize the MCC for efficient mitotic exit. The dual function of the IC20BD thus elegantly couples SAC signaling or silencing to the localization of BubR1.

METHODS

Cloning and stable cell lines

All constructs were cloned into pcDNA5/FRT/TO FLAG Venus (N-terminal tagging). Cdc20 and BubR1 and fragments thereof were amplified by PCR and inserted into the BamHI and NotI sites of the vector using the primers specified in Supplementary table I. Mutations into these constructs were done using whole plasmid PCR or 2-step PCR. All constructs were verified by sequencing.

The generation of stable HeLa cell lines was done using the Flp-In system (Invitrogen) and the resulting clones were kept under selection by supplementing the growth media with 200 µg/ml Hygromycin B and 5 µg/ml Blasticidin S.

Antibodies

The following antibodies were used for western blot, immunofluorescence or immunoprecipitation as indicated: α -tubulin 11H10 (rabbit, Cell Signaling) 1:100 IF, β -actin AC-15 (mouse, Abcam) 1:5000 WB, APC1 A301-653A (rabbit, Bethyl) 1:500 WB, APC3 35/CDC27 (mouse, BD Biosciences) 1:250 WB, APC4 (mouse, raised against a C-terminal peptide) 5µg IP, APC7 A302-551A (rabbit, Bethyl) 1:1000 WB, Bub3 clone 31 (mouse, BD Transduction Laboratories) 1:500 WB, BubR1 A300-995A (rabbit, Bethyl) 1:500 WB, BubR1 A300-386A (rabbit, Bethyl) 1:200 IF, Cdc20 AR12 (mouse, Millipore) 1:200 IF/5 µg IP, Cdc20 E-7 (mouse, Santa Cruz Biotechnology) 1:500 WB, Cdc20 A301-180A (rabbit, Bethyl) 1:1000 WB, CREST (human, Antibodies Inc.) 1:400 IF, FLAG M2 (mouse, Sigma) 1:5000 WB, GFP clones 7.1 and 13.1

(mouse, Roche) 1:5000 WB /1:200 IF, Mad2 from G. Kops (rabbit) 1:200 IF, Mad2 A300-301A (rabbit, Bethyl) 1:1000 WB, p31 rabbit antibody was generated using full-length p31 as antigen and affinity purified.

RNAi depletion of proteins

RNAi depletion of proteins was performed for 48 hours, except for p31 that was performed twice at 72 and 48 hours before analysis. DMEM media was replaced with OPTIMEM media right before adding a mix containing 100 nM RNAi oligo (Sigma) and 4 ul/ml RNAimax (Invitrogen) diluted in OPTIMEM. The media was changed after 5 hours and replaced with DMEM supplemented with 10% FBS. siRNAs used are: BubR1 5'-GAUGGUGAAUUGUGGAAUA-3'; Cdc20 5'-CGGAAGACCUGCCGUUACAUU-3'; KNL1 5'-AAGAUCUGAUUAAGGAUCCACGAAA-3'; Luciferase 5'-CGUACGCGGAAUACUUCGA-3'; Mad2 5'-GGAAGAGUCGGGACCACAG-3'; p31 5'-GGCUGCUGUCAGUUUACUU-3'

Immunofluorescence analysis

For immunofluorescence microscopy, cells were synchronized with 2.5 mM thymidine, 10 μ M MG132 and 660 mM nocodazole as shown in figure 1A. Cells were pre-fixed with 4% PFA in PHEM buffer (50 mM Pipes, 25mM HEPES, 10 mM EGTA, 8.5 mM MgSO₄, pH 7.0) for 20 seconds, permeabilized in 0.5% TritonX-100 in PHEM buffer for 5 minutes and fixed in 4% PFA in PHEM buffer for 20 minutes. Coverslips were quenched with 25 mM Glycin in PBS for 20 minutes, blocked with 3% BSA in PBS-T (0.1% Tween in PBS) for

30 minutes, incubated with primary antibodies for 1 hour, with Alexa Fluor goat secondary antibodies for 45 minutes and mounted on slides using ProLong Gold Antifade mounting media (Invitrogen).

Images were acquired taking z stacks of 200 nm using a 100X/1.4NA objective on a DeltaVision Elite Microscope (GE Healthcare). Images were analyzed after deconvolution using SoftWoRx (GE Healthcare). Figures were generated by maximum intensity projection of entire cells using Softworx and ImageJ.

Live cell imaging

Live cell analysis was performed on a Deltavision Elite system using a 40x or 60x objective (GE Healthcare). Cells were seeded in 8 well Ibidi dishes (Ibidi) in advance and prior to filming the media was changed to Leibovitz's L-15 (Life technologies). Appropriate channels were recorded for 18 hours and data analyzed using Softworx (GE Healthcare). MEFs were transfected with Venus-Cdc20 proteins by electroporation using the Neon Transfection System according to manufacturer instructions. 1 μ M 4-OHT and 200 nM taxol were added where indicated 24 hours and 2 hours before filming respectively. For experiments in figure 2B images were acquired on a Perkin Elmer UltraView Vox-Spinning Disk CSU-X1 using a 60x1.4 oil objective.

Immunopurification of complexes

Immunoprecipitation of Venus proteins from stable HeLa cell lines was performed from cells synchronized with 2.5 mM thymidine and arrested in mitosis with either 660 nM nocodazole or 200 nM taxol. The mitotic cells

were harvested by shake off, washed with ice-cold PBS and lysed for 30 minutes on ice in lysis buffer (150 mM NaCl, 50 mM Tris-HCl pH 7.5, 1 mM EDTA, 1% NP40, 1 mM DTT, 1x protease inhibitor cocktail (Roche), 1x phosphatase inhibitor cocktail (Roche)). Lysates were clarified by centrifugation at 20,000 rcf at 4°C for 15 minutes and the resulting cell extracts were incubated with 20 µl of GFP-Trap A beads (ChromoTek) at 4°C with gentle agitation for 30 minutes. The beads were then washed 3 times with ice-cold lysis buffer and the complexes eluted in 2x SDS sample buffer. Low salt/high salt IP was performed using a different lysis buffer (50 mM NaCl, 25 mM Tris-HCl pH 7.5, 0.1% NP40, 1 mM DTT, 1x protease inhibitor cocktail (Roche), 1x phosphatase inhibitor cocktail (Roche)) and washing buffer (50 mM Tris-HCl pH 7.5, 1 mg/ml BSA, 20% glycerol, 0.5 mM DTT, 1x protease inhibitor cocktail (Roche), 1x phosphatase inhibitor cocktail (Roche)) supplemented with 300 mM NaCl only for the high salt condition. IP of endogenous proteins was performed using Dynabeads Protein G (Invitrogen) and 5 µg of antibodies.

***In vitro* binding experiments with BubR1 and Cdc20.**

FLAG-HA-BubR1 516-715 was expressed in the E. coli BL21 Rosetta2 (DE3) R3 T1 strain by induction with 0.5 mM IPTG for 20 hours at 18°C. The bacterial pellet was resuspended in buffer L (50 mM NaP, 300 mM NaCl, 10% glycerol, 0.5 mM TCEP, protease inhibitors) and lysed using a high-pressure homogenizer at 1,000 Bar. After clarification by centrifugation (18,000 g for 30 minutes) the protein was purified by affinity (1 ml Ni column (GE healthcare), loaded in buffer L with 10 mM imidazole) and gel filtration chromatography

(Superdex 200 PG 16/60 equilibrated with SEC buffer (50 mM NaP, 150 mM NaCl, 0.5 mM TCEP, 10% glycerol, pH 7.5)). Strep-His-Cdc20 and mutants thereof were expressed in HEK293 6E cell lines by transfection with 100 µg/ml Polyethylenimine "MAX"(PEI) (polysciences). After 3 days the Cdc20 proteins were affinity purified using a Strep-tag/Strep-Tactin purification system (IBA) according to manufactures description. 10 µg of FLAG-HA-BubR1 516-715 protein was incubated with 20 µl of anti-FLAG M2 affinity gel (Sigma) in PBS for 1 hour at room temperature with gentle mixing followed by 3 washes with washing buffer (PBS, 0.1% TritonX-100). The affinity gel was then resuspended in binding buffer (150 mM NaCl, 50 mM Tris-HCl pH 7.5, 0.3% Triton X-100, 1x protease inhibitor cocktail (Roche), 1x phosphatase inhibitor cocktail (Roche)) and incubated with 2, 5 or 10 µg of Strep-His-Cdc20 proteins for 1 hour at room temperature with gentle mixing. After 4 washes with the binding buffer the bound proteins were released from the affinity gel by addition of 2x SDS sample buffer and incubation at 95°C. The eluates were analyzed by Coomassie staining and western blotting against BubR1 (FLAG) and Cdc20 (E7, Santa Cruz).

***In vitro* ubiquitination assays with APC/C.**

APC/C-Cdc20 complexes were purified from mitotic cells using an APC4 antibody. Cells were either control treated or depleted of Cdc20 by RNAi and the expression of the different Venus-Cdc20 proteins induced by doxycycline. *In vitro* ubiquitination assays with *in vitro* translated Cyclin B1 1-86 was carried out as previously described ⁴¹.

Binding experiments in HEK293 cells.

For experiments presented in Figure S3A-B HEK293T cell were transfected with Venus-BubR1 proteins and depleted of endogenous BubR1 using Lipofectamine 2000 (Invitrogen), 500 ng/ml plasmid and 100 nM RNAi oligo. Cells were subsequently synchronized with 2.5 mM thymidine for 24 hours and treated with 200 ng/ml nocodazole after the thymidine release. 10 μ M MG132 was added after 6 hours from the thymidine release for 3 hours. Cells were then harvested and washed with ice-cold PBS.

Peptide binding experiments.

Two peptides were used for this assay: Biotin-YSVPFSIFDEFLLESEKKNKS (WT peptide) and Biotin-YSVPFSIAKKAASEKKNKS (mut peptide) (BIOSYNTAN). 2 μ g of purified Strep-His-Cdc20 was incubated with peptide-streptavidin agarose complex (Sigma) for 1 hour at room temperature. The binding buffer contains 50 mM Tris-HCl, pH= 7.5, 150 mM NaCl, 0.3% TritonX-100 with protease inhibitors. After 4 times of washing with the binding buffer, the bound proteins were eluted by 1x Laemmli buffer and released by incubating at 95°C. The eluates were analyzed by western blot against Cdc20 (E7, Santa Cruz).

Size exclusion chromatography.

Cell lines were arrested in nocodazole and treated for 1 hour with reversine and MG132 before harvesting. The cells were resuspended in lysis buffer (150 mM NaCl, 25 mM TRIS pH 8.0, 0.1% NP-40, 1 mM DTT, 1x protease inhibitor cocktail (Roche) and 1x phosphatase inhibitor cocktail (Roche)).

Following lysis the extract was clarified by spinning for 10 minutes at 20000 g and subsequently for 10 minutes at 186000 g in a TLA-55 rotor (Beckman-Coulter). 500 ul extract at 6 ug/ul was loaded on a Superose 6 10/300 GL column equilibrated with lysis buffer. 500 ul fractions where collected and analyzed.

REFERENCES

1. Musacchio, A. Spindle assembly checkpoint: the third decade. *Philos. Trans. R. Soc. Lond., B, Biol. Sci.* **366**, 3595–3604 (2011).
2. Lara-Gonzalez, P., Westhorpe, F. G. & Taylor, S. S. The spindle assembly checkpoint. *Curr. Biol.* **22**, R966–80 (2012).
3. Rieder, C. L., Schultz, A., Cole, R. & Sluder, G. Anaphase onset in vertebrate somatic cells is controlled by a checkpoint that monitors sister kinetochore attachment to the spindle. *J. Cell Biol.* **127**, 1301–1310 (1994).
4. Rieder, C. L., Cole, R. W., Khodjakov, A. & Sluder, G. The checkpoint delaying anaphase in response to chromosome monoorientation is mediated by an inhibitory signal produced by unattached kinetochores. *J. Cell Biol.* **130**, 941–948 (1995).
5. Pines, J. Cubism and the cell cycle: the many faces of the APC/C. *Nat. Rev. Mol. Cell Biol.* **12**, 427–438 (2011).
6. Tang, Z., Bharadwaj, R., Li, B. & Yu, H. Mad2-Independent inhibition of APCCdc20 by the mitotic checkpoint protein BubR1. *Dev. Cell* **1**, 227–237 (2001).
7. Fang, G. Checkpoint protein BubR1 acts synergistically with Mad2 to inhibit anaphase-promoting complex. *Mol. Biol. Cell* **13**, 755–766 (2002).
8. Sudakin, V., Chan, G. K. & Yen, T. J. Checkpoint inhibition of the APC/C in HeLa cells is mediated by a complex of BUBR1, BUB3, CDC20, and MAD2. *J. Cell Biol.* **154**, 925–936 (2001).
9. Kim, S. H., Lin, D. P., Matsumoto, S., Kitazono, A. & Matsumoto, T. Fission yeast Slp1: an effector of the Mad2-dependent spindle checkpoint. *Science* **279**, 1045–1047 (1998).
10. Hwang, L. H. *et al.* Budding yeast Cdc20: a target of the spindle checkpoint. *Science* **279**, 1041–1044 (1998).
11. Fang, G., Yu, H. & Kirschner, M. W. The checkpoint protein MAD2 and the mitotic regulator CDC20 form a ternary complex with the anaphase-promoting complex to control anaphase initiation. *Genes Dev.* **12**, 1871–1883 (1998).
12. Nilsson, J., Yekezare, M., Minshull, J. & Pines, J. The APC/C maintains the spindle assembly checkpoint by targeting Cdc20 for destruction. *Nat. Cell Biol.* **10**, 1411–1420 (2008).
13. Kulukian, A., Han, J. S. & Cleveland, D. W. Unattached kinetochores catalyze production of an anaphase inhibitor that requires a Mad2 template to prime Cdc20 for BubR1 binding. *Dev. Cell* **16**, 105–117 (2009).
14. Han, J. S. *et al.* Catalytic Assembly of the Mitotic Checkpoint Inhibitor BubR1-Cdc20 by a Mad2-Induced Functional Switch in Cdc20. *Mol. Cell* **51**(1):92-104. (2013).
15. Hardwick, K. G., Johnston, R. C., Smith, D. L. & Murray, A. W. MAD3 encodes a novel component of the spindle checkpoint which interacts with Bub3p, Cdc20p, and Mad2p. *J. Cell Biol.* **148**, 871–882 (2000).
16. Taylor, S. S., Ha, E. & McKeon, F. The human homologue of Bub3 is required for kinetochore localization of Bub1 and a Mad3/Bub1-related protein kinase. *J. Cell Biol.* **142**, 1–11 (1998).
17. Lara-Gonzalez, P., Scott, M. I. F., Diez, M., Sen, O. & Taylor, S. S.

- BubR1 blocks substrate recruitment to the APC/C in a KEN-box-dependent manner. *J. Cell. Sci.* 124(Pt 24):4332-45 (2011).
18. Elowe, S. *et al.* Uncoupling of the spindle-checkpoint and chromosome-congression functions of BubR1. *J. Cell. Sci.* **123**, 84–94 (2010).
 19. Chao, W. C. H., Kulkarni, K., Zhang, Z., Kong, E. H. & Barford, D. Structure of the mitotic checkpoint complex. *Nature* **484**, 208–213 (2012).
 20. Tipton, A. R. *et al.* BUBR1 and closed MAD2 (C-MAD2) interact directly to assemble a functional mitotic checkpoint complex. *J. Biol. Chem.* **286**, 21173–21179 (2011).
 21. Westhorpe, F. G., Tighe, A., Lara-Gonzalez, P. & Taylor, S. S. p31comet-mediated extraction of Mad2 from the MCC promotes efficient mitotic exit. *J. Cell. Sci.* **124**, 3905–3916 (2011).
 22. King, E. M. J., van der Sar, S. J. A. & Hardwick, K. G. Mad3 KEN boxes mediate both Cdc20 and Mad3 turnover, and are critical for the spindle checkpoint. *PLoS ONE* **2**, e342 (2007).
 23. Sczaniecka, M. *et al.* The spindle checkpoint functions of Mad3 and Mad2 depend on a Mad3 KEN box-mediated interaction with Cdc20-anaphase-promoting complex (APC/C). *J. Biol. Chem.* **283**, 23039–23047 (2008).
 24. Burton, J. L. & Solomon, M. J. Mad3p, a pseudosubstrate inhibitor of APCCdc20 in the spindle assembly checkpoint. *Genes Dev.* **21**, 655–667 (2007).
 25. Tian, W. *et al.* Structural analysis of human Cdc20 supports multisite degron recognition by APC/C. *Proc. Natl. Acad. Sci. U.S.A.* 109(45):18419-24 (2012).
 26. Davenport, J., Harris, L. D. & Goorha, R. Spindle checkpoint function requires Mad2-dependent Cdc20 binding to the Mad3 homology domain of BubR1. *Exp. Cell Res.* **312**, 1831–1842 (2006).
 27. Malureanu, L. A. *et al.* BubR1 N terminus acts as a soluble inhibitor of cyclin B degradation by APC/C(Cdc20) in interphase. *Dev. Cell* **16**, 118–131 (2009).
 28. Suijkerbuijk, S. J. E. *et al.* Molecular causes for BUBR1 dysfunction in the human cancer predisposition syndrome mosaic variegated aneuploidy. *Cancer Res.* **70**, 4891–4900 (2010).
 29. Meraldi, P., Draviam, V. M. & Sorger, P. K. Timing and checkpoints in the regulation of mitotic progression. *Dev. Cell* **7**, 45–60 (2004).
 30. Howell, B. J. *et al.* Spindle checkpoint protein dynamics at kinetochores in living cells. *Curr. Biol.* **14**, 953–964 (2004).
 31. Kallio, M. J., Beardmore, V. A., Weinstein, J. & Gorbsky, G. J. Rapid microtubule-independent dynamics of Cdc20 at kinetochores and centrosomes in mammalian cells. *J. Cell Biol.* **158**, 841–847 (2002).
 32. Shah, J. V. *et al.* Dynamics of centromere and kinetochore proteins; implications for checkpoint signaling and silencing. *Curr. Biol.* **14**, 942–952 (2004).
 33. Hein, J. B. & Nilsson, J. Stable MCC binding to the APC/C is required for a functional spindle assembly checkpoint. *EMBO Rep.* 15(3):264-72 (2014).
 34. Li, D., Morley, G., Whitaker, M. & Huang, J.-Y. Recruitment of Cdc20 to the kinetochore requires BubR1 but not Mad2 in *Drosophila*

- melanogaster. *Mol. Cell. Biol.* **30**, 3384–3395 (2010).
35. Manchado, E. *et al.* Targeting mitotic exit leads to tumor regression in vivo: Modulation by Cdk1, Mastl, and the PP2A/B55 α,δ phosphatase. *Cancer Cell* **18**, 641–654 (2010).
 36. Acquaviva, C., Herzog, F., Kraft, C. & Pines, J. The anaphase promoting complex/cyclosome is recruited to centromeres by the spindle assembly checkpoint. *Nat. Cell Biol.* **6**, 892–898 (2004).
 37. Herriott, A., Sweeney, M., Whitaker, M., Taggart, M. & Huang, J.-Y. Kinetochore localized Mad2 and Cdc20 is itself insufficient for triggering the mitotic checkpoint when Mps1 is low in *Drosophila melanogaster* neuroblasts. *Cell Cycle* **11**, (2012).
 38. Xu, P., Raetz, E. A., Kitagawa, M., Virshup, D. M. & Lee, S. H. BUBR1 recruits PP2A via the B56 family of targeting subunits to promote chromosome congression. *Biol Open* **2**, 479–486 (2013).
 39. Kruse, T. *et al.* Direct binding between BubR1 and B56-PP2A phosphatase complexes regulate mitotic progression. *J. Cell. Sci.* **126**(Pt 5):1086-92. (2013).
 40. Suijkerbuijk, S. J. E., Vleugel, M., Teixeira, A. & Kops, G. J. P. L. Integration of Kinase and Phosphatase Activities by BUBR1 Ensures Formation of Stable Kinetochore-Microtubule Attachments. *Dev. Cell* **23**, 745–755 (2012).
 41. Sedgwick, G. G. *et al.* Mechanisms controlling the temporal degradation of Nek2A and Kif18A by the APC/C-Cdc20 complex. *EMBO J.* **32**(2):303-14 (2013).

END NOTES

ACKNOWLEDGEMENTS

We thank Geert Kops for providing the rabbit Mad2 antibody, Stephen Taylor for providing the HeLa FRT TRex cell line and Marcos Malumbres for the Cdc20 conditional knockout MEFs. We thank Jon Pines and Barbara Di Fiore for sharing unpublished results, Silke Hauf and Andrea Musacchio for fruitful discussions, Werner W. Streicher for help with the peptide arrays, Giuseppe Cazzamali and Michael Williamson for producing purified Cdc20 proteins, Jutta Bulkescher for help with spinning disk confocal microscopy and Dan Hayward and Marie Larsen for comments on the manuscript. This project was supported by a grant from the Novo Nordisk Foundation, the Lundbeck Foundation and a grant from the Danish Council for Independent Research-Natural Sciences to JN.

AUTHOR CONTRIBUTION

TL performed all experiments except for the experiments in Figure 3B-E and Figure S3B-C, which were performed by GZ and Figure S2B which was performed by GGS. VBG helped with generating structural homology models of human Cdc20 prior to their direct structural determination. JN performed cloning work and wrote the paper.

CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

FIGURE LEGENDS

Figure 1. Recruitment of Cdc20 to kinetochores requires BubR1.

A) Schematic of protocol used to synchronize cells and deplete specific proteins using RNAi. B) Cells depleted of the indicated proteins were stained for CREST and Cdc20 and BubR1 or Mad2 to check for depletion efficiency. Scale bar represents 5 μm . C) Quantification of Cdc20, BubR1 and Mad2 kinetochore levels in cells treated with the indicated RNAi oligoes. The fluorescence from the three z-stacks (200 nm apart) encompassing the bulk kinetochore fluorescent intensity was used and normalized to the CREST signal. Cells were co-stained for Mad2 or BubR1 to ensure that only cells with efficient depletion were analyzed. At least 80 kinetochore pairs from 8 cells were quantified and the mean and standard error of mean is indicated. D) Stable HeLa cell line expressing Venus-Cdc20 was treated with the indicated RNAi oligoes and the localization of Venus-Cdc20 followed by time-lapse microscopy. Scale bar represents 10 μm . E) Quantification of Venus-Cdc20 at kinetochores in the movies in D). The signal was quantified from a single z-section and in the frame right after NEBD. A total number of 50 kinetochores from 10 cells were analyzed for each condition and the mean and standard error of mean indicated. An unpaired t-test was performed for statistical analysis (**** is $p < 0.0001$) and the comparison is to the control treated cells.

Figure 2. A novel Cdc20 mutant specifically defective in BubR1 binding.

A) The structure of human Cdc20 bound to the BubR1 KEN-box motif (blue) (PDB: 4GGD) with the residues mutated in Cdc20 4A in red. B) Stable HeLa cell lines expressing the indicated Cdc20 proteins were analyzed by live cell

spinning disk confocal microscopy and representative still images are shown. Scale bar represents 10 μm . C) Quantification of Cdc20 intensity at the kinetochores from the experiment shown in B). A total of 50 kinetochores from 10 different cells were analyzed for each condition. Cdc20 kinetochore signal was determined using the ImageJ software and mean and standard error of mean are shown. An unpaired t-test was used for statistical analysis. (**** is $p < 0.0001$). D) The Cdc20 intensity in the cytoplasm was measured from the images acquired with the spinning disk from the 10 cells used for the analysis in C). An unpaired t-test was used for statistical analysis. E) Stable HeLa cell lines expressing the indicated Cdc20 proteins were arrested by nocodazole and MG132, and stained for CREST and Cdc20 using a GFP specific antibody. Scale bar represents 5 μm . F) Stable HeLa cell lines expressing the indicated Cdc20 proteins were arrested in nocodazole and MG132 and cells collected by mitotic shake-off. Venus-Cdc20 proteins were purified using a GFP affinity resin and washed with either a low salt buffer or high salt buffer as indicated. The composition of proteins associated with Cdc20 was determined by blotting for the indicated proteins. G) Quantification of the proteins bound to Venus-Cdc20 proteins was determined. The mean and standard error of means of two independent experiments are shown. H) Stable HeLa cells were treated with a control RNAi oligo (Luciferase) or depleted of Cdc20 and then complemented with the different Venus-Cdc20 proteins as indicated. Each circle represents a single cell analyzed (at least 60 cells were analyzed for each condition) and the red line indicates the median ($m =$). A Mann-Whitney test was performed for statistical analysis. I) Mouse embryonic fibroblasts (MEFs) were depleted of Cdc20 by addition of 4-

OHT 24 hours prior to imaging and complemented with the indicated Venus-Cdc20 proteins. At least 25 cells were analyzed per condition and a Mann Whitney test was used for statistical analysis. Median time (m=) is indicated for each condition.

Figure 3. The internal Cdc20 binding site of BubR1 recruits Cdc20 to kinetochores.

A) Schematic of human BubR1 and the location of Cdc20 binding sites and Bub3 binding site as well as the pseudo-kinase domain. Alignment of the region encompassing residues 530-535 of human BubR1 is shown on top and the different constructs used are indicated below. The BubR1 KEN/AAA has the first KEN-box mutated to AAA. B) Stable HeLa cell lines expressing the different Venus-BubR1 siRNA resistant constructs were used to determine the domains in BubR1 required for Cdc20 kinetochore localization. Briefly, cells were treated with a control RNAi oligo (Luciferase) or a BubR1 RNAi oligo and then arrested in mitosis using nocodazole treatment and the proteasome inhibitor, MG132. BubR1 RNAi treated cells were complemented with the indicated Venus-BubR1 constructs. Cells were stained for BubR1, CREST and Cdc20. Scale bar represents 5 μ m C-D). The kinetochore levels of BubR1 (C) and Cdc20 (D) normalized to CREST in control and BubR1 depleted cells. At least 80 kinetochore pairs from 8 different cells were analyzed and the mean and standard error of mean is indicated. E) The level of Cdc20 at kinetochores in cells complemented with the indicated Venus-BubR1 constructs was determined. Only cells with endogenous levels of BubR1 at kinetochores were used for this analysis. At least 80 kinetochore pairs from 8

different cells were analyzed and the mean and standard error of mean is indicated. F) Binding of Cdc20 and Cdc20 4A to BubR1 516-715 was determined by binding 10 μ g recombinant FLAG-HA-BubR1 516-715 to FLAG affinity beads and incubating these with increasing concentrations of Strep-His tagged Cdc20 or Cdc20 4A expressed and purified from HEK293 cells. The beads were washed and bound proteins were eluted and analyzed by western blot. G) Quantification of western blot in F) using Licor technology. Experiment in F-G is representative of two independent experiments.

Figure 4. The internal Cdc20 binding site of BubR1 contributes to SAC signaling during a taxol-induced arrest.

A) Mitotic progression in HeLa cells stably expressing similar levels of Venus-BubR1 or Venus-BubR1 Δ 490-560 and depleted of endogenous BubR1 was determined by time-lapse microscopy. This was compared to Luciferase RNAi and BubR1 RNAi treated cells. Each circle represents a single cell analyzed (at least 50 cells were analyzed per condition) and the red line indicates the median ($m=$). A Mann Whitney test was used for statistical analysis. B) Still images from the time-lapse movies of Venus-BubR1 and Venus-BubR1 Δ 490-560. Scale bar represents 10 μ m. C) Similar to A) but cells was challenged with either 200 nM taxol or 100 nM nocodazole as indicated. The red line indicates the median ($m=$) and each circle represents a single cell analyzed (at least 50 cells were analyzed per condition). A Mann Whitney test was used for statistical analysis. D) The indicated Venus-BubR1 proteins were purified from taxol arrested cells using a GFP affinity resin and washed with different salt conditions, as indicated. The binding to the indicated proteins was

analyzed by western blotting. E) HeLa cells stably expressing similar levels of Venus-BubR1, Venus-BubR1 Δ Bub3 or Venus-BubR1 Δ Bub3/ Δ 490-560 and depleted of endogenous BubR1 were treated with 200 nM taxol and their mitotic progression was followed by time-lapse microscopy. Each circle represents a single cell analyzed (at least 40 cells were analyzed per condition) and the red bars indicate the medians (m=). A Mann Whitney test was used for statistical analysis. F) HeLa cells were depleted of BubR1 and/or p31^{comet} as indicated. Luciferase knock down was included as control. BubR1 depletion was complemented with Venus-BubR1 or Venus-BubR1 Δ 490-560 as indicated and 200 nM taxol was added prior filming. Each circle represents a single cell analyzed and at least 50 cells were analyzed per condition. Medians (m=) are shown in red and a Mann Whitney test was used for the statistical analysis.

Figure 5. The IC20BD is required for SAC silencing.

A) Stable HeLa cells were depleted of endogenous BubR1 or treated with control RNAi oligo (Luciferase). BubR1 knock down was complemented by expression of Venus-BubR1, Venus-BubR1 Δ Bub3 or Venus-BubR1 Δ Bub3/ Δ 490-560 as indicated and mitotic progression was followed by time-lapse microscopy. Each circle represents a single cell analyzed (at least 40 cells were analyzed for each condition) and the median (m=) is shown as a red bar. Statistical analysis was performed using a Mann Whitney test. B) Stable HeLa cell lines were depleted of endogenous BubR1 and complemented by expression of Venus-BubR1 or Venus-BubR1 Δ 490-560 as indicated. Cells were arrested in mitosis with 100 nM nocodazole prior filming

and subsequently treated with 0.5 μ M reversine to silence the SAC. Each circle represents a single cell analyzed and at least 120 cells were analyzed per condition. Red bars indicate means ($m=$) and a t-test was used for statistical analysis. C) Representative still images from B. Scale bar represents 10 μ m. D-E) Venus-Cdc20 was purified from a stable HeLa cell line arrested in mitosis with nocodazole. The beads were incubated with increasing concentrations of recombinant FLAG-HA-BubR1 516-715 at room temperature for 1 hour and afterwards washed. The binding of endogenous BubR1 and Mad2 to Venus-Cdc20 was analyzed by western blot analysis and quantified using Licor technology. Representative of two independent experiments. F) HeLa cells were transfected with the indicated BubR1 constructs and before filming 200 nM taxol was added. The time from NEBD to mitotic exit was measured by analyzing the time-lapse movies and at least 50 cells were analyzed per condition. Medians ($m=$) are shown as red bars and a Mann-Whitney test was used for statistical analysis.

Figure 6. Dual role of the IC20BD in regulating the SAC.

At the kinetochore the IC20BD recruits Cdc20 to facilitate its interaction with Mad2 and potentially the formation of a partly assembled MCC. In the cytoplasm the IC20BD contributes to SAC silencing likely by competing for binding to Cdc20 with the N-terminal Mad3 homology region of BubR1. These two opposing activities of the IC20BD could make the SAC more dynamic and also couples MCC production or disassembly to BubR1 localization.

Figure 1.




Figure 2.




Figure 3.




Figure 4.




Figure 5.




Figure 6.

