

RESEARCH ARTICLE

Bioactive secondary metabolites from new endophytic fungus *Curvularia*. sp isolated from *Rauwolfia macrophylla*

Fatma Kaaniche^{1,2,3}, Abdelaaty Hamed^{1,4}, Ahmed S. Abdel-Razek^{1,5}, Daniel Wibberg⁶, Negera Abdissa⁷, Imene Zendah El Euch¹, Noureddine Allouche², Lotfi Mellouli³, Mohamed Shaaban^{1,8}, Norbert Sewald^{1*}

1 Department of Chemistry, Organic and Bioorganic Chemistry, Bielefeld University, Bielefeld, Germany, **2** Laboratory of Organic Chemistry, Natural Substances Team, Faculty of Sciences of Sfax, University of Sfax, Sfax, Tunisia, **3** Laboratory of Microorganisms and Biomolecules of the Centre of Biotechnology of Sfax-Tunisia, Sfax, Tunisia, **4** Chemistry Department, Faculty of Science, Al-Azhar University, Nasr City-Cairo, Egypt, **5** Microbial Chemistry Department, Genetic Engineering and Biotechnology Research Division, National Research Centre, Dokki-Giza, Egypt, **6** Center of Biotechnology(CeBiTec), Bielefeld University, Bielefeld, Germany, **7** Department of Chemistry, Jimma University, Jimma, Ethiopia, **8** Chemistry of Natural Compounds Department, Pharmaceutical and Drug Industries Research Division, National Research Centre, Dokki-Cairo, Egypt

* Norbert.sewald@uni-bielefeld.de


 OPEN ACCESS

Citation: Kaaniche F, Hamed A, Abdel-Razek AS, Wibberg D, Abdissa N, El Euch IZ, et al. (2019) Bioactive secondary metabolites from new endophytic fungus *Curvularia*. sp isolated from *Rauwolfia macrophylla*. PLoS ONE 14(6): e0217627. <https://doi.org/10.1371/journal.pone.0217627>

Editor: Vijai Gupta, Tallinn University of Technology, ESTONIA

Received: December 12, 2018

Accepted: May 15, 2019

Published: June 27, 2019

Copyright: © 2019 Kaaniche et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the manuscript and Supporting Information files.

Funding: This research work has been financed by the German Academic Exchange Service (DAAD) with funds from the German Federal Foreign Office in the frame of the Research Training Network "Novel Cytotoxic Drugs from Extremophilic Actinomycetes" (Project ID57166072), website: <https://www.daad.de/en/>. The Deutsche

Abstract

Over the last decades, endophytic fungi represent a new source of pharmacologically active secondary metabolites based on the underlying assumption that they live symbiotically within their plant host. In the present study, a new endophytic fungus was isolated from *Rauwolfia macrophylla*, a medicinal plant from Cameroon. The fungus showed a highest homology to *Curvularia* sp. based on complete nucleotide sequence data generated from the internal transcribed spacer (ITS) of ribosomal DNA region. Large scale fermentation, working-up and separation of the strain extract using different chromatographic techniques afforded three bioactive compounds: 2'-deoxyribolactone (**1**), hexylitaconic acid (**2**) and ergosterol (**3**). The chemical structures of compounds **1–3** were confirmed by 1 and 2D NMR spectroscopy and mass spectrometry, and comparison with corresponding literature data. Biologically, the antimicrobial, antioxidant activities and the acetylcholinesterase inhibitory of the isolated compounds were studied.

Introduction

Rauwolfia macrophylla (Apocynaceae) is a tree of about 12–15 m high occurring in Upper Guinea, Southern Nigeria, and Cameroon [1]. The stem and root barks were commonly used to treat malaria and other parasitic diseases in African indigenous medicine [2, 3]. Endophytes are microorganisms that inhabit the inner tissue of their hosts and perform various ecological relationships without showing visible host infection symptoms [4–6]. Isolation of three endophytic fungi namely *Aspergillus* sp., *Fusarium* sp., and *Purpureocillium lilacinum* from the

Forschungsgemeinschaft and the Open Access Publication Fund of Bielefeld University support for the Article Processing Charge.

Competing interests: The authors have declared that no competing interests exist.

stem bark of *Rauwolfia macrophylla* were recently reported [7]. The last endophytic fungus (*Purpureocillium lilacinum*) was among them specifically studied due to its potent antiprotozoal activity against *Leishmania donovani* (IC_{50} $0.174 \mu\text{g mL}^{-1}$); which was attributed to its production of purpureone, an ergochrome moiety [7]. It should be considered that different factors such as the plant organs, genotypic and geographic location influence the endophytic fungal community structure [8]. Endophytic fungi *Curvularia* sp. including *Curvularia trifolii* [9] and *Curvularia lunata* [10] provide a broad variety of bioactive secondary metabolites including polyketides and steroids. The genus *Curvularia* includes more than 40 species of saprophytes, endophytes and pathogens [11]. Most of these species are plant pathogens and give rise to losses in agricultural production [12].

In a continuing search for new endophytic fungi from medicinal plants as source of bioactive secondary metabolites, we isolated for the first time an endophytic fungus belonging to *Curvularia* sp. which has been characterized as *Curvularia* sp. T12 based on ITS sequencing (99% sequence identity with *Curvularia sorghina* BRIP 15900) from the stem bark of *Rauwolfia macrophylla* (Apocynaceae). An investigation of the fungus metabolites after its fermentation on solid-rice medium, working up and purification afforded three diverse classes of bioactive compounds. Identification of their structures on the bases of NMR and mass spectrometry assigned them as 2'-deoxyribolactone (1), hexylitaconic acid, (2) and ergosterol (3). The antibacterial, antioxidant, and acetylcholinesterase inhibitory activities of the compounds 1–3 were intensively reported as well.

Materials and methods

General

NMR spectra (^1H NMR, ^{13}C NMR, DEPT, COSY, HMQC, and HMBC) were measured on Bruker Avance DRX 500 spectrometer using standard pulse sequences and referenced to residual solvent signals. Column chromatography was carried out on silica gel 60 (0.040–0.063 mm, Merck, Darmstadt, Germany) and Sephadex LH-20 (GE Healthcare, Uppsala, Sweden) as stationary phases. ESI-MS was done on a Micromass AC-TOF micro mass spectrometer (Micromass, Agilent Technologies 1200 series, Tokyo, Japan). Analytical TLC was performed with pre-coated Merck silica gel 60 $_{\text{PF}254+366}$ (Merck, Darmstadt, Germany). R_f values of the strain extract bioactive compounds and visualization of their chromatograms was carried out with UV light (254 and 366 nm) followed by spraying with anisaldehyde-sulfuric acid and heating. Acetylcholine iodide, AChE from electric eel (type VI-S lyophilized powder), 2,2-diphenyl-1-picrylhydrazyl (DPPH), 2,2' azino-bis-(3-ethylbenzthiazoline-6-sulfonic acid (ABTS⁺), 5,5-dithiobis-2-nitrobenzoic acid (DTNB), BHA (butylated hydroxyanisole) and BHT (butylated hydroxytoluene), galanthamine were obtained from Sigma-Aldrich (Germany).

Strain material

The fungal strain T12 was isolated from the stem bark of *Rauwolfia macrophylla* (Apocynaceae) collected from Mount Kalla in Cameroon. The stem surface of the plant was sterilized using the methods described by Pimentel et al. (2006) [13]. The stems were cleaned with distilled and sterilised water for 10 min to remove impurities and surface debris. After air-drying, the cleaned stems were cut into small pieces and then sterilized under aseptic conditions using 70% ethanol for 30 s, 2.4% sodium hypochlorite solution for 4 min, and then 70% ethanol for 30 s. The plant samples were finally washed (3 ×) with sterile distilled water for 1 min. The surface-sterilized samples were then further cut into smaller pieces (1 cm^2) and aseptically placed in petri dishes containing sterile potato dextrose agar (PDA medium), supplemented with chloramphenicol (250 mg L^{-1}) to inhibit bacterial growth. All plates were incubated at room

temperature until mycelium grew out hyphal tips that were cut and transferred to PDA. The isolated strain was stored on PDA agar slants at 4 °C.

To identify the isolated endophytic fungal strain, the ITS region of the ribosomal DNA was analyzed by applying Sanger sequencing [14]. For DNA isolation, fungal cultures were grown in liquid potato dextrose broth (PDB) (24 g L⁻¹, pH 7) for 5–7 days at room temperature. After centrifugation (12 000 ×g, 10 min, at room temperature), the mycelium pellet was shock frozen and ground with a pestle in liquid nitrogen. The fungal DNA was isolated using the DNeasy Plant Mini kit (Qiagen, Hilden, Germany). PCR mixtures to amplify the ITS regions contained 2.5 μL of 10 × PCR buffer (Promega, Fitchburg, MA, USA), 0.5 μL of dNTP mixture (10 mmol·L⁻¹ each, NEB, Ipswich, MA, USA), 0.5 μL of each primer (10 μmol L⁻¹, ITS1 (5′-TCC GTA GGT GAA CCT GCG G-3′) and ITS4 (5′-TCC TCC GCT TAT TGA TAT GC-3′) (White et al., 1990) [15], 0.5 μL of *Pfu* DNA polymerase (3 U μL⁻¹, Promega), and 2 μL of isolated DNA in a total volume of 25 μL [15]. PCRs were carried out and the resulting products were purified using the QIAquick PCR purification kit (Qiagen) and sequenced on an Applied Biosystems Abi-Prism 3730 sequencer using BigDye-terminator chemistry at the Center for Biotechnology (CeBiTec, Bielefeld, Germany). For taxonomic classification, the resulting sequence was compared with the NT database (NCBI) by means of blastn and default settings. Sequences of the best hits were used for the construction of a phylogenetic tree. Phylogenetic analyses were conducted using MEGA version 7.0.2 [15, 16]. The phylogenetic tree was constructed by means of neighbourjoining (NJ) algorithm [17] using Kimura 20-parameter distance. The robustness of the inferred tree was evaluated by bootstrap (1000 replications).

Fermentation, extraction and isolation

Inocula of the endophytic fungus *Curvularia* sp. T12 were introduced to five 1L-flasks, each containing sterilized solid-rice medium (100 g rice, 100 mL distilled water, followed by sterilization in an autoclave). The flasks were incubated at room temperature (25 °C) for 21 days. After fermentation, the obtained cultures were soaked in methanol (3 × 0.5 L per each flask for 24 h), followed by filtration. The filtered methanol extract was concentrated under vacuum at 40 °C to afford 60 g of brown crude extract.

The extract was then applied to fractionation by silica gel column chromatography (80 × 10 cm) and eluted with petroleum ether-EtOAc-MeOH gradient with gradually increasing the polarity, affording six major fractions according to TLC monitoring: FI (2.9 g), FII (2.1 g), FIII (6.0 g), FIV (3.1 g), FV (5.0 g), FVI (8.1 g). Application of FI to silica gel column (60×1.5 cm) using cyclohexane-DCM gradient afforded ergosterol (**3**, 10 mg) as colorless solid. Further purification of fraction III was done on silica gel column chromatography eluted with petroleum ether/EtOAc (1:1) and followed by gel filtration using Sephadex LH-20 (DCM/MeOH, 1:1) resulting in colourless solids of 2′-deoxyriboflavone (**1**, 1.3 mg) and hexylitaconic acid (**2**, 2.2 mg) (Fig 1).

2′-Deoxyriboflavone (1): colourless solid, UV-non-absorbing, showed yellow-green colour on spraying with anisaldehyde/sulfuric acid; $R_f = 0.52$ (DCM/5% MeOH); ¹H NMR (500 MHz, CD₃OD) and ¹³C NMR (125 MHz, CD₃OD) (S1 Table, S1–S6 Figs); (+)-ESI-MS $m/z = 155$ ([M+Na]⁺); (-)-ESI-MS $m/z = 131$ ([M-H]⁻).

Hexylitaconic acid (2): colourless solid, UV-absorbing (254 nm), showing brown colouration on spraying with anisaldehyde/sulphuric acid and heating; $R_f = 0.39$ (5% MeOH in DCM); ¹H NMR (500 MHz, CDCl₃) and ¹³C NMR (125 MHz, CDCl₃) (S2 Table, S7–S12 Figs); (+)-ESI-MS m/z (%) = 237 ([M+Na]⁺); (-)-ESI-MS m/z (%) = 213 ([M-H]⁻).

Ergosterol (3): Colourless amorphous solid, UV-absorbing (254 nm), purple to dark blue colour on spraying with anisaldehyde/sulfuric acid; $R_f = 0.48$ (3% MeOH in DCM); ¹H NMR




Fig 1. Structure of the compounds (1–3).

<https://doi.org/10.1371/journal.pone.0217627.g001>

(500 MHz, CDCl_3 , [S13 Fig](#)) δ = 5.50 (dd, J = 5.7, 2.5 Hz, 1H), 5.31 (dt, J = 5.6, 2.8 Hz, 1H), 5.16 (dd, J = 15.2, 7.0 Hz, 1H), 5.10 (dd, J = 15.3, 7.7 Hz, 1H), 3.57 (tt, J = 11.2, 4.0 Hz, 1H), 2.40 (ddd, J = 14.3, 4.8, 2.4 Hz, 1H), 2.22 (ddt, J = 14.2, 11.8, 2.3 Hz, 1H), 2.03–1.93 (m, 2H), 1.91 (dddd, J = 11.9, 7.2, 4.6, 2.2 Hz, 1H), 1.80 (tdd, J = 13.9, 12.8, 6.4, 4.6 Hz, 4H), 1.73–1.48 (m, 4H), 1.48–1.36 (m, 2H), 1.36–1.13 (m, 6H), 0.97 (d, J = 6.6 Hz, 3H), 0.88 (s, 3H), 0.85 (d, J = 6.9 Hz, 3H), 0.76 (t, J = 7.2 Hz, 6H), 0.56 (s, 3H). ^{13}C NMR (125 MHz, CDCl_3 , [S14 Fig](#)) δ 141.3, 139.8, 135.6, 132.0, 119.6, 116.3, 70.4, 55.7, 54.6, 46.2, 42.8, 40.8, 40.4, 39.1, 38.4, 37.0, 33.1, 32.0, 28.3, 23.0, 21.1, 20.0, 19.6, 17.6, 16.3, 12.0. (+)-ESI-MS m/z (%) = 419 ($[\text{M}+\text{Na}]^+$); (-)-ESI-MS m/z (%) = 395 ($[\text{M}-\text{H}]^-$).

***In vitro* antibacterial assay**

The antimicrobial activity of the isolated compounds was studied using micro-dilution method with some modification [18]. The bacterial strains, *Escherichia coli* (DSMZ1058), *Micrococcus luteus* (DSMZ1605), *Pseudomonas agarici* (DSMZ11810), and *Staphylococcus warneri* (DSMZ20036) were grown on a nutrient agar medium (3 g L^{-1} beef extract, 10 g L^{-1} peptone, and 20 g L^{-1} agar) and the pH was adjusted to 7.2. The test was performed by inoculating a suspension of the overnight tested microorganism ($\text{DO}_{600} = 0.05 \sim 0.1$) on the nutrient agar medium. The investigated compounds were diluted to 100 μL of broth in the first well of 96 wells of microtiter plate. The concentrations of the compounds were finally in the range of 250 mg mL^{-1} to 0.01 mg mL^{-1} . Microtiter plates containing *Escherichia coli* and *Staphylococcus warneri* were incubated at 37 °C and those containing *Micrococcus luteus* and *Pseudomonas agarici* were incubated at 30 °C for 24 h. The minimum inhibition concentration (MIC) was calculated according to our previous work [19].

***In vitro* acetylcholinesterase inhibition assay**

AChE inhibitory activity was measured according to the spectrophotometric method of Ellman with slight modifications [20]. Electric eel AChE was used, while acetylthiocholine iodide (ATCI) was employed as substrate. 5,5'-Dithiobis-(2-nitrobenzoic acid) (DTNB) was used for the determination of released thiocholine. Briefly, in this method, 100 μL of Tris buffer at 50 mM (pH 8.0), 30 μL of sample (100 $\mu\text{g}/\text{mL}$) and 5 μL of AChE enzyme (0.5 U/mL) were added in a 96 well microplate and incubated for 10 min at 25 °C. Then, 27 μL of DTNB (3 mM) and 23 μL of substrate (1.16 mM) were added. Hydrolysis of ATCI was monitored by the formation of the yellow 5-thio-2-nitrobenzoate anion as a result of the reaction of DTNB with thiocholines, at 405 nm utilizing a 96 well microplate reader (Thermo Scientific/Varioskan Flash, Germany). The reaction progress was monitored until the AChE activity decreased, and then the reaction has been stopped. Inhibition percentage was calculated according to Michaelis–Menten model by using *EZ-Fit*. The IC_{50} values were determined by monitoring the inhibition effects of various concentrations of under investigation compounds and calculated *EZ-Fit*, Enzyme Kinetics Program (Perrella Scientific In., Amherst, USA). Galanthamine dissolved in 10% DMSO in Tris buffer were used as standard drugs at 10 $\mu\text{g mL}^{-1}$ and 1 mg mL^{-1} concentrations.

***In vitro* antioxidant assays**

DPPH and (ABTS⁺) free radical scavenging activities. The DPPH and ABTS⁺ assays [21, 22] were used to evaluate the free radical scavenging activities of the compounds. Briefly, the compounds were dissolved in 10% DMSO and diluted at different concentrations of 25 to 1 $\mu\text{g mL}^{-1}$. For the DPPH assay, 500 μL of a 4% (w/v) solution of DPPH radical in methanol was mixed with 500 μL of samples under investigation. The mixture was incubated for 30 min in the dark at room temperature. The scavenging capacity was determined spectrophotometrically by monitoring the decrease in absorbance at 517 nm against a blank. For the ABTS⁺ assay, the reaction mixture was containing ABTS solution (7 mM) with 2.45 mM $\text{K}_2\text{S}_2\text{O}_8$. The mixture was let to stand for 15 minutes in the dark at room temperature. The solution was diluted with ethanol to obtain the absorbance of 0.7 ± 0.2 units at 734 nm. In order to measure the antiradical activity of extracts and standards, 20 μL of each sample at various concentrations was added to 180 μL of ABTS⁺ and vortexed for 30 s, and then incubated for 6 min in the dark at room temperature. The absorbance was measured spectrophotometrically at 734 nm. The percentage of antioxidant activity had been calculated as follows:

$$\text{Antioxidant activity (\%)} = [(A_{\text{control}} - A_{\text{sample}}) / A_{\text{control}}] \times 100.$$

BHA and BHT were used as positive controls.

Data analyses

Curve-fitting analyses was made using GraphPad Prism version 4.03 (GraphPad Software, San Diego, CA, USA) [23].

Results and discussion

Taxonomical characterization of the producing fungus

An endophytic fungus was isolated from *Rauwolfia macrophylla* (Apocynaceae). For taxonomic classification of this endophyte, its ITS sequences were amplified by applying PCR, followed by Sanger sequencing and a BLAST-based approach. The obtained sequences showed hits against the databases with high identities (99%–100%) and sequence coverage (97%–100%). Best hits




Fig 2. Phylogenetic tree of the endophyte strain T12.

<https://doi.org/10.1371/journal.pone.0217627.g002>

were selected to construct a phylogenetic tree (Fig 2). Based on the phylogenetic and the BLAST analyses, the fungal strain named T12 was assigned as *Curvularia* sp., since the closest available strains in the NCBI database are belonging to *Curvularia* spp, and the most similar one is *Curvularia sorghina* (99% sequence identity).

Fermentation and structural identification

The extract produced by the endophytic fungus *Curvularia* sp. T12 was found to exhibit an antimicrobial inhibition zones (IZ) against *Escherichia coli* (IZ = 14.7 ± 1.0 mm), *Micrococcus luteus* (IZ = 14.0 ± 0.5 mm), *Pseudomonas agarici* (IZ = 12.3 ± 0.6 mm), and *Staphylococcus warneri* (IZ = 17.0 ± 1.5 mm), which represent a strong antimicrobial activities against these tested microbes. Accordingly, the fungus T12 was subjected to large scale fermentation to produce reasonable amounts of crude extract, After then, the crude extract has been applied for working up and purification of its desired metabolites using a series of chromatographic techniques, starting with silica gel column chromatography and ended by Sephadex LH-20. Based on that, three diverse bioactive compounds (1–3) were obtained.

The chemical structures of the isolated metabolites were determined as 2'-deoxyribolactone (1) [24–27], hexylitaconic acid (2) [28–30], ergosterol (3) [31–33] based on comparing their

1D & 2D nuclear magnetic resonance and ESI-MS data with those previously reported in literature (see experimental section and supplementary data).

Ribolactone derivatives. Synthetically developed and natural ribolactones derivatives are included as building blocks of several antiviral regimens [34]. The diversity and availability of these relatively cheap chiral compounds has led to their use as starting materials for the design and syntheses of naturally occurring compounds and biologically important molecules [35, 36].

Hexylitaconic acid derivatives. Itaconic acid derivatives (IA) represent promising compounds that have a wide range of applications and can be obtained in large scale using fermentation processes. One of the most important uses of these biomonomer is the environmentally sustainable production of biopolymers. IA is used worldwide in the industrial synthesis of resins such as polyesters, plastics and artificial glass and in the preparation of bioactive compounds in the agriculture, pharmacy and medicine sectors. IA also provides possibilities for selective enzymatic transformations to create useful polyfunctional building blocks [37]. IAs were found to possess significant anti-inflammatory and analgesic activities as well [38]. Moreover, hexylitaconic acid (2) has been reported as inhibitor of p53-HDM2 interaction [39].

Ergosterol and structural analogues. Ergosterol is the major product of sterol biosynthesis in fungi, and also in some trypanosomes. Most of the current antifungal agents interfere with ergosterol-function in some way, either through inhibition of various steps in ergosterol biosynthesis (allylamines, azoles, morpholines) or by complexing directly with membrane ergosterol (polyenes). Biologically, ergosterol was reported to have several effective activities, for example it has an inhibitory effect on the activity of mucosal-type mast cells [40], anti-*Trypanosoma cruzi* activity against trypomastigotes (IC₅₀ 51.3 µg/mL) [41], and antifungal activity as well. More recently, ergosterol derivatives were reported as Anti-tumor and Anti-angiogenic agents [42].

Biological activity studies

The isolated compounds 1–3 were evaluated for antimicrobial activity against four bacterial strains (*Escherichia coli*, *Micrococcus luteus*, *Pseudomonas agarici* and *Staphylococcus warneri*). The antibacterial activity test (Table 1) indicated that compounds 1 and 2 have antibacterial activities against all the tested bacterial strains with MIC ranging between 0.17 µg mL⁻¹ and 0.58 µg ml⁻¹.

The isolated compounds (1–3) were evaluated also as inhibitors of acetylcholinesterase. They showed good inhibitory potential towards acetylcholinesterase (Table 2) with IC₅₀ values of 1.93, 1.54 and 1.52 µM, respectively, while the selected reference galanthamine showed an IC₅₀ value of 0.5 µM and 0.01 µM for tacrine.

Antioxidant activity of the three compounds was measured according to DPPH and ABTS⁺ scavenging methods [43]. The results (Table 3) showed that compounds 1–3 were active with

Table 1. MIC (µg ml⁻¹) of the isolated compounds (1–3).

	<i>Pseudomonas agarici</i>	<i>Escherichia coli</i>	<i>Staphylococcus warneri</i>	<i>Micrococcus luteus</i>
	MIC (µg/ml)			
1	0.19	0.30	0.17	0.32
2	0.29	0.58	0.58	0.29
3	-	-	-	-
Gentamicin ^a	0.01	0.01	0.01	0.01

MIC = Minimal inhibition concentration, Minimal, (*) = European Pharmacopoeia (EP) Reference Standard.

<https://doi.org/10.1371/journal.pone.0217627.t001>

Table 2. Acetylcholinesterase inhibition of compounds (1–3).

	1	2	3	Galanthamine*	Tacrine*
IC ₅₀ (μM)	1.93	1.54	1.52	0.5	0.01

IC₅₀ = Concentration of acetylcholinesterase inhibitor drug that gives half maximal response, (*) = European Pharmacopoeia (EP) Reference Standard.

<https://doi.org/10.1371/journal.pone.0217627.t002>

Table 3. Antioxidant and radical scavenging capacity of the compounds (1–3).

	1	2	3	BHA*	BHT*
	EC ₅₀				
Antioxidant activity (DPPH*)	0.66 ± 0.02	0.56 ± 0.02	1.09 ± 0.02	0.13	0.46
Antiradical activity (ABTS*)	0.49 ± 0.02	0.88 ± 0.02	0.83 ± 0.02	0.01	0.06

EC₅₀ = Concentration of an antioxidant drug that gives half maximal response, (*) = European Pharmacopoeia (EP) Reference Standard.

<https://doi.org/10.1371/journal.pone.0217627.t003>

EC₅₀ values 0.66, 0.56 and 1.09 μM, respectively in DPPH assay and 0.49, 0.88 and 0.83 μM in ABTS assay. Compounds **1** and **2** exhibited close EC₅₀ values to those of the reference BHT (EC₅₀ = 0.46 μM in DPPH assay), reflecting the high scavenging ability of the compounds.

Numerous diverse bioactive compounds have been recently isolated from the endophyte *Curvularia* spp. indicating their importance as talented and major sources of bioactive molecules [9, 44]. However, the fungal metabolites have increased the interest in biomolecules activity studies [10].

Conclusion

The new endophytic fungus *Curvularia* sp. T12 was isolated from the medicinal plant *Rauwolfia macrophylla*. Its Large scale fermentation and working up of the crude extract, led to the isolation of 2'-deoxyribolactone (**1**), hexylitaconic acid (**2**) and ergosterol (**3**). Compounds **1–3** displayed antimicrobial, antioxidant and acetylcholinesterase inhibition activities. To our knowledge, these biological activities have never been assayed for these compounds. These findings are greatly highlighting the importance of endophytic fungi as source of bioactive secondary metabolites.

Supporting information

S1 Table. ¹³C (125 MHz) and ¹H (500 MHz) NMR data of 2'-deoxyribolactone (**1**) in CD₃OD.
(PDF)

S2 Table. ¹³C (125 MHz) and ¹H (500 MHz) NMR data of hexylitaconic acid (**2**) in CDCl₃.
(PDF)

S1 Fig. ¹H NMR spectrum (CD₃OD, 500 MHz) of 2'-deoxyribolactone (**1**).
(PDF)

S2 Fig. ¹³C NMR spectrum (CD₃OD, 125MHz) of 2'-deoxyribolactone (**1**).
(PDF)

S3 Fig. DEPT spectrum (CD₃OD, 125MHz) of 2'-deoxyribolactone (**1**).
(PDF)

S4 Fig. H,H-COSY spectrum (CD₃OD, 500 MHz) of 2'-deoxyribolactone (1).
(PDF)

S5 Fig. HMQC spectrum (CD₃OD, 500 MHz) of 2'-deoxyribolactone (1).
(PDF)

S6 Fig. HMBC spectrum (CD₃OD, 500 MHz) of 2'-deoxyribolactone (1).
(PDF)

S7 Fig. ¹H NMR spectrum (CDCl₃, 500 MHz) of Hexylitaconic acid (2).
(PDF)

S8 Fig. ¹³C NMR spectrum (CDCl₃, 125MHz) of Hexylitaconic acid (2).
(PDF)

S9 Fig. DEPT spectrum (CDCl₃, 125MHz) of Hexylitaconic acid (2).
(PDF)

S10 Fig. H,H-COSY spectrum (CDCl₃, 500 MHz) of Hexylitaconic acid (2).
(PDF)

S11 Fig. HMQC spectrum (CDCl₃, 500 MHz) of Hexylitaconic acid (2).
(PDF)

S12 Fig. HMBC spectrum (CDCl₃, 500 MHz) of Hexylitaconic acid (2).
(PDF)

S13 Fig. ¹H NMR spectrum (CDCl₃, 500 MHz) of ergosterol (3).
(PDF)

S14 Fig. ¹³C NMR spectrum (CDCl₃, 125MHz) of ergosterol (3).
(PDF)

S1 File. Raw data.
(RAR)

Acknowledgments

The authors are thankful to the NMR and MS Departments in Bielefeld University for the spectral measurements. We thank Prof. Dr. Bruno Lenta Ndjakou, Chemistry Department-, Yaoundé University, Cameroon for donating the fungal extracts as well as Mr. Sebastian Gohlke for his assistance regarding HPLC chromatography analysis. We thank also Carmela Michalek for biological activity tests and Marco Wißbrock for informatic technical assistance. The bioinformatics support of the BMBF-funded project "Bielefeld-Gießen Resource Center for Microbial Bioinformatics-BiGi (Grant number 031A533)" within the German Network for Bioinformatics Infrastructure (de.NBI) is gratefully acknowledged. This research work has been financed by the German Academic Exchange Service (DAAD) with funds from the German Federal Foreign Office in the frame of the Research Training Network "Novel Cytotoxic Drugs from Extremophilic Actinomycetes" (Project ID57166072).

Author Contributions

Conceptualization: Fatma Kaaniche, Abdelaaty Hamed, Daniel Wibberg.

Data curation: Fatma Kaaniche, Abdelaaty Hamed, Daniel Wibberg.

Formal analysis: Abdelaaty Hamed, Daniel Wibberg.

Funding acquisition: Imene Zendah El Euch, Nobert Sewald.

Investigation: Abdelaaty Hamed, Ahmed S. Abdel-Razek.

Methodology: Fatma Kaaniche.

Project administration: Lotfi Mellouli, Mohamed Shaaban, Nobert Sewald.

Software: Daniel Wibberg.

Supervision: Negera Abdissa, Imene Zendah El Euch, Noureddine Allouche, Lotfi Mellouli, Mohamed Shaaban, Nobert Sewald.

Validation: Fatma Kaaniche, Abdelaaty Hamed, Ahmed S. Abdel-Razek.

Visualization: Fatma Kaaniche.

Writing – original draft: Fatma Kaaniche, Abdelaaty Hamed, Ahmed S. Abdel-Razek, Daniel Wibberg, Negera Abdissa.

Writing – review & editing: Mohamed Shaaban, Nobert Sewald.

References

1. Amer MMA, Court WE. Stem Bark Alkaloids of *Rauwolfia-Macrophylla*. *Planta Medica*. 1980; 40(S1):8–12. <https://doi.org/10.1055/s-2008-1074994>
2. Ndjakou Lenta B, Vonthron-Sénécheau C, Fongang Soh R, Tantangmo F, Ngouela S, Kaiser M, et al. In vitro antiprotozoal activities and cytotoxicity of some selected Cameroonian medicinal plants. *Journal of Ethnopharmacology*. 2007; 111(1):8–12. <https://doi.org/10.1016/j.jep.2006.10.036>. <https://doi.org/10.1016/j.jep.2006.10.036> PMID: 17141994
3. Sabrin RMI, Gamal AM, Rwaida AAH, Amal AE-K, Mohamed FZ. Potential Anti-Malarial Agents from Endophytic Fungi: A Review. *Mini-Reviews in Medicinal Chemistry*. 2018; 18(13):1110–32. <http://dx.doi.org/10.2174/1389557518666180305163151>. <https://doi.org/10.2174/1389557518666180305163151> PMID: 29512460
4. Xia Y, Amna A, Opiyo SO. The culturable endophytic fungal communities of switchgrass grown on a coal-mining site and their effects on plant growth. *PLoS One*. 2018; 13(6):e0198994. <https://doi.org/10.1371/journal.pone.0198994> PMID: 29902231
5. Hardoim PR, van Overbeek LS, Berg G, Pirttilä AM, Compant S, Campisano A, et al. The Hidden World within Plants: Ecological and Evolutionary Considerations for Defining Functioning of Microbial Endophytes. *Microbiology and Molecular Biology Reviews*. 2015; 79(3):293. <https://doi.org/10.1128/MMBR.00050-14> PMID: 26136581
6. Kaul S, Gupta S, Ahmed M, Dhar MK. Endophytic fungi from medicinal plants: a treasure hunt for bioactive metabolites. *Phytochemistry Reviews*. 2012; 11(4):487–505. <https://doi.org/10.1007/s11101-012-9260-6>
7. Lenta Bruno N, Ngatchou J, Frese M, Ladoh-Yemeda F, Voundi S, Nardella F, et al. Purpureone, an antileishmanial ergochrome from the endophytic fungus *Purpureocillium lilacinum*. *Zeitschrift für Naturforschung B*. 2016; 71(11):1159–67. <https://doi.org/10.1515/znb-2016-0128>
8. Silva FdA, Liotti RG, Boleti APdA, Reis ÉdM, Passos MBS, dos Santos EL, et al. Diversity of cultivable fungal endophytes in *Paullinia cupana* (Mart.) Ducke and bioactivity of their secondary metabolites. *PLoS One*. 2018; 13(4):e0195874. <https://doi.org/10.1371/journal.pone.0195874> PMID: 29649297
9. Samanthi K, Wickramarachchi S, Wijeratne E, Paranagama P. Two new bioactive polyketides from *Curvularia trifolii*, an endolichenic fungus isolated from *Usnea* sp., in Sri Lanka. *Journal of the National Science Foundation of Sri Lanka*. 2015; 43(3):217–24. <http://doi.org/10.4038/jnsfsr.v43i3.7950>.
10. Avinash KS, Ashwini HS, Babu HNR, Krishnamurthy YL. Antimicrobial Potential of Crude Extract of *Curvularia lunata*, an Endophytic Fungi Isolated from *Cymbopogon caesius*. *Journal of Mycology*. 2015; 2015:4. <https://doi.org/10.1155/2015/185821>
11. Zhou L, Zhao J, Shan T, Cai X, Peng Y. Spirobisnaphthalenes from Fungi and their Biological Activities. *Mini-Reviews in Medicinal Chemistry*. 2010; 10(10):977–89. <http://dx.doi.org/10.2174/138955710792007178>. PMID: 20491650

12. Brun R, Don R, Jacobs RT, Wang MZ, Barrett MP. Development of novel drugs for human African trypanosomiasis. *Future Microbiology*. 2011; 6(6):677–91. <https://doi.org/10.2217/fmb.11.44> PMID: 21707314
13. Pimentel IC, Glienke-Blanco C, Gabardo J, Stuart RM, Azevedo JL. Identification and colonization of endophytic fungi from soybean (*Glycine max* (L.) Merrill) under different environmental conditions. *Brazilian Archives of Biology and Technology*. 2006; 49:705–11.
14. Hidayat I, Radiastuti N, Rahayu G, Achmadi S, Okane I. Endophytic fungal diversity from *Cinchona calisaya* based on phylogenetic analysis of the ITS ribosomal DNA sequence data. *Current Research in Environmental & Applied Mycology*. 2016; 6(2):132–42. <https://doi.org/10.5943/cream/6/2/7>
15. Neubauer PR, Widmann C, Wibberg D, Schröder L, Frese M, Kottke T, et al. A flavin-dependent halogenase from metagenomic analysis prefers bromination over chlorination. *PLoS One*. 2018; 13(5): e0196797. <https://doi.org/10.1371/journal.pone.0196797> PMID: 29746521
16. Kumar S, Stecher G, Tamura K. MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0 for Big-ger Datasets. *Molecular Biology and Evolution*. 2016; 33(7):1870–4. <https://doi.org/10.1093/molbev/msw054> PMID: 27004904
17. Saitou N, Nei M. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Molecular Biology and Evolution*. 1987; 4(4):406–25. <https://doi.org/10.1093/oxfordjournals.molbev.a040454> PMID: 3447015
18. Sellem I, Kaaniche F, Chakchouk A, Mellouli L. Anti-oxidant, antimicrobial and anti-acetylcholinesterase activities of organic extracts from aerial parts of three Tunisian plants and correlation with polyphenols and flavonoids contents. *Bangladesh Journal of Pharmacology*. 2016; 11(2):531–44. <https://doi.org/10.3329/bjp.v11i2.26001>
19. Hamed A, Abdel-Razek AS, Omran DA, El-Metwally MM, El-Hosari DG, Frese M, et al. Terretinin O: a new meroterpenoid from *Aspergillus terreus*. *Natural Product Research*. 2019:1–10. <https://doi.org/10.1080/14786419.2018.1544977> PMID: 30602325
20. Ellman GL, Courtney KD, Andres V, Featherstone RM. A new and rapid colorimetric determination of acetylcholinesterase activity. *Biochemical Pharmacology*. 1961; 7(2):88–95. [https://doi.org/10.1016/0006-2952\(61\)90145-9](https://doi.org/10.1016/0006-2952(61)90145-9).
21. Molyneux P. The use of the stable free radical diphenylpicrylhydrazyl (DPPH) for estimating antioxidant activity. *Songklanakarinn Journal of Science and Technology*. 2004; 26(2):211–9.
22. Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice Evans C. Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radical Biology & Medicine*. 1999; 26(9–10):1231–7. [https://doi.org/10.1016/s0891-5849\(98\)00315-3](https://doi.org/10.1016/s0891-5849(98)00315-3)
23. Abdissa N, Gohlke S, Frese M, Sewald N. Cytotoxic Compounds from *Aloe megalacantha*. *Molecules*. 2017; 22(7). <https://doi.org/10.3390/molecules22071136> PMID: 28686200
24. Tarawneh AH, León F, Radwan MM, Rosa LH, Cutler SJ. Secondary metabolites from the fungus *Emericella nidulans*. *Nat Prod Commun*. 2013; 8(9):1285–8. PMID: 24273867.
25. Liu X, Dong M, Chen X, Jiang M, Lv X, Zhou J. Antimicrobial activity of an endophytic *Xylaria* sp. YX-28 and identification of its antimicrobial compound 7-amino-4-methylcoumarin. *Applied Microbiology and Biotechnology*. 2008; 78(2):241–7. <https://doi.org/10.1007/s00253-007-1305-1> PMID: 18092158
26. Arunpanichlert J, Rukachaisirikul V, Phongpaichit S, Supaphon O, Sakayaroj J. Xylariphilone: a new azaphilone derivative from the seagrass-derived fungus *Xylariales* sp. PSU-ES163. *Natural Product Research*. 2016; 30(1):46–51. <https://doi.org/10.1080/14786419.2015.1032282> PMID: 25880133
27. Salat K, Librowski T, Moniczewski A, Stanisz-Wallis K, Wieckowski K, Malawska B. Analgesic, antioedematous and antioxidant activity of γ -butyrolactone derivatives in rodents. *Behavioural Pharmacology*. 2012; 23(4):407–16. <https://doi.org/10.1097/FBP.0b013e3283566042> PMID: 22785386
28. Isogai A, Washizu M, Kondo K, Murakoshi S, Suzuki A. Isolation and Identification of (+)-Hexylitaconic Acid as a Plant Growth Regulator. *Agricultural and Biological Chemistry*. 1984; 48(10):2607–9. <https://doi.org/10.1271/bbb1961.48.2607>
29. Klemke C, Kehraus S, Wright AD, König GM. New Secondary Metabolites from the Marine Endophytic Fungus *Apiospora montagnei*. *Journal of Natural Products*. 2004; 67(6):1058–63. <https://doi.org/10.1021/np034061x> PMID: 15217297
30. Nakahashi A, Miura N, Monde K, Tsukamoto S. Stereochemical studies of hexylitaconic acid, an inhibitor of p53–HDM2 interaction. *Bioorganic & Medicinal Chemistry Letters*. 2009; 19(11):3027–30. <https://doi.org/10.1016/j.bmcl.2009.04.057>.
31. Torres S, Cajas D, Palfner G, Astuya A, Aballay A, Pérez C, et al. Steroidal composition and cytotoxic activity from fruiting body of *Cortinarius xiphidipus*. *Natural Product Research*. 2017; 31(4):473–6. <https://doi.org/10.1080/14786419.2016.1185717> PMID: 27198920

32. Lee NK, Aan B-Y. Optimization of ergosterol to vitamin D(2) synthesis in *Agaricus bisporus* powder using ultraviolet-B radiation. *Food Sci Biotechnol*. 2016; 25(6):1627–31. <https://doi.org/10.1007/s10068-016-0250-0> PMID: 30263454.
33. Mattila P, Lampi A-M, Ronkainen R, Toivo J, Piironen V. Sterol and vitamin D2 contents in some wild and cultivated mushrooms. *Food Chemistry*. 2002; 76(3):293–8. [https://doi.org/10.1016/S0308-8146\(01\)00275-8](https://doi.org/10.1016/S0308-8146(01)00275-8).
34. De Schutter C, Sari O, Coats SJ, Amblard F, Schinazi RF. Synthesis of (2S)-2-Chloro-2-fluororibolactone via Stereoselective Electrophilic Fluorination. *The Journal of Organic Chemistry*. 2017; 82(24):13171–8. <https://doi.org/10.1021/acs.joc.7b02245> PMID: 29140703
35. Witczak ZJ. Thio sugars: biological relevance as potential new therapeutics. *Current Medicinal Chemistry*. 1999; 6(2):165–78. PMID: 10189230.
36. Pal P, Shaw AK. The evolution of comprehensive strategies for furanoid glycal synthesis and their applications. *RSC Advances*. 2017; 7(42):25897–963. <https://doi.org/10.1039/C6RA28598H>
37. Okabe M, Lies D, Kanamasa S, Park EY. Biotechnological production of itaconic acid and its biosynthesis in *Aspergillus terreus*. *Applied Microbiology and Biotechnology*. 2009; 84(4):597–606. <https://doi.org/10.1007/s00253-009-2132-3> PMID: 19629471
38. Antia BS, Aree T, Kasetrathat C, Wiyakrutta S, Ekpa OD, Ekpe UJ, et al. Itaconic acid derivatives and diketopiperazine from the marine-derived fungus *Aspergillus aculeatus* CRI322-03. *Phytochemistry*. 2011; 72(8):816–20. <https://doi.org/10.1016/j.phytochem.2011.02.013> PMID: 21397285
39. Tsukamoto S, Yoshida T, Hosono H, Ohta T, Yokosawa H. Hexylitaconic acid: A new inhibitor of p53–HDM2 interaction isolated from a marine-derived fungus, *Arthrinium* sp. *Bioorganic & Medicinal Chemistry Letters*. 2006; 16(1):69–71. <https://doi.org/10.1016/j.bmcl.2005.09.052>.
40. Kageyama-Yahara N, Wang P, Wang X, Yamamoto T, Kadowaki M. The Inhibitory Effect of Ergosterol, a Bioactive Constituent of a Traditional Japanese Herbal Medicine Saireito on the Activity of Mucosal-Type Mast Cells. *Biological and Pharmaceutical Bulletin*. 2010; 33(1):142–5. <https://doi.org/10.1248/bpb.33.142> PMID: 20045953
41. Alexandre TR, Lima ML, Galuppo MK, Mesquita JT, do Nascimento MA, Dos Santos AL, et al. Ergosterol isolated from the basidiomycete *Pleurotus salmoneostramineus* affects *Trypanosoma cruzi* plasma membrane and mitochondria. *J Venom Anim Toxins Incl Trop Dis*. 2017; 23:30-. <https://doi.org/10.1186/s40409-017-0120-0> PMID: 28572816.
42. Chen S, Yong T, Zhang Y, Su J, Jiao C, Xie Y. Anti-tumor and Anti-angiogenic Ergosterols from *Ganoderma lucidum*. *Front Chem*. 2017; 5:85. <https://doi.org/10.3389/fchem.2017.00085> PMID: 29164102.
43. Lü J-M, Lin PH, Yao Q, Chen C. Chemical and molecular mechanisms of antioxidants: experimental approaches and model systems. *Journal of Cellular and Molecular Medicine*. 2010; 14(4):840–60. <https://doi.org/10.1111/j.1582-4934.2009.00897.x> PMID: 19754673
44. Mondol MAM, Farthouse J, Islam MT, Schüffler A, Laatsch H. Metabolites from the Endophytic Fungus *Curvularia* sp. M12 Act as Motility Inhibitors against *Phytophthora capsici* Zoospores. *Journal of Natural Products*. 2017; 80(2):347–55. <https://doi.org/10.1021/acs.jnatprod.6b00785> PMID: 28195475