

---

1 **A taxonomic revision of *Noriopterus complicidens* (Young, 1973)**  
2 **and Asian members of the Dsungaripteridae**

3

4 **\*D.W.E. Hone<sup>1</sup>**

5 **S. Jiang<sup>2</sup>**

6 **X. Xu<sup>2</sup>**

7

8 1. Queen Mary University of London, Mile End Road, London, E14NS, UK.

9 2. Institute of Vertebrate Paleontology and Paleoanthropology, Xizhimenwai

10 Dajie, 100044, Beijing, China.

11

12 \*Correspondence: [d.hone@qmul.ac.uk](mailto:d.hone@qmul.ac.uk)

13

14

15

16 Taxonomy of Asian Dsungaripteridae

17

18

19

20 **Abstract:**

21 After being inaccessible for a number of years, the holotype and other

22 specimens of the dsungaripterid pterodactyloid pterosaur *Noriopterus*

23 *complicidens* (Young, 1973) are again available for study. Numerous taxa

24 assigned to the Dsungaripteridae have been described since the erection of

---

1 *Noriopteris*, but with limited comparisons to this genus. Based on the information  
2 from Young's original material here we revise the taxonomic identity of *N.*  
3 *complicidens* and that of other Asian dsungaripterids. We conclude that *N.*  
4 *complicidens* is likely distinct from the material recovered from Mongolia and  
5 this latter material should be placed in a separate genus.

6

7

8 **Keywords:** Pterosauria, *Noriopteris*, *Dsungaripteris*, Dsungaripteroidea, China,  
9 Mongolia

10

11

12

13 The dsungaripterid pterosaurs are a group of derived pterodactyloids  
14 that are characterized by having toothless jaw tips (Kellner, 2003; Unwin,  
15 2003). A number of taxa also show expansions of the bone around the tooth  
16 alveoli such that the jaw is swollen at the bases of the teeth, or the teeth may  
17 even be covered with bone entirely (e.g. see Martill et al., 2000). Dsungaripterids  
18 also have characteristically thick bone cortices, such that their long bones are  
19 more dense than similar sized pterosaurs (Fastnacht, 2005).

20 Described by C.C. Young (1964), the dsungaripterids remain a clade with

21 few taxa assigned to them (see Lü et al., 2009a). The more inclusive clade

22 Dsungaripteroidea may or may not include the somewhat problematic

23 *Germanodactylus cristatus* from the Late Jurassic of the Solnhofen limestones of

24 Germany. This species has been recovered as both a basal dsungaripteroid (e.g.

25 Unwin, 2003; Lü et al., 2009a) of close to the Ctenochasmatidae (e.g. Kellner,

---

1 2003). Currently its affinities remain uncertain, but the recent rediscovery of the  
2 missing counterplate to the *G. cristatus* holotype in Dublin, Ireland (Hone, 2010)  
3 may yet help solve this issue. Here, we follow Lü et al., (2009a, b) in considering  
4 this species a dsungaripteroid and also follow their definition of the clade (see  
5 also Unwin, 2003; Witton, 2013, p.201, and for an alternate definition see  
6 Kellner, 2003).

7         Dsungaripteroids have a wide distribution (Witton, 2013., p. 203) but the  
8 dsungaripterids are known primarily from the Cretaceous of Asia with  
9 *Dsungaripterus*, *Noripterus* and *Longchognathosaurus* all being found in the  
10 Junggar Basin of western China (Lü et al., 2009b) and further material coming  
11 from Mongolia (Bakhurina, 1986; Lü et al., 2009b). Other specimens referred to  
12 the Dsungaripteroidea herald from South America (e.g. Martill et al., 2000) and  
13 Europe (in the form of *Germanodactylus*) although the identification of a number  
14 of these as dsungaripteroid is questionable and many are fragmentary.

15         The holotype and referred material of *Noripterus* that was discovered and  
16 described by Young has not featured in the literature to our knowledge since the  
17 original description (Young, 1973), despite the description of new material  
18 referred to this genus (Lü et al., 2009b). While reasonably well illustrated, much  
19 of Young's original paper was devoted to new material of *Dsungaripterus*, is  
20 written in Chinese, and not easy to obtain. Indeed the holotype of *Noripterus* has  
21 not been available for at least a decade and was thought lost for a time. Part of  
22 the holotype and two referred specimens have however now returned to the  
23 IVPP and this material is now available for study. The referral of new material to  
24 *Noripterus* (Lü et al., 2009b), and putative synonymies of some dsungaripterid  
25 taxa (Maish et al., 2004) makes this material important for pterosaurian

---

1 taxonomy. Here we present a revision of the taxonomy of this genus and other  
2 Asian dsungaripterids.

3

4 **Institutional Abbreviations:**

5 BSPG, Bavarian State Collection of Palaeontology, Munich, Germany

6 IVPP, Institute of Vertebrate Paleontology and Paleoanthropology, Chinese

7 Academy of Sciences, Beijing, China.

8

9 **Specimens:**

10 *Identification of specimens:*

11 Multiple specimens referred to *Noriopteris complicidens* are housed at the  
12 IVPP following their collection by Young. In his description, Young (1973)  
13 suggests that he collected elements of approximately eight individuals. However,  
14 not all of them were given different field numbers or museum accession  
15 numbers, and only a limited number were illustrated or measured in the  
16 description. This makes it difficult at this point to refer each element correctly to  
17 Young's (1973) identifications. Furthermore, examination of the *Noriopteris*  
18 specimens currently at the IVPP reveals the holotype to be incomplete but also  
19 that the specimen numbers and field numbers do not match across specimens.  
20 This suggests some confusion in assignment of material to formal IVPP collection  
21 numbers.

22 The holotype is IVPP V 4062 bears the field number 64045. Young (1973)  
23 considered two specimens with field numbers 64041-7 and 64043-3 to be  
24 paratypes and a fourth specimen was unnumbered.

---

1           According to Young's (1973) description of the holotype, IVPP V 4062  
2 should consist of a partial lower jaw, several cervical vertebrae, several dorsal  
3 vertebrae, the distal part of coracoid, the proximal part of a humerus, the distal  
4 part of an ulna, proximal wrist elements, a partial 4<sup>th</sup> metacarpal, a partial wing  
5 phalanx, the distal part of both femora and some further bone fragments. Of this  
6 material, only the fused partial dentaries (Figure 1) can be identified and this  
7 does bear the field number 64045, but the other elements are missing and may  
8 now be lost. One large wing metacarpal (which appears far too large to be  
9 associated with the jaws) also bears the field number 640[numeral missing]5 but  
10 Young (1973) considered this too large to belong to the holotype and we agree  
11 with this assessment. However, this does imply that multiple individuals were  
12 collected under a single field number.

13           Fortunately although little of this material has been illustrated, Young  
14 (1973) did measure many of the elements described in his paper and thus the  
15 identity of a number of specimens can be determined by their published  
16 dimensions and occasional field numbers. Sorting of the available material  
17 reveals the presence of a minimum of five individuals that can currently be  
18 identified:

19

20 1. The holotype IVPP V 4062 (field number 64045) – a pair of partial, fused  
21 dentaries.

22

23 2. Unnumbered paratype specimen (field number 64043-3) - consisting of one  
24 near complete left wing: humerus, radius and ulna, wrist including partial  
25 pteroid, manus and two complete wing phalanges and a broken third. A second

---

1 incomplete wing metacarpal and first wing phalanx are also present  
2 (presumably the right). A complete left hindlimb is also preserved. This  
3 specimen was figures in Young's (1973, plate V) though is incorrectly described  
4 as being field number 64041-7. This list of material matches the description of  
5 material given this field number in Young's description and thus this specimen  
6 appears to be complete as originally recovered. This specimen has now been  
7 given an IVPP designation and is hereafter referred to as IVPP RV 73001 (Figure  
8 2).

9

10 3. IVPP V 4059 (field number 64041-7) – a partial skeleton consisting of two  
11 cranial pieces that are thought to be from the dentary rami, one near complete  
12 cervical, a second very partial cervical, two fused vertebral centra that based on  
13 their size are part of the notarium, a scapulacoracoid, a humerus, two proximal  
14 ulnae and one distal ulna (or radius), one partial wrist complex (proximal and  
15 distal syncarpals), the midshaft of metacarpal IV with parts of two other  
16 metacarpals attached, two manual phalanges of digits I-III, two proximal parts of  
17 the first wing phalanges and parts of three other wing phalanges, a near  
18 complete pelvis and sacrum (with three proximal caudal centra in association),  
19 two femoral heads and two distal femoral ends, two tibial shafts and two distal  
20 tibial ends, a number of isolated metatarsals and pedal phalanges and a small  
21 block of matrix of numerous pedal elements. These pieces were mounted on a  
22 board in their approximate anatomical positions and it is clear that many of the  
23 now fragmentary longbones were originally complete based on marks in the  
24 underlying dust where the shafts are now absent (Figure 3). The pelvis and

---

1 hindlimbs are those illustrated as the specimen on the left side of Young's (1973)  
2 plate IV.

3

4 4. Unnumbered specimens consisting of a distal wing metacarpal, a large carpal  
5 (assuming this wrist elements belongs with the metacarpal). The larger  
6 metacarpal is the one with the field number of the holotype but that is here  
7 considered a separate individual.

8

9       Originally Young (1973) mentions a number of other elements but  
10 without illustrations or any measurements, and as not all elements described  
11 above have field numbers they cannot be positively referred to his description.  
12 The fact that Young gave them separate field numbers does suggest that the  
13 specimens came from different localities. Some field numbers in Young's paper  
14 include suffixes (e.g. 64041-10 as opposed to 64041) and are presumed to refer  
15 to different specimens recovered at a single locality, although none of the  
16 numbers written on specimens contain the suffix values even when they were  
17 used in the paper and so cannot necessarily be aligned to a specific specimen.  
18 These specimens are: 64041, distal part of the wing metacarpal; 64041-10, a  
19 humerus; 64044, a fragment of cervical vertebra (this might now be included  
20 with IVPP V 4059).

21       Almost all of the material is in good condition and despite breaks and  
22 damage appears to have undergone little or no distortion and has also suffered  
23 little erosion. Measurements and subtle anatomical features can therefore be  
24 treated as correct. Fusion of various elements (e.g. fusion of the scapula to the  
25 coracoid, fusion of the extensor tendon process to the first wing phalanx, fusion

---

1 of the pelvic elements) across several specimens suggest that most if not all of  
2 these animals were close to osteological maturity. Although there is variation in  
3 the sizes of various elements that occur in multiple specimens (e.g. the humeri,  
4 femoral heads) the apparently smaller specimens still show the above fused  
5 elements and thus the whole collection is tentatively treated as being of similar  
6 osteological maturity.

7         The material collected by Young came from four different localities but all  
8 of them were close together (most quarries were less than 10 km apart (Dong,  
9 1973)) and there is little reason to think that these were not comparable  
10 localities of the same or similar horizons. The fact that multiple specimens that  
11 are all a close match in size, shape, morphology and preservational condition /  
12 colour where they overlap is weak, but supporting evidence that all of the  
13 material is of one taxon. Here we therefore follow Young (1973) in considering  
14 all of this material to belong to *N. complicidens*, despite the current lack of  
15 overlap between the holotype and other material. The material is also  
16 consistently different to that referred to *Dsungaripterus* (e.g. the ratios of the  
17 limbs – see below) while consistent between specimens again also suggesting  
18 that all of this material represents a single taxon, though identification of the  
19 original quarries and a specimen with a skull would greatly help strengthen this  
20 case.

21

## 22 **Systematic Palaeontology:**

### 23 *Diagnosis of Noripterus:*

24         Young's (1973) original diagnosis is largely redundant in the context of  
25 modern taxonomic characters. He listed the following characters to distinguish



---

1 *Noriopterus complicidens* from other pterosaurs / *Dsungaripterus*: 1) smaller than  
2 two thirds of the size of *Dsungaripterus weii*; 2) having teeth on the anterior tips  
3 of the mandible, unlike the toothless tip in *Dsungaripterus weii*; 3) narrow and  
4 elongated cervical vertebrae; 4) the angle formed by scapula and coracoid is not  
5 big, and the distal part of coracoid may not connect with sternum directly; 5) the  
6 diaphysis of humerus is straight, without a hatchet-like deltopectoral crest; 6)  
7 the proximal carpals form a triangle, and the ratio of ulna to metacarpal IV is  
8 69%; 7) the forelimbs and hind limbs are thin, the formula of phalanges are 2, 3,  
9 4, 4, 0, and the formula of pes phalanges is 2, 4, 4, 5, 0.

10 Of these, characters 1, 3 and 4 are vague and thus not diagnostic as they  
11 cannot therefore be easily compared to other pterosaurs. For character 6, the  
12 ratio of the ulna to the wing metacarpal is 74% in at least one specimen Young  
13 collected, making the value for this ratio of 69% questionable. The description of  
14 the proximal carpals as triangular in general form is correct but the shape is  
15 unknown in *Dsungaripterus*. The description of the limbs as 'thin' in character 7  
16 is vague and the phalangeal formulae of the manus and pes is the same for all  
17 pterodactyloid pterosaurs and thus not diagnostic.

18 Characters 2 and 5 are not immediately problematic, but comparisons of  
19 the specimens to other pterosaurs reveals issues with them. However, the  
20 description of the teeth as reaching the tips of the mandible is unclear since the  
21 rostral end is missing and thus it is not clear if the teeth actually did extend to  
22 the tips of the jaws. However, they certainly do appear to be closer to the tip of  
23 the jaw than in *Dsungaripterus* and so this does suggest a potential difference  
24 between them. The shaft of the humerus is straight in *Noriopterus*, but this is also  
25 the case in many other pterodactyloid pterosaurs and is therefore not diagnostic.

---

1 It is not clear how the non-hatchet-like deltopectoral crest is supposed to be  
2 diagnostic and so remains unclear. Collectively therefore this definition is  
3 problematic and this is difficult to justify as a diagnosis that is currently valid  
4 (though this in itself does not invalidate the taxon). More recently, Lü et al.  
5 (2009b) provided a new diagnosis for *Noripterus* based on newly recovered  
6 material from Mongolia.

7         The definition of Lü et al. (2009b) is as follows: “Skull with a developed  
8 saggital crest, which begins above the interval between the 7th and the 8th tooth  
9 position (from anterior to posterior) of the upper jaw, extending posteriorly  
10 along the midline of the skull and terminating above at the level of the middle of  
11 the dorsal rim of the orbit; Anterior toothless parts of both jaws straight; Ratio of  
12 the length of the mandibular symphysis to that of the lower jaw approximately  
13 0.54; Deep groove on the midline of the dorsal surface of the dentition [sic] part  
14 of the mandibular symphysis; Teeth laterally compressed with sharp tips; Thirty  
15 teeth on the upper jaw and 20 teeth on the lower jaw; Six teeth on upper jaw  
16 below the margin of the nasoantorbital opening; The alveoli are not expanded  
17 into protuberances; The dentition in the upper jaw extends about one-third  
18 further posteriorly than that of lower jaw; Ratio of tibia to femur length is  
19 approximately 1.7.”

20         Although this is a significant improvement on the original diagnosis of  
21 Young, this is also problematic. Assuming that this material does relate to  
22 *Noripterus complicitens* (see below) only three of the characters of Lü et al.  
23 (2009b) can be seen in the holotype of *Noripterus*, and two of these are not  
24 apomorphic. Among dsungaripterids, the character “anterior toothless parts of  
25 both jaws straight” is also true of the holotypes of *Germanodactylus cristatus*

---

1 (BSPG 1982 IV 1) and *Longchoganthosaurus* (Maish et al., 2004 – though see the  
2 discussion below on the validity of this genus). The character “alveoli are not  
3 expanded into protuberances” is also present in *Germanodactylus cristatus* and  
4 may be present here – despite some damage to the base of the teeth, at least one  
5 tooth in the holotype jaw of *Noriopteris* has a slight expansion of bone around the  
6 alveolus in the holotype and more may be present (Figures 1, 4). The third  
7 character “teeth laterally compressed with sharp tips” can be partly inferred in  
8 the holotype by the shape of the alveoli being laterally compressed and the sole  
9 tooth present, while damaged, is sharp tipped. However, again this is also true of  
10 *Germanodactylus cristatus* which appears to have rather laterally compressed  
11 teeth and these are certainly pointed. Thus under Lü et al.’s (2009b) revised  
12 diagnosis, the holotype of Young (1973) is not necessarily a specimen of  
13 *Noriopteris*, and a revised diagnosis is therefore provided below.

14

15 Pterosauria (Kaup, 1834)

16 Pterodactyloidea (Pleninger, 1901)

17 Dsungaripteridae (Young, 1964)

18 *Noriopteris complicidens* (Young, 1973)

19

20 *Definition and diagnosis:*

21 Dsungaripterid pterosaur that can be diagnosed by presence the following

22 characteristics: only mild expansion of bone around the base of the alveoli;

23 possesses a relatively short wing metacarpal (ratio to first wing phalanx close to

24 0.8). It can be further diagnosed by the following combination of characteristics:

25 straight jaw-tips with a deep midline groove on the dentary symphysis.

1

2 **Discussion:**3 *Dsungaripterid Taxonomy:*

4 *Noriopteris* has been repeatedly assigned to the Dsungaripteridae or  
5 recovered as a part of this group in phylogenetic analyses (e.g. Wellnhofer, 1978;  
6 Kellner, 2003; Unwin, 2003; Maish, 2004) based on Young's material and  
7 description. However, none of the diagnostic characters listed by Unwin (2003)  
8 to define the dsungaripterids can be seen in the remaining material of the  
9 holotype of *N. complicidens*. However, Unwin's (2003) characters of limb bones  
10 with relatively thick walls and a strongly bowed femur are both clearly present  
11 in the other material collected by Young that are referred to this taxon. A number  
12 of longbone elements are broken and the cortex thickness can be measured, and  
13 these are between 0.75 and 1.25 mm, for elements that are 5.4 and 4.3 mm in  
14 diameter (ulna and tibia respectively). These are close to those ratios reported  
15 considered diagnostic for the Dsungaripteridae (Fastnacht, 2005) and are above  
16 the values recorded for most other pterosaurs.

17 One character from Kellner (2003) diagnoses the other available material  
18 as belonging to the Tapejaroidea (i.e. the dsungaripterids + azhdarchoids): a  
19 massive medial crest on the humerus with a developed proximal ridge. However  
20 the presence of teeth therefore supports this taxon as a dsungaripterid alone as  
21 all azhdarchoids are toothless. A second character of Kellner (2003) - teeth with  
22 proximal oval base - is also seen here in the holotype dentaries and supports the  
23 referral to Dsungaripteridea (sensu Kellner, 2003). Witton (2013, p 2-8) also  
24 notes that the humeri of dsungaripterids lack pneumatopores and also have a  
25 large deflected deltopectoral crest as seen here in Young's material (Figure 2).

---

1           *Noriopteris* is then a dsungaripteroid pterosaur and also can be assigned  
2 to the Dsungaripteridae. The straight tips to the mandible, and presence of only  
3 very mild expansion of the bone around the base of the teeth clearly separate  
4 *Noriopteris* from *Dsungaripteris* (Young, 1964) and its overall size at osteological  
5 maturity is considerably smaller than that of specimens of *Dsungaripteris*. Thus,  
6 despite the fact that much of the holotype of *Noriopteris* cannot be accounted for,  
7 what remains is diagnostic and the taxon is valid. The additional material  
8 representing the paratypes and other specimens help further separate  
9 *Noriopteris* from *Dsungaripteris*. Although in some details (e.g. the pelvis and  
10 wing phalanx morphology) the two are very similar there are differences.  
11 *Dsungaripteris* has a proportionally much shorter humerus (or longer femur)  
12 compared to *Noriopteris* (humerus to femur ratio of 0.57, based on IVPP V 2776  
13 from Elgin, 2014, compared to 0.81 in IVPP RV 73001).

14           As the second named dsungaripterid, *Noriopteris* must then be considered  
15 a valid taxon. The question remains however, as to whether or not other more  
16 recently described taxa are synonymous with *N. complicidens*.

17           As part of the revision of the genus and description of new material, Lü et  
18 al., (2009b) synonymised '*Phobetor*' (Bakhurina, 1986 – the name is preoccupied  
19 and thus required replacement – Bakhurina & Unwin, 1995) with *Noriopteris*.  
20 Much material has been assigned to '*Phobetor*' (Bakhurina & Unwin, 1995) and  
21 this is a small, straight-jawed dsungaripterid known from Mongolia (Bakhurina,  
22 1986) and thus clearly bears at least some resemblance to *Noriopteris*.

23           As described above, the new and largely complete specimen described by  
24 Lü et al. (2009b) was shown to be a very close match for that described as  
25 '*Phobetor*' by Bakhurina (1986; Bakhurina & Unwin, 1995) and some of the

---

1 details also match the holotype and referred material presented here. A detailed  
2 description of the remains of '*Phobetor*' has yet to be produced by either group,  
3 so detailed comparisons between this and the *Noripterus* material cannot be  
4 made, however there are some notable differences between the two taxa.

5         The sole diagnostic character given by Bakhurina (1982) was the shape of  
6 the facets on the proximal tibia and these at least appears to be very similar to  
7 that of IVPP RV 73001 (although this is partially obscured by the proximal  
8 tarsals). Unwin and Bakhurina (2000) suggested that the limbs of the limited and  
9 fragmentary '*Phobetor*' holotype were indistinguishable from their counterparts  
10 in *Dsungaripterus* and *Noripterus* but that other material confirmed the validity  
11 of the Mongolian taxon. However, the femora of IVPP RV 73001 and V 4059 have  
12 a pronounced anterior-posterior curvature along the shaft – a characteristic  
13 shared by *Dsungaripterus* (Young, 1964) this does not appear to be present in  
14 the referred '*Phobetor*' material of Lü et al., (2009b) as the femur figured has  
15 instead a slight lateral curve. The condition of this referred specimen suggests it  
16 has undergone little or no taphonomic distortion and thus this may yet be a  
17 significant difference between '*Phobetor*' and other taxa.

18         The ratio of the tibia to the femur is also distinct – it is very high (> 1.8) in  
19 IVPP RV 73001 (Figure 2), but only c. 1.7 in the material described by Lü et al.,  
20 (2009b). Similarly the ratio of the length of the wing metacarpal to the first wing  
21 phalanx is 0.83 in *Noripterus* but 0.88 in the referred material. Given the similar  
22 sizes of these animals (humeral lengths of 77 and 84 mm respectively) at the fact  
23 that both are likely osteologically mature and these differences are quite marked.  
24 Other characters also potentially separate this material from *Noripterus*. For  
25 example, the humerus illustrated by Lü et al., (2009b – their figure 2) shows a

---

1 deltopectoral crest that extends further from the shaft of the humerus but is less  
2 dorsoventrally tall compared to that of *Noriopterus*.

3         The coding of Lü et al. (2009a) for a major phylogenetic analysis that  
4 covered numerous pterosaurs, includes several differences between *Noriopterus*  
5 (apparently coded from Young's material) and what they refer to as the 'Tatal  
6 pterosaur' (which we infer as their material that was referred to *Noriopterus* by  
7 Lü et al., 2009b). Two characters refer to major proportions between long bone  
8 elements (ratio of humerus to ulna, and of metatarsal III to the tibia) and thus  
9 are additional differences to those we identify above. The third scored difference  
10 in the datamatrix gives the Tatal pterosaur laterally compressed teeth, but this is  
11 scored as absent in *Noriopterus* (though as noted above, we would also consider  
12 *Noriopterus* to possess this trait). In short, the material of Lü et al. (2009b) may be  
13 synonymous with '*Phobetor*', but both sets of specimens (the '*Phobetor*' material,  
14 and in particular the Tatal material) have a number of notable differences with  
15 *Noriopterus* and suggest that they are distinct taxa.

16         The diagnosis of another dsugaripterid from the Early Cretaceous Tugulu  
17 group, *Longchognathosaurus* (Maish et al., 2004), features characters of the  
18 cranium which cannot be observed in the *Noriopterus* holotype (or currently  
19 available material). Only two characters can be compared to *Noriopterus* and one  
20 of these is present in the holotype and thus cannot be considered an apomorphy  
21 of *Longchognathosaurus*.

22         Maish et al.'s (2004) character of "alveoli not bulbously expanded but  
23 surrounded by a low ring of bone" can be seen in at least one alveolus of the  
24 *Noriopterus* holotype and is thus not diagnostic for *Longchognathosaurus*. This  
25 feature also varies within specimens as can be seen with the anterior most

---

1 alveoli in the holotype *Longchognathosaurus* that lack any kind of bone  
2 expansion around them (Figure 4C), but those alveoli that are more posteriorly  
3 located do show some bony expansions. This change may represent a continuum  
4 along the tooth row from the anterior to posterior teeth and may point to  
5 differential use of the jaws in biting. A stronger bite is typically possible at the  
6 rear of the jaws and therefore it would make sense that durophagus animals  
7 might develop more robust dentition or support for their teeth in this part of the  
8 jaw. It is notable that although badly damaged, the jaw of *Noriopterus* also seems  
9 to show variation in the degree on bony expansions at the alveoli and thus  
10 caution should be used with characters based on bone expansions in these taxa.

11         The second of Maish et al.'s (2004) characters is "Teeth widely spaced  
12 (distance between individual tooth positions always more than distomesial  
13 length of tooth)" is also true of *Noriopterus*. Although the two taxa have different  
14 parts preserved (dentary vs maxilla) the teeth in dsungaripterids are similar in  
15 size, shape or spacing between the upper and lowers jaws, so these should be  
16 broadly comparable here. *Noriopterus* has a range of 1.6-2.4 tooth lengths to  
17 spaces between adjoining teeth and this is near identical in  
18 *Longchognathosaurus* being 1.7 to 2.4 tooth lengths. These both lie in sharp  
19 contrast to *Dsungaripterus* that has teeth larger than their successive spaces. The  
20 referred 'Phobetor' material cannot be easily measured from the figures of Lü et  
21 al. (2009a), but they do appear to have relatively large spaces that are  
22 comparable to *Noriopterus* (Figure 4).

23         The proportional length to width of the teeth in these taxa are also very  
24 similar measuring between 1.4-1.6 in *Noriopterus* and between 1.6-1.8 in  
25 *Longchognathosaurus* (cf. *Dsungaripterus* mandible measures on an unnumbered



---

1 IVPP specimens as 1.2-1.5). Although *Longchognathosaurus* is based on maxillae,  
2 and the part of *Noriopteris* preserving teeth are dentaries, this does leave  
3 *Longchognathosaurus* with nothing to distinguish it from *Noriopteris*. As a result,  
4 this taxon is here suggested to be potentially synonymous with *Noriopteris*.

5       Thus a more detailed and comprehensive comparison is required  
6 between the holotypes of *Noriopteris*, *Longchognathosaurus*, and '*Phobetor*' as  
7 well as the undescribed material of '*Phobetor*' and the new material recently  
8 referred to *Noriopteris*. Provisionally we suggest that '*Phobetor*' is indeed a valid  
9 genus that is distinct from *Noriopteris* (assuming that Bakhurina's Total material  
10 is the same taxon as that of Lü et al.) based on the very different limb  
11 proportions, and that *Longchognathosaurus* is likely synonymous with  
12 *Noriopteris* (although Andres et al., 2010 have also suggested it may be  
13 synonymous with *Dsungaripteris* which seems unlikely given the differences in  
14 tooth morphology and spacing, and the straight tips of the premaxillae).  
15 However, we refrain from making formal revisions here while much of the  
16 *Noriopteris* holotype remains missing, and the original material of '*Phobetor*'  
17 awaits detailed description.

18       Although a number of specimens of the *Dsungaripteridae* have been  
19 recovered from Asia, few have been described or even illustrated in any detail to  
20 date making comparisons between specimens and putative taxa difficult.  
21 However, the renewed access to C.C. Young's material of *Noriopteris complicidens*  
22 reveals important character information that helps resolve some issues in the  
23 taxonomy of the members of this group, and gives a much improved  
24 understanding of this intriguing taxon (Figure 5).

25

---

1 **Acknowledgements:**

2 DWEH wishes to dedicate this manuscript to Wann Langston who sent him on  
3 the path to hunt down *Noriopterus* and led to this manuscript being created. Our  
4 thanks to Xiaoqing Ding for the reparation of parts of the specimens. We  
5 thank Dave Unwin, Natasha Bakhurina, Lü Junchang, Taissa Rodrigues, Brian  
6 Andres, Liz Martin-Silverstone and Mark Witton for discussions of these  
7 specimens and providing images of key specimens. Our thanks to Rebecca  
8 Gelernter for her superb rendering of the reconstructed skeleton used in Figure  
9 5. XX's work is supported by the National Natural Science Foundation of China  
10 (grant number: 41120124002).

11

12 **References:**

13 Andres, B., Clark, J.M., & Xu, X. 2010. A new rhamphorhynchid pterosaur from the  
14 Upper Jurassic of Xinjiang, China, and the phylogenetic relationships of basal  
15 pterosaurs. *Journal of Vertebrate Paleontology*, **30**, 163–187.

16

17 Bakhurina, N.N. 1982. A pterodactyl from the Lower Cretaceous of Mongolia.  
18 *Paleontological Journal*, **1982**, 105–109.

19

20 Bakhurina, N.N. 1986. Letayushchiye yashchery. ["Flying saurians."] [in Russian].  
21 *Priroda*, **1986**, 27–36.

22

23 Bakhurina, N.N. & Unwin, D.M. 1995. A survey of pterosaurs from the Jurassic  
24 and Cretaceous of the former Soviet Union and Mongolia. *Historical Biology*,  
25 **10**,197–245.

- 
- 1  
2 Dong, Z. 1973. Cretaceous Stratigraphy of Wuerho district, Dsungar Basin.  
3 Reports of Paleontological Expedition to Sinkiang (II). Pterosaurian Fauna from  
4 Wuerho, Sinkiang. Memoirs of the Institute of Vertebrate Palaeontology and  
5 Paleoanthropology, Academy Sinica, **11**, 1–7.  
6  
7 Fastnacht, M. 2005. The first dsungaripterid pterosaur from the Kimmeridgian of  
8 Germany and the biomechanics of pterosaur long bones. *Acta Palaeontologica*  
9 *Polonica*, **50**, 273–288.  
10  
11 Hone, D.W.E. 2010. A short note on modifications to Nineteenth Century  
12 pterosaur specimens held in the National Museum of Ireland – Natural History,  
13 Dublin. *Geological Curator*, **9**, 261–265.  
14  
15 Kellner, A.W.A. 2003. Pterosaur phylogeny and comments on the evolutionary  
16 history of the group. In: Buffetaut, E. & Mazin, J.-M. (eds) *Evolution and*  
17 *Palaeobiology of Pterosaurs*. Geological Society Special Publications, **217**, 105–  
18 137.  
19  
20 Lü, J., Unwin, D. M., Jin, X., Liu, Y., and Ji, Q. 2009a. Evidence for modular evolution  
21 in a long-tailed pterosaur with a pterodactyloid skull. *Proceedings of the Royal*  
22 *Society of London, Series B*, rspb20091603  
23  
24 Lü, J., Azuma, Y., Dong, Z., Barsbold, R., Kobayashi, Y., & Lee, Y.N. 2009b. New  
25 material of dsungaripterid pterosaurs (Pterosauria: Pterodactyloidea) from

- 
- 1 western Mongolia and its palaeoecological implications. *Geological Magazine*,  
2 **146**, 690–700.
- 3
- 4 Maisch, M.W., Matzke, A.T. & Sun, G. 2004. A new dsungaripteroid pterosaur from  
5 the Lower Cretaceous of the southern Junggar Basin, north-west China.  
6 *Cretaceous Research*, **25**, 625–634.
- 7
- 8 Martill, D.M., Frey, E., Chong, D.G. & Bell, C.M. 2000. Reinterpretation of a Chilean  
9 pterosaur and the occurrence of Dsungaripteridae in South America. *Geology*  
10 *Magazine*, **137**, 19–25.
- 11
- 12 Unwin, D.M. 2003. On the phylogeny and evolutionary history of pterosaurs. In:  
13 Buffetaut, E. & Mazin, J.-M. (eds) *Evolution and Palaeobiology of Pterosaurs*.  
14 *Geological Society Special Publications*, **217**, 139–190.
- 15
- 16 Unwin, D.M. & Bakhurina, N.N. 2000. Pterosaurs from Russia, Middle Asia and  
17 Mongolia. In: Benton, M.J., Shishikin, M.A., Unwin, D.M. & Kurochkin, E.N. (eds)  
18 *The age of dinosaurs in Russia and Mongolia*. Cambridge University Press. pp.420-  
19 433.
- 20
- 21 Wellnhofer, P. 1978. Pterosauria. *Handbuch der Paläoherpetologie*, vol. 19. 83pp  
22 CIP-Kurztitelaufnahme der Deutschen Biblioth Sttutgart.
- 23
- 24 Witton, M.P. 2013. *Pterosaurs: natural history, evolution, anatomy*. Princeton  
25 University Press. **ADD PP**

1

2 Young, C.C. 1964. On a new pterosaurian from Sinkiang, China. *Vertebrata*3 *PalAsiatica*, **8**, 221–255.

4

5 Young, C.C. 1973. Reports of Paleontological Expedition to Sinkiang (II).

6 *Pterosaurian Fauna from Wuerho, Sinkiang*. *Memoirs of the Institute of*7 *Vertebrate Palaeontology and Paleoanthropology, Academy Sinica*, **11**, 18–35.

8

9

10

11 **Figures:**

12



13


14 Figure 1. The currently available holotype material of *Noripterus complicidens*

15 (IVPP V 4062) – partial dentaries with some intact teeth. The teeth are widely-

16 spaced and show slight expansion of bone around the base of some towards the

17 rear of the jaw. Scale bar is 20 mm.

18




1

2 Figure 2. The near complete fore and hindlimbs of IVPP RV 73001. Elements still  
3 bear Young's (1973) original fieldnumber for the specimen – 64043-3.


4 Abbreviations as follows: Cp, carpal block; D, manual digits; Fb, fibula; Fm,  
5 femur; Hu, humerus; Mc, metacarpals; Pes, foot (including tarsals); Ph, wing  
6 phalanges; Rd, radius; Tb, tibia; Ul, ulna.

7



1 Figure 3. The material belonging to specimen IVPP V 4059 of *Noriopterus* (field  
2 number 64041-7) as found in the collections with material mounted on a  
3 wooden board – note that in many cases longbones are broken and parts are  
4 missing but their original position and size can often be identified based on the  
5 cleaner parts of the board. Abbreviations as in Figure 2 with the following  
6 additions: Cv, cervical vertebra; MD, manual and / or pedal elements; Pv, pelvis;  
7 SC, scapulocoracoid. Not all elements are identified or labeled here.

8




9



1 Figure 4. Tracings of the jaws of Asian dsungaripterid pterosaurs to show tooth  
 2 size and spacing (teeth or alveoli are in grey). A The holotype dentaries of  
 3 *Noriopteris*, B the dentaries of the Mongolian material referred to '*Phobetor*' by Lü  
 4 et al., 2009 (modified from their figure 4e), C the holotype maxillae of  
 5 *Longchognathosaurus* (the premaxillae are known but not drawn here), D  
 6 unnumbered IVPP specimen of *Dsungaripteris*. Scales bars A, B, and C, 20 mm; D,  
 7 50 mm.

8



9

10 Figure 5. Reconstruction of *Noriopteris complicidens* based on the C.C. Young  
 11 material (image by Rebecca Gelernter). Elements known are in white, unknown  
 12 elements are greyed out. Missing parts are restored based on Lü et al., (2009b)  
 13 and Witton, (2013, p. 207).