

Tightness and Line Ensembles for Brownian Polymers Under Geometric Area Tilts

Pietro Caputo, Dmitry Ioffe and Vitali Wachtel

Abstract We prove tightness and limiting Brownian-Gibbs description for line ensembles of non-colliding Brownian bridges above a hard wall, which are subject to geometrically growing self-potentials of tilted area type. Statistical properties of the resulting ensemble are very different from that of non-colliding Brownian bridges without self-potentials. The model itself was introduced in order to mimic level lines of $2 + 1$ discrete Solid-On-Solid random interfaces above a hard wall.

Keywords Level lines of low temperature interfaces · Brownian Gibbs line ensembles · Geometric area tilts · Tightness

1 Introduction, Notation and Results

1.1 Level Lines of SOS Surfaces and Related Ensembles of Non-intersecting Paths

In this paper we investigate large scale behavior of ensembles of non-intersecting Brownian polymers under geometric area tilts. Our main motivation comes from a

We dedicate this paper to Anton Bovier on the occasion of his 60th birthday.

DI was supported by the Israeli Science Foundation grants 1723/14 and 765/18.

VW was supported by the Humboldt Foundation.

P. Caputo

Dipartimento di Matematica e Fisica, Roma Tre University, Rome, Italy

e-mail: caputo@mat.uniroma3.it

D. Ioffe (✉)

The Faculty of Industrial Engineering and Management, Technion, Haifa 32000, Israel

e-mail: ieioffe@ie.technion.ac.il

V. Wachtel

Institut für Mathematik, Universität Augsburg, 86135 Augsburg, Germany

e-mail: vitali.wachtel@math.uni-augsburg.de

desire to understand and model limiting properties of microscopic large level lines of the $2 + 1$ SOS (solid-on-solid) random interfaces conditioned to stay above a flat wall. We refer to [3] for a careful exposition of this connection with references to earlier works.

Models of a (single) Brownian polymer constrained to stay above a barrier were introduced in [13]. By Girsanov's transform an equivalent formulation is in terms of positive Brownian bridges under area tilts [21]. These models are of an independent interest and, in particular, they have a rich limiting variational and fluctuation structure. We would like to mention recent works [11, 22, 26] and references therein.

Gibbsian structure of (single $\mathbf{C}(\mathbb{R}, \mathbb{R})$ -valued) Brownian polymers with self-potentials and self-interactions was introduced and explored in [20, 25]. Space-time Gibbsian states on $\mathbf{C}(\mathbb{R}, \mathbb{R})^{\mathbb{Z}^d}$ were introduced and constructed in [9, 23] in the weak interaction case, both as Gibbs states and as systems of infinite dimensional interacting diffusions. Hard core interactions between ordered paths in one spatial dimension do not fall into this framework. Variants of infinite volume dynamics whose equilibrium states are determinantal point fields were constructed in [24, 27], see also various refinements with references to the latter works.

Recent developments which are closely related to the model we consider here are [2, 12]. However, since geometrically growing area tilts preclude using Karlin-McGregor formula, the determinantal structure is apparently lost, and one has to deal with a rather different situation. For instance no rescaling as the number of paths grow is needed. Yet, the Brownian-Gibbsian framework introduced in [6, 7] fits in extremely well, and we shall rely on the philosophy and the ideas which were developed in these remarkable papers.

1.2 Organization of the Paper

The Brownian notation and the models of Brownian polymers under geometric area tilts are introduced in Sects. 1.3–1.5. The main confinement results from [3] which enables a uniform control over one-point distributions of the top path is recalled in Sect. 1.6. Our main results here which are formulated in Sect. 1.7 yield full tightness of the ensembles we consider, both with free and zero boundary conditions, and, furthermore, we establish Brownian-Gibbs property of limiting infinite dimensional ensembles. Whether there is a unique such limiting ensemble or not remains an open question, although monotonicity arguments imply unicity of the limit for polymer measures with zero boundary conditions. The proofs of full tightness appear in Sect. 2. Limiting line ensembles are discussed in Sect. 3. Finally, future research directions and some open problems are outlined in the concluding Sect. 3.5.

1.3 Brownian Motion and Brownian Bridges

In the sequel we shall use the same notation for path measures of underlying Brownian motion and Brownian bridges and for expectations with respect to these path measures. For $\ell < r$ and $x \in \mathbb{R}$, let $\mathbf{P}_{\ell,r}^x$ be the path measure of the Brownian motion X on $[\ell, r]$ which starts at x at time ℓ ; $X(\ell) = x$. We can record $\mathbf{P}_{\ell,r}^x$ as follows:

$$\mathbf{P}_{\ell,r}^x(F(X)) = \int \mathbf{B}_{\ell,r}^{x,y}(F(X)) dy \quad (1.1)$$

where $\mathbf{B}_{\ell,r}^{x,y}$ the *unnormalized* path measure of the Brownian bridge X on $[\ell, r]$ which starts at x at time ℓ and ends at y at time r ; $X(\ell) = x$, $X(r) = y$. In this way the total mass of $\mathbf{B}_{\ell,r}^{x,y}$ is given by

$$q_{r-\ell}(x, y) := \mathbf{B}_{\ell,r}^{x,y}(1) = \frac{1}{\sqrt{2\pi(r-\ell)}} e^{-\frac{(y-x)^2}{2(r-\ell)}}. \quad (1.2)$$

The corresponding normalized Brownian bridge probabilities will be denoted as $\Gamma_{\ell,r}^{x,y}$, that is $\Gamma_{\ell,r}^{x,y}(\cdot) = \mathbf{B}_{\ell,r}^{x,y}(\cdot)/q_{r-\ell}(x, y)$.

For an n -tuple $\underline{x} = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$, set

$$\mathbf{P}_{n;\ell,r}^{\underline{x}} = \mathbf{P}_{\ell,r}^{x_1} \otimes \mathbf{P}_{\ell,r}^{x_2} \otimes \dots \otimes \mathbf{P}_{\ell,r}^{x_n}.$$

Similarly for n -tuples $\underline{x}, \underline{y} \in \mathbb{R}^n$, set

$$\mathbf{B}_{n;\ell,r}^{\underline{x},\underline{y}} = \mathbf{B}_{\ell,r}^{x_1,y_1} \otimes \mathbf{B}_{\ell,r}^{x_2,y_2} \otimes \dots \otimes \mathbf{B}_{\ell,r}^{x_n,y_n}, \quad (1.3)$$

and

$$q_{n,r-\ell}(\underline{x}, \underline{y}) = \prod_{i=1}^n q_{n,r-\ell}(x_i, y_i), \quad \Gamma_{n,\ell,r}^{\underline{x},\underline{y}} = \frac{\mathbf{B}_{n;\ell,r}^{\underline{x},\underline{y}}}{q_{n,r-\ell}(\underline{x}, \underline{y})}. \quad (1.4)$$

1.4 Polymer Measures with Geometric Area Tilts

Given a function h , the signed h -area under the trajectory of X is defined as

$$\mathcal{A}_{\ell,r}^h(X) = \int_{\ell}^r h(t)X(t)dt. \quad (1.5)$$

We shall drop the superscript if $h \equiv 1$ and use $\mathcal{A}_{\ell,r}(X)$ accordingly. For $n \in \mathbb{N}$ define

$$\mathbb{A}_n^+(M) = \{\underline{x} \in \mathbb{R}^n : M > x_1 > \dots > x_n > 0\} \quad \text{and} \quad \mathbb{A}_n^+ = \mathbb{A}_n^+(\infty). \quad (1.6)$$

We also define the set

$$\bar{\mathbb{A}}_n^+ = \{\underline{x} \in \mathbb{R}^n : x_1 \geq \cdots \geq x_n \geq 0\}. \quad (1.7)$$

Polymer measures which we consider in the sequel are always concentrated on the set $\Omega_{n;\ell,r}^+$ of n -tuples \underline{X} ,

$$\Omega_{n;\ell,r}^+ = \{\underline{X} : \underline{X}(t) \in \mathbb{A}_n^+ \quad \forall t \in (\ell, r)\}. \quad (1.8)$$

Given $n \geq 1$, $a > 0$, $\lambda > 1$ and $\underline{x}, \underline{y} \in \mathbb{A}_n^+$, consider the partition function

$$Z_{n;\ell,r}^{\underline{x},\underline{y}}(a, \lambda) := \mathbf{B}_{\ell,r}^{\underline{x},\underline{y}} \left(\mathbb{1}_{\Omega_{n;\ell,r}^+} e^{-\sum_1^n a \lambda^{i-1} \mathcal{A}_{\ell,r}(X_i)} \right), \quad (1.9)$$

and the associated probability measure $\mathbb{P}_{n;\ell,r}^{\underline{x},\underline{y}}[\cdot | a, \lambda]$ defined by

$$\mathbb{P}_{n;\ell,r}^{\underline{x},\underline{y}}[F(\underline{X}) | a, \lambda] := \frac{1}{Z_{n;\ell,r}^{\underline{x},\underline{y}}(a, \lambda)} \mathbf{B}_{\ell,r}^{\underline{x},\underline{y}} \left(F(\underline{X}) \mathbb{1}_{\Omega_{n;\ell,r}^+} e^{-\sum_1^n a \lambda^{i-1} \mathcal{A}_{\ell,r}(X_i)} \right), \quad (1.10)$$

where F is any bounded measurable function over the set of n -tuples of continuous functions from $[\ell, r]$ to \mathbb{R} . The measure $\mathbb{P}_{n;\ell,r}^{\underline{x},\underline{y}}[\cdot | a, \lambda]$ will be referred to as the n -polymer measure with (a, λ) -geometric area tilts with boundary conditions $(\underline{x}, \underline{y})$ on the interval $[\ell, r]$.

We remark that $\mathbb{P}_{n;\ell,r}^{\underline{x},\underline{y}}[\cdot | a, \lambda]$ is well defined for all $\underline{x}, \underline{y} \in \mathbb{A}_n^+$. Indeed, if $\underline{x}, \underline{y} \in \mathbb{A}_n^+$ one has

$$\mathbf{B}_{\ell,r}^{\underline{x},\underline{y}}(\Omega_{n;\ell,r}^+) > 0, \quad (1.11)$$

and therefore $Z_{n;\ell,r}^{\underline{x},\underline{y}}(a, \lambda) \in (0, \infty)$. While this does not apply to all $\underline{x}, \underline{y} \in \bar{\mathbb{A}}_n^+$, it is still possible to define $\mathbb{P}_{n;\ell,r}^{\underline{x},\underline{y}}[\cdot | a, \lambda]$ in these cases by a limiting procedure, see e.g. [6, Definition 2.13] for the case $a = 0$. In particular, we shall often deal with the case of *zero boundary conditions*

$$\mathbb{P}_{n;\ell,r}^0[\cdot | a, \lambda] := \mathbb{P}_{n;\ell,r}^{0,0}[\cdot | a, \lambda].$$

It is also natural to consider the polymer measure with *free boundary conditions* defined by

$$\mathbb{P}_{n;\ell,r}[F(\underline{X}) | a, \lambda] := \frac{1}{\mathcal{Z}_{n;\ell,r}(a, \lambda)} \int_{\mathbb{A}_n^+} \int_{\mathbb{A}_n^+} \mathbf{B}_{\ell,r}^{\underline{x},\underline{y}} \left(F(\underline{X}) \mathbb{1}_{\Omega_{n;\ell,r}^+} e^{-\sum_1^n a \lambda^{i-1} \mathcal{A}_{\ell,r}(X_i)} \right) d\underline{x} d\underline{y} \quad (1.12)$$

where

$$\mathcal{Z}_{n;\ell,r}(a, \lambda) := \int_{\mathbb{A}_n^+} \int_{\mathbb{A}_n^+} Z_{n;\ell,r}^{\underline{x},\underline{y}}(a, \lambda) d\underline{x} d\underline{y}, \quad (1.13)$$

and \underline{dx} , \underline{dy} denote Lebesgue measures on \mathbb{R}^n . For a proof that $\mathbb{P}_{n;\ell,r}[\cdot | a, \lambda]$ is well defined, that is $\mathcal{Z}_{n;\ell,r}(a, \lambda) \in (0, \infty)$, for all $a > 0$, $\lambda > 1$, see [3, Appendix A].

1.5 A General Class of Polymers with Area Tilts

Let us say that two functions f and g on $I \subset \mathbb{R}$ satisfy $f < g$ if $f(t) \leq g(t)$ for any $t \in I$. By construction, if $\underline{X} \in \Omega_{n;\ell,r}^+$, then $0 < X_n < X_{n-1} < \dots < X_1$ on $[l, r]$. For every $n \in \mathbb{N}$ and $\ell < r$, consider the following general class $\mathbb{P}_{n;\ell,r}^{\underline{x}, \underline{y}}[\cdot | h_-, h_+, \underline{\rho}]$ of polymer measures which is parametrized by:

- a:** Boundary conditions $\underline{x}, \underline{y} \in \mathbb{A}_n^+$.
- b:** A pair $\underline{h} = (h_-, h_+)$ of non-negative continuous functions, called respectively the floor and the ceiling, satisfying $h_- < h_+$ on $[\ell, r]$.
- c:** An n -tuple of (not necessarily ordered) nonnegative continuous functions $\underline{\rho} = \{\rho_1, \dots, \rho_n\}$, called the area tilts.

Then, setting

$$\Omega_{n;\ell,r}^{\underline{h}} = \Omega_{n;\ell,r}^+ \cap \{h_- < X_n\} \cap \{X_1 < h_+\}, \quad (1.14)$$

define:

$$\mathbb{P}_{n;\ell,r}^{\underline{x}, \underline{y}}[\underline{dX} | h_-, h_+, \underline{\rho}] \propto e^{-\sum_1^n \mathcal{A}_{\ell,r}^{\rho_i}(X_i)} \mathbb{1}_{\Omega_{n;\ell,r}^{\underline{h}}} \mathbf{B}_{\ell,r}^{\underline{x}, \underline{y}}(\underline{dX}). \quad (1.15)$$

The corresponding partition function is denoted $Z_{n;\ell,r}^{\underline{x}, \underline{y}}(h_-, h_+, \underline{\rho})$. Clearly, the polymer measure $\mathbb{P}_{n;\ell,r}^{\underline{x}, \underline{y}}[\cdot | h_-, h_+, \underline{\rho}]$ coincides with the $\mathbb{P}_{n;\ell,r}^{\underline{x}, \underline{y}}[\cdot | a, \lambda]$ defined in (1.10) in the case of geometric tilts $\rho_i \equiv a\lambda^{i-1}$ and trivial floor and ceilings $(h_-, h_+) \equiv (0, +\infty)$. Similarly, we shall employ the following notation for Brownian bridge measures conditioned to $\Omega_{n;\ell,r}^{\underline{h}}$,

$$\mathbf{B}_{n;\ell,r}^{\underline{x}, \underline{y}}[\cdot | h_-, h_+] := \Gamma_{n;\ell,r}^{\underline{x}, \underline{y}}[\cdot | h_-, h_+] := \Gamma_{n;\ell,r}^{\underline{x}, \underline{y}}(\cdot | \Omega_{n;\ell,r}^{\underline{h}}). \quad (1.16)$$

There is a straightforward extension of (1.8) and (1.14) to sets of paths over more general subsets $I \subset \mathbb{R}$ (in the sequel we shall need to work with finite union of intervals). Namely,

$$\Omega_{n;I}^+ = \{\underline{X} : \underline{X}(t) \in \mathbb{A}_n^+ \quad \forall t \in I\} \quad (1.17)$$

and

$$\Omega_{n;I}^{\underline{h}} = \Omega_{n;I}^+ \cap \{h_- < X_n\} \cap \{X_1 < h_+\} \quad (1.18)$$

Moreover, there is a version of (1.15) which permits more general boundary conditions: Let $\underline{\nu}$ and $\underline{\eta}$ be n -tuples of functions on \mathbb{R}_+ . For $\underline{x} \in \mathbb{A}_n^+$ set $\underline{\nu}(\underline{x}) = \sum_1^n \nu_i(x_i)$, and $\underline{\eta}(\underline{x}) = \sum_1^n \eta_i(x_i)$. Similarly, set $\mathcal{A}_{\ell,r}^{\underline{\rho}}(\underline{X}) = \sum_1^n \mathcal{A}_{\ell,r}^{\rho_i}(X_i)$. Then,

$$\mathbb{P}_{n;\ell,r}^{\underline{\nu},\underline{\eta}} \left[d\underline{X} | h_-, h_+, \underline{\rho} \right] \propto \int_{\mathbb{A}_n^+} \int_{\mathbb{A}_n^+} e^{-A_T^\ell(\underline{X})} \mathbb{1}_{\Omega_{n;\ell,r}^{\underline{\nu}}}(\underline{X}) e^{-\underline{\nu}(x)} \mathbf{B}_{\ell,r}^{x,y}(\underline{dX}) e^{-\underline{\eta}(y)} d\underline{x} d\underline{y}. \tag{1.19}$$

The corresponding partition function is denoted $\mathcal{Z}_{n;\ell,r}^{\underline{\nu},\underline{\eta}}(h_-, h_+, \underline{\rho})$.

Below we shall tacitly assume that boundary conditions $\underline{\nu}, \underline{\eta}$ are chosen in such a way that the corresponding polymer measures are well defined, this is justified in all the relevant cases in Appendix A of [3].

Reduced notation. We shall, unless this creates a confusion, employ the following reduced notation: If $\underline{\eta}, \underline{\nu}$ are identically zero, we shall drop them from the notation. We refer to this as the case of free (or empty) boundary conditions. Similarly we shall drop from the notation the floor h_- whenever $h_- \equiv 0$ and the ceiling h_+ whenever $h_+ \equiv \infty$.

In the case of pure geometric tilts we shall write a, λ instead of $\underline{\rho}$ whenever

$$\underline{\rho} = \{a, a\lambda, \dots, a\lambda^{n-1}, \dots\} \tag{1.20}$$

and, furthermore, we shall drop a, λ from the notation whenever this creates no confusion.

In the sequel we shall drop sub-index n unless we would like to stress the number of polymers in a stack, and whenever this will cause no confusion.

In the case of symmetric time intervals we shall write

$$\Omega_{n,T}^+ = \Omega_{n;-T,T}^+, \quad Z_{n,T}^{x,y} = Z_{n;-T,T}^{x,y}, \quad \mathbf{B}_{n,T}^{x,y} = \mathbf{B}_{n;-T,T}^{x,y}, \quad \mathbb{P}_{n,T}^{x,y} = \mathbb{P}_{n;-T,T}^{x,y},$$

and so on.

1.6 A Confinement Statement and Uniform Control of Curved Maxima

The main result of [3] could be formulated as follows:

Theorem 1.1 *For any fixed $a > 0, \lambda > 1$ and $\chi > 0$ the family of (one-dimensional) distributions of the height $X_1(0)$ of the top path at the origin under the free boundary condition field $\{\mathbb{P}_{n;\ell,r}[\cdot | a, \lambda]\}_{n \in \mathbb{N}, \ell \leq -\chi, r \geq \chi}$ is tight. In other words the top path does not fly away as the number of polymers and the length of their horizontal span grow. Moreover, the same statement holds for the zero-boundary condition field $\mathbb{P}_{n;\ell,r}^0[\cdot | a, \lambda]$.*

The statement of Theorem 1.1 follows from uniform control of the expectations of a certain curved maxima. Let us describe the latter notion.

Let $\varphi_\alpha(t) = |t|^\alpha$ with $\alpha \in (0, \frac{1}{2})$. Given a continuous function h on $[\ell, r]$ define (see Fig. 1)




Fig. 1 The curved maximum $\xi_\alpha^{\ell,r}(h)$, in the case where $[\ell, r]$ is a symmetric interval around the origin

$$\xi_\alpha^{\ell,r}(h) = \min \{y \geq 0 : y + \varphi_\alpha > h\} = \max_{t \in [\ell, r]} (h(t) - |t|^\alpha)_+, \quad (1.21)$$

where $(\cdot)_+$ denotes the positive part. Informally, $\xi_\alpha^{\ell,r}(h)$ is the minimal amount to lift φ_α so that it will stay above h . We think of $\xi_\alpha^{\ell,r}(h)$ in terms of the curved maximum of h on $[\ell, r]$.

Then, Theorem 1.1 is an immediate consequence of the following result [3]:

Theorem 1.2 For any $a > 0$, $\lambda > 1$, $\chi > 0$ and $\alpha \in (0, \frac{1}{2})$:

$$\begin{aligned} \xi_\alpha^* &:= \sup_{\ell \leq -\chi} \sup_{r \geq \chi} \sup_n \mathbb{E}_{n;\ell,r} [\xi_\alpha^{\ell,r}(X_1) \mid a, \lambda] \\ &= \sup_{\ell \leq -\chi} \sup_{r \geq \chi} \sup_n \mathbb{E}_{n;\ell,r} \left[\max_{t \in [\ell, r]} (X_1(t) - |t|^\alpha)_+ \mid a, \lambda \right] < \infty. \end{aligned} \quad (1.22)$$

The fact that Theorem 1.2 also holds for the zero boundary condition field is a consequence of stochastic domination, see Sect. 2.1.

1.7 Full Tightness, Non-intersecting Line Ensembles and Brownian-Gibbs Property

The main contribution of this paper is a strengthening of the above mentioned results, with the extension of the one-dimensional tightness to a full tightness statement on path space. To formulate this we need to introduce some more notation. Fix $a > 0$ and $\lambda > 1$. Given $\gamma > 0$ and $k \in \mathbb{N}$, let $n > k$, $[\ell, r] \supseteq [-\gamma, \gamma]$, and consider the

free boundary condition field $\mathbb{P}_{n;\ell,r}[\cdot | a, \lambda]$. We shall use $\mu_{n;\ell,r}^{k,\gamma}$ for the induced distribution of k top paths $\underline{X} = (X_1(\cdot), \dots, X_k(\cdot))$ over the interval $[-\gamma, \gamma]$. Similarly, we write $\mu_{n;\ell,r}^{0,k,\gamma}$ for the distribution of k top paths under the zero boundary condition field $\mathbb{P}_{n;\ell,r}^0[\cdot | a, \lambda]$. In this way $\mu_{n;\ell,r}^{k,\gamma}, \mu_{n;\ell,r}^{0,k,\gamma}$ are distributions on the set of continuous functions from $[-\gamma, \gamma]$ to $\bar{\mathbb{A}}_k^+$, denoted $\mathbf{C}([-\gamma, \gamma]; \bar{\mathbb{A}}_k^+)$. The latter is equipped with the topology of uniform convergence.

Theorem 1.3 *For any $k \in \mathbb{N}$ and $\gamma > 0$ the family*

$$\left\{ \mu_{n;\ell,r}^{k,\gamma}, [\ell, r] \supseteq [-\gamma, \gamma], n > k \right\} \tag{1.23}$$

is tight on $\mathbf{C}([-\gamma, \gamma]; \bar{\mathbb{A}}_k^+)$. The same holds for the family

$$\left\{ \mu_{n;\ell,r}^{0,k,\gamma}, [\ell, r] \supseteq [-\gamma, \gamma], n > k \right\}.$$

We turn to the analysis of the consequences of the above tightness results regarding convergence to a limiting polymer measure describing infinitely many non-intersecting random lines. We start by recalling the definition of Brownian-Gibbs line ensembles. We refer to [6, 7, 10] for a more comprehensive setup, as well as to earlier papers [23–25] on the subject. Here we formulate the concepts that are relevant for our setting.

Define the spaces

$$\mathbb{A}_\infty^+ = \left\{ \underline{x} \in (0, \infty)^\mathbb{N} : x_1 > x_2 > \dots > 0 \right\} \subset \bar{\mathbb{A}}_\infty^+ = \left\{ \underline{x} \in [0, \infty)^\mathbb{N} : x_1 \geq x_2 \geq \dots \geq 0 \right\}.$$

The sample space is $\Omega = \mathbf{C}(\mathbb{R}, \bar{\mathbb{A}}_\infty^+)$, the set of continuous functions $f : \mathbb{R} \mapsto \bar{\mathbb{A}}_\infty^+$, equipped with the topology of uniform convergence of any finite number of paths on compact subsets, and with the corresponding Borel σ -field \mathcal{B} . The coordinate maps $\underline{X} \in \Omega \mapsto X_i(t)$ are viewed as position of i th particle at time t . Following [6], for each $n \in \mathbb{N}$, and time interval $[\ell, r] \subset \mathbb{R}$, define the internal and external σ -algebras

$$\mathcal{B}_{n;\ell,r}^i = \sigma(X_i(t) : t \in (\ell, r) \text{ and } i \leq n) \text{ and } \mathcal{B}_{n;\ell,r}^e = \sigma(X_i(t) : \text{either } t \notin (\ell, r) \text{ or } i > n). \tag{1.24}$$

Recall that $\mathbb{E}_{n;\ell,r}^{\underline{x}, \underline{y}}[\cdot | h, a, \lambda]$ stands for the expectation w.r.t. n -polymer measure with (a, λ) -geometric area tilts with boundary conditions $(\underline{x}, \underline{y})$ and floor h on the interval $[\ell, r]$.

Definition 1 A probability measure \mathbb{P} on Ω is said to have the *Brownian-Gibbs property* with respect to (a, λ) -geometric area tilts (1.20) if for any bounded measurable $F : \Omega \mapsto \mathbb{R}$,

$$\mathbb{E} \left(F \mid \mathcal{B}_{n;\ell,r}^e \right) = \mathbb{E}_{n;\ell,r}^{\underline{X}^{(\leq n)}(\ell), \underline{X}^{(\leq n)}(r)} \left[F(\cdot, \underline{X}^{(>n)}) \mid X_{n+1}, a, \lambda \right], \tag{1.25}$$

\mathbb{P} -a.s for any $-\infty < \ell < r < \infty$ and $n \in \mathbb{N}$. In (1.25), we use the notation $\underline{X}^{(\leq n)} = (X_1, \dots, X_n)$ and $\underline{X}^{(>n)} = (X_{n+1}, X_{n+2}, \dots)$. A probability measure \mathbb{P} on Ω with the above Brownian-Gibbs property is called a Brownian-Gibbs measure or a Brownian-Gibbs line ensemble with respect to (a, λ) -geometric area tilts. The line ensemble is said to be *non-intersecting* if \mathbb{P} is concentrated on $\mathbf{C}(\mathbb{R}, \mathbb{A}_\infty^+) \subset \mathbf{C}(\mathbb{R}, \bar{\mathbb{A}}_\infty^+)$.

Notice that measures $\mathbb{P}_{m;S,T}^{x,y}$ and $\mathbb{P}_{m;S,T}$ describing m lines in the time interval $[S, T]$, as defined in Sect. 1.4, are trivially extendable to the whole of Ω by setting $X_i \equiv 0$ for $i > m$ and $\underline{X}(t) \equiv \underline{X}(T)$ for $t > T$ (respectively, $\underline{X}(s) \equiv \underline{X}(S)$ for $s < S$). Moreover, these measures satisfy the Brownian-Gibbs property (1.25) whenever $S \leq \ell < r \leq T$ and $n \leq m$.

Theorem 1.3 says that any sequence from $\{\mu_{n;\ell,r}^{k,\gamma}, [\ell, r] \supseteq [-\gamma, \gamma], n > k\}$ has a weakly converging subsequence in the space of probability measures on $\mathbf{C}([-\gamma, \gamma]; \bar{\mathbb{A}}_k^+)$, for all fixed $k \in \mathbb{N}$ and $\gamma > 0$. Moreover, by using an extraction argument together with consistency of marginals along $\gamma \rightarrow \infty$ and $k \rightarrow \infty$, Theorem 1.3 also implies that any sequence from $\{\mathbb{P}_{n;\ell,r}\}$ or from $\{\mathbb{P}_{n;\ell,r}^0\}$ has a subsequence that converges weakly to a probability measure \mathbb{P} on $\mathbf{C}(\mathbb{R}, \bar{\mathbb{A}}_\infty^+)$. Concerning the nature of the limit points, the main consequences of our analysis can be summarized as follows.

Theorem 1.4 *Any sub-sequential limit (as $n \rightarrow \infty, \ell \rightarrow -\infty$ and $r \rightarrow \infty$) \mathbb{P} of $\{\mathbb{P}_{n;\ell,r}\}$ or $\{\mathbb{P}_{n;\ell,r}^0\}$ has the following properties:*

- (1) \mathbb{P} is Brownian-Gibbs with respect to (a, λ) -geometric area tilts.
- (2) \mathbb{P} is non-intersecting.
- (3) *Control of maxima: For any $\gamma > 0$, there exists $C = C(\gamma) > 0$ such that for all $k \in \mathbb{N}$, and $M > 0$:*

$$\mathbb{P} \left(\max_{t \in [-\gamma, \gamma]} X_k(t) > \lambda^{-(k-1)/3} M \right) \leq \frac{C}{M}. \tag{1.26}$$

- (4) *Absolute continuity with respect to independent Brownian bridges: For any $k \in \mathbb{N}, M > 0$ and $\gamma > 0$ there exists $\beta = \beta(k, M, \gamma) > 0$ such that*

$$\mathbb{P}(E) \leq \frac{C(\gamma)}{M} + \frac{1}{\beta} \sup_{v, w \in \mathbb{A}_n^+(M)} \Gamma_{k,\gamma}^{v,w}(E), \tag{1.27}$$

for any $E \in \mathcal{B}_{k,\gamma}^i$.

Finally, let us remark that Theorem 1.4 does not imply unicity of limiting points, and therefore it does not allow us to conclude that $\mathbb{P}_{n;\ell,r}$ or $\mathbb{P}_{n;\ell,r}^0$ actually converge as $n \rightarrow \infty, \ell \rightarrow -\infty$ and $r \rightarrow \infty$. Moreover, Theorem 1.4 does not address the issue of ergodicity or even time stationarity of limiting points. Neither it addresses the issue of representation of limiting points as infinite dimensional diffusions.

However, thanks to monotonicity, we shall obtain the desired convergence and time stationarity, at least for the zero-boundary condition field. We believe that the same holds for free boundary conditions, and that the limiting ensembles coincide.

Theorem 1.5 *There is a unique (as $n \rightarrow \infty$, $\ell \rightarrow -\infty$ and $r \rightarrow \infty$) limit \mathbb{P}^0 of the family $\mathbb{P}_{n;\ell,r}^0$. Besides the properties listed in Theorem 1.4, \mathbb{P}^0 is time stationary.*

2 Tightness of Ensembles of Top Paths

We follow the scheme developed in the seminal [6]. Since, however, geometric area tilts preclude exact solutions and, somehow, complicate matters, we had to modify, adjust and eventually simplify the arguments involved.

2.1 Stochastic Domination

As in [3, 6] stochastic domination plays an important technical role in our approach. Equip $\Omega_{n;\ell,r}^+$ with the partial order \prec , defined by

$$\underline{X} \prec \underline{Y} \text{ iff } X_i \prec Y_i, \text{ for all } i = 1, \dots, n,$$

and let \prec^{FKG} denote the associated notion of stochastic domination of probability measures. Recall (1.16). As it was proved in [6].

Lemma 2.1 *For any $n, \ell, r, h_- \prec g_-$ and $h_+ \prec g_+$ the following holds. If, $\underline{x} \prec \underline{u}$ and $\underline{y} \prec \underline{v}$, then*

$$\mathbf{B}_{\ell,r}^{\underline{x},\underline{y}} [\cdot | h_-, h_+] \stackrel{\text{FKG}}{\prec} \mathbf{B}_{\ell,r}^{\underline{u},\underline{v}} [\cdot | g_-, g_+]. \quad (2.1)$$

Our generalization [3] to measures with geometric area tilts reads as:

Lemma 2.2 *For any $n, \ell, r, h_- \prec g_-$, $h_+ \prec g_+$ and $\underline{\rho} \succ \underline{\kappa}$ the following holds. If, $\underline{x} \prec \underline{u}$ and $\underline{y} \prec \underline{v}$, then*

$$\mathbb{P}_{\ell,r}^{\underline{x},\underline{y}} [\cdot | h_-, h_+, \underline{\rho}] \stackrel{\text{FKG}}{\prec} \mathbb{P}_{\ell,r}^{\underline{u},\underline{v}} [\cdot | g_-, g_+, \underline{\kappa}]. \quad (2.2)$$

Moreover, for an n -tuple $\underline{\chi} = \{\chi_1, \dots, \chi_n\}$ of smooth boundary condition let $\underline{\chi}'$ be the n -tuple of corresponding first derivatives. Then,

$$\mathbb{P}_{\ell,r}^{\underline{\xi},\underline{\zeta}} [\cdot | h_-, h_+, \underline{\rho}] \stackrel{\text{FKG}}{\prec} \mathbb{P}_{\ell,r}^{\underline{\nu},\underline{\eta}} [\cdot | g_-, g_+, \underline{\kappa}], \quad (2.3)$$

whenever, $h_- \prec g_-$, $h_+ \prec g_+$, $\underline{\rho} \succ \underline{\kappa}$ and, both $\underline{\xi}' \succ \underline{\nu}'$ and $\underline{\zeta}' \succ \underline{\eta}'$. In particular, (2.3) holds if $\underline{\xi} = \underline{\nu}$ and $\underline{\zeta} = \underline{\eta}$ (by approximation without any assumptions on smoothness).

2.2 Proof of Theorem 1.3

To facilitate the exposition we shall restrict attention to symmetric intervals $[\ell, r] = [-T, T] \supseteq [-\gamma - 2, \gamma + 2]$. The choice of the constant 2 here is purely conventional, and any positive number could be handled with minor modifications.

Following the standard compactness criterion (see [1, Theorem 8.10]), and thanks to the single time tightness statement that follows from Theorem 1.2, the proof of Theorem 1.3 will be reduced to the control of the maximal modulus of continuity

$$m^{k,\gamma}(\underline{X}, \delta) := \max_{1 \leq i \leq k} \sup_{\substack{s, t \in [-\gamma, \gamma]: \\ |s-t| < \delta}} |X_i(s) - X_i(t)|. \quad (2.4)$$

Thus, Theorem 1.3 is a consequence of the following technical estimate.

Proposition 2.1 *Fix $\gamma > 0$, $k \in \mathbb{N}$. For any $\eta, \varepsilon > 0$, there exists $\delta > 0$ such that*

$$\sup_{T \geq \gamma+2, n > k} \mu_{n;T}^{k,\gamma} (m^{k,\gamma}(\underline{X}, \delta) \geq \eta) \leq \varepsilon \quad (2.5)$$

The estimate above holds as well for the zero boundary distribution $\mu_{n;T}^{0,k,\gamma}$.

2.3 Scheme of Proof of Proposition 2.1

We fix $\gamma > 0$, and $k \in \mathbb{N}$. All estimates to be derived below implicitly depend on these two parameters. On the other hand, they will be uniform in $n > k$ and $T \geq \gamma + 2$. We give the details of the proof in the case of free boundary conditions only, since the case of zero boundary conditions requires only cosmetic changes.

Consider the event $E = E(\delta, \eta)$ from (2.5):

$$E = \{m^{k,\gamma}(\underline{X}, \delta) \geq \eta\}. \quad (2.6)$$

Clearly, this event belongs to the σ -algebra $\sigma(X_i(t), t \in [-\gamma, \gamma], i = 1, \dots, k)$. Thus, we want to prove that

$$\mu_{n;T}^{k,\gamma} (E) = \mathbb{P}_{n;T}(E) \leq \varepsilon. \quad (2.7)$$

We shall rely on the fact that the probability of the event E can be made suitably small in the case of k independent Brownian bridges, uniformly in the boundary conditions, provided the latter are taken in a bounded set.

Lemma 2.3 *For any $\eta, \varepsilon > 0$, and $M > 0$, there exists $\delta > 0$ such that*

$$\sup_{\underline{x}, \underline{y} \in \mathbb{A}_k^+(M)} \Gamma_{k,\gamma}^{\underline{x}, \underline{y}} (m^{k,\gamma}(\underline{X}, \delta) \geq \eta) \leq \varepsilon \quad (2.8)$$

Proof This follows by reduction to the case of k independent Brownian bridges from 0 to 0, see e.g. [6, p. 463]. \square

The proof of Proposition 2.1 will be achieved by controlling the probability $\mathbb{P}_{n;T}(E)$ in terms of the probability $\Gamma_{k,\gamma}^{x,y}(E)$, uniformly in $n > k, T \geq \gamma + 2$. Our proof comprises several intermediate steps that can be roughly summarized as:

- Step 1. Control of probabilities of large excursion events for top and bottom paths X_1, X_{k+1} .
- Step 2. Reduction to probabilities without area tilts.
- Step 3. Use of heat kernel bounds for Brownian bridges in conical domains.
- Step 4. Control of boundary values $X_k(\pm\gamma)$ and of $\min_{j=1}^{k-1} (X_j(\pm\gamma) - X_{j+1}(\pm\gamma))$.

2.4 Control of Maxima

For each $M > 0$, define the event

$$H^i(M) := \left\{ \max_{t \in [-\gamma-2, \gamma+2]} X_i(t) \leq \lambda^{-(i-1)/3} M \right\}. \tag{2.9}$$

The main estimates concerning large excursions to be used below are gathered in the next lemma. Recall the notation (1.16). In particular, $\mathbf{B}_{n;T}^{x,y}[\cdot | h]$ denotes the polymer measure with no area tilts and with floor h .

Lemma 2.4 *The following holds for all fixed $k \in \mathbb{N}$ and $\gamma > 0$:*

- (1) *For all $\varepsilon > 0$, there exists $L = L(k, \gamma, \varepsilon) > 0$ such that for all $M > 0, \underline{x}, \underline{y} \in \mathbb{A}_k^+(M)$ and $h < M$, with $x_k > h(-\gamma - 2)$ and $y_k > h(\gamma + 2)$:*

$$\mathbf{B}_{k;\gamma+2}^{x,y} [H^1(M + L) | h] \geq 1 - \varepsilon. \tag{2.10}$$

- (2) *For all $i = 1, \dots, n, \alpha \in (0, 1/2)$ and $M > 0$:*

$$\mathbb{P}_{n,T} (H^i(M) | a, \lambda) \geq 1 - \frac{\xi_\alpha^* + (\gamma + 2)^\alpha}{M}. \tag{2.11}$$

where ξ_α^* is as in Theorem 1.2.

Proof Using Lemma 2.1 and a recursive argument (see e.g. [3, Proposition 2.1]), it is not hard to see that the maximum of X_1 under $\mathbf{B}_{k;T}^{x,y}[\cdot | h]$ is stochastically dominated by M plus the sum of k independent copies of the maximum of the Brownian bridge of length $2T$ between 0 and 0 constrained to be nonnegative. This implies (2.10).

Inequality (2.11) for $i = 1$ is implied by Markov inequality and Theorem 1.2. The case of general i is obtained using Brownian scaling and the stochastic domination Lemma 2.2, see [3, Remark 2] for the details. \square

2.5 Reduction to Probabilities Without Area Tilts

Estimate (2.11) applied to $i = 1$ and $i = k + 1$ implies that by taking M sufficiently large (depending on ε)

$$\mathbb{P}_{n;T}(E) \leq \varepsilon/2 + \mathbb{P}_{n;T}(E \cap H^1(M) \cap H^{k+1}(M)). \quad (2.12)$$

Recall the notation (1.14). Since E depends only on top k paths, it follows that

$$\mathbb{P}_{n;T}(E \cap H^1(M) \cap H^{k+1}(M)) \leq \sup_{f < \lambda^{-k/3}M} \mathbb{P}_{n;T}\left(E \cap \Omega_{k,\gamma+2}^{f,M} \mid \Omega_{k,\gamma+2}^f\right). \quad (2.13)$$

Clearly, given $f < \lambda^{-k/3}M$,

$$\mathbb{P}_{n;T}\left(E \cap \Omega_{k,\gamma+2}^{f,M} \mid \Omega_{k,\gamma+2}^f\right) \leq \mathbb{P}_{n;T}\left(E \cap \Omega_{k,\gamma+2}^{f,M} \mid \underline{X}(\pm(\gamma+2)) \in \mathbb{A}_n^+(M); \Omega_{k,\gamma+2}^{f,2M}\right).$$

By the Brownian-Gibbs property, the last expression is bounded by

$$\sup_{\underline{x}, \underline{y} \in \mathbb{A}_k^+(M)} \frac{\mathbf{B}_{k,\gamma+2}^{\underline{x}, \underline{y}}\left(\Phi(\underline{X}); E \cap \Omega_{k,\gamma+2}^{f,M}\right)}{\mathbf{B}_{k,\gamma+2}^{\underline{x}, \underline{y}}\left(\Phi(\underline{X}); \Omega_{k,\gamma+2}^{f,2M}\right)},$$

where $\Phi(\underline{X}) = e^{-\sum_0^{k-1} a\lambda^j \mathcal{A}_{\gamma+2}(X_i)}$. In the numerator we estimate $\Phi(\underline{X}) \leq 1$ while in the denominator we may estimate

$$\Phi(\underline{X}) \geq e^{-2aM(2\gamma+4)\sum_0^{k-1}\lambda^j} =: \frac{1}{C(M)}.$$

Taking M appropriately large, (2.10) shows that

$$\mathbf{B}_{k,\gamma+2}^{\underline{x}, \underline{y}}\left(\Omega_{k,\gamma+2}^{f,2M}\right) \geq \frac{1}{2}\mathbf{B}_{k,\gamma+2}^{\underline{x}, \underline{y}}\left(\Omega_{k,\gamma+2}^f\right)$$

uniformly in $f < \lambda^{-k/3}M$ and $\underline{x}, \underline{y} \in \mathbb{A}_k^+(M)$. Hence, we infer that the probability term on the right hand side of (2.12) is bounded above by

$$2C(M) \sup_{f < \lambda^{-k/3}M} \sup_{\underline{x}, \underline{y} \in \mathbb{A}_k^+(M)} \mathbf{B}_{k,\gamma+2}^{\underline{x}, \underline{y}}\left(E \cap \Omega_{k,\gamma+2}^{f,M} \mid \Omega_{k,\gamma+2}^f\right). \quad (2.14)$$

Above it is understood that the conditional probability is taken to be zero if either $x_k < f(-\gamma - 2)$ or $y_k < f(\gamma + 2)$.

Recall that for simplicity of notation, we set $\ell = -\gamma - 2$ and $r = \gamma + 2$. Moreover, we call \mathcal{F}_M^k the family of functions $f \geq 0$ on $[\ell, r]$ satisfying $\sup_t f(t) \leq \lambda^{-k/3}M$. Then, probabilities in (2.14) are now written as

$$\sup_{f \in \mathcal{F}_M^k} \sup_{\underline{x}, \underline{y} \in \mathbb{A}_k^+(M)} \mathbf{B}_{k;\ell,r}^{\underline{x}, \underline{y}} \left(E \cap \Omega_{k;\ell,r}^{f,M} \mid \Omega_{k;\ell,r}^f \right). \quad (2.15)$$

Following Sect. 5 in [6] let us set

$$\Omega_{k;\ell,r}^f = \Omega_{k;\gamma}^f \cap \Omega_{k;[\ell,r] \setminus [-\gamma, \gamma]}^f, \quad (2.16)$$

where we use the notation (1.18). Then, since $\Omega_{k;\ell,r}^{f,M} \subset \{X_1(\pm\gamma) \leq M\}$,

$$\begin{aligned} \mathbf{B}_{k;\ell,r}^{\underline{x}, \underline{y}} \left(E \cap \Omega_{k;\ell,r}^{f,M} \mid \Omega_{k;\ell,r}^f \right) &\leq \mathbf{B}_{k;\ell,r}^{\underline{x}, \underline{y}} \left(E; X_1(\pm\gamma) \leq M \mid \Omega_{k;\gamma}^f \Omega_{k;[\ell,r] \setminus [-\gamma, \gamma]}^f \right) \\ &\leq \frac{\mathbf{B}_{k;\ell,r}^{\underline{x}, \underline{y}} \left(E \mid X_1(\pm\gamma) \leq M; \Omega_{k;[\ell,r] \setminus [-\gamma, \gamma]}^f \right)}{\mathbf{B}_{k;\ell,r}^{\underline{x}, \underline{y}} \left(\Omega_{k;\gamma}^f \mid \Omega_{k;[\ell,r] \setminus [-\gamma, \gamma]}^f \right)}, \end{aligned} \quad (2.17)$$

where we relied on elementary inequality

$$\mathbb{P}(AB|CD) = \frac{\mathbb{P}(ABCD)}{\mathbb{P}(CD)} \leq \frac{\mathbb{P}(A|BD)\mathbb{P}(D)}{\mathbb{P}(CD)} = \frac{\mathbb{P}(A|BD)}{\mathbb{P}(C|D)}.$$

For the numerator in (2.17) we use the fact that E depends only on the time interval $[-\gamma, \gamma]$, and the spatial Markov property for k independent Brownian bridges to obtain

$$\mathbf{B}_{k;\ell,r}^{\underline{x}, \underline{y}} \left(E \mid X_1(\pm\gamma) \leq M; \Omega_{k;[\ell,r] \setminus [-\gamma, \gamma]}^f \right) \leq \sup_{\underline{v}, \underline{w} \in \mathbb{A}_k^+(M)} \Gamma_{k,\gamma}^{\underline{v}, \underline{w}}(E). \quad (2.18)$$

The last expression can be made arbitrarily small thanks to the estimate of Lemma 2.3.

In conclusion, to finish the proof of Proposition 2.1 we need to show that for any γ, M and k there exists $\beta = \beta(k, M, \gamma) > 0$, such that

$$\inf_{f \in \mathcal{F}_M^k} \inf_{\underline{x}, \underline{y} \in \mathbb{A}_k^+(M)} \mathbf{B}_{k;\ell,r}^{\underline{x}, \underline{y}} \left(\Omega_{k;\gamma}^f \mid \Omega_{k;[\ell,r] \setminus [-\gamma, \gamma]}^f \right) \geq \beta. \quad (2.19)$$

Here it is understood that $\underline{x}, \underline{y} \in \mathbb{A}_k^+(M)$ also satisfy $x_k > f(\ell), y_k > f(r)$.

2.6 Main Technical Estimates About Constrained Brownian Bridges

In the sequel $k \in \mathbb{N}$ is fixed and we employ the reduced notation $\Omega_{\ell,r}^f, \Omega_\gamma^f, \Omega_{[\ell,r] \setminus [-\gamma, \gamma]}^f$ for the sets of k -tuples of ordered paths.

Let us take a look at our target (2.19). By the spatial Markov property of Brownian bridges,

$$\mathbf{B}_{\ell,r}^{\underline{x},\underline{y}}\left(\Omega_{\gamma}^f \mid \Omega_{[\ell,r]\setminus[-\gamma,\gamma]}^f\right) = \mathbf{B}_{k;\ell,r}^{\underline{x},\underline{y}}\left(\Gamma_{\gamma}^{\underline{X}(-\gamma),\underline{X}(\gamma)}(\Omega_{\gamma}^f) \mid \Omega_{[\ell,r]\setminus[-\gamma,\gamma]}^f\right). \quad (2.20)$$

Definition 2 Given $f \in \mathcal{F}_M^k$ and $\delta > 0$ let us define the set $\mathcal{G}_{k,\gamma}^f(\delta)$ of pairs of boundary conditions $(\underline{x}, \underline{y}) \in \mathbb{A}_k^+ \times \mathbb{A}_k^+$, such that

$$\mathbf{B}_{\gamma}^{\underline{x},\underline{y}}(\Omega_{\gamma}^f) \geq \delta. \quad (2.21)$$

In view of (2.20) our target (2.19) would follow if we show that there exist $\delta > 0$ and $\chi > 0$ such the following estimate holds:

$$\inf_{f \in \mathcal{F}_M^k} \inf_{\underline{x}, \underline{y} \in \mathbb{A}_k^+(M)} \mathbf{B}_{\ell,r}^{\underline{x},\underline{y}}\left(\left(\underline{X}(-\gamma), \underline{X}(\gamma)\right) \in \mathcal{G}_{k,\gamma}^f(\delta) \mid \Omega_{[\ell,r]\setminus[-\gamma,\gamma]}^f\right) \geq \chi. \quad (2.22)$$

It remains to prove the following two lemmas:

Lemma 2.5 Consider sets

$$\mathcal{S}_k(\eta, L) = \left\{ \underline{x} : x_k > 1 + M\lambda^{-k/3}, x_1 < L \right\} \bigcap_{j=1}^{k-1} \{x_j - x_{j+1} > \eta\}. \quad (2.23)$$

Fix $k \in \mathbb{N}$ and $\gamma > 0$. Then for any $\eta > 0$ and $L > 1 + M\lambda^{-k/3} + k\eta$ there exists $\delta > 0$ such that

$$\mathcal{S}_k(\eta, L) \times \mathcal{S}_k(\eta, L) \subset \mathcal{G}_{k,\gamma}^f(\delta) \quad (2.24)$$

for any $f \in \mathcal{F}_M^k$.

Lemma 2.6 There exists $\chi > 0$, $\eta > 0$ and $L > 1 + M\lambda^{-k/3} + k\eta$ such that

$$\inf_{f \in \mathcal{F}_M^k} \inf_{\underline{x}, \underline{y} \in \mathbb{A}_k^+(M)} \mathbf{B}_{\ell,r}^{\underline{x},\underline{y}}\left(\left(\underline{X}(-\gamma), \underline{X}(\gamma)\right) \in \mathcal{S}_k(\eta, L) \times \mathcal{S}_k(\eta, L) \mid \Omega_{[\ell,r]\setminus[-\gamma,\gamma]}^f\right) \geq \chi. \quad (2.25)$$

2.7 Proof of Lemma 2.5

Since $f(t) \leq \lambda^{-k/3}M$ for all $t \in [-\gamma, \gamma]$, and $\underline{x}, \underline{y} \in \mathcal{S}_k(\eta, L)$:

$$\mathbf{B}_{\gamma}^{\underline{x},\underline{y}}\left(\Omega_{k;\gamma}^f\right) \geq \mathbf{B}_{\gamma}^{\underline{x},\underline{y}}\left(\Omega_{k;\gamma}^{\lambda^{-k/3}M}\right).$$

Let $p_{S,T}(\underline{x}, \underline{z})$ denote the density of the measure $\mathbf{P}_{S,T}^{\underline{x}}(X(T) \in d\underline{z}; \Omega_{k,S,T}^+)$. Then, by the translation invariance and by the symmetry of the Brownian motion,

$$\begin{aligned} \mathbf{B}_{\gamma}^{\underline{x}, \underline{y}}\left(\Omega_{k;\gamma}^{\lambda^{-k/3}M}\right) &= \int_{\mathbb{A}_k^+} p_{0,\gamma}(\underline{x} - \lambda^{-k/3}M, \underline{z}) p_{0,\gamma}(\underline{y} - \lambda^{-k/3}M, \underline{z}) d\underline{z} \\ &\geq \int_{\mathbb{A}_k^+(L)} p_{0,\gamma}(\underline{x} - \lambda^{-k/3}M, \underline{z}) p_{0,\gamma}(\underline{y} - \lambda^{-k/3}M, \underline{z}) d\underline{z}. \end{aligned}$$

Above we use $\underline{c} = (c, c, \dots, c)$ for the k -tuple with constant entries $c = \lambda^{-k/3}M$.

Fix $t_0 > 0$. According to (0.3.2) in [28], for any L there exists a non-decreasing function $t \mapsto C(t, L)$ on $[t_0, \infty)$ such that

$$\frac{1}{C(t, L)} U(\underline{x})U(\underline{z}) \leq p_{0,t}(\underline{x}, \underline{z}) \leq C(t, L)U(\underline{x})U(\underline{z}), \quad x_1, z_1 < L. \quad (2.26)$$

where

$$U(\underline{x}) := \prod_{j=1}^k x_j \prod_{i < j} (x_i^2 - x_j^2)$$

is the positive harmonic function in \mathbb{A}_k^+ . It is then clear that if $\underline{x} \in \mathcal{S}_k(\eta, L)$ then

$$U(\underline{x} - \lambda^{-k/3}M) \geq (2\eta)^{k(k-1)/2}.$$

Therefore,

$$\mathbf{B}_{\gamma}^{\underline{x}, \underline{y}}\left(\Omega_{k;\gamma}^f\right) \geq \frac{1}{C^2(\gamma, L)} (2\eta)^{k(k-1)} \int_{\mathbb{A}_k^+(L)} U^2(\underline{z}) d\underline{z}$$

and the proof is complete.

2.8 Proof of Lemma 2.6

The proof comprises two steps:

Step 1. Control of $X_k(\pm(\gamma + 1))$. One first checks that there exists $\epsilon = \epsilon(\gamma, M, k) > 0$, such that for all $f \in \mathcal{F}_M^k$,

$$\inf_{\underline{x}, \underline{y} \in \mathbb{A}_k^+(M)} \mathbf{B}_{\ell,r}^{\underline{x}, \underline{y}}\left(L > X_1(\pm(\gamma + 1)) > X_k(\pm(\gamma + 1)) \geq 1 + M\lambda^{-k/3} \mid \Omega_{[\ell,r] \setminus [-\gamma, \gamma]}^f\right) \geq \epsilon. \quad (2.27)$$

Let us observe that if L is sufficiently large, then the bound (2.27) is a consequence of

$$\inf_{\underline{x}, \underline{y} \in \mathbb{A}_k^+(M)} \mathbf{B}_{\ell, r}^{\underline{x}, \underline{y}} \left(X_1(\pm(\gamma + 1)) > X_k(\pm(\gamma + 1)) \geq 1 + M\lambda^{-k/3} \mid \Omega_{[\ell, r] \setminus [-\gamma, \gamma]}^0 \right) \geq \epsilon. \quad (2.28)$$

Indeed, set $A = \{L > X_1(\pm(\gamma + 1))\}$ and $B = \{X_k(\pm(\gamma + 1)) \geq 1 + M\lambda^{-k/3}\}$. Both A^c and B are monotone non-decreasing. Hence, Lemma 2.1 implies that

$$\mathbf{B}_{\ell, r}^{\underline{x}, \underline{y}} \left(AB \mid \Omega_{[\ell, r] \setminus [-\gamma, \gamma]}^f \right) \geq \mathbf{B}_{\ell, r}^{\underline{x}, \underline{y}} \left(B \mid \Omega_{[\ell, r] \setminus [-\gamma, \gamma]}^0 \right) - \mathbf{B}_{\ell, r}^{\underline{x}, \underline{y}} \left(A^c \mid \Omega_{[\ell, r] \setminus [-\gamma, \gamma]}^{M\lambda^{-k/3}} \right). \quad (2.29)$$

Moreover, for any $\epsilon > 0$,

$$\mathbf{B}_{\ell, r}^{\underline{x}, \underline{y}} \left(A^c \mid \Omega_{[\ell, r] \setminus [-\gamma, \gamma]}^{M\lambda^{-k/3}} \right) < \epsilon/2 \quad (2.30)$$

for all L large enough, uniformly in $\underline{x}, \underline{y} \in \mathbb{A}_k^+(M)$. The estimate (2.30) follows from (2.10) and the observation that, by monotonicity (2.1),

$$\mathbf{B}_{\ell, r}^{\underline{x}, \underline{y}} \left(X_1(\pm(\gamma + 1)) \geq L \mid \Omega_{[\ell, r] \setminus [-\gamma, \gamma]}^{M\lambda^{-k/3}} \right) \leq \mathbf{B}_{\ell, r}^{\underline{x}, \underline{y}} \left(X_1(\pm(\gamma + 1)) \geq L \mid \Omega_{[\ell, r]}^{M\lambda^{-k/3}} \right).$$

The above shows that (2.28) implies (2.27) with ϵ replaced by $\epsilon/2$.

(2.28) will be derived in Sect. 2.9. \square

Step 2. Once bottom boundary conditions at $\pm(\gamma + 1)$ satisfy $x_k, y_k \geq 1 + M\lambda^{-k/3}$ and $x_1, y_1 < L$, one can steer into $\mathcal{S}_k(\eta, L) \times \mathcal{S}_k(\eta, L)$ at $\pm\gamma$.

Lemma 2.7 *Recall (2.16). Fix L and η such that $L > 1 + M\lambda^{-k/3} + k\eta$, in particular such that $\mathcal{S}_k(\eta, L)$ has a positive Lebesgue measure. There exists $\chi = \chi(k, L, \eta) > 0$ such that*

$$\mathbf{B}_{\gamma+1}^{\underline{x}, \underline{y}} \left(\left(\underline{X}(-\gamma), \underline{X}(\gamma) \right) \in \mathcal{S}_k(\eta, L) \times \mathcal{S}_k(\eta, L) \mid \Omega_{[-\gamma-1, \gamma+1] \setminus [-\gamma, \gamma]}^f \right) \geq \chi \quad (2.31)$$

for all $f \in \mathcal{F}_M^k$ and for all $\underline{x}, \underline{y} \in \mathbb{A}_k^+$ satisfying $x_k, y_k \geq 1 + M\lambda^{-k/3}$ and $x_1, y_1 < L$.

Proof Set, for brevity $\widehat{\Omega}_\gamma^f := \Omega_{[-\gamma-1, \gamma+1] \setminus [-\gamma, \gamma]}^f$ and

$$A = A(\eta, L) := \left\{ \left(\underline{X}(-\gamma), \underline{X}(\gamma) \right) \in \mathcal{S}_k(\eta, L) \times \mathcal{S}_k(\eta, L) \right\}.$$

Then, since $0 \leq f \leq M\lambda^{-k/3}$,

$$\mathbf{B}_{\gamma+1}^{\underline{x}, \underline{y}} \left(A \mid \widehat{\Omega}_\gamma^f \right) = \frac{\mathbf{B}_{\gamma+1}^{\underline{x}, \underline{y}} \left(A; \widehat{\Omega}_\gamma^f \right)}{\mathbf{B}_{\gamma+1}^{\underline{x}, \underline{y}} \left(\widehat{\Omega}_\gamma^f \right)} \geq \frac{\mathbf{B}_{\gamma+1}^{\underline{x}, \underline{y}} \left(A; \widehat{\Omega}_\gamma^{M\lambda^{-k/3}} \right)}{\mathbf{B}_{\gamma+1}^{\underline{x}, \underline{y}} \left(\widehat{\Omega}_\gamma^0 \right)}. \quad (2.32)$$

It is straightforward to check that (2.26) implies the following upper bound on $\mathbf{B}_{\gamma+1}^{\underline{x}, \underline{y}} \left(\widehat{\Omega}_\gamma^0 \right)$: There exists $\widehat{C} = \widehat{C}(\gamma, L)$ such that

$$\frac{U(\underline{x})U(\underline{y})}{\widehat{C}(\gamma, L)} \leq \mathbf{B}_{\gamma+1}^{\underline{x}, \underline{y}}(\widehat{\Omega}_\gamma^0) \leq \widehat{C}(\gamma, L)U(\underline{x})U(\underline{y}), \quad (2.33)$$

uniformly in $\underline{x}, \underline{y} \in \mathbb{A}_k^+(L)$. Furthermore, for every \underline{x} with $x_1 < L$ we have

$$U(\underline{x}) < L^k (2L)^{k(k-1)/2} \prod_{i < j} (x_i - x_j).$$

This implies that

$$\mathbf{B}_{\gamma+1}^{\underline{x}, \underline{y}}(\widehat{\Omega}_\gamma^0) \leq \widehat{C}(\gamma, L)(2L)^{k(k+1)} \prod_{i < j} (x_i - x_j) \prod_{i < j} (y_i - y_j). \quad (2.34)$$

For the probability in the numerator in the right-most expression in (2.32) we have

$$\mathbf{B}_{\gamma+1}^{\underline{x}, \underline{y}}\left(A; \widehat{\Omega}_\gamma^{\lambda^{-k/3}M}\right) = \int_{\mathcal{S}_k(\eta, L)} \int_{\mathcal{S}_k(\eta, L)} p_{0,1}(\tilde{\underline{x}}, \tilde{\underline{u}}) q_{2\gamma}(\tilde{\underline{u}}, \tilde{\underline{v}}) p_{0,1}(\tilde{\underline{v}}, \tilde{\underline{y}}) d\underline{u} d\underline{v}, \quad (2.35)$$

where $\tilde{\underline{z}} := \underline{z} - \lambda^{-k/3}M$, $q_{2\gamma}(\tilde{\underline{u}}, \tilde{\underline{v}})$ is the unconstrained quantity defined in (1.4), and $p_{0,1}$ is the density defined before (2.26).

Since $\underline{u}, \underline{v}$ are in $\mathcal{S}_k(\eta, L) \subset \mathbb{A}_k(L)$,

$$q_{2\gamma}(\tilde{\underline{u}}, \tilde{\underline{v}}) \geq \frac{1}{(4\pi\gamma)^{k/2}} e^{-\frac{kL^2}{4\gamma}}.$$

as it readily follows from (1.2). Consequently,

$$\mathbf{B}_{\gamma+1}^{\underline{x}, \underline{y}}\left(A; \widehat{\Omega}_\gamma^{M\lambda^{-k/3}}\right) \geq \frac{(2\eta)^{k(k-1)} e^{-\frac{kL^2}{4\gamma}}}{(4\pi\gamma)^{k/2}} \int_{\mathcal{S}_k(\eta, L)} p_{0,1}(\tilde{\underline{x}}, \tilde{\underline{u}}) d\underline{u} \int_{\mathcal{S}_k(\eta, L)} p_{0,1}(\tilde{\underline{v}}, \tilde{\underline{y}}) d\underline{v}.$$

Using (2.26) once again, we obtain

$$\int_{\mathcal{S}_k(\eta, L)} p_{0,1}(\tilde{\underline{x}}, \tilde{\underline{u}}) d\underline{u} \geq \frac{1}{\widehat{C}(\gamma, L)} U(\tilde{\underline{x}}) \int_{\mathcal{S}_k(\eta, L)} U(\tilde{\underline{u}}) d\underline{u}.$$

We have already shown that $U(\tilde{\underline{u}}) \geq (2\eta)^{k(k-1)/2}$ for $\underline{u} \in \mathcal{S}_k(\eta, L)$. Furthermore, if $x_k > 1 + \lambda^{-k/3}M$ then

$$U(\tilde{\underline{x}}) \geq \prod_{i < j} (x_i - x_j).$$

Therefore

$$\int_{\mathcal{S}_k(\eta, L)} p_{0,1}(\tilde{\underline{x}}, \tilde{\underline{u}}) d\underline{u} \geq \frac{1}{\widehat{C}(\gamma, L)} (2\eta)^{k(k-1)/2} \prod_{i < j} (x_i - x_j)$$

and, consequently,

$$\mathbf{B}_{\gamma+1}^{\underline{x}, \underline{y}} \left(A; \widehat{\Omega}_{\gamma+1}^{M\lambda^{-k/3}} \right) \geq \frac{(2\eta)^{k(k-1)} e^{-\frac{kL^2}{4\gamma}}}{(4\pi\gamma)^{k/2} C^2(\gamma, L)} \prod_{i < j} (x_i - x_j) \prod_{i < j} (y_i - y_j). \quad (2.36)$$

Plugging (2.34) and (2.36) into (2.32), we get the desired estimate. \square

2.9 Proof of (2.28)

By the Markov property, (2.26) and (2.33),

$$\begin{aligned} \mathbf{B}_{\ell, r}^{\underline{x}, \underline{y}} \left(X_k(\pm(\gamma + 1)) \geq 1 + M\lambda^{-k/3}, \Omega_{[\ell, r] \setminus [-\gamma, \gamma]}^0 \right) \\ \geq \int_{\mathcal{S}_k(\eta, 2M) \times \mathcal{S}_k(\eta, 2M)} p_{\ell, -\gamma-1}(\underline{x}, \underline{u}) \mathbf{B}_{\gamma+1}^{\underline{u}, \underline{v}} \left(\widehat{\Omega}_{\gamma}^0 \right) p_{\gamma+1, r}(\underline{v}, \underline{y}) d\underline{u} d\underline{v} \\ \geq \frac{1}{C^2(2, 2M) \widehat{C}(\gamma, 2M)} U(\underline{x}) U(\underline{y}) \int_{\mathcal{S}_k(\eta, 2M) \times \mathcal{S}_k(\eta, 2M)} U^2(\underline{u}) U^2(\underline{v}) d\underline{u} d\underline{v} \\ =: C_1(\gamma, \eta, M) U(\underline{x}) U(\underline{y}). \end{aligned}$$

Furthermore, using upper bounds in (2.26) and (2.33), we infer that there exists $C_2(\gamma, M)$ such that

$$\begin{aligned} \mathbf{B}_{\ell, r}^{\underline{x}, \underline{y}} \left(\Omega_{[\ell, r] \setminus [-\gamma, \gamma]}^0 \right) &= \int p_{\ell, -\gamma-1}(\underline{x}, \underline{u}) \mathbf{B}_{\gamma+1}^{\underline{u}, \underline{v}} \left(\widehat{\Omega}_{\gamma}^0 \right) p_{\gamma+1, r}(\underline{v}, \underline{y}) d\underline{u} d\underline{v} \\ &\leq C_2(\gamma, M) U(\underline{x}) U(\underline{y}). \end{aligned}$$

Therefore,

$$\inf_{\underline{x}, \underline{y} \in \mathbb{A}_n^+(M)} \mathbf{B}_{\ell, r}^{\underline{x}, \underline{y}} \left(X_k(\pm(\gamma + 1)) \geq 1 + M\lambda^{-k/3} \mid \Omega_{[\ell, r] \setminus [-\gamma, \gamma]}^0 \right) \geq \chi_1 \quad (2.37)$$

holds with $\chi_1 := C_1(\gamma, \eta, M)/C_2(\gamma, M)$. This concludes the proof of (2.28).

3 Limiting Line Ensemble

In this section we prove Theorems 1.4 and 1.5. The key technical tool will be a uniform control of the minimal gaps between lines.

3.1 Minimal Gaps

As in [6], for the purpose of analyzing the limiting line ensemble, an important quantity to be controlled is the minimal gap between the paths, defined as

$$g^{k,\gamma}(\underline{X}) := \min_{1 \leq i \leq k-1} \inf_{s \in [-\gamma, \gamma]} |X_{i+1}(s) - X_i(s)|. \quad (3.1)$$

The main result concerning this quantity is the following

Proposition 3.1 *For any $\varepsilon > 0$, there exists $\delta > 0$ such that*

$$\sup_{T \geq \gamma+2, n > k} \mu_{n;T}^{k,\gamma} (g^{k,\gamma}(\underline{X}) \leq \delta) \leq \varepsilon \quad (3.2)$$

The estimate above holds as well for the zero boundary distribution $\mu_{n;T}^{0,k,\gamma}$.

We remark that (3.2) shows that any limiting point of $\{\mu_{n;T}^{k,\gamma}, T \geq \gamma + 2, n > k\}$, must be concentrated on $\mathbf{C}([-\gamma, \gamma]; \mathbb{A}_k^+)$. Since this holds for all γ and k , it follows that if \mathbb{P} is a limiting point of the free or the zero boundary measures as in Theorem 1.4, then \mathbb{P} must be concentrated on $\mathbf{C}(\mathbb{R}, \mathbb{A}_\infty^+)$, that is \mathbb{P} is non-intersecting.

3.2 Proof of Proposition 3.1

In view of the strategy which we have already employed to prove Proposition 2.1, we will first reduce the problem to the case of Brownian measures without area tilts. Once this is accomplished, one could adapt the proof in [6, Proposition 5.6] to obtain the desired result. However, below we will actually provide an alternative, perhaps more direct proof of the same estimate. We provide the proof in the case of free boundary conditions only, since the same argument applies with no modifications to the case of zero boundary conditions.

Let $E = E(\delta, k, \gamma)$ denote the event

$$E = \{g^{k,\gamma}(\underline{X}) \leq \delta\}. \quad (3.3)$$

As in (2.12)–(2.13) we may restrict ourselves to an upper bound on

$$\sup_{f < \lambda^{-k/3} M} \mathbb{P}_{n;T} \left(E \cap \Omega_{k,\gamma+2}^{f,M} \right).$$

Repeating the argument leading to (2.15), it suffices to show that for all $\varepsilon > 0$, $M > 0$ there exists $\delta > 0$ such that

$$\sup_{f \in \mathcal{F}_M^k} \sup_{\underline{x}, \underline{y} \in \mathbb{A}_k^+(M)} \mathbf{B}_{k;\ell,r}^{\underline{x},\underline{y}} \left(E \cap \Omega_{k;\ell,r}^{f,M} \mid \Omega_{k;\ell,r}^f \right) \leq \varepsilon, \quad (3.4)$$

where $\ell = -\gamma - 2$ and $r = \gamma + 2$.

We now prove (3.4). Fix $\gamma > 0$, and $f \in \mathcal{F}_M^k$. Fix boundary conditions $\underline{x}, \underline{y} \in \mathbb{A}_k^+(M)$ satisfying $x_k > f(\ell)$, $y_k > f(r)$. By (2.19),

$$\mathbf{B}_{k;\ell,r}^{\underline{x},\underline{y}} \left(\Omega_{k;\ell,r}^f \right) \geq \beta \mathbf{B}_{k;\ell,r}^{\underline{x},\underline{y}} \left(\Omega_{k;[\ell,r] \setminus [-\gamma,\gamma]}^f \right) \quad (3.5)$$

Thus, it would be enough to check that regardless of $f, \underline{x}, \underline{y}$ fixed as above,

$$\mathbf{B}_{k;\ell,r}^{\underline{x},\underline{y}} \left(\Omega_{k;\ell,r}^{f,M} \cap E \right) \leq \nu \mathbf{B}_{k;\ell,r}^{\underline{x},\underline{y}} \left(\Omega_{k;[\ell,r] \setminus [-\gamma,\gamma]}^f \right) \quad (3.6)$$

where $E = E(\delta, k, \gamma)$ are the events defined in (3.3), whereas $\nu = \nu(\delta, k, \gamma)$ satisfies:

$$\lim_{\delta \rightarrow 0} \nu(\delta, k, \gamma) = 0. \quad (3.7)$$

Now, by (2.25) there exists $\xi = \xi(k, \gamma) > 0$, such that

$$\mathbf{B}_{k;\ell,r}^{\underline{x},\underline{y}} \left(\Omega_{k;[\ell,r] \setminus [-\gamma,\gamma]}^f \right) \geq \xi \mathbf{P}_{k;\ell}^{\underline{x}} \left(\Omega_{k;\ell,-\gamma}^f \right) \widehat{\mathbf{P}}_{k;r}^{\underline{y}} \left(\Omega_{k;\gamma,r}^f \right), \quad (3.8)$$

where $\mathbf{P}_{k;\ell}^{\underline{x}}$ denotes the measure associated to k independent Brownian motions started at \underline{x} at time ℓ and $\widehat{\mathbf{P}}_{k;r}^{\underline{y}}$ is the notation for the time-reversed k -tuple of independent Brownian motions started at time r at \underline{y} .

On the other hand,

$$\mathbf{B}_{k;\ell,r}^{\underline{x},\underline{y}} \left(\Omega_{k;\ell,r}^{f,M} \cap E \right) \leq \mathbf{P}_{k;\ell}^{\underline{x}} \left(\Omega_{k;\ell,-\gamma}^f \right) \widehat{\mathbf{P}}_{k;r}^{\underline{y}} \left(\Omega_{k;\gamma,r}^f \right) \max_{\underline{u}, \underline{v} \in \mathbb{A}_k^+(M)} \mathbf{B}_{k;\gamma}^{\underline{u},\underline{v}} \left(\Omega_{k;\gamma}^0 \cap E \right) \quad (3.9)$$

If we show that

$$\max_{\underline{u}, \underline{v} \in \mathbb{A}_k^+(M)} \mathbf{B}_{k;\gamma}^{\underline{u},\underline{v}} \left(\Omega_{k;\gamma}^0 \cap E(\delta, k, \gamma) \right) \leq \nu(\delta, k, \gamma), \quad (3.10)$$

with ν satisfying (3.7), then (3.6) will hold with $\nu' = \nu/\xi$.

We first note that

$$\mathbf{B}_{k;\gamma}^{\underline{u},\underline{v}} \left(\Omega_{k;\gamma}^0 \cap E(\delta, k, \gamma) \right) \leq \sum_{j=1}^{k-1} \mathbf{B}_{k;\gamma}^{\underline{u},\underline{v}} \left(\inf_{|t| \leq \gamma} (X_j(t) - X_{j+1}(t)) \in (0, \delta) \right)$$

Then, by the scaling property of the Brownian motion,

$$\max_{\underline{u}, \underline{v} \in \mathbb{A}_k^+(M)} \mathbf{B}_{k;\gamma}^{\underline{u}, \underline{v}}(\Omega_{k;\gamma}^0 \cap E(\delta, k, \gamma)) \leq (k-1) \max_{x, y > 0} \mathbf{B}_{2\gamma}^{x, y} \left(\inf_{|t| \leq 2\gamma} X(t) \in (0, \delta) \right)$$

If $x \leq \sqrt{2}\delta$ then

$$\begin{aligned} \mathbf{B}_{2\gamma}^{x, y} \left(\inf_{|t| \leq 2\gamma} X(t) \in (0, \delta) \right) &\leq \int_0^\infty \mathbf{P}_{-2\gamma, 0}^x \left(\inf_{t \in [-2\gamma, 0]} X(t) > 0, X(0) \in dz \right) \mathbf{B}_{0, 2\gamma}^{z, y} (1) \\ &\leq \frac{1}{2\sqrt{\pi\gamma}} \mathbf{P}_{-2\gamma, 0}^x \left(\inf_{t \in [-2\gamma, 0]} X(t) > 0 \right) \\ &\leq \frac{x}{\sqrt{2\pi\gamma}} \leq \frac{\delta}{\pi\gamma}, \end{aligned}$$

in the last step we have used the reflection principle. By the symmetry,

$$\mathbf{B}_{2\gamma}^{x, y} \left(\inf_{|t| \leq 2\gamma} X(t) \in (0, \delta) \right) \leq \frac{\delta}{\pi\gamma},$$

for all $y \leq \sqrt{2}\delta$.

Consider now the remaining case $x, y > \sqrt{2}\delta$. Set

$$\tau = \inf\{t : X(t) = \sqrt{2}\delta\}.$$

Then we have

$$\begin{aligned} \mathbf{B}_{2\gamma}^{x, y} \left(\inf_{|t| \leq 2\gamma} X(t) \in (0, \delta) \right) &= \mathbf{B}_{2\gamma}^{x, y} \left(\inf_{|t| \leq 2\gamma} X(t) > 0, \tau < 2\gamma \right) \\ &= \mathbf{B}_{2\gamma}^{x, y} \left(\inf_{|t| \leq 2\gamma} X(t) > 0, \tau < 0 \right) + \mathbf{B}_{2\gamma}^{x, y} \left(\inf_{|t| \leq 2\gamma} X(t) > 0, \tau \in (0, 2\gamma) \right). \end{aligned}$$

By the strong Markov property and by the reflection principle,

$$\begin{aligned} \mathbf{B}_{2\gamma}^{x, y} \left(\inf_{|t| \leq 2\gamma} X(t) > 0, \tau < 0 \right) &= \int_{-2\gamma}^0 \mathbf{P}_{-2\gamma, 0}^x(\tau \in ds) \mathbf{B}_{s, 2\gamma}^{\sqrt{2}\delta, y} \left(\inf_{t \in (s, 2\gamma)} X(t) > 0 \right) \\ &\leq \frac{2\delta}{\pi\gamma} \mathbf{P}_{-2\gamma, 0}^x(\tau < 0). \end{aligned}$$

Therefore, by the symmetry,

$$\mathbf{B}_{2\gamma}^{x, y} \left(\inf_{|t| \leq 2\gamma} X(t) > 0, \tau \in (0, 2\gamma) \right) \leq \frac{2\delta}{\pi\gamma}.$$

As a result,

$$\mathbf{B}_{2\gamma}^{x,y} \left(\inf_{|t| \leq 2\gamma} X(t) \in (0, \delta) \right) \leq \frac{4\delta}{\pi\gamma}.$$

This completes the proof of (3.4).

3.3 Proof of Theorem 1.4

As already noted, the fact that \mathbb{P} is non-intersecting follows from Proposition 3.1. It is also immediate to check that the estimate on the maxima stated in the theorem follows from the control of maxima established in Lemma 2.4. Notice that Lemma 2.4 applies to the zero boundary case as well, as a consequence of the stochastic domination from Lemma 2.2.

Once we have the control of the minimal gaps from Proposition 3.1, to prove the Brownian-Gibbs property of the limiting line ensemble we may use exactly the same coupling argument of [6, Proposition 3.7]. Indeed, it is not hard to check that all the basic properties of Brownian bridges used in that argument can be extended with only minor modifications to the case of Brownian bridges with area tilts.

The absolute continuity statement (1.27) follows by a literal repetition of arguments employed for the proof of Proposition 2.1.

3.4 Proof of Theorem 1.5

In view of Theorem 1.3, we need only show convergence of finite dimensional distributions of $\mathbb{P}_{n:\ell,r}^0$ as $n \rightarrow \infty$, $\ell \rightarrow -\infty$, and $r \rightarrow \infty$. To this end, fix $m \in \mathbb{N}$, let $\mathcal{S} = \{s_1, \dots, s_m\} \subset \mathbb{R}^m$, $\mathcal{I} = \{i_1, \dots, i_m\} \in \mathbb{N}^m$, and let $\mathcal{T} = \{t_1, \dots, t_m\} \in \mathbb{R}_+^m$. Consider the event

$$E(\mathcal{S}, \mathcal{I}, \mathcal{T}) = \{\underline{X} \in \Omega : X_{i_j}(s_j) > t_j, j = 1, \dots, m\}.$$

It suffices to show that for each choice of $\mathcal{S}, \mathcal{I}, \mathcal{T}$ there exists $u(\mathcal{S}, \mathcal{I}, \mathcal{T})$ such that

$$\mathbb{P}_{n:\ell,r}^0(E(\mathcal{S}, \mathcal{I}, \mathcal{T})) \rightarrow u(\mathcal{S}, \mathcal{I}, \mathcal{T}) \tag{3.11}$$

as $n \rightarrow \infty$, $\ell \rightarrow -\infty$, and $r \rightarrow \infty$, regardless of the order of the limits. The main observation here is that, since the event $E(\mathcal{S}, \mathcal{I}, \mathcal{T})$ is increasing, the Brownian-Gibbs property on sub-intervals of $[\ell, r]$ and the stochastic domination from Lemma 2.2 imply monotonicity of the above probabilities:

$$\mathbb{P}_{n:\ell,r}^0(E(\mathcal{S}, \mathcal{I}, \mathcal{T})) \leq \mathbb{P}_{n':\ell',r'}^0(E(\mathcal{S}, \mathcal{I}, \mathcal{T})), \tag{3.12}$$

whenever $n' \geq n$, $r' \geq r$ and $\ell' \leq \ell$. Clearly, (3.12) implies (3.11), and the proof of convergence is complete.

Concerning time-translation invariance, let $\mathcal{S}_t = \{s_1 + t, \dots, s_m + t\}$, $t \in \mathbb{R}$. Then,

$$\mathbb{P}_{n:\ell,r}^0(E(\mathcal{S}_t, \mathcal{I}, \mathcal{T})) = \mathbb{P}_{n:\ell-t,r-t}^0(E(\mathcal{S}, \mathcal{I}, \mathcal{T})). \quad (3.13)$$

Monotonicity (3.12) implies that any two sequences $(n_k^{(1)}, \ell_k^{(1)}, r_k^{(1)})$, $(n_k^{(2)}, \ell_k^{(2)}, r_k^{(2)})$, such that $n_k^{(i)} \leq n_{k+1}^{(i)}$, $\ell_k^{(i)} \geq \ell_{k+1}^{(i)}$, $r_k^{(i)} \leq r_{k+1}^{(i)}$, $(n_k^{(i)}, \ell_k^{(i)}, r_k^{(i)}) \rightarrow (\infty, -\infty, \infty)$; $i = 1, 2$, must satisfy

$$\lim_{k \rightarrow \infty} \mathbb{P}_{n_k^{(1)}; \ell_k^{(1)}, r_k^{(1)}}^0(E(\mathcal{S}, \mathcal{I}, \mathcal{T})) = \lim_{k \rightarrow \infty} \mathbb{P}_{n_k^{(2)}; \ell_k^{(2)}, r_k^{(2)}}^0(E(\mathcal{S}, \mathcal{I}, \mathcal{T})) = u(\mathcal{S}, \mathcal{I}, \mathcal{T}). \quad (3.14)$$

Therefore $\mathbb{P}_{n:\ell-t,r-t}^0(E(\mathcal{S}, \mathcal{I}, \mathcal{T}))$ and $\mathbb{P}_{n:\ell,r}^0(E(\mathcal{S}, \mathcal{I}, \mathcal{T}))$ have the same limit

$$u(\mathcal{S}_t, \mathcal{I}, \mathcal{T}) = u(\mathcal{S}, \mathcal{I}, \mathcal{T}), \quad t \in \mathbb{R}, \quad (3.15)$$

so that \mathbb{P}^0 is invariant by time translations.

3.5 Directions of Further Research and Open Questions

In this concluding Subsection we briefly summarize open problems and future research directions.

3.5.1 Unicity, Time-Ergodicity and Structure of Limiting Line Ensembles

As it has been already mentioned Theorem 1.4 does not imply unicity of limiting line ensembles. There are two possible venues to address this issue, both require insights. Indeed, although there are no ready determinantal formulas for finite dimensional distributions at hand (and hence, unlike [6] one cannot characterize the unique limiting ensemble from tightness considerations), the simple geometric structure of tilts gives hope that there is an appropriate algebraic characterization to be uncovered. On the more analytic part, although mixing estimates of [18] do not directly apply in the case of random floors, one may hope that there is an appropriate adjustment, which would also imply ergodicity without resorting to exact solutions as in [8].

3.5.2 Scaling Limits as $\lambda \downarrow 1$

It would be interesting to explore, how this scaling regime is related to the scaling regimes described in [2].

3.5.3 Random Walks and Level Lines Ensembles

As it was stated in the very beginning, the original motivation was to understand the large scale structure of level lines of low temperature $(2 + 1)$ -dimensional SOS interfaces above a hard wall [4, 5, 17, 19]. A natural intermediate step is to study ensembles of ordered random walks under properly normalized geometric area tilts, and to recover limiting line ensembles in the 1:2:3 diffusive rescaling. In view of unbounded number of paths and in view of geometrically growing tilts such an endeavor clearly requires insights beyond [18]. In the case of level lines one has to employ renormalization procedures of e.g. [15, 16] for an effective finite scale description of level lines in terms of random walks. Growing number of macroscopic level lines and multi-body interactions between them pose additional challenges [14, 15].

References

1. Billingsley, P.: *Convergence of Probability Measures*. Wiley, New York (1968)
2. Bornemann, F.: On the scaling limits of determinantal point processes with kernels induced by Sturm-Liouville operators. *SIGMA Symmetry Integr. Geom.: Methods Appl.* **12**(83), 1–20 (2016)
3. Caputo, P., Ioffe, D., Wachtel, V.: Confinement of Brownian polymers under geometric area tilts. *Electron. J. Probab.* **24**(37), 1–21
4. Caputo, P., Lubetzky, E., Martinelli, F., Sly, A., Toninelli, F.L.: Dynamics of $(2 + 1)$ -dimensional SOS surfaces above a wall: slow mixing induced by entropic repulsion. *Ann. Probab.* **42**(4), 1516–1589 (2014)
5. Caputo, P., Lubetzky, E., Martinelli, F., Sly, A., Toninelli, F.L.: Scaling limit and cube-root fluctuations in SOS surfaces above a wall. *J. Eur. Math. Soc.* **18**(5), 931–995 (2016)
6. Corwin, I., Hammond, A.: Brownian Gibbs property for Airy line ensembles. *Invent. Math.* **195**(2), 441–508 (2014)
7. Corwin, I., Hammond, A.: KPZ line ensemble. *Probab. Theory Relat. Fields* **166**(1–2), 67–185 (2016)
8. Corwin, I., Sun, X., et al.: Ergodicity of the Airy line ensemble. *Electron. Commun. Probab.* **19** (2014)
9. Dai Pra, P., Roelly, S.: An existence result for infinite-dimensional Brownian diffusions with non-regular and non-Markovian drift. *WIAS* (2002)
10. Dauvergne, D., Virág, B.: Basic properties of the Airy line ensemble (2018). [arXiv:1812.00311](https://arxiv.org/abs/1812.00311)
11. Dobrokhotov, S.Y., Minenkov, D.S., Shlosman, S.B.: Asymptotics of wave functions of the stationary Schrödinger equation in the Weyl chamber. *Theor. Math. Phys.* **197**(2), 1626–1634 (2018)
12. Duits, M.: On global fluctuations for non-colliding processes. *Ann. Probab.* **46**(3), 1279–1350 (2018)
13. Ferrari, P.L., Spohn, H., et al.: Constrained Brownian motion: fluctuations away from circular and parabolic barriers. *Ann. Probab.* **33**(4), 1302–1325 (2005)
14. Ioffe, D., Shlosman, S.: Formation of facets for an effective model of crystal growth (2017). [arXiv:1704.06760](https://arxiv.org/abs/1704.06760)
15. Ioffe, D., Shlosman, S., Toninelli, F.L.: Interaction versus entropic repulsion for low temperature Ising polymers. *J. Stat. Phys.* **158**(5), 1007–1050 (2015)
16. Ioffe, D., Velenik, Y.: Ballistic phase of self-interacting random walks. *Analysis and Stochastics of Growth Processes and Interface Models*, pp. 55–79 (2008)

17. Ioffe, D., Velenik, Y.: Low-temperature interfaces: prewetting, layering, faceting and Ferrari-Spohn diffusions. *Markov Processes Relat. Fields* **24**, 487–537 (2018)
18. Ioffe, D., Velenik, Y., Wachtel, V.: Dyson Ferrari–Spohn diffusions and ordered walks under area tilts. *Probab. Theory Relat. Fields*, 11–47 (2017)
19. Lacoin, H.: Wetting and layering for Solid-on-Solid I: Identification of the wetting point and critical behavior. *Commun. Math. Phys.*, 1–42 (2017)
20. Lőrinczi, J., Minlos, R.A.: Gibbs measures for Brownian paths under the effect of an external and a small pair potential. *J. Stat. Phys.* **105**(3–4), 605–647 (2001)
21. Maillard, P., Zeitouni, O.: Slowdown in branching Brownian motion with inhomogeneous variance. *Annales de l’Institut Henri Poincaré, Probabilités et Statistiques* **52**(3), 1144–1160 (2016)
22. Meerson, B., Smith, N.R.: Geometrical optics of constrained Brownian motion: three short stories (2019). [arXiv:1901.04209](https://arxiv.org/abs/1901.04209)
23. Minlos, R., Røelly, S., Zessin, H.: Gibbs states on space-time. *Potential Analysis* **13**(4), 367–408 (2000)
24. Osada, H.: Non-collision and collision properties of Dyson’s model in infinite dimension and other stochastic dynamics whose equilibrium states are determinantal random point fields. In: *Stochastic Analysis on Large Scale Interacting Systems*, pp. 325–343. Mathematical Society of Japan (2004)
25. Osada, H., Spohn, H.: Gibbs measures relative to Brownian motion. *Ann. Probab.*, 1183–1207 (1999)
26. Smith, N.R., Meerson, B.: Geometrical optics of constrained Brownian excursion: from the KPZ scaling to dynamical phase transitions (2018). [arXiv:1811.01565](https://arxiv.org/abs/1811.01565)
27. Spohn, H.: Interacting Brownian particles: a study of Dyson’s model. In: *Hydrodynamic Behavior and Interacting Particle Systems*, pp. 151–179. Springer (1987)
28. Varopoulos, N.T.: Potential theory in conical domains. *Math. Proc. Camb. Philos. Soc.* **125**(2), 335–384 (1999)