



## RESEARCH NOTE/NOTA CIENTÍFICA

### FIRST RECORD OF *ANISAKIS SP.* (NEMATODA, ANISAKIDAE) L3 INFECTING THE BODY CAVITY OF *ATLANTORAJA PLATANA* (CHONDRICHTHYES, RAJIDAE)

### PRIMER REGISTRO DE *ANISAKIS SP.* (NEMATODA, ANISAKIDAE) L3 EN LA CAVIDAD CORPORAL DE *ATLANTORAJA PLATANA* (CHONDRICHTHYES, RAJIDAE)

A.C. Moya<sup>1,2,4</sup>, E.J. Galíndez<sup>1,2</sup>, E.E. Di Giacomo<sup>4,5</sup> & R.D. Tanzola<sup>2,3</sup>

1 Laboratorio de Citología, Histología y Embriología Animal, DBByF, UNS, San Juan 670, 8000, Bahía Blanca, Argentina.

2 Instituto de Investigaciones Biológicas y Biomédicas del Sur (INBIOSUR)-CONICET (Consejo Nacional de Investigaciones Científicas y Tecnológicas).

3 Laboratorio de Patología de Organismos Acuáticos, DBByF, UNS, San Juan 670, 8000, Bahía Blanca, Argentina.

4 CONDROS-Laboratorio de Recursos Ícticos. Instituto de Biología Marina y Pesquera "Alte. Storni", Güemes 1030, CP8520, San Antonio Oeste, Río Negro.

5 Universidad Nacional del Comahue, San Martín 247, CP 8520, San Antonio Oeste, Río Negro, Argentina. Tel/Fax: 54 – 2934 – 21002.

Corresponding-authors: [acmoya83@gmail.com](mailto:acmoya83@gmail.com) ; [rtanzola@uns.edu.ar](mailto:rtanzola@uns.edu.ar)

Neotropical Helminthology, 2015, 9(2), jul-dec: 359-365.

## ABSTRACT

This communication is the first record of the presence of a third stage larva of *Anisakis sp.* infecting *Atlantoraja platana*. The hosts were collected from fishery landings at processing plants of San Antonio Oeste (40° 44' S 64° 57' O) and San Antonio Este ports (40° 49' S 64° 57' O), Río Negro province, Argentina. They were found in the visceral cavity near the epigonal organ, a lymphomyeloid tissue closely associated with gonads and only in cartilaginous fish. The high concentrations of urea in the body fluid and tissues of elasmobranch hosts made an inhospitable environment to the colonization of helminthes. The results produced by this work constitute the first report of L<sub>3</sub> of *Anisakis sp.* in the body cavity of an elasmobranch, in particular *A. platana*, and show the capability of this anisakid to survive in the visceral mass of these hosts.

**Keywords:** *Anisakis sp.* - *Atlantoraja platana* - Epigonal organ - L<sub>3</sub> - Nematodes.

## RESUMEN

En la presente nota se registra por primera vez el tercer estadio larval de *Anisakis sp.* infectando a *Atlantoraja platana* (Günther, 1880). Los hospederos fueron obtenidos en desembarques pesqueros de plantas de procesado de los puertos de San Antonio Oeste (40° 44' S 64° 57' O) y San Antonio Este (40° 49' S 64° 57' O), Provincia de Río Negro, Argentina. Las larvas fueron colectadas en la cavidad visceral de los peces, cerca del órgano epigonal, un tejido linfomieloide estrechamente asociado a las gónadas y exclusivo de los peces cartilagosos. Hay evidencias documentadas que las altas concentraciones de urea en tejidos y fluidos corporales tornan inhabitable el medioambiente celómico para ser colonizado por helmintos. Los resultados expuestos en este trabajo constituyen el primer reporte de L<sub>3</sub> de *Anisakis sp.* en la cavidad corporal de un elasmobranquio, en particular *A. platana*, y demuestran la capacidad de este anisákido para sobrevivir en la masa visceral de estos hospederos.

**Palabras clave:** *Anisakis sp.* L<sub>3</sub> - *Atlantoraja platana* - Epigonal organ - Nematodes.

## INTRODUCTION

The anisakid nematodes parasitize elasmobranches as larvae or adults and are generally internal parasites (Caira, 1990) besides they can be found at integumentary level. The location where larval nematodes can be found range from the uterus (Benz *et al.*, 1987) and ovaries (Rosa-Molinar *et al.*, 1983; Aragort *et al.*, 2002) to superficial body tissues (Ruyck & Chabaud, 1960).

Anisakids use fish as paratenic hosts to close their complex life cycles (Guagliardo *et al.*, 2009). There are relatively few records of nematodiasis in rays (Mc Vicar 1977; Romera, 1993; Tanzola *et al.*, 1998; Borucinska & Heger, 1999; Sanmartín *et al.*, 2000, Knoff *et al.*, 2001; Aragort *et al.*, 2002; Santos *et al.*, 2004; Alvarez *et al.*, 2006; Díaz Andrade *et al.*, 2008). The present communication is the first report of the third stage larva (L<sub>3</sub>) of *Anisakis sp.* infecting the body cavity of *Atlantoraja platana* (Günther, 1880). This skate is an endemic species from the Southwest Atlantic Ocean and one of the most commonly taken as bycatch in the coastal fisheries at the San Matías Gulf (Northern Patagonia, Argentina). Its IUCN Red List Category is “Vulnerable A4bd” (<http://www.iucnredlist.org/details/63110/0>).

## MATERIAL AND METHODS

Males specimens of *A. platana* were collected from fishery landings at processing plants of San Antonio Oeste (40° 44' S 64° 57' O) and San Antonio Este ports (40° 49' S 64° 57' O), Rio Negro province, Argentina. The specimens were taxonomically identified following Coller (2012). Two of fifteen (13.3% prevalence) result parasitized by anisakid larvae. These findings proceed from histopathological examination.

Small pieces of the male reproductive system were fixed in Bouin's solution in seawater for at least 24 h. Afterwards, all material was dehydrated through a graded series of ethanol and embedded in Paraplast®. Sections of 4-5 µm thick were stained by Masson's trichromic stain and hematoxylin-eosin. Selected sections were photographed using an Olympus BX51 light microscope equipped with an Olympus C-7070 digital camera.

Following the criteria of Oshima (1972) the larvae were identified as third stage (L<sub>3</sub>) of *Anisakis sp.* and were characterized by a cuticle with slight ripples, number of muscular cells (41-56 muscular cells per quadrant) and hypodermic cords with “Y-shape” associated with the excretory gland (Figs. 1 y 2). The maximum diameter of the larvae were between 0.35 – 0.37 mm.

## RESULTS AND DISCUSSION

The parasites were found in visceral cavity of two males of *A. platana* (one juvenile and other sexually mature). In the juvenile specimen, larvae were found near the epigonal tissue adjacent to the testes. In this case, inflammatory tissue was evident (Fig. 3). In the mature host, larvae were found near the epigonal organ bordering the genital ducts and no inflammatory tissue was detected (Fig. 4). Despite the presence of larvae, the cytoarchitecture of reproductive and lymphomyeloid structures of the hosts was preserved.

The larvae stages of *Anisakis sp.* present in fish tissues can be determined, in cross section, by the structure of the digestive tract, the morphology of the hypodermic cords and the association of these structures with the excretory gland (Oshima, 1972). However, few characters of the adults are present in larvae, so the specific determination of the

parasite, based on the larvae, is complex (Dick & Choudhury, 1995; Guagliardo *et al.*, 2009).


The general pathology of fish specify that the major diseases caused by nematodes have targeted organs as intestine, liver, heart, body cavity, muscle, gonads (especially ovary) and swim bladder (Williams & Jones, 1994; Dick & Choudhury, 1995). Borucinska & Heger (1999) associated the presence of granulomas scattered throughout different organs (spleen, stomach, spiral valve, kidney and gill septa) in *Isurus oxyrinchus* (Rafinesque, 1891) with nematoda larvae and were attributed likely to drancunculoids.

Santos *et al.* (2004) reported the occurrence of a *Pseudoanisakis* (Ascaridida: Acanthocheilidae) from the intestines of two rajids: *Rioraja agassizii* (Müller & Henle, 1841) and *Psammobatis extenta* (Garman,

1913) from Brazilian southwestern Atlantic waters. Alvarez *et al.* (2006) recorded six species of nematode, among them *A. simplex*, all of them in the lumen of stomach and intestine.

Knoff *et al.*, (2001) reported L<sub>3</sub> of *Anisakis* sp. in the stomach and spiral valve of 263 elasmobranchs of Brazil.

Guagliardo *et al.* (2009) found L<sub>3</sub> of *Anisakis* sp. in a bony fish *Serirolella porosa* (Guichenot, 1848) and Tanzola & Guagliardo (2004) reported L<sub>3</sub> of several anisakids in the intestinal lumen of cartilaginous fish, but no invasion of these in the body cavity. The *Anisakis* larvae found in the present study are morphologically similar and share the range of measurements with those parasitize silver warehou, *S. porosa*, from the same geographical area, San Matias Gulf (Guagliardo *et al.*, 2009).




**Figure 1.** High magnification of the anterior part of a third larva stage of *Anisakis* sp. Arrow shows the cuticle. Asterisks indicate the hypodermic cordons. **Mc:** muscular cells; **Oe:** oesophagus; **Eg:** excretory gland. Scale bar: 70  $\mu$ m.




The body cavity of elasmobranch is rarely parasitized by nematodes (Caira & Healy, 2004). Moravec & Little (1988) reported two species of micropleurid nematodes in bull sharks and Caira & Healy (2004) found larvae of ascarid nematodes in elasmobranchs from the Gulf of California. Williams (1964) and Williams *et al.* (1970) stated that the high concentrations of urea in the body fluid and tissues of elasmobranch hosts made an inhospitable environment to the colonization of helminthes, such as acanthocephalans. This

statement could be extrapolate to explain the few records of larval helminths parasitizing the body cavity and tissues of cartilaginous fishes to date (Díaz Andrade *et al.*, 2008).


Base on this background, the results exposed in this work constitute the first report of L<sub>3</sub> of *Anisakis sp.* in the body cavity of an elasmobranch, in particular *A. platana*, and show the capability of this anisakid to survive in the visceral mass of these hosts.



**Figure 2.** High magnification of the middle part of a third larva stage of *Anisakis sp.* Asterisks indicate hypodermic cords. **In:** intestine. Scale bar: 60  $\mu$ m.



**Figure3.** Transverse section of a third larva stage of *Anisakis sp.* in visceral tissue of a juvenile male of *Atlantoraja platana*. Asterisk depicts inflammatory tissue. **Ct:** connective tissue. Scale bar: 130  $\mu$ m.



**Figure4.** General view of a third larval stage of *Anisakis sp.* in visceral tissue of a mature male of *A. platana*. **Eo:** epigonal organ; **Ct:** connective tissue; **Gd:** genital ducts. Scale bar: 150  $\mu$ m.

## ACKNOWLEDGEMENT

We thank to San Antonio Oeste and San Antonio Este fish-processing plants for providing us the samples. This work was supported by the SGCyT-UNS, PGI: 24/B173.

## BIBLIOGRAPHIC REFERENCES

- Aragot, WS, Alvarez, F, Iglesias, R, Leiro, J & Sanmartín, ML. 2002. *Histodytes microocellatus gen. et sp. nov. (Dracunculoidea: Guyaneimidae), a parasite of Raja microocellata on the European Atlantic coast (north-Western Spain)*. Parasitology Research, vol. 88, pp. 932-940.
- Álvarez MF, Aragort, W, Leiro, JM & Sanmartín, ML. 2006. *Macroparasites of five species of ray (genus Raja) on the northwest coast of Spain*. Diseases of Aquatic Organisms, vol 70, pp. 93-100.
- Benz, GW, Pratt, HL & Adamson, ML. 1987. *Larval philometrid nematodes (Philometridae) from the uterus of a sandbar shark, Carcharhinus plumbeus*. Proceedings of the Helminthological Society of Washington, vol. 54, pp. 154-155.
- Borucinska, JD & Heger, K. 1999. *Disseminated granulomas associated with nematode larvae in a shortfin mako shark*. Journal of Wildlife Diseases, vol. 35, pp. 98-100.
- Caira, JN. 1990. *Metazoan parasites as indicators of elasmobranch biology*. In: Pratt Jr., HL, Gruber, SH & Taniuchi, T (eds.). *Elasmobranchs as Living Resources: Advances in the Biology, Ecology, Systematics, and the Status of the Fisheries*. NOAA Technical Report.
- Caira, JN & Healy, CJ. 2004. *Elasmobranchs as Hosts of Metazoan Parasites*. In: Carrier, JC, Musick, JA & Heithaus, MR (eds.). *Biology of Sharks and their Relatives*. CRC Marine Biology Series, Florida.
- Coller, NM. 2012. *Biología, ecología y explotación de la raya platana Atlantoraja platana (Günther, 1880), (Chondrichthyes, Rajidae) del Golfo San Matías*. Tesis Doctoral de la Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata, Argentina, 179 pp.
- Diaz Andrade, MC, Galíndez, EJ, Estecondo, S & Tanzola, RD. 2008. *Hepatic infection by Terranova sp. (Nematoda, Anisakidae) larvae in Sympterygia acuta (Chondrichthyes, Rajidae)*. Bulletin European Association of Fish Pathologists, vol. 28, pp. 144-147.
- Dick TA & Choudhury A. 1995. *Phylum Nematoda*. In: Wood, PTK (ed). *Fish Diseases and Disorders Vol. 1. Protozoan and metazoan infections*. CAB International.
- Guagliardo, SE, De Salvo, MN, Schwerdt, CB, Galeano, NA & Tanzola, RD. 2009. *Anisákidos del savorín, Seriolella porosa (Pisces: Centrolophidae). Análisis de la interacción parásito-hospedador*. BioScriba, vol. 2, pp. 106-114.
- Knoff M, De São Clemente, SC, Magalhães Pinto, R & Corrêa Gomes, D. 2001. *Nematodes of elasmobranch fishes from the Southern Coast of Brazil*. Memórias do Instituto Oswaldo Cruz, vol. 96, pp. 81-87.
- Mc Vicar, AH. 1977. *Intestinal helminth parasites of the ray Raja naevus in British waters*. Journal of Helminthology, vol. 51, pp. 11-21.
- Moravec, F & Little, MD. 1988. *Granulinema gen. n. a new dracunculoid genus with two new species (G. carcharhini sp. n. and G. smile sp. n.) from the bull shark, Carcharhinus leucas (Valenciennes), from Louisiana, USA*. Folia Parasitologica, vol. 35, pp. 113-120.
- Oshima, T. 1972. *Anisakis and anisakiasis in*



- Japan and adjacent area. Progress of Medical Parasitology in Japan, vol. 4, pp. 301-393.
- Romera, SA. 1993. *Proleptus acutus* (Nematoda: Physalopteridae), a parasite from an Argentinian Skate, *Sympterygia bonapartei* (Pisces: Rajidae). Journal of Parasitology, vol. 79, pp. 620-623.
- Rosa-Molinar, E, Williams, CS & Lichtenfels, JR. 1983. Larval nematodes (Philometridae) in granulomas in ovaries of black-tip sharks, *Carcharhinus limbatus* (Valenciennes). Journal of Wildlife Disease, vol. 19, pp. 275-277.
- Ruyck, R & Chabaud, AG. 1960. Un cas de parasitisme attribuable a des larves de Phlyctainophora lamnae Steiner chez un sélacien, et cycle évolutif probable de ce nematode. Vie Milieu, vol. 11, pp. 386-389.
- Sanmartín, ML, Alvarez, MS, Peris, D, Iglesias, R, Leiro, J. 2000. Parasite community study of the undulate ray *Raja undulata* in the Ría of Muros (Galicia, northwest Spain). Aquaculture, vol. 184, pp. 189-201.
- Santos, CP, Lent, H & Gibson, DI. 2004. A new species of *Pseudanisakis* Layman & Borovkova, 1926 (Nematoda: Ascaridida) from *Rioraja agassizii* and *Psammobatis extenta* (Rajidae) in Brazilian southwestern Atlantic waters. Systematic Parasitology, vol. 57, pp. 229-235.
- Tanzola, RD, Guagliardo, SE, Brizzola, SM, Arias, MV & Botte, SE. 1998. Parasite assemblage of *Sympterygia bonapartei* (Pisces: Rajidae), an endemic skate of the Southwest Atlantic. Helminthologia, vol. 35, pp. 123-129.
- Tanzola, RD & Guagliardo, SE. 2004. Nematodes anisákidos presentes en peces del área de Bahía Blanca y el riesgo potencial de anisakidosis humana. Revista Científica de la AMBB, vol. 14, pp. 67-73.
- Williams, HH. 1964. Observations on the helminths of *Raja*. Parasitology, vol. 54, p. 5.
- Williams, HH, McVicar, AH & Ralph, R. 1970. The alimentary canal of fish as an environment for helminth parasites. Symposia of the British Society for Parasitology, vol. 8, pp. 43-77.
- Williams, H & Jones, A. 1994. Parasitic worms of fish. Taylor and Francis Ltd. London, 593 pp.

Received June 29, 2015.  
Accepted September 26, 2015.