

Analysis of antennal sensilla patterns of *Rhodnius prolixus* from Colombia and Venezuela

Lyda Esteban/⁺, Víctor Manuel Angulo, M Dora Feliciangeli*, Silvia Catalá**

Centro de Investigaciones en Enfermedades Tropicales, Universidad Industrial de Santander, Bucaramanga, Colombia *Sección de Entomología Médica, Universidad de Carabobo, Maracay, Venezuela **Centro Regional de Investigación y Transferencia Tecnológica de la Rioja, Crilar, Argentina

Antennal sensilla patterns were used to analyze population variation of domestic Rhodnius prolixus from six departments and states representing three biogeographical regions of Colombia and Venezuela. Discriminant analysis of the patterns of mechanoreceptors and of three types of chemoreceptors on the pedicel and flagellar segments showed clear differentiation between R. prolixus populations east and west of the Andean Cordillera. The distribution of thick and thin-walled trichoids on the second flagellar segment also showed correlation with latitude, but this was not seen in the patterns of other sensilla. The results of the sensilla patterns appear to be reflecting biogeographic features or population isolation rather than characters associated with different habitats and lend support to the idea that domestic R. prolixus originated in the eastern region of the Andes

Key words: Chagas disease - sensilla patterns - *Rhodnius prolixus* - Triatominae - Colombia - Venezuela

Triatominae (Hemiptera, Reduviidae), the insect vectors of Chagas disease (American trypanosomiasis) have been recorded from 30 of the 32 departments of Colombia, from the coast to 2000 m.a.s.l. (Molina et al. 2000). Over 20 species have been reported, of which *Rhodnius prolixus* (Stål) is the most frequently encountered in domestic habitats, representing 95.4% of domestic Triatominae collected, with house infestation rates above 20% in many municipalities (Corredor et al. 1990, Angulo et al. 1997). *R. prolixus* is also widespread in Venezuela where, in spite of extensive control campaigns since 1966, it continues to be the most important domestic vector of Chagas disease (Aché & Matos 2001, Feliciangeli et al. 2003).

Historical reconstruction, combined with biogeographical studies and available genetic comparisons, suggests that *R. prolixus* may have been first domesticated in the savanna-like areas (llanos) of Venezuela, and subsequently spread by accidental human intervention (Dujardin et al. 1998, Schofield & Dujardin 1999, Zeledón 2004). In general however, populations of *R. prolixus* tend to show low levels of genetic or allozyme variability (Dujardin et al. 1998, Monteiro et al. 2000, 2003, Jaramillo et al. 2001), and the aim of the present study was to examine phenetic variability between Colombian and Venezuelan populations using antennal sensilla patterns, which have been shown to be informative characters in studies of population structuring in other species of Triatominae (Gracco & Catalá 2000, Catalá & Dujardin 2001, Carbajal et al. 2002).

MATERIALS AND METHODS

Areas of study - Adults *R. prolixus* were collected from domestic habitats in six departments from three biogeographical regions of Colombia selected on the basis of previous reports and the national entomological survey (Angulo et al. 1997, Molina et al. 2000, Ministerio de Salud 2000). These represented the Andean zone (departments of Santander and Tolima), the Sierra Nevada (departments of Magdalena and Guajira), and the llanos (department of Arauca). *R. prolixus* was also collected from houses in the Venezuelan state of Barinas, which neighbours the department of Arauca (Fig. 1) and from where acute cases of Chagas disease have recently been reported (Añez et al. 1999).

Santander and Tolima in the Andean zone west of the Cordillera are mountainous woodland regions with altitudes from 1000 to 5500 m.a.s.l. The average annual temperature in Santander is 18°C, compared to 28°C in Tolima. Magdalena and Guajira in the Sierra Nevada de Santa Marta, mountain at the Caribbean coast in the North of Colombia, isolated from the rest of the Andean range, with a wide diversity of habitats. The lower slopes are used for agriculture, cattle-ranching and tourism and the upper slopes are inhabited by the indigenous Arhuaco and Kogi cultures. The llanos of Arauca and Barinas are part of an extensive savanna-like plain extending east of the Andes in Colombia and Venezuela. The climate is humid-tropical, with average temperatures of 22-28°C and extensive woodlands with numerous stands of palm trees.

The majority of rural houses in the studied areas are of adobe or 'bahareque' (woven stick and mud) with roofs of palm thatch or corrugated metal or fibre-cement sheets. *R. prolixus* was collected from infested houses during 2003-2005, by timed manual collection (using forceps and a torch to see into crevices in the house structure) (Table I). The collected insects were maintained live or placed into 70% ethanol for transportation to our laboratory. From each collection, adults were selected at random until at

Financial support: Colciencias in Colombia (project 1102-04-13009), Fonacit in Venezuela (project 2000001888)

⁺Corresponding author. E-mail: cintrop@uis.edu.co

Received 30 June 2005

Accepted 10 November 2005

least 10 with complete antennae could be included from each of the departments. Magdalena and Guajira were taking as the same department and included in the same biogeographical regions (Fig. 1).

Analysis of antennal sensilla - Antennae were cut from the head using fine forceps, and stored in 70% ethanol prior to diaphanisation in 4% NaOH. After clearing, they were individually slide-mounted in glycerine and examined at 40 \times . Sensilla were classified according to Catalá

and Schofield (1994), and counted along the entire length of the pedicel (P) and the two flagellar segments (F1 and F2). Only one antenna was examined from each specimen. Four sensilla types were used for analysis: BR, mechanoreceptive bristles, BA, basiconics, TH, thin walled trichoids, TK, thick-walled trichoids; these sensilla have shown to be significant in differentiating Triatominae populations (Catalá & Dujardin 2001, Catalá et al. 2004, 2005).

TABLE I
Source of *Rhodnius prolixus* populations used in the analysis

Municipalities	Department/State	Latitude (deg/min)	Longitude (deg/min)	Altitude (masl)
Santa Marta	Magdalena	11°09'	73°18'	300
Dibulla	Guajira	11°09'	73°18'	297
Albania	Santander	5°45'	73°55'	1900
Sucre	Santander	5°55'	73°47'	1640
Jesús Maria	Santander	5°52'	73°48'	1950
Coyaima	Tolima	3°47'	75°12'	350
Arauca	Arauca	6°98'	71°12'	130
Bolívar	Barinas	8°47'	70°30'	990
Barinas	Barinas	7°51'	71°15'	460
Ezequiel Zamora	Barinas	7°52'	71°14'	160




Fig. 1: map of Colombia and Venezuela showing the collection places of *Rhodnius prolixus* in Tolima, Santander, Magdalena, Guajira, and Arauca in Colombia and Barinas in Venezuela.

A total of 53 insects (27 males and 26 females) was included in the analysis. For each population, mean sensilla density (\pm SD) was calculated for each segment (Table II). Logs of these variables were compared by ANOVA or Kruskal-Wallis, with significance of differences estimated by Newman-Keuls with Bonferroni correction for continuity (Sokal & Rohlf 1997) using STATISTICA v6. Discriminant analysis were made between the five populations, and also between populations from the three biogeographical regions, using PADWIN v65 (<http://www.mpl.ird.fr/morphometrics>). Latitudinal variation, which has been observed in populations of *Triatoma dimidiata* (Catalá et al. 2005), was also examined by linear correlation with TH and TK densities on the flagellar segments.

RESULTS

ANOVA between sexes showed no significant differences between sexes for any sensilla pattern in any of the departments studied (data not shown) so that subsequent analysis were made on groups of 10 (5 males and 5 females) or 11 insects (5 males and 6 females) for each department (Table II).

Univariate analysis showed significant differences in densities of pedicel mechanoreceptors (PBR) and flagellar chemoreceptors (F1BA, F1TK, F2TK, F1TH, F2TH), especially those of the first flagellar segment (F1) (Table III). Insects from the Andean region differed from those of the Sierra Nevada only in numbers of F2TH. However,

insects from the llanos differed significantly from those of the Andes and those from la Sierra Nevada by densities of five receptor types (Table III).

Multivariate discriminant analysis using all variables except F2BA and F1BR (due to heteroscedascity) correctly reclassified 86% of all individuals to their respective department. The population from Magdalena-Guajira was 100% correctly reclassified, followed by those of Barinas (90%), Arauca (88%), Santander (80%), and Tolima (72%). Of the four discriminant functions obtained, the first two contributed 94% of the observed variation (63% for the first, and 31% for the second), and the two variables with highest contribution to variability were PBR and F1TH. Plotting individuals within the discriminant space defined by the first two discriminant functions revealed clear groupings according to biogeographical region (Fig. 2). Similar levels of significance were shown for the Mahalanobis distances between these biogeographical groupings, but with insignificant differences between those of Tolima-Santander or Arauca-Barinas; the Magdalena-Guajira population showed no significant grouping with either of these two.

Regression analysis of sensilla density on latitude showed significant increase in TH of the second flagellar segment in accordance with increasing north latitude (Fig. 3A) with a similar decrease in TK (Fig. 3B). However, no such correlation was found for other sensilla types or for TH and TK on the first flagellar segment (data not shown).

TABLE II
Mean (standard deviation) of antennal sensilla of domestic *Rhodnius prolixus* from Colombia and Venezuela

Segment	Sensilla type ^a	Number of specimens				
		Magdalena Guajira	Santander	Tolima	Arauca	Barinas
		(11)	(10)	(11)	(11)	(10)
Pedicel	PBR	137 (13,36)	128 (15,68)	135 (10,27)	176 (20,51)	166 (24,14)
1st flagellar segment	F1BR	30 (4,85)	27 (5,10)	27 (2,16)	38 (22,62)	36 (5,32)
	F1TH	104 (18,95)	77 (15,74)	92 (20,62)	133 (18,64)	131 (23,10)
	F1TK	155 (21,13)	173 (30,86)	163 (21,14)	212 (42,78)	178 (34,41)
	F1BA	26 (6,50)	31 (7,81)	31 (7,59)	39 (7,60)	35 (10,83)
2nd flagellar segment	F2BR	14 (8,09)	11 (2,01)	11 (2,02)	21 (2,60)	15 (3,06)
	F2TH	81 (12,55)	47 (12,74)	43 (12,27)	57 (12,45)	66 (13,59)
	F2TK	203 (33,38)	225 (27,87)	242 (39,44)	265 (42,02)	210 (30,89)
	F2BA	34 (8,98)	38 (5,65)	38 (6,80)	46 (19,78)	35 (12,52)

^a: mecanoreceptors: PBR, F1BR and F2BR; chemoreceptors: thin-walled trichoids (TH), thick-walled trichoids (TK), basiconics (BA). Following Catalá and Schofield (1994).

TABLE III
Univariate analysis of *Rhodnius prolixus* sensilla patterns

ANOVA ^a	Pedicel	1st flagellar segment				2nd flagellar segment			
	PBR	F1BR	F1TH	F1TK	F1BA	F2BR	F2TH	F2TK	F2BA
Between departments	0,000*	0,0318	0,000*	0,002*	0,004*	0,134	0,000*	0,003*	0,563
Llanos vs Andes	0,000*	0,000*	0,000*	0,031	0,041	0,003*	0,000*	0,784	0,705
Andes vs Sierra Nevada	0,249	0,130	0,038	0,170	0,073	0,307	0,000*	0,019	0,176
Sierra Nevada vs llanos	0,000*	0,028	0,001*	0,003*	0,000*	0,330	0,001*	0,072	0,396

^a: significant value after being examined with Bonferroni, were marked with *.




Fig. 2: discriminant analysis using PBR, F1TH, F1TK, F1BA, F2BR, F2TH, F2TK. Each point represents one individual insect on the canonical axes (FD1, FD2), with polygons encompassing the individuals from Arauca (Ar), Barinas (Ba), Santander (Sa), Tolima (To), and Magdalena-Guajira (Mg).

DISCUSSION

The lack of overt sexual dimorphism in *Rhodnius*, and the relatively simple sensilla patterns, for example shown by the absence of chemoreceptors on the pedicel has been suggested to relate to the abundance of these insects in stable habitats such as domestic environments and palm tree crowns (Schofield 1988, Catalá 1997, Carbajal de la Fuente 2002). As shown by Dujardin et al. (1999), Triatominae tend to show reduced sexual dimorphism in domestic populations, compared to those occupying silvatic habitats. In the case of *R. prolixus*, the high densities attained in domestic habitats (Rabinovich et al. 1995, Sandoval et al. 2000) presumably facilitate encounters between sexes, reducing any necessity for sexual dimorphism or for specific means of sexual identity.

Populations of Triatominae occupying similar biogeographical regions will presumably evolve similarly, so that their antennal sensilla patterns can be expected to show similar features (Catalá & Dujardin 2001, Carbajal & Catalá 2002, Jaramillo et al. 2002). In the present study, the high degree of correct reclassification of *R. prolixus* popula-




Fig. 3: latitudinal variation of the thin walled trichoids TH (A) and the thick walled trichoids (TK) (B) of the flagellum 2 in both sexes of populations of *Rhodnius prolixus* from Colombia and Venezuela (for Barinas, the highest latitude was taken from Table I).

tions in accordance with their biogeographical origin supports the idea that sensilla patterns can act as specific indicators for particular populations. However, since all populations examined were taken from similar domestic habitats, the sensilla patterns appear to be reflecting biogeographic features or population isolation rather than characters associated with different habitats.

The analysis of sensilla patterns indicates clear differentiation between *R. prolixus* populations to the east (Arauca and Barinas) and west (Santander and Tolima) of the Andean Cordillera, as has also been indicated by comparative head morphometrics (Esteban et al. 2000). Populations from Magdalena-Guajira do not group directly with

either the west or eastern forms, but appear to be more similar to the western forms in the sense that they are differentiated from the western forms only by the distribution of TH on the second flagellar segment (F2TH). These groupings lend support to the idea that the original forms of domestic *R. prolixus* developed to the east of the Andes, in the region of the llanos of Venezuela and Colombia, being subsequently spread westwards presumably by accidental human intervention along a route following the pass between the Sierra Nevada and the northern tip of the Andean Cordillera (Schofield & Dujardin 1999). The direction of this presumed dispersal is further supported by the fact that for all sensilla types on all segments (except for the F2TH of the Magdalena-Guajira populations); density is reduced from east to west.

The latitudinal variation in the number of F2TK and F2TH, reported for the first time by *R. prolixus*, was also observed in first flagellar segment in populations of *T. dimidiata*, and explained by the authors in relation to the greater housing capacity to Ecuador (Catalá et al. 2005)

If this interpretation is correct, that domestic *R. prolixus* has been dispersed westwards into Colombia presumably by accidental human intervention and that the Andean Cordillera remains as a biogeographical barrier between the eastern populations of the llanos and the western populations of the Magdalena valley, then this could suggest that control interventions would be more successful against the western populations, since these are no longer within the original biogeographical region of the species. There is evidence that *R. prolixus* populations in Central America were also carried there from a Venezuelan source (Dujardin et al. 1998, Zeledón 2004). By contrast, in the llanos of Venezuela and eastern Colombia which are characterized by extensive palm tree stands where silvatic forms of *R. prolixus* have been reported (Gamboa 1963) the biology of domestic *R. prolixus* may require additional studies to reorientate binational control interventions.

ACKNOWLEDGEMENTS

To Dr Gustavo Vallejo, University of Tolima, and the communities of Magdalena, Arauca, Santander, and Barinas, for their assistance in collecting the insects. To Dr JP Dujardin and Dr CJ Schofield for assistance with the analysis. This work benefited from international collaboration through the ECLAT network.

REFERENCES

- Aché A, Matos JA 2001. Interrupting Chagas disease transmission in Venezuela. *Rev Inst Med Trop São Paulo* 43: 37-43.
- Añez N, Carrasco H, Parada H, Crisante G, Rojas A, Gonzalez N, Ramirez IL, Guevara P, Rivero C, Borges R, Scorza JV 1999. Acute Chagas disease in western Venezuela: a clinical, seroparasitologic, and epidemiologic study. *Am J Trop Med Hyg* 60: 215-222.
- Angulo VM, Tarazona Z, Arismendi MJ, Joya MI, Sandoval CM 1997. Distribución de triatomíneos (Hemiptera: Reduviidae) domiciliarios en 26 municipios de Santander, Colombia. "IX Congreso Colombiano de Parasitología y Medicina Tropical". *Biomédica* 17 (Supl. 7): 162-163.
- Carbajal de la Fuente AN, Catalá S 2002. Relationship between antennal sensilla pattern and habitat in six species of Triatominae. *Mem Inst Oswaldo Cruz* 97: 1121-1125.
- Catalá S 1997. Antennal sensilla of Triatominae (Hemiptera: Reduviidae): a comparative study of five genera. *Int J Insect Morphol & Embryol* 26: 67-73.
- Catalá S, Schofield CJ 1994. The antennal sensilla of *Rhodnius*. *J Morphol* 21: 193-203.
- Catalá S, Dujardin JP 2001. Antennal sensilla patterns indicate geographic and ecotopic variability among *Triatoma infestans* populations (Hemiptera: Reduviidae). *J Med Entomol* 38: 423-428.
- Catalá S, Maida DM, Caro-Riaño H, Jaramillo N, Moreno J 2004. Changes associated with laboratory rearing in antennal sensilla patterns of *Triatoma infestans*, *Rhodnius prolixus*, and *Rhodnius pallescens* (Hemiptera, Reduviidae, Triatominae). *Mem Inst Oswaldo Cruz* 99: 25-30.
- Catalá S, Sachetto C, Moreno M, Rosales R, Salazar-Schettino PM, Gorla D 2005. The antennal phenotype of *Triatoma dimidiata* populations and its relationship with species of the *phyllosoma* and *protracta* complexes. *J Med Entomol* 42: 719-725.
- Corredor A, Santacruz MM, Paez S, Guatame LA 1990. *Distribución de los Triatomíneos Domiciliarios en Colombia*, Instituto Nacional de Salud, Bogotá, 144 pp.
- Dujardin JP, Muñoz M, Chavez T, Ponce C, Moreno J, Schofield CJ 1998. The origin of *Rhodnius prolixus* in Central America. *Med Vet Entomol* 12: 113-115.
- Dujardin JP, Steindel M, Chavez T, Machane M, Schofield CJ 1999. Changes in the sexual dimorphism of Triatominae in the transition from natural to artificial habitats. *Mem Inst Oswaldo Cruz* 94: 565-569.
- Esteban L, Sandoval CM, Dujardin JP, Angulo VM 2000. Caracterización morfométrica de *Rhodnius prolixus* provenientes de dos áreas endémicas para la enfermedad de Chagas en Colombia. Memorias, Curso Taller Internacional Biología, Epidemiología y Control de la Tripanosomiasis Americana y Leishmaniosis, Universidad del Tolima, Colombia, p. 73-78.
- Feliciangeli MD, Campbell-Lendrum D, Martinez C, Gonzalez D, Coleman P, Davies C 2003. Chagas disease control in Venezuela: lessons for the Andean region and beyond. *Trends Parasitol* 19: 44-49.
- Gamboa CJ 1963. Comprobación de *Rhodnius prolixus* extradomiciliarios en Venezuela (Comunicación preliminar). *Bol Ofic Sanit Panam* 54: 18-25.
- Gracco M, Catalá S 2000. Inter-specific and developmental differences on the array of antennal chemoreceptors in four species of Triatominae. (Hemiptera: Reduviidae). *Mem Inst Oswaldo Cruz* 95: 67-74.
- Jaramillo C, Montana MF, Castro LR, Vallejo GA, Guhl F 2001. Differentiation and genetic analysis of *Rhodnius prolixus* and *Rhodnius colombiensis* by rDNA and RAPD amplification. *Mem Inst Oswaldo Cruz* 96: 1043-1048.
- Jaramillo N, Calle DA, Caro Riaño H, Calle J 2002. Diferencias morfométricas de *Rhodnius pallescens* provenientes de siete localidades de Colombia y Panamá. Taller Técnico de Estudio Sobre *Rhodnius pallescens*, Su Vigilancia y Control OPS/DPC/CD/273/03, Panamá, Panamá, Sep 16-17.

- Ministerio de Salud 2000. Informe Programa Nacional de Prevención y Control de la Enfermedad de Chagas y la Cardiopatía Infantil, Colombia.
- Molina JA, Gualdrón L, Brochero HL, Olano V, Barrios D, Ghul F 2000. Distribución actual e importancia epidemiológica de las especies de triatomíneos (Reduviidae: Triatominae) en Colombia. *Biomédica* 20: 344-360.
- Monteiro FA, Barrett T, Fitzpatrick S, Cordon-Rosales C, Feliciangeli D, Beard B 2003. Molecular phylogeography of the Amazonian Chagas disease vectors *Rhodnius prolixus* and *R. robustus*. *Mol Ecol* 12: 997-1006.
- Monteiro FA, Wesson DM, Dotson EM, Schofield CJ, Beard CB 2000. Phylogeny and molecular taxonomy of the Rhodniini derived from mitochondrial and nuclear DNA sequences. *Am J Trop Med Hyg* 62: 460-465.
- Rabinovich JE, Gurtler RE, Leal JA, Feliciangeli MD 1995. Density estimates of the domestic vector of Chagas disease, *Rhodnius prolixus* Stål (Hemiptera: Reduviidae), in rural houses in Venezuela. *Bull WHO* 73: 347-357.
- Sandoval CM, Gutiérrez R, Luna S, Amaya A, Esteban L, Ariza H, Angulo VM 2000. High density of *Rhodnius prolixus* in a rural house in Colombia. *Am J Trop Med Hyg* 94: 372-373.
- Schofield CJ 1988. Biosystematics of Triatominae. In MW Service, *Biosystematics of Haematophagous Insects*, Clarendon Press, Oxford, p. 287-312.
- Schofield CJ, Dujardin JP 1999. Theories on the evolution of *Rhodnius*. *Actualidades Biológicas* (Medellin) 21: 183-197.
- Sokal RR, Rohlf FJ 1995. *Biometry*, 3rd ed., WH Freeman, New York, 887 pp.
- Zeledón R 2004. Some historical facts and recent issues related to the presence of *Rhodnius prolixus* (Stål, 1859) (Hemiptera: Reduviidae) in Central America. *Entomol y Vect* 11: 233-246.