

From DNF Compression to Sunflower Theorems via Regularity

Shachar Lovett

University of California, San Diego, CA, USA
slovett@ucsd.edu

Noam Solomon

MIT, Cambridge, MA, USA
noam.solom@gmail.com

Jiapeng Zhang

University of California, San Diego, CA, USA
jpeng.zhang@gmail.com

Abstract

The sunflower conjecture is one of the most well-known open problems in combinatorics. It has several applications in theoretical computer science, one of which is DNF compression, due to Gopalan, Meka and Reingold (Computational Complexity, 2013). In this paper, we show that improved bounds for DNF compression imply improved bounds for the sunflower conjecture, which is the reverse direction of the DNF compression result. The main approach is based on regularity of set systems and a structure-vs-pseudorandomness approach to the sunflower conjecture.

2012 ACM Subject Classification Theory of computation → Randomness, geometry and discrete structures

Keywords and phrases DNF sparsification, sunflower conjecture, regular set systems

Digital Object Identifier 10.4230/LIPIcs.CCC.2019.5

Funding *Shachar Lovett*: Supported by NSF grant CCF-1614023.

Jiapeng Zhang: Supported by NSF grant CCF-1614023.

Acknowledgements We thank anonymous reviewers for insightful suggestions.

1 Introduction

The sunflower conjecture is one of the most well-known open problems in combinatorics. An r -sunflower is a family of r sets S_1, \dots, S_r where all pairwise intersections are the same. A w -set system is a collection of sets where each set has size at most w . Erdős and Rado [3] asked how large can a w -set system be, without containing an r -sunflower. They proved an upper bound of $w!(r-1)^w$, and conjectured that the bound can be improved.

► **Conjecture 1** (Sunflower conjecture, [3]). *Let $r \geq 3$. There is a constant c_r such that any w -set system \mathcal{F} of size $|\mathcal{F}| \geq c_r^w$ contains an r -sunflower.*

60 years later, only lower order improvements have been achieved, and the best bounds are still of the order of magnitude of about w^w for any fixed r , same as in the original theorem of Erdős and Rado. A good survey on the current bounds is [6].

Sunflowers have been useful in various areas in theoretical computer science. Some examples include monotone circuit lower bounds [9, 10], barriers for improved algorithms for matrix multiplication [1] and faster deterministic counting algorithms via DNF compression [5]. The focus on this paper is on this latter application, in particular DNF compression.



© Shachar Lovett, Noam Solomon, and Jiapeng Zhang;
licensed under Creative Commons License CC-BY
34th Computational Complexity Conference (CCC 2019).
Editor: Amir Shpilka; Article No. 5; pp. 5:1–5:14



Leibniz International Proceedings in Informatics
Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany



A DNF (Disjunctive Normal Form) is disjunction of conjunctive terms. The *size* of a DNF is the number of terms, and the *width* of a DNF is the maximal number of literals in a term. It is a folklore result that any DNF of size s can be approximated by another DNF of width $O(\log s)$, by removing all terms of larger width. The more interesting direction is whether DNFs of small width can be approximated by DNFs of small size. Namely - can DNFs of small width be “compressed” while approximately preserving their computational structure?

A beautiful result of Gopalan, Meka and Reingold [5] shows that DNFs of small width can be approximated by small size DNFs. Their proof relies on the sunflower theorem (more precisely, a variant thereof due to Rossman [10] that we will discuss shortly). Before stating their result, we introduce some necessary terminology. We say that two functions $f, g : \{0, 1\}^n \rightarrow \{0, 1\}$ are ε -close if $\Pr[f(x) \neq g(x)] \leq \varepsilon$ over a uniformly chosen input. We say that f is a lower bound of g , or that g is an upper bound of f , if $f(x) \leq g(x)$ for all x .

► **Theorem 2** (DNF compression using sunflowers, sandwiching bounds [5]). *Let f be a width- w DNF. Then for every $\varepsilon > 0$ there exist two width- w DNFs, f_{lower} and f_{upper} such that*

- (i) $f_{\text{lower}}(x) \leq f(x) \leq f_{\text{upper}}(x)$ for all x .
- (ii) f_{lower} and f_{upper} are ε -close.
- (iii) f_{lower} and f_{upper} have size $(w \log(1/\varepsilon))^{O(w)}$.

Recently, Lovett and Zhang [8] improved the dependence of the size of the lower bound DNF on w (but with a worse dependence on ε). In particular, the proof avoids the use of the sunflower theorem.

► **Theorem 3** (DNF compression without sunflowers, lower bound [8]). *Let f be a width- w DNF. Then for every $\varepsilon > 0$ there exists a width- w DNFs f_{lower} such that*

- (i) $f_{\text{lower}}(x) \leq f(x)$ for all x .
- (ii) f_{lower} and f are ε -close.
- (iii) f_{lower} has size $(1/\varepsilon)^{O(w)}$.

It is natural to speculate that a similar bound holds for upper bound DNFs.

► **Conjecture 4** (Improved upper bound DNF compression). *Let f be a width- w DNF. Then for every $\varepsilon > 0$ there exists a width- w DNF f_{upper} such that*

- (i) $f(x) \leq f_{\text{upper}}(x)$ for all x .
- (ii) f_{upper} and f are ε -close.
- (iii) f_{upper} has size $(1/\varepsilon)^{O(w)}$.

To study the connection between DNF compression and sunflowers, we would need an analog of Conjecture 4 for monotone DNFs.

► **Conjecture 5** (Improved upper bound monotone DNF compression). *In Conjecture 4, if f is a monotone DNF, then f_{upper} can also be taken to be a monotone DNF.*

The main result of this paper is that Conjecture 5 implies an improved bound for the sunflower conjecture, with a bound of $(\log w)^{O(w)}$ instead of the current bound of $O(w)^w$. Thus, the connection between sunflower theorems and DNF compression goes both ways. We note that the proof of [5] is also true for monotone DNF compression.

To simplify the presentation, we assume from now on that $w \geq 2$. This will allow us to assume that $\log w > 0$. In any case, for $w = 1$ the sunflower conjecture is trivial, as any 1-set system of size r is an r -sunflower.

► **Theorem 6** (Main theorem). *Assume that Conjecture 5 holds. Then for any $r \geq 3$ there exists a constant c_r such that the following holds. Any w -set system \mathcal{F} of size $|\mathcal{F}| \geq (\log w)^{c_r w}$ contains an r -sunflower.*

In fact, Theorem 6 holds even with a slightly weaker conjecture instead of Conjecture 5, where the size bound can be assumed to be $((\log w)/\varepsilon)^{O(w)}$ instead of $(1/\varepsilon)^{O(w)}$.

1.1 Proof overview

The proof of Erdős and Rado [3] is by a simple case analysis which we now recall. Let \mathcal{F} be a w -set system. Then either \mathcal{F} contains r disjoint sets, which are in particular an r -sunflower; or at most $r - 1$ sets whose union intersects all other sets. In the latter case, there is an element that belongs to a $\frac{1}{(r-1)w}$ fraction of the sets in \mathcal{F} . If we restrict to these sets, and remove the common element, then we reduced the problem to a $(w - 1)$ -set system of size $\frac{|\mathcal{F}|}{(r-1)w}$. The proof concludes by induction.

Our approach is to refine this via a structure-vs-pseudorandomness approach. Either there is a set T of elements that belong to many sets in \mathcal{F} (concretely, at least $|\mathcal{F}|/\kappa^{|T|}$, for an appropriately chosen κ), or otherwise the set system \mathcal{F} is pseudo-random, in the sense that no set T is contained in too many sets in \mathcal{F} . The main challenge is showing that by choosing κ large enough, this notion of pseudo-randomness is useful. This will involve introducing several new concepts and tying them to the sunflower problem.

The following proof overview follows the same structure as the sections in the paper, to ease readability.

Section 2: DNFs and set systems

First, we note that set systems are one-to-one correspondence to monotone DNFs. Formally, we identify a set system $\mathcal{F} = \{S_1, \dots, S_m\}$ with the monotone DNF $f_{\mathcal{F}}(x) = \bigvee_{S \in \mathcal{F}} \bigwedge_{i \in S} x_i$. This equivalence will be useful in the proof, as at different stages one of these viewpoints is more convenient.

The notions of “lower bound DNF” f_{lower} and “upper bound DNF” f_{upper} used in Theorem 2, Theorem 3 and Conjecture 5 have analogs for set systems, which we refer to as *proper* lower bound and upper bound DNFs (or set systems). For the purpose of this high level overview, we ignore this distinction here.

Section 3: Approximate sunflowers

The notion of approximate sunflowers was introduced by Rossman [10]. It relies on the notion of *satisfying set systems*.

Let \mathcal{F} be a set system on a universe X . We say that \mathcal{F} is (p, ε) -*satisfying* if $\Pr_{x \sim X_p}[f_{\mathcal{F}}(x) = 1] > 1 - \varepsilon$, where $f_{\mathcal{F}}$ is the corresponding monotone DNF for \mathcal{F} , and X_p is the p -biased distribution on X . The importance of satisfying set systems in our context is that a $(1/r, 1/r)$ -satisfying set system contains r disjoint sets (Claim 14).

Let $K = \bigcap_{S \in \mathcal{F}} S$ be the intersection of all sets in \mathcal{F} . We say that \mathcal{F} is a (p, ε) -*approximate sunflower* if the set system $\{S \setminus K : S \in \mathcal{F}\}$ is (p, ε) -satisfying. An interesting connection between approximate sunflowers and sunflowers is that a $(1/r, 1/r)$ -approximate sunflower contains an r -sunflower (Corollary 15).

Section 4: Regular set systems

Let \mathcal{D} be a distribution over subsets of X . We say that \mathcal{D} is *regular* if when sampling $S \sim \mathcal{D}$, the probability that S contains any given set T is exponentially small in the size of T . Formally, \mathcal{D} is κ -regular if for any set $T \subseteq X$ it holds that $\Pr_{S \sim \mathcal{D}}[T \subseteq S] \leq \kappa^{-|T|}$.

A set system \mathcal{F} is κ -regular if there exists a κ -regular distribution supported on sets in \mathcal{F} . We show that if \mathcal{F} is κ -regular, then the same holds for any upper bound set system (Claim 18) and any “large enough” lower bound set system (Claim 19). These facts will turn out to be useful later.

Section 5: Regular set systems are $(1/2, 1/2)$ -satisfying

In this section, we focus on regular set systems \mathcal{F} , or equivalently regular DNFs $f = f_{\mathcal{F}}$. We show that, assuming Conjecture 5 (or the slightly weaker Conjecture 21), any κ -regular DNF of width w , where $\kappa = (\log w)^{O(1)}$, is $(1/2, 1/2)$ -satisfying. Namely, $\Pr[f(x) = 1] \geq 1/2$, where x is uniformly chosen. In particular, this implies that \mathcal{F} contains two disjoint sets. However, our goal is to prove that \mathcal{F} contains an r -sunflower for $r \geq 3$, so we are not done yet.

Section 6: Intersecting regular set systems

Let $\alpha(w, r)$ denote the maximal κ such that there exists a κ -regular w -set system without r disjoint sets. It is easy to prove that the sunflower theorem holds for any set system of size $|\mathcal{F}| > \alpha(w, r)^w$ (Claim 29). However, our discussion so far only allows us to bound $\beta(w) = \alpha(w, 2)$; concretely, assuming Conjecture 5 we have $\beta(w) \leq (\log w)^{O(1)}$.

We show (Lemma 30) that nontrivial upper bounds on $\beta(w)$ imply related upper bounds on $\alpha(w, r)$ for every r . Concretely, if $\beta(w) \leq (\log w)^{O(1)}$ then $\alpha(w, r) \leq (\log w)^{c_r}$ where $c_r > 0$ are constants. This concludes the proof, as we get that any w -set system of size $|\mathcal{F}| \geq (\log w)^{c_r w}$ must contain an r -sunflower.

2 DNFs and set systems

A DNF is *monotone* if it contains no negated variables. Monotone DNFs are in one-to-one correspondence with set systems. Formally, if \mathcal{F} is a set system then the corresponding monotone DNF is

$$f_{\mathcal{F}}(x) = \bigvee_{S \in \mathcal{F}} \bigwedge_{i \in S} x_i.$$

In the other direction, if $f = \bigvee_{j \in [m]} \bigwedge_{i \in S_j} x_i$ is a monotone DNF then its corresponding set system is

$$\mathcal{F}_f = \{S_1, \dots, S_m\}.$$

Observe that a w -set system corresponds to a width- w monotone DNF, and vice versa. If X is the set of elements over which \mathcal{F} is defined then we write $\mathcal{F} \subseteq \mathcal{P}(X)$.

To recall, we consider both lower bound and upper bound DNFs. As our main motivation is to better understand sunflowers, we restrict attention to monotone DNFs from now on; however, all the definitions can be easily adapted for general DNFs.

We next define *proper* upper and lower bound DNFs. Proper lower bound DNFs are obtained by removing terms from the DNF, and proper upper bound DNFs are obtained by removing variables from terms in the DNF. We describe both in terms of the corresponding set systems.

► **Definition 7** (Proper lower bound DNF / set system). *Let \mathcal{F} be a set system. A proper lower bound set system for \mathcal{F} is simply a sub set system $\mathcal{F}' \subseteq \mathcal{F}$. Observe that indeed*

$$f_{\mathcal{F}'}(x) \leq f_{\mathcal{F}}(x) \quad \forall x.$$

► **Definition 8** (Proper upper bound DNF / set system). *Let \mathcal{F} be a set system. A proper upper bound set system for \mathcal{F} is a set system \mathcal{F}' that satisfies the following: for each $S \in \mathcal{F}$ there exists $S' \in \mathcal{F}'$ such that $S' \subseteq S$. Observe that indeed*

$$f_{\mathcal{F}'}(x) \geq f_{\mathcal{F}}(x) \quad \forall x.$$

For monotone DNFs, upper bounds and proper upper bounds are the same.

▷ **Claim 9.** Let $\mathcal{F}, \mathcal{F}'$ be set systems over the same universe, such that

$$f_{\mathcal{F}'}(x) \geq f_{\mathcal{F}}(x) \quad \forall x.$$

Then \mathcal{F}' is a proper upper bound set for \mathcal{F} .

Proof. Assume not. Then there exists $S \in \mathcal{F}$ such that there is no $S' \in \mathcal{F}'$ with $S' \subseteq S$. Let $x = 1_S$ be the indicator vector for S . Then $f_{\mathcal{F}}(x) = 1$ but $f_{\mathcal{F}'}(x) = 0$, a contradiction. ◁

► **Corollary 10.** *In Conjecture 5, we may assume that f_{upper} is a proper upper bound DNF for f .*

We note that the lower and upper bound DNFs in [5] are in fact proper lower and upper bounds, and the same holds for the lower bound DNF in [8].

3 Approximate sunflowers

We introduce the notion of *approximate sunflowers*, first defined by Rossman [10]. We first need some notation. Given a finite set X and $0 < p < 1$, we denote by X_p the p -biased distribution over X , where $W \sim X_p$ is sampled by including each $x \in X$ in W independently with probability p . The definition of approximate sunflowers relies on the notion of a *satisfying set system*.

► **Definition 11** (Satisfying set system). *Let $\mathcal{F} \subseteq \mathcal{P}(X)$ be a set system and let $0 < p, \varepsilon < 1$. We say that \mathcal{F} is (p, ε) -satisfying if*

$$\Pr_{W \sim X_p} [\exists S \in \mathcal{F} : S \subseteq W] > 1 - \varepsilon.$$

Equivalently, if $f_{\mathcal{F}} : \{0, 1\}^X \rightarrow \{0, 1\}$ is the DNF corresponding to \mathcal{F} , then \mathcal{F} is (p, ε) -satisfying if

$$\Pr_{x \sim X_p} [f_{\mathcal{F}}(x) = 1] > 1 - \varepsilon.$$

An approximate sunflower is a set system which is satisfying if we first remove the common intersection of all the sets in the set system.

► **Definition 12** (Approximate sunflower). *Let $\mathcal{F} \subseteq \mathcal{P}(X)$ be a set system and let $0 < p, \varepsilon < 1$. Let $K = \bigcap_{S \in \mathcal{F}} S$. Then \mathcal{F} is a (p, ε) -approximate sunflower if the set system $\{S \setminus K : S \in \mathcal{F}\}$ is (p, ε) -satisfying.*

5:6 From DNF Compression to Sunflower Theorems via Regularity

Rossman proved an analog of the sunflower theorem for approximate sunflowers. Li, Lovett and Zhang [7] reproved this theorem by using a connection to randomness extractors.

► **Theorem 13** (Approximate sunflower lemma [10]). *Let \mathcal{F} be a w -set system and let $\varepsilon > 0$. If $|\mathcal{F}| \geq w! \cdot (1.71 \log(1/\varepsilon)/p)^w$ then \mathcal{F} contains a (p, ε) -approximate sunflower.*

To conclude this section, we show that satisfying set systems contain many disjoint sets, and hence approximate sunflowers contain sunflowers.

▷ **Claim 14.** Let $r \geq 2$ and \mathcal{F} be a $(1/r, 1/r)$ -satisfying set system. Then \mathcal{F} contains r pairwise disjoint sets.

Proof. Let $\mathcal{F} \subseteq \mathcal{P}(X)$. Consider a uniform random coloring of X with r colors. A coloring induces a partition of X into $X = W_1 \cup \dots \cup W_r$, where W_c is the set of all elements that attain the color c . Given a color $c \in [r]$, a set $S \in \mathcal{F}$ is c -monochromatic if all its elements attain the color c . Observe that for each color c ,

$$\Pr[\exists S \in \mathcal{F}, S \text{ is } c\text{-monochromatic}] = \Pr[\exists S \in \mathcal{F}, S \subseteq W_c].$$

The marginal distribution of each W_c is $(1/r)$ -biased. By our assumption that \mathcal{F} is $(1/r, 1/r)$ -satisfying, the probability that W_c contains some $S \in \mathcal{F}$ is more than $1 - 1/r$. So by the union bound,

$$\Pr[\forall c \in [r] \exists S \in \mathcal{F}, S \text{ is } c\text{-monochromatic}] > 0.$$

In particular, there exists a coloring where this event happens. Let S_1, \dots, S_r be the sets for which S_c is c -monochromatic. Then S_1, \dots, S_r must be pairwise disjoint. ◀

► **Corollary 15.** *Let \mathcal{F} be a $(1/r, 1/r)$ -approximate sunflower. Then \mathcal{F} contains an r -sunflower.*

Proof. Let $K = \bigcap_{S \in \mathcal{F}} S$. Apply Claim 14 to the set system $\mathcal{F}' = \{S \setminus K : S \in \mathcal{F}\}$ which by assumption is $(1/r, 1/r)$ -satisfying. We obtain that \mathcal{F}' contains r pairwise disjoint sets $S_1 \setminus K, \dots, S_r \setminus K$. This implies that S_1, \dots, S_r form an r -sunflower. ◀

4 Regular set systems

The notion of regularity of a set system is pivotal in this paper. At a high level, a set system is regular if no element belongs to too many sets, no pair of elements belongs to too many sets, and so on. It is closely related to the notion of block min-entropy studied in the context of lifting theorems in communication complexity [4].

► **Definition 16** (Regular distribution). *Let X be a finite set, and let \mathcal{D} be a distribution on subsets $S \subseteq X$. The distribution \mathcal{D} is κ -regular if for any set $T \subseteq X$ it holds that*

$$\Pr_{S \sim \mathcal{D}}[T \subseteq S] \leq \kappa^{-|T|}.$$

► **Definition 17** (Regular set system). *A set system \mathcal{F} is κ -regular if there exists a κ -regular distribution \mathcal{D} supported on the sets in \mathcal{F} .*

The following claims show that if \mathcal{F} is a κ -regular set system then any proper upper bound set system for it is also κ -regular, and any “large” proper lower bound set system is approximately κ -regular.

▷ **Claim 18.** Let \mathcal{F} be a κ -regular set system. Let \mathcal{F}' be a proper upper bound set system for \mathcal{F} . Then \mathcal{F}' is also κ -regular.

Proof. Let \mathcal{D} be a κ -regular distribution supported on \mathcal{F} . Let $\varphi : \mathcal{F} \rightarrow \mathcal{F}'$ be a map such that $\varphi(S) \subseteq S$ for all $S \in \mathcal{F}$. Define a distribution \mathcal{D}' on \mathcal{F}' as follows:

$$\mathcal{D}'(S') = \sum_{S \in \varphi^{-1}(S')} \mathcal{D}(S).$$

Then for any set T ,

$$\Pr_{S' \sim \mathcal{D}'}[T \subseteq S'] = \sum_{S' \in \mathcal{F}': T \subseteq S'} \mathcal{D}'(S') = \sum_{S \in \mathcal{F}: T \subseteq \varphi(S)} \mathcal{D}(S) \leq \sum_{S \in \mathcal{F}: T \subseteq S} \mathcal{D}(S) = \Pr_{S \sim \mathcal{D}}[T \subseteq S]$$

Since \mathcal{D} is a regular distribution, the claim then follows. ◁

▷ **Claim 19.** Let \mathcal{F} be a κ -regular set system, and \mathcal{D} be a κ -regular distribution supported on \mathcal{F} . Let $\mathcal{F}' \subseteq \mathcal{F}$ be a proper lower bound set system for \mathcal{F} , and let $\alpha = \mathcal{D}(\mathcal{F}')$. Then \mathcal{F}' is $(\kappa\alpha)$ -regular.

Proof. Define a distribution \mathcal{D}' on \mathcal{F}' by $\mathcal{D}'(S) = \alpha^{-1}\mathcal{D}(S)$. Then for any non-empty set T ,

$$\Pr_{S \sim \mathcal{D}'}[T \subseteq S] \leq \alpha^{-1} \Pr_{S \sim \mathcal{D}}[T \subseteq S] \leq \alpha^{-1}\kappa^{-|T|} \leq (\kappa\alpha)^{-|T|}. \blacktriangleleft$$

5 Regular set systems are $(1/2, 1/2)$ -satisfying

In this section we use Conjecture 5 to prove that regular enough DNFs are $(1/2, \varepsilon)$ -satisfying, where in light of Claim 14 we care about $\varepsilon = 1/2$. To recall the definitions, a DNF f is $(1/2, \varepsilon)$ -satisfying if for a uniformly chosen x ,

$$\Pr_x[f(x) = 1] > 1 - \varepsilon.$$

Define

$$\gamma(w) = \sup\{\kappa : \exists \kappa\text{-regular } w\text{-set system which is not } (1/2, 1/2)\text{-satisfying}\}.$$

We start by giving a lower bound on $\gamma(w)$, where the motivation is to help the reader gain intuition.

▷ **Claim 20.** $\gamma(w) \geq \log w - O(1)$.

Proof. We construct a κ -regular w -set system which is not $(1/2, 1/2)$ -satisfying, for $\kappa = \log w - O(1)$. Let X_1, \dots, X_w be disjoint sets, each of size $\kappa = \log w - c$ for a constant $c > 0$ to be determined. Let $X = X_1 \cup \dots \cup X_w$. Let $\mathcal{F} \subseteq \mathcal{P}(X)$ be the w -set system of all sets S that contain exactly one element from each set X_i . It is simple to verify that the uniform distribution over \mathcal{F} is κ -regular, and hence \mathcal{F} is κ -regular. Let $W \sim X_{1/2}$. Then

$$\Pr[\exists S \in \mathcal{F}, S \subseteq W] = \Pr[\forall i \in [w], |X_i \cap W| \geq 1] = (1 - 2^{-\kappa})^w = (1 - c/w)^w \leq \exp(-c).$$

In particular, for $c \geq 1$ we get that \mathcal{F} is not $(1/2, 1/2)$ -satisfying. ◁

As we shall soon see, Conjecture 5 implies that the lower bound is not far from tight:

$$\gamma(w) \leq (\log w)^{O(1)}.$$

It will be sufficient to assume a slightly weaker version of Conjecture 5, where we allow the size of f_{upper} to be somewhat bigger.

► **Conjecture 21** (Weaker version of Conjecture 5). *Let $w \geq 2, \varepsilon > 0$. For any monotone width- w DNF f there exists a monotone width- w DNF f_{upper} such that*

- (i) f_{upper} is a proper upper bound DNF for f .
- (ii) f_{upper} and f are ε -close.
- (iii) f_{upper} has size at most $((\log w)/\varepsilon)^{cw}$ for some absolute constant $c > 1$.

► **Lemma 22.** *Assume Conjecture 21 holds. Then there exists a constant $c_0 > 1$ such that the following holds. For $w \geq 2, \varepsilon > 0$ let $\kappa_0(w, \varepsilon) = ((\log w)/\varepsilon)^{c_0}$. Let \mathcal{F} be a w -set system which is $\kappa_0(w, \varepsilon)$ -regular. Then \mathcal{F} is $(1/2, \varepsilon)$ -satisfying.*

► **Corollary 23.** $\gamma(w) \leq \kappa_0(w, 1/2) = (\log w)^{O(1)}$.

We prove Lemma 22 in the remainder of this section. We start with some simple claims that would serve as a base case for Lemma 22 for $w = O(1)$.

▷ **Claim 24.** Let $r \geq 2$. Let \mathcal{F} be a κ -regular w -set system, where $\kappa > w \binom{r}{2}$. Then \mathcal{F} contains r pairwise disjoint sets.

Proof. Let \mathcal{D} be a κ -regular distribution over \mathcal{F} . Sample independently $S, S' \sim \mathcal{D}$. The probability that S, S' intersect is at most

$$\Pr[|S \cap S'| \geq 1] \leq \sum_{i \in S} \Pr[i \in S'] \leq w/\kappa.$$

Let $S_1, \dots, S_r \sim \mathcal{D}$ be chosen independently. Then by the union bound, the probability that two of them intersect is at most $\binom{r}{2} w/\kappa < 1$. In particular, there exist r pairwise disjoint sets in \mathcal{F} . ◁

▷ **Claim 25.** Let $\varepsilon > 0$. Let \mathcal{F} be a κ -regular w -set system, where $\kappa = w(2^w \log(1/\varepsilon))^2$. Then \mathcal{F} is $(1/2, \varepsilon)$ -satisfying.

Proof. Assume $\mathcal{F} \subseteq \mathcal{P}(X)$. Claim 24 implies that \mathcal{F} contains $r = 2^w \log(1/\varepsilon)$ disjoint sets S_1, \dots, S_r . Let $W \sim X_{1/2}$. Then

$$\Pr[\exists S \in \mathcal{F}, S \subseteq W] \geq \Pr[\exists i \in [r], S_i \subseteq W] = 1 - (1 - 2^{-w})^r > 1 - \varepsilon. \quad \blacktriangleleft$$

Proof of Lemma 22. We will need several properties from κ_0 in the proof. To simplify notations, we shorthand $\kappa_0(w/2, \varepsilon)$ for $\kappa_0(\lfloor w/2 \rfloor, \varepsilon)$ throughout. The constant $c > 1$ below is the absolute constant from Conjecture 5. We need a constant $c' > 1$ so that the following conditions are satisfied:

- (i) $\kappa_0(w, \varepsilon) \geq w(2^w \log(1/\varepsilon))^2$ for $w = 1, 2$ and $\varepsilon > 0$.
- (ii) $\kappa_0(w, \varepsilon) \geq ((\log w)/\varepsilon)^{12c}$ for $w \geq 3, \varepsilon > 0$.
- (iii) $\kappa_0(w, \varepsilon) \geq \kappa_0(w/2, \varepsilon(1 - 1/\log w)) + 1$ for $w \geq 3, \varepsilon > 0$.

One can check that the function $\tau(w, \varepsilon) = (\log w)/\varepsilon$ satisfies $\tau(w/2, \varepsilon(1 - 1/\log w)) \leq \tau(w, \varepsilon)$, with equality when w is even. Thus taking $\kappa_0(w, \varepsilon) = ((\log w)^2/\varepsilon)^{c'}$ satisfies the conditions for a large enough $c' \geq 12c$. We then take $c_0 = 2c'$.

The proof of lemma Lemma 22 is by induction on w . The base cases are $w = 1$ and $w = 2$ which follow from Claim 25 and condition (i) on κ_0 . Thus, we assume from now that $w \geq 3$. We need to prove that for $f = f_{\mathcal{F}}$ we have

$$\Pr[f(x) = 0] < \varepsilon.$$

Let $\gamma = \varepsilon/\log w$ and assume that \mathcal{F} is κ -regular for $\kappa = \kappa_0(w, \varepsilon)$. Let $\mathcal{F}_1 = \{S \in \mathcal{F} : |S| \geq w/2\}$ and let $f_1 = f_{\mathcal{F}_1}$ be the corresponding DNF. Applying Conjecture 21 to f_1 with error parameter γ , we obtain that there exists a γ -approximate proper upper bound DNF f_2 for f_1 of size $s = ((\log w)/\gamma)^{cw} \leq ((\log w)/\varepsilon)^{2cw}$. Let \mathcal{F}_2 be the corresponding set system to f_2 , and observe that \mathcal{F}_2 is a proper upper bound set system for \mathcal{F}_1 . Let $\mathcal{F}_3 = (\mathcal{F} \setminus \mathcal{F}_1) \cup \mathcal{F}_2$ and let $f_3 = f_{\mathcal{F}_3}$ be the corresponding DNF. Then

$$\Pr[f(x) = 0] \leq \Pr[f_3(x) = 0] + (\Pr[f_2(x) = 0] - \Pr[f_1(x) = 0]) \leq \Pr[f_3(x) = 0] + \gamma.$$

Next, observe that \mathcal{F}_3 is a proper upper bound set system for \mathcal{F} . As we assume that \mathcal{F} is κ -regular, then by Claim 18 we obtain that \mathcal{F}_3 is also κ -regular. Let \mathcal{D} be a κ -regular distribution supported on \mathcal{F}_3 . Let $\mathcal{F}_4 = \{S \in \mathcal{F}_3 : |S| \geq w/2\}$, where $\mathcal{F}_4 \subseteq \mathcal{F}_2$. As each set $S \in \mathcal{F}_4$ has size $|S| \geq w/2$ then, since \mathcal{D} is κ -regular, we have

$$\mathcal{D}(S) \leq \kappa^{-w/2}.$$

Summing over all $S \in \mathcal{F}_4$ we obtain that

$$\mathcal{D}(\mathcal{F}_4) \leq |\mathcal{F}_4| \cdot \kappa^{-w/2} \leq |\mathcal{F}_2| \cdot \kappa^{-w/2} \leq \left(\left(\frac{\log w}{\varepsilon} \right)^{2c} \kappa^{-1/2} \right)^w.$$

We would need that $\mathcal{D}(\mathcal{F}_4) \leq 1/\kappa$. As $w \geq 3$, this follows from condition (ii) on κ_0 . Let $\mathcal{F}_5 = \mathcal{F}_3 \setminus \mathcal{F}_4$. Then \mathcal{F}_5 is a $(w/2)$ -set system. By Claim 19 \mathcal{F}_5 is κ' -regular for

$$\kappa' = \kappa \cdot \mathcal{D}(\mathcal{F}_5) = \kappa(1 - \mathcal{D}(\mathcal{F}_4)) \geq \kappa - 1.$$

Let $\varepsilon' = \varepsilon(1 - 1/\log w)$. Assumption (iii) on κ_0 gives that $\kappa_0(w/2, \varepsilon') \leq \kappa_0(w, \varepsilon) - 1$. Thus, \mathcal{F}_5 is $\kappa_0(w/2, \varepsilon')$ -regular. Applying the induction hypothesis, if we denote by f_5 the corresponding DNF for \mathcal{F}_5 , then

$$\Pr[f_5 = 0] < \varepsilon'.$$

Finally, as $\mathcal{F}_5 \subseteq \mathcal{F}_3$ we have $\Pr[f_3(x) = 0] \leq \Pr[f_5(x) = 0]$. Putting these together we obtain that

$$\begin{aligned} \Pr[f(x) = 0] &\leq \Pr[f_3(x) = 0] + \gamma \\ &\leq \Pr[f_5(x) = 0] + \gamma < \varepsilon' + \gamma = \varepsilon(1 - 1/\log w) + \varepsilon/\log w = \varepsilon. \end{aligned} \quad \blacktriangleleft$$

6 Intersecting regular set systems

As we showed in Claim 14, if \mathcal{F} is a $(1/r, 1/r)$ -satisfying set system, then it contains an r -sunflower. However we only proved that a regular enough set system is $(1/2, 1/2)$ -satisfying so far. In this section, we prove that this is enough to show the existence of an r -sunflower for any constant r , and with a comparable condition of regularity. Our proof is based on a the study of regular intersecting set systems.

► **Definition 26** (Intersecting set system). *A set system is intersecting if any two sets in it intersect. In other words, it does not contain two disjoint sets.*

► **Definition 27.** *For $w \geq 1, r \geq 2$ define*

$$\alpha(w, r) = \sup\{\kappa : \exists \kappa\text{-regular } w\text{-set system without } r \text{ pairwise disjoint sets}\}.$$

It will be convenient to shorthand $\beta(w) = \alpha(w, 2)$, which can equivalently be defined as

$$\beta(w) = \sup\{\kappa : \exists \kappa\text{-regular intersecting } w\text{-set system}\}.$$

5:10 From DNF Compression to Sunflower Theorems via Regularity

▷ **Claim 28.** $\alpha(w+1, r) \geq \alpha(w, r)$ and $\alpha(w, r+1) \geq \alpha(w, r)$ for all $w \geq 1, r \geq 2$.

Proof. The first claim follows by our definition that a w -set system is a set system where all sets have size at most w . In particular, any w -set system is also a $(w+1)$ -set system and hence $\alpha(w+1, r) \geq \alpha(w, r)$. The second claim holds since a set system that does not contain r disjoint sets, also does not contain $r+1$ disjoint sets. ◁

We start by showing that upper bounds on $\alpha(w, r)$ directly translate to upper bounds on sunflowers. This is reminiscent to the original proof of Erdős and Rado [3].

▷ **Claim 29.** Let \mathcal{F} be a w -set system of size $|\mathcal{F}| > \alpha(w, r)^w$. Then \mathcal{F} contains an r -sunflower.

Proof. The proof is by induction on w . If \mathcal{F} contains r pairwise disjoint sets then we are done. Otherwise, \mathcal{F} is not κ -regular for any $\kappa > \alpha(w, r)$. In particular, the uniform distribution over \mathcal{F} is not κ -regular. This implies that there exists a nonempty set T of size $|T| = t \geq 1$ such that

$$\mathcal{F}' = \{S \setminus T : S \in \mathcal{F}, T \subseteq S\}$$

has size $|\mathcal{F}'| \geq |\mathcal{F}| \kappa^{-t} > \alpha(w, r)^{w-t} \geq \alpha(w-t, r)^{w-t}$. By induction, \mathcal{F}' contains an r -sunflower $S_1 \setminus T, \dots, S_r \setminus T$. Hence S_1, \dots, S_r is a sunflower in \mathcal{F} . ◁

The main lemma we prove in this section is that upper bounds on β imply upper bounds on α .

► **Lemma 30.** For all $w \geq 1, r \geq 3$ it holds that $\alpha(w, r) \leq r2^{r+1}\beta(wr)^r$.

Before proving Lemma 30, we first prove some upper and lower bounds on $\beta(w)$. Although these are not needed in the proof of Lemma 30, we feel that they help gain intuition on $\beta(w)$.

▷ **Claim 31.** $\beta(w) \leq w$.

Proof. Apply Claim 24 for $r = 2$. ◁

It is easy to construct examples that show that $\beta(w) > 1$; for example, the family of all sets of size w in a universe of size $2w-1$ is intersecting and $((2w-1)/w)$ -regular. The following example shows that $\beta(w)$ is super-constant.

▷ **Claim 32.** $\beta(w) \geq \Omega\left(\frac{\log w}{\log \log w}\right)$.

Proof. We construct an example of an intersecting w -set system for $\kappa = \Omega(\log w / \log \log w)$. Let $t \leq w/2$ to be optimized later and set $m = w - t + 1$. Let X_1, \dots, X_m be disjoint sets of size t each, and let $X = X_1 \cup \dots \cup X_m$. Consider the set system \mathcal{F} of all sets $S \subseteq X$ of the following form:

$$\mathcal{F} = \{S \subseteq X : \exists i \in [m], X_i \subseteq S, \forall j \neq i, |X_j \cap S| = 1\}.$$

Observe that \mathcal{F} is an intersecting w -set system.

Let \mathcal{D} be the uniform distribution over \mathcal{F} . We show that \mathcal{D} is κ -regular, and hence \mathcal{F} is κ -regular. There are two extreme cases: for sets T of size $|T| = 1$ we have

$$\Pr_{S \sim \mathcal{D}} [T \subseteq S] = \frac{1}{m} + \left(1 - \frac{1}{m}\right) \frac{1}{t} \leq \frac{2}{t}.$$

For sets $T = X_i$ we have

$$\Pr_{S \sim \mathcal{D}} [X_i \subseteq S] = \frac{1}{m}.$$

One can verify that these are the two extreme cases which control the regularity, and hence \mathcal{F} is κ -regular for

$$\kappa = \min(t/2, m^{1/t}).$$

Setting $t = \Theta(\log w / \log \log w)$ gives $\kappa = \Theta(\log w / \log \log w)$. \triangleleft

We conjecture that this is essentially tight. In fact, by Claim 14 we have that

$$\beta(w) \leq \gamma(w)$$

As we proved, Conjecture 21 implies $\gamma(w) = (\log w)^{O(1)}$, thus it also implies $\beta(w) = (\log w)^{O(1)}$.

Proof of Lemma 30. For $w, r \geq 1$ define

$$\eta(w, r) = r2^{r+1}\beta(wr)^r.$$

We will first prove that

$$\alpha(w, 2r) \leq \max(\eta(w, r), 2\alpha(w, r))$$

and then that this implies the bound

$$\alpha(w, r) \leq \eta(w, r).$$

Let \mathcal{F} be a κ -regular w -set system \mathcal{F} which does not contain $2r$ pairwise disjoint sets, where $\kappa > \max(\eta(w, r), 2\alpha(w, r))$. We will show that this leads to a contradiction.

Let \mathcal{D} be the corresponding κ -regular distribution on \mathcal{F} . Let $\mathcal{F}' \subseteq \mathcal{F}$ be any sub set-system with $\mathcal{D}(\mathcal{F}') \geq 1/2$. Claim 19 then implies that \mathcal{F}' is $(\kappa/2)$ -regular. By our choice of κ , $\kappa/2 > \alpha(w, r)$, and hence \mathcal{F}' contains r pairwise disjoint sets.

More generally, consider the following setup. Let $\mathcal{D}' : \mathcal{F} \rightarrow \mathbb{R}_{\geq 0}$ with $\mathcal{D}'(S) \leq \mathcal{D}(S)$ for all $S \in \mathcal{F}$. Define $\mathcal{F}' = \{S : \mathcal{D}'(S) > 0\}$ and $\mathcal{D}'(\mathcal{F}) = \sum \mathcal{D}'(S)$. As long as $\mathcal{D}'(\mathcal{F}) \geq 1/2$ we are guaranteed that \mathcal{F}' is $(\kappa/2)$ -regular, and hence contains r pairwise disjoint sets. Consider the following process:

1. Initialize $D_0(S) = \mathcal{D}(S)$ for all $S \in \mathcal{F}$ and $i = 0$.
2. As long as $D_i(\mathcal{F}) \geq 1/2$ do:
 - a. Let $\mathcal{F}_i = \{S : D_i(S) > 0\}$.
 - b. Find r pairwise disjoint sets $S_{i,1}, \dots, S_{i,r} \in \mathcal{F}_i$.
 - c. Let $w_i = \min(D_i(S_{i,1}), \dots, D_i(S_{i,r}))$.
 - d. Set $D_{i+1}(S) = D_i(S) - w_i$ if $S \in \{S_{i,1}, \dots, S_{i,r}\}$, and $D_{i+1}(S) = D_i(S)$ otherwise.
 - e. Set $i \leftarrow i + 1$

Assume that the process terminates after m steps. Let $W_i = S_{i,1} \cup \dots \cup S_{i,r}$, which by construction is a set of size wr . Note that as we assume that \mathcal{F} does not contain $2r$ pairwise disjoint sets, we obtain that W_1, \dots, W_m must be an intersecting set system (possibly with some repeated sets). Let $w = \sum w_i$. As $D_{i+1}(\mathcal{F}) = D_i(\mathcal{F}) - w_i r$, and as we terminate when $D_m(\mathcal{F}) < 1/2$, we have

$$w \geq 1/2r.$$

5:12 From DNF Compression to Sunflower Theorems via Regularity

Let $\mathcal{F}^* = \{W_1, \dots, W_m\}$, namely taking each set exactly once. As it may be the case that W_1, \dots, W_m are not all distinct, we only know that $|\mathcal{F}^*| \leq m$. Consider the distribution D^* on \mathcal{F}^* given by $D^*(W) = \frac{1}{w} \sum_{i: W_i=W} w_i$. Then as \mathcal{F}^* is an intersecting set system, we obtain that D^* cannot be β -regular for $\beta = \beta(wr)$.

Thus, there exists a nonempty set T of size $|T| = t \geq 1$ such that

$$\sum_{W \in \mathcal{F}^*: T \subseteq W} D^*(W) \geq \beta^{-t}.$$

This implies that if we denote $I = \{i \in [m] : T \subseteq W_i\}$ then

$$\sum_{i \in I} w_i \geq w\beta^{-t} \geq \frac{1}{2r\beta^t}.$$

Next, consider some $i \in I$. Recall that W_i is the union of pairwise disjoint sets $S_{i,1}, \dots, S_{i,r} \in \mathcal{F}$. In particular, there must exist $j_i \in [r]$ such that $|T \cap S_{i,j_i}| \geq |T|/r$. We denote $T_i = T \cap S_{i,j_i}$. As the number of possible subsets of T is $2^{|T|}$, there must exist $T^* \subseteq T$ such that

$$\sum_{i \in I: T_i = T^*} w_i \geq 2^{-t} \sum_{i \in I} w_i \geq \frac{1}{2r(2\beta)^t}.$$

In particular, $|T^*| \geq |T|/r$ and

$$\sum_{i \in I: T^* \subseteq S_{i,j_i}} w_i \geq \frac{1}{2r(2\beta)^t}.$$

It may be that the list of S_{i,j_i} contains repeated sets (namely, that $S_{i,j_i} = S_{i',j_{i'}}$ for some $i \neq i'$). For each $S \in \mathcal{F}$ let $I(S) = \{i \in I : S_{i,j_i} = S\}$. In particular, $I(S)$ is not empty only for sets S with $T^* \subseteq S$. We can rewrite the sum as

$$\sum_{i \in I: T^* \subseteq S_{i,j_i}} w_i = \sum_{S \in \mathcal{F}: T^* \subseteq S} \sum_{i \in I(S)} w_i.$$

Next, fix some $S \in \mathcal{F}$ with $T^* \subseteq S$ and consider the internal sum. Recall that $w_i = D_i(S) - D_{i+1}(S)$, and hence the sum is a telescopic sum and can be bounded by

$$\sum_{i \in I(S)} w_i \leq D_0(S) - D_m(S) \leq D(S).$$

We thus obtain that

$$\sum_{S \in \mathcal{F}: T^* \subseteq S} \mathcal{D}(S) \geq \sum_{i \in I: T^* \subseteq S_{i,j_i}} w_i \geq \frac{1}{2r(2\beta)^t}.$$

Recall that \mathcal{D} is κ -regular. We can upper bound κ by

$$\kappa \leq (2r(2\beta)^t)^{1/|T^*|} \leq (2r(2\beta)^t)^{r/t} \leq 2r(2\beta)^r = \eta(w, r).$$

Putting everything together, we get

$$\alpha(w, 2r) \leq \max(\eta(w, r), 2\alpha(w, r)).$$

To conclude the proof, note that if r is a power of two then by induction and our choice of η we have

$$\alpha(w, 2r) \leq \max(\eta(w, r), 2\eta(w, r/2), 4\eta(w, r/4), \dots) = \eta(w, r).$$

Thus for a general r , if $r \leq s \leq 2r$ is the smallest power of two that upper bounds r then

$$\alpha(w, r) \leq \alpha(w, s) \leq \eta(w, s/2) \leq \eta(w, r). \quad \blacktriangleleft$$

Proof of Theorem 6. We put all the pieces together. Let $w \geq 2$. Assume Conjecture 21 holds. Lemma 22 gives that

$$\gamma(w) \leq (\log w)^c$$

for some constant $c \geq 1$. Claim 24 then gives that

$$\beta(w) \leq \gamma(w)$$

and Lemma 30 gives that

$$\alpha(w, r) \leq r2^{r+1}\beta(wr)^r \leq r2^{r+1}(\log(wr))^{cr} \leq (\log w)^{c_r}$$

for some constant $c_r \geq 1$. Finally, Claim 29 shows that if \mathcal{F} is a w -set system of size $|\mathcal{F}| \geq (\log w)^{c_r w}$ then \mathcal{F} contains an r -sunflower. \blacktriangleleft

7 Further discussions

Recall that $\beta(w)$ is the maximal κ such that there exists an intersecting κ -regular w -set system.

► **Conjecture 33.** $\beta(w) \leq (\log w)^{O(1)}$.

We would like to point out that in Conjecture 33, the assumption that the set system is intersecting cannot be replaced by a weaker assumption that it is almost intersecting, namely that most pairs of sets intersect. To see that, consider the following example.

► **Example 34.** Let \mathcal{F} be the family of all sets of size w in a universe of size $n = cw^2$. By choosing an appropriate constant $c > 0$, we get that 99% of the sets $S, S' \in \mathcal{F}$ intersect. However, \mathcal{F} is (w/c) -regular.

The following is an interesting family of examples, that might help shed light on Conjecture 33.

► **Example 35.** Let \mathbb{F}_p be a finite field and $n \geq 1$. Let $V \subset \mathbb{F}_p^n$ be a linear subspace of dimension k . Given a set of coordinates $I \subseteq [n]$, define $V_I = \{(v_i)_{i \in I} : v \in V\}$ to be the subspace obtained by restricting vectors $v \in V$ to coordinates I . We say that V is α -large if

$$\dim(V_I) \geq \alpha|I| \quad \forall I \subseteq [n].$$

In particular, this implies that $k \geq \alpha n$.

Next, we define a set system corresponding to a subspace. Let $X = \{(i, a) : i \in [n], a \in \mathbb{F}_p\}$. For any vector $v \in \mathbb{F}_p^n$ define its corresponding set

$$S(v) = \{(i, v_i) : i \in [n]\} \subset X.$$

For a subspace $V \subset \mathbb{F}_p^n$ define the set system

$$\mathcal{F}(V) = \{S(v) : v \in V\}.$$

Observe that:

- (i) $\mathcal{F}(V)$ is an n -set system of size p^k .
- (ii) For any $T \subseteq X$ it holds that $|\{S \in \mathcal{F}(V) : T \subseteq S\}| \leq p^{-\alpha|T|}|\mathcal{F}|$. Hence $\mathcal{F}(V)$ is κ -regular for $\kappa = p^\alpha$.
- (iii) $\mathcal{F}(V)$ is intersecting iff any $v \in V$ contains at least one zero coordinate.

If Conjecture 33 holds and $p \geq (\log n)^c$ for some absolute constant $c > 0$, then it must hold that V contains a vector with no zero coordinates. This motivates the following problem.

► **Problem 36.** *Let $V \subset \mathbb{F}_p^n$ be a α -large subspace. Prove that if $p \geq (\log n)^c$, for some $c = c(\alpha)$, then V must contain a vector with no zero coordinates.*

A previous version of this paper gave a more restricted version of Example 35, corresponding to the case when V spans an MDS code. Namely, $\dim(V_I) = |I|$ for all $I \subseteq [n]$ with $|I| \leq k$. Ryan Alweiss [2] proved the analog of Problem 36 for this case, in fact where $p \geq p_0(n/k)$.

References

- 1 Noga Alon, Amir Shpilka, and Christopher Umans. On sunflowers and matrix multiplication. *computational complexity*, 22(2):219–243, 2013.
- 2 Ryan Alweiss. Personal communication, 2019.
- 3 Paul Erdős and Richard Rado. Intersection theorems for systems of sets. *Journal of the London Mathematical Society*, 35(1):85–90, 1960.
- 4 Mika Goos, Shachar Lovett, Raghu Meka, Thomas Watson, and David Zuckerman. Rectangles are nonnegative juntas. *SIAM Journal on Computing*, 45(5):1835–1869, 2016.
- 5 Parikshit Gopalan, Raghu Meka, and Omer Reingold. DNF sparsification and a faster deterministic counting algorithm. *Computational Complexity*, 22(2):275–310, 2013.
- 6 Alexandr V Kostochka. Extremal Problems on Δ -Systems. In *Numbers, Information and Complexity*, pages 143–150. Springer, 2000.
- 7 Xin Li, Shachar Lovett, and Jiapeng Zhang. Sunflowers and Quasi-Sunflowers from Randomness Extractors. In *Approximation, Randomization, and Combinatorial Optimization. Algorithms and Techniques (APPROX/RANDOM 2018)*. Schloss Dagstuhl-Leibniz-Zentrum fuer Informatik, 2018.
- 8 Shachar Lovett and Jiapeng Zhang. DNF sparsification beyond sunflowers. In *Electronic Colloquium on Computational Complexity (ECCC)*, 2018.
- 9 Alexander A Razborov. Lower bounds for the monotone complexity of some Boolean functions. In *Soviet Math. Dokl.*, volume 31, pages 354–357, 1985.
- 10 Benjamin Rossman. The monotone complexity of k-clique on random graphs. In *Proceedings of the 2010 IEEE 51st Annual Symposium on Foundations of Computer Science*, pages 193–201. IEEE Computer Society, 2010.