



OC31 – 24988 - EXTRACTION OF CAROTENOIDS SUPPORTED BY OHMIC HEATING AND CHARACTERIZATION OF BIOLOGICAL PROPERTIES AND STABILITY THROUGHOUT GASTROINTESTINAL TRACT

Marta Coelho¹; Ricardo Pereira²; António Rodrigues³; José Teixeira²; Manuela Pintado¹

1 - Universidade Católica Portuguesa, CBQF - Centro de Biotecnologia e Química Fina – Laboratório Associado, Escola Superior de Biotecnologia, Rua Arquitecto Lobão Vital, 172, 4200-374 Porto, Portugal; 2 - CEB - Centre of Biological Engineering, University of Minho, 4710-057 Braga, Portugal.; 3 - Centre for Toxicogenomics and Human Health, Genetics, Oncology and Human Toxicology, NOVA Medical School/Faculdade de Ciências Médicas, Universidade Nova de Lisboa, Lisbon, Portugal

E-mail: mrtcoelho@hotmail.com

Keywords: Ohmic heating, carotenoids, gastrointestinal tract, carotenoids-derived aroma compounds, prebiotic, tomato by-products, anti-inflammatory

Abstract

Carotenoids are lipophilic isoprenoid compounds with a polyene backbone that contains a variable number of conjugated double bonds, being precursors of aroma compounds. They are important in human nutrition since they are a source of vitamin A (β -carotene). Several studies have linked the regular consumption of carotenoids to prevent human diseases, such as cardiovascular, cholesterol, neurodegenerative diseases, antioxidant and anti-cancer. Very little is known about release of entrapped carotenoids from complex matrixes and their impact on the gut microbiota. Extractable carotenoids were obtained from tomato by-products through green efficient process using ohmic heating (OH). The extracts were submitted to simulated gastrointestinal tract and digested samples were subjected to fermentation using faecal matter from 5 different controlled donors (male and female alike). The total carotenoids were assessed by spectrophotometric method. Individual carotenoids and carotenoids-derived aroma compounds during each step of digestive tract and from the microbiota assay were also analysed by LC-MS and LC-HR-QTOF-MS, respectively. The short-chain fatty acids (SCFA) and lactic acid and sugars from microbiota assay were analysed by HPLC. The human faecal microbiota was assessed by real-time quantitative PCR. Prebiotic, anti-hypertensive activity, anti-inflammatory activities were also evaluated. In the OH extract, the major carotenoids identified were lycopene and β -carotene (corresponding to 92% of total carotenoids). After digestion process and fermentation only carotenoids-derived aroma compounds and other metabolites resulting the carotenoids were found, including camphenol, linalool acetate, (+)-sabinol, acetovanillone and hexadecanedioic acid. The results suggest that



fermentation of OH tomato by-product extract demonstrated a prebiotic effect through the increasing of the number of *Bifidobacterium animalis* and improving the production of SCFA, approving a potential modelatory effect upon the gut microbiota and consequently providing a prevention of numerous diseases. Complementary, the OH extract also proved anti-inflammatory activity and moderate anti-hypertensive activity.

Acknowledgements:

The authors would like to thank to the project Co-promoção nº 016403, "MULTIBIOREFINERY", supported by Programa Operacional Competitividade e Internacionalização e pelo Programa Operacional Regional de Lisboa, na sua componente FEDER, e pela Fundação para a Ciência e Tecnologia and project UID/Multi/50016/2013, administrated by FCT. Furthermore, his study was supported by the Portuguese Foundation for Science and Technology (FCT) under the scope of the strategic funding of UID/BIO/04469/2019 unit and BioTecNorte operation (NORTE-01-0145-FEDER-000004) funded by the European Regional Development Fund under the scope of Norte2020 - Programa Operacional Regional do Norte. The author Marta Coelho would like to acknowledge FCT for your PhD grant with the reference [grant number SFRH/BD/111884/2015]