

BMJ Open Effects of dietary antioxidant vitamins on lung functions according to gender and smoking status in Korea: a population-based cross-sectional study

Ji Young Hong,^{1,2} Chang Youl Lee,^{1,2} Myung Goo Lee,^{1,2} Young Sam Kim³

To cite: Hong JY, Lee CY, Lee MG, *et al.* Effects of dietary antioxidant vitamins on lung functions according to gender and smoking status in Korea: a population-based cross-sectional study. *BMJ Open* 2018;**8**:e020656. doi:10.1136/bmjopen-2017-020656

► Prepublication history for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2017-020656>).

Received 15 November 2017
Revised 16 January 2018
Accepted 6 February 2018



¹Division of Pulmonary and Critical Care Medicine, Department of Medicine, Chuncheon Sacred Heart Hospital, Hallym University Medical Center, Chuncheon, South Korea

²Lung Research Institute of Hallym, University College of Medicine, Chuncheon, South Korea

³Division of Pulmonology, Department of Internal Medicine, Yonsei University College of Medicine, Seoul, Republic of Korea

Correspondence to

Professor Young Sam Kim;
yksamkim@yuhs.ac

ABSTRACT

Objective Cigarette smoke-induced oxidative stress plays an important role in the pathogenesis of chronic obstructive pulmonary disease (COPD). Dietary antioxidants are thought to prevent smoke-induced oxidative damage. The aim of this study was to investigate associations between lung function and the consumption of antioxidant vitamins in Korean adults.

Methods In total, 21 148 participants from the Korean National Health and Nutrition Examination Survey (2007–2014) were divided into four groups based on smoking history and gender. Multivariate regression models were used to evaluate associations between lung function and intake of dietary antioxidants.

Results Subjects in the highest intake quintile (Q5) of vitamin A, carotene and vitamin C intake had mean forced expiratory volume in 1 s (FEV₁) measurements that were 30 mL, 32 mL and 36 mL higher than those of individuals in the lowest intake quintile (Q1), respectively (p for trend; p=0.008, p=0.010 and p<0.001, respectively). The risks of COPD for male smokers in Q1 increased 7.60-fold (95% CI 5.92 to 9.76), 7.16-fold (95% CI 5.58 to 9.19) and 7.79-fold (95% CI 6.12 to 9.92), for vitamin A, carotene and vitamin C, respectively, compared with those of female non-smokers in Q5. Among patients with COPD, men who smoked >20 pack-years had mean FEV₁ measurements that were 192 mL, 149 mL and 177 mL higher than those of patients in Q1 (p for trend; p=0.018, p=0.024 and p=0.043, for vitamin A, carotene and vitamin C, respectively).

Conclusions These findings indicate that the influence of antioxidant vitamins on lung function depends on gender and smoking status in the Korean COPD population.

INTRODUCTION

Chronic obstructive pulmonary disease (COPD) causes morbidity and mortality.¹ Smoking is a primary risk factor for COPD; however, other factors also contribute as only 10%–20% of smokers develop airflow limitations.²

Dietary antioxidants protect against oxidative stress caused by smoking,³ and multiple studies have revealed associations between the intake of antioxidant vitamins or fibres

Strengths and limitations of this study

- This study revealed that the influence of antioxidant vitamins on lung function depends on gender and smoking status in Korean patients with chronic obstructive pulmonary disease.
- This is a cross-sectional study with a large sample size collected from a national health survey.
- Main limitations include a possible recall bias and no further verification of nutritional intake.

and respiratory diseases.^{4–8} However, evidence supporting the benefits of vitamin supplement therapy is lacking.^{9,10}

Because micronutrient status is affected by dietary intake and metabolic turnover, which are regulated by oxidative stress, the benefits of antioxidant vitamins may vary by gender and smoking status. Multiple studies have shown that different antioxidants exhibit different effects based on smoking status. Morabia *et al* reported an association between airway obstruction and vitamin A intake in smokers compared with former smokers, whereas Hu *et al* reported that carotene was less strongly associated with FEV₁ in smokers compared with former smokers and non-smokers.^{11,12}

This study used Korean National Health and Nutritional Examination Survey (KNHANES) data to investigate whether dietary antioxidant vitamins were independently associated with pulmonary function and COPD in the Korean population. This study also evaluated whether the effects of antioxidant vitamins on pulmonary function differed based on gender or smoking status.

PATIENTS AND METHODS

Study population

Participants were sampled from KNHANES (2007–2014) IV–VI, a nationwide survey

designed to be representative of the population that is used to establish health policies. KNHANES contains a massive database with information about demographic characteristics, comorbidities, lung function, nutritional status and health (<https://knhanes.cdc.go.kr/knhanes>).

A two-stage stratified systemic sampling method was used to select 65 973 individuals to survey between February 2007 and December 2014. Of the chosen individuals, 34 278 participants over 40 years of age responded to questionnaires regarding diet and smoking history and underwent a medical examination. After excluding subjects who omitted lung function or nutrition data, we analysed data from 21 148 (8804 men and 12 344 women) in this study. This study was approved by the Institutional Review Board of the Korean Centres for Disease Control and Prevention. All participants provided informed written consent.

PROTOCOL

KNHANES collects survey data through health questionnaire surveys, screening surveys and nutrition surveys. Health questionnaires were divided into household survey, health interview survey and health behaviour survey. The health interview survey examined the use of medical services, activity limitations, education and economic activities and physical activity by interview method. The health behaviour survey examined smoking status, drinking, mental health and safety consciousness by self-filling method. The screening consisted of physical measurement, blood pressure and pulse measurement, blood and urine test, oral examination, pulmonary function test, visual and refractive examination, colour vision test, hearing test and muscle strength test. Nutrition surveys consisted of dietary behaviours, dietary supplements, nutritional knowledge and the contents of food intake (24-hour recall method) a day before the survey.

Spirometry and airflow obstruction definitions

The pulmonary function test was performed using dry-rolling seal volume spirometers (Model 2130; Sensor Medics, Yorba Linda, California, USA) and standardised according to the American Thoracic Society/European Respiratory Society criteria.¹³ Qualified technicians and principal investigators assessed the spirometry data for acceptability and reproducibility. The predictive equations for the forced expiratory volume in 1 s (FEV_1) and the forced vital capacity (FVC) were derived from survey data on non-smokers who had normal chest X-rays and no history of respiratory diseases.¹⁴ COPD was defined as an $FEV_1/FVC < 70\%$.¹⁵

Dietary assessments

Food intake data were obtained using the 24-hour recall method, in which participants were asked to report the foods and amounts thereof consumed during the previous 24 hours. Total energy (kJ/d (kcal/d)) and total intake of antioxidant vitamins were calculated using the Korean

Food Composition Table¹⁶ as the reference. Antioxidant vitamin consumption was adjusted for total energy intake.

Potential confounders

Data regarding demographic information, education level, household income, smoking status, smoking amount, alcohol intake, place of residence, body mass index (BMI) and comorbid diseases were obtained. Educational level was categorised as elementary school or lower, completion of middle school, completion of high school and college or higher. Household income was divided by quartile. Place of residence was divided to rural and urban.

Smokers were subjects who smoked more than 100 cigarettes in their lifetime.¹⁷ Participants were categorised in terms of smoking status as follows: smoker, ex-smoker or never smoked. Those who answered in the negative to the question 'Do you currently smoke?' were defined as ex-smokers.

The smoking amount was determined in pack-years, which was calculated by multiplying the duration of smoking (years) by the number of packs of cigarettes smoked. Comorbid diseases included hypertension, stroke, cardiovascular disease, arthritis, tuberculosis, asthma, diabetes mellitus, thyroid disorders, renal failure, liver disease and malignancy.

Statistical analysis

A total of 21 148 subjects participated in this study (figure 1). The relationship between antioxidant vitamin intake and lung function was analysed using multiple linear regression analyses. We analysed the energy-adjusted antioxidant vitamin intake by quintiles. The adjustment factors were age, sex, BMI, educational level, household income, total energy intake, place of residence, number of comorbid diseases, smoking history, alcohol intake and pack-years.^{4 11 18} Assessments of linear trends across increasing antioxidant vitamin quintiles were also performed.

We estimated the ORs of COPD using multivariate logistic regression analyses of quintiles after adjusting for confounding factors. Participants were divided into four groups based on gender and smoking status (male smokers, male non-smokers, female smokers, female non-smokers) to determine whether the relationship between COPD risk and antioxidant vitamin intake is related to gender and smoking status. For combined analyses between the effects of antioxidant vitamin intake, gender and smoking status on the risk of COPD, interaction tests were performed. Patients with COPD and male patients with COPD were analysed separately. We attempted to determine whether the association of antioxidant vitamins and lung function varies with gender and smoking status in patients with COPD. Multiple linear regression analyses were performed after categorising patients with COPD by smoking status and amount.




Figure 1 The study population framework. COPD, chronic obstructive pulmonary disease; FVC, forced vital capacity; FEV₂, forced expiratory volume in 2 s; KNHANES, Korean National Health and Nutritional Examination Survey .

Statistical analyses were performed using PASW Statistics V.20 (SPSS) and SAS V.9.4 (SAS Institute) software. $P < 0.05$ was considered statistically significant.

RESULTS

The baseline characteristics of the 21 148 participants are shown in [table 1](#). All subjects were classified into four groups based on smoking history and gender. Of the 7986 smokers, 7178 were male (mean age, 57.8 ± 11.0 years) and 808 were female (mean age, 57.4 ± 12.5 years). Of the 13162 individuals who had never smoked, 1626 were male (mean age, 57.9 ± 11.3 years) and 11 536 were female (mean age, 57.1 ± 10.8 years). Among all subjects, 3005 were diagnosed with COPD. The prevalence of COPD was highest in male smokers (26.4%) and lowest in female non-smokers (6.4%).

The four groups differed regarding age, BMI, educational level, household income and alcohol usage ($p < 0.001$). Energy intake was significantly higher in men

than women (men, 2256.5 kcal; women, 1648.3 kcal; $p < 0.001$). The levels of vitamin A, carotene and vitamin C were highest in the male non-smoker group and lowest in the female smoker group. Korean male non-smokers are predisposed to COPD compared with female non-smokers (incidence rate of 15.7% vs 6.4%). Age and the percentage of alcohol intake were higher in Korean male non-smokers than female non-smokers.

[Table 2](#) showed the association between lung function (FEV₁, FVC) and dietary antioxidant vitamin levels. Participants in the highest quintile (Q5) of vitamin A intake had 30 mL higher FEV₁ (p for trend across quintiles=0.008) and 33 mL higher FVC (p for trend across quintiles=0.007) compared with participants in the lowest quintile (Q1). Participants in Q5 for carotene intake had 32 mL higher FEV₁ (p for trend across quintiles=0.010) and 36 mL higher FVC (p for trend across quintiles=0.005) measurements compared with participants in Q1. Participants in Q5 of vitamin C intake had 36 mL higher FEV₁

Table 1 Study population characteristics

	Total (n=21 148)	Male smokers (n=7178)	Male non-smokers (n=1.626)	Female smokers (n=808)	Female non-smokers (n=11 536)	P values
Age*	57.4 (10.9)	57.8 (11.0)	57.9 (11.3)	57.4 (12.5)	57.1 (10.8)	<0.001
40–49	6048 (28.6)	1998 (27.8)	464 (28.5)	273 (33.8)	3313 (28.7)	<0.001
50–59	6131 (29.0)	1981 (27.6)	431 (26.5)	199 (24.6)	3520 (30.5)	
60–69	5387 (25.5)	1913 (26.7)	430 (26.4)	158 (19.6)	2866 (25.0)	
70–	3582 (16.9)	1286 (17.9)	301 (18.5)	178 (22.0)	1817 (15.8)	
BMI*	24.2 (3.0)	24.2 (2.8)	24.3 (2.8)	23.8 (3.6)	24.2 (3.2)	0.007
Education						<0.001
Elementary	7229 (34.2)	1763 (24.6)	321 (19.7)	381 (47.2)	4764 (41.3)	
Middle school	3315 (15.7)	1216 (16.9)	267 (16.4)	112 (13.9)	1720 (14.9)	
High school	6427 (30.4)	2366 (33.0)	458 (28.2)	228 (28.2)	3375 (29.3)	
More than college	4169 (19.7)	1831 (25.5)	580 (35.7)	87 (10.8)	1671 (14.5)	
Household income						<0.001
First quartile	4763 (22.5)	1440 (20.1)	289 (17.8)	315 (39.0)	2719 (23.6)	
Second quartile	5427 (25.7)	1874 (26.1)	391 (24.1)	223 (27.6)	2939 (25.5)	
Third quartile	5162 (24.4)	1869 (26.1)	414 (25.5)	145 (17.9)	2734 (23.7)	
Fourth quartile	5780 (27.3)	1988 (27.7)	530 (32.6)	125 (15.5)	3137 (27.2)	
Comorbidity*	0.9 (1.1)	0.9 (1.0)	0.8 (0.9)	1.0 (1.2)	1.0 (1.1)	<0.001
Pack-years*	4.7 (13.6)	13.3 (20.3)	0.2 (2.2)	3.3 (11.9)	0.0 (0.0)	<0.001
Alcohol	17 554 (83.0)	6877 (95.8)	1399 (86.0)	714 (88.4)	8564 (74.2)	<0.001
Energy intake (kcal/day)*	1901.5 (797.8)	2266.5 (869.7)	2212.6 (855.9)	1538.2 (653.4)	1656.0 (630.0)	<0.001
Vitamin A (µg RE/day)*	822.5 (1118.5)	881.9 (1067.5)	925.5 (1095.2)	600.3 (644.2)	786.6 (1173.9)	<0.001
Carotene (µg/day)*	4337.3 (6206.0)	4596.2 (5557.8)	4803.8 (5506.6)	3143.5 (3682.6)	4194.1 (6780.6)	<0.001
Vitamin C (mg/day)*	111.9 (107.6)	111.8 (97.9)	128.8 (107.1)	84.8 (96.9)	111.5 (113.5)	<0.001
FEV ₁ (mL)*	2.60 (0.67)	3.02 (0.68)	3.09 (0.66)	2.23 (0.56)	2.30 (0.46)	<0.001
FVC (mL)*	3.38 (0.84)	4.07 (0.72)	4.04 (0.73)	2.88 (0.62)	2.89 (0.51)	<0.001
FEV ₁ /FVC (%)*	77.3 (7.9)	73.9 (9.1)	76.6 (7.9)	77.2 (8.0)	79.5 (6.1)	<0.001
COPD	3005 (14.2)	1893 (26.4)	256 (15.7)	119 (14.7)	737 (6.4)	<0.001

*Numbers represent mean percentages (SD).

BMI, body mass index; COPD, chronic obstructive pulmonary disease; FVC, forced vital capacity; FEV₁, forced expiratory volume in 1 s; RE, retinol equivalent.

(p for trend across quintiles<0.001) and 35 mL higher FVC (p for trend across quintiles=0.014) measurements compared with participants in Q₁. A statistically significant dose–response relationship was observed (all, p for trend across quintiles<0.005), but participants in Q₃ of vitamin A and carotene had comparable lung function to those in Q₅.

The effects of gender, smoking and dietary antioxidant vitamins on the risk of COPD are summarised in table 3. The risk of COPD for male smokers in Q₁ for vitamin A, carotene and vitamin C intake increased by 7.60-fold (95% CI 5.92 to 9.76), 7.16-fold (95% CI 5.58 to 9.19) and 7.79-fold (95% CI 6.12 to 9.92), respectively, which was greater than that observed for female non-smokers in Q₅ for antioxidant vitamin intake. Interestingly, the

risk of COPD for male non-smokers in Q₅ for vitamin A, carotene and vitamin C intake increased by 3.26-fold (95% CI 2.24 to 4.75), 3.35-fold (95% CI 2.31 to 4.86) and 3.28-fold (95% CI 2.27 to 4.73), respectively, compared with female non-smokers in Q₅ for antioxidant vitamin intake. The risk of COPD for male non-smokers in Q₁ for vitamin A, carotene and vitamin C intake increased by 2.80-fold (95% CI 1.90 to 4.12), 3.25-fold (95% CI 2.21 to 4.78) and 3.17-fold (95% CI 2.04 to 4.91), respectively, compared with female non-smokers in Q₁ for antioxidant vitamin intake. These results suggest that men may have other causes of COPD as well as smoking, compared with women who took similar amounts of antioxidant vitamins.

The interaction exists between the antioxidant vitamin intake and gender/smoking status on the risk of COPD

Table 2 Mean values of adjusted lung function measurements across quintiles of vitamin A, carotene and vitamin C intake

	Q1	Q2	Q3	Q4	Q5	Difference between Q5 and Q1 (95% CI)	P values for trend
Vitamin A							
Mean intake ($\mu\text{g RE}$)	151.2	353.6	573.1	893.9	2140.8		
FEV ₁ (mL)	2379	2389	2410	2397	2409	30 (10 to 50)	0.008
FVC (mL)	3119	3136	3158	3148	3152	33 (10 to 57)	0.007
Predicted FEV ₁ (%)	91.37	91.44	91.91	91.45	91.94	0.57 (-0.08 to 1.22)	0.185
Predicted FVC (%)	90.93	91.06	91.45	91.22	91.48	0.55 (0.00 to 1.10)	0.195
Carotene							
Mean intake (μg)	691.1	1747.4	2938.9	4736.1	11574.1		
FEV ₁ (mL)	2347	2363	2376	2370	2379	32 (12 to 52)	0.010
FVC (mL)	3088	3117	3127	3119	3124	36 (13.59)	0.005
Predicted FEV ₁ (%)	91.55	92.03	92.26	91.85	92.31	0.76 (0.12 to 1.39)	0.096
Predicted FVC (%)	91.02	91.68	91.78	91.41	91.82	0.80 (0.26 to 1.33)	0.015
Vitamin C							
Mean intake (mg)	24.2	53.6	84.2	128.8	268.9		
FEV ₁ (mL)	2411	2423	2436	2441	2453	36 (16.56)	<0.001
FVC (mL)	3117	3122	3132	3140	3154	35 (12.58)	0.014
Predicted FEV ₁ (%)	91.3	91.5	91.9	91.99	92.21	0.91 (0.27 to 1.55)	0.050
Predicted FVC (%)	91.29	91.33	91.58	91.77	92.0	0.71 (0.17 to 1.26)	0.118

Data were adjusted for age, sex, body mass index, energy intake, number of comorbid diseases, alcohol consumption, place of residence, smoking history, pack-years (smoking amount), household income and education level. P values were determined using tests for linear trends across increasing quintiles (means) of antioxidant vitamin intake.

FVC, forced vital capacity; FEV₁, forced expiratory volume in 1 s; RE, retinol equivalent; Q1, lowest quintile; Q5, highest quintile.

Table 3 Association between vitamin A, carotene and vitamin C intake and COPD according to gender and smoking status

	Intake		COPD		OR		P interaction
	Q5	Q1	Q5	Q1	Q5	Q1	
Vitamin A							
Female non-smokers	2096	2564	105	242	Ref	1.16 (0.89, 1.49)	<0.001
Female smokers	109	264	16	53	3.90 (2.12, 7.17)	2.42 (1.63, 3.58)	
Male non-smokers	394	225	53	47	3.26 (2.24, 4.75)	3.15 (2.10, 4.72)	
Male smokers	1630	1176	320	444	5.54 (4.28, 7.16)*	7.60 (5.92, 9.76)*	
Carotene							
Female non-smokers	2118	2529	108	226	Ref	1.10 (0.85, 1.42)	<0.001
Female smokers	104	268	15	49	3.47 (1.86, 6.47)	2.16 (1.45, 3.23)	
Male non-smokers	397	243	55	50	3.35 (2.31, 4.86)	3.24 (2.18, 4.82)	
Male smokers	1610	1189	321	425	5.83 (4.51, 7.53)*	7.16 (5.58, 9.19)*	
Vitamin C							
Female non-smokers	2303	2466	112	465	Ref	1.00 (0.77, 1.30)	<0.001
Female smokers	107	294	12	35	2.37 (1.20, 4.71)	2.27 (1.55, 3.34)	
Male non-smokers	401	191	55	55	3.28 (2.27, 4.73)	3.24 (2.07, 5.06)	
Male smokers	1419	1278	317	204	6.20 (4.82, 7.98)*	7.79 (6.12, 9.92)*	

OR was determined following adjustment for age, body mass index, energy intake, number of comorbid diseases, alcohol consumption, place of residence, household income and education level.

*The risk for COPD was significantly different between Q1 and Q5.

COPD, chronic obstructive pulmonary disease; Q1, lowest quintile; Q5, highest quintile.




Figure 2 OR for the association between antioxidant vitamin intake and COPD among (A) male and (B) female smokers and non-smokers. OR were adjusted for age, body mass index, energy intake, number of comorbid diseases, alcohol consumption, place of residence, household income and education level.

(all p values < 0.001). The effect of the antioxidant vitamin intake depends on the gender/smoking status. When assessing the risk of COPD following reduction of antioxidant intake from Q5 to Q1, only male smokers showed significant difference in risk of COPD, but other three groups did not. Figure 2 shows that the risk of COPD was influenced by dietary antioxidant vitamin levels in male smokers in detail. In male smokers, the risk of COPD in subjects in Q5 for antioxidant vitamins intake was significantly lower than that for subjects in Q1 (vitamin A, OR=0.77, 95% CI 0.63 to 0.94, $p=0.009$; carotene, OR=0.81, 95% CI 0.67 to 0.99, $p=0.041$; vitamin C, OR=0.74, 95% CI 0.61 to 0.91, $p=0.004$). The dose-dependent effect of vitamin C was observed between COPD risk and dietary antioxidant vitamin levels, but it was not for vitamin A and carotene. Although not significant, Q3 group of carotene had increased risk to develop COPD than Q1 group of carotene.

The prevalence of COPD did not increase significantly as the intake of dietary antioxidant vitamins increased in male non-smokers, female smokers or female non-smokers. No significant interaction between the effects of antioxidant vitamins on COPD and smoking status was observed. The correlation between the risk of COPD and antioxidant vitamin intake was stronger in male smokers who smoked less than 20 pack-years (not shown).

We investigated the association between dietary antioxidant vitamin intake and lung function after limiting the analyses to individuals with COPD. The changes in FEV₁ were not statistically significant based on the levels of dietary antioxidant vitamins in subjects with COPD.

Similar to the previous results, only male smokers in subjects with COPD, exhibited a beneficial association between dietary antioxidant vitamin intake and FEV₁ (figure 3). Male smokers with COPD in Q5 for vitamin A intake had a 71 mL higher FEV₁ (p for trend across quintiles=0.019) compared with those in Q1. Male smokers with COPD in Q5 of carotene and vitamin C intake had 71 mL higher FEV₁ (p for trend across quintiles=0.037) and a 109 mL higher FEV₁ (p for trend across quintiles=0.026), respectively, compared with individuals in Q1.

Additional analyses were performed to determine if lung function was reduced by smoking amount or smoking status in male smoker patients with COPD. Male patients with COPD who had smoked ≥ 20 pack-years exhibited a beneficial association between dietary antioxidant vitamin intake and FEV₁ (figure 4). Male patients with COPD in Q5 of vitamin A intake who had smoked ≥ 20 pack-years had a 192 mL higher FEV₁ (p for trend across quintiles=0.018) compared with individuals in Q1. Patients with COPD in Q5 of carotene intake who had smoked ≥ 20 pack-years had a 149 mL higher FEV₁ (p for trend across quintiles=0.024) compared with patients in Q1. Patients with COPD in Q5 of vitamin C intake who had smoked > 20 pack-years had a 177 mL higher FEV₁ (p for trend across quintiles=0.043) compared with patients in Q1.

DISCUSSION

This study examined the association between the intake of antioxidant vitamins and lung functions in the Korean




Figure 3 Mean values of adjusted FEV₁ measurements across quintiles of vitamin A, carotene and vitamin C intake (energy adjusted) in subjects with COPD. Adjusted for age, body mass index, energy intake, number of comorbid diseases, alcohol consumption, place of residence, household income and education level. P values were determined using tests for linear trends across increasing quintiles (means) of antioxidant vitamin intake. COPD, chronic obstructive pulmonary disease; FEV₁, forced expiratory volume in 1 s.

population. Previous studies showed that antioxidant vitamins, including vitamin C, were protective of the human lung,^{5 12} whereas high levels of vitamin A and carotene were also associated with increased lung functions in multiple studies.^{11 19–23} In a randomised controlled trial, Keranis *et al* reported that increasing the intake of antioxidants improved lung function.²⁴

Cigarette smoking is the primary cause of COPD as it increases oxidative stress in the lungs and activates inflammatory responses.²⁵ Notably, one inhalation from a cigarette generates more than 10¹⁵ free radicals and other oxidants.²⁶

Antioxidants protect against the damage caused by smoking in multiple ways.³ For example, as it is




Figure 4 Mean values of adjusted FEV₁ measurements across quintiles of vitamin A, carotene and vitamin C intake (energy adjusted) in male patients with COPD according to smoking status. Values were adjusted for age, BMI, energy intake, number of comorbid diseases, alcohol consumption, place of residence, household income and education level. P values were determined using tests for linear trends across increasing quintiles (median) of antioxidant vitamin intake. BMI, body mass index; COPD, chronic obstructive pulmonary disease; FEV₁, forced expiratory volume in 1 s.

water soluble, vitamin C scavenges free radicals in the cytoplasm. Koike *et al* reported that vitamin C diminished smoke-induced oxidative stress and corrected emphysematous lungs *in vivo*.²⁷

Carotenoids quench singlet oxygen and inhibit lipid peroxidation.³ In an animal study, the respiratory epithelial of retinol-deficient animals had atrophied ciliated cells and modified lipid contents.²⁸ The pathological features of the retinol-deficient animals were similar to those of human smokers.²⁹

Smokers exhibit nicotine-induced reductions in intestinal absorption and elevated metabolic turnover.³⁰ The metabolism or destruction of antioxidant vitamins increases in inflammatory environments,^{31–34} which suggests that smokers with COPD require larger amounts of antioxidant vitamins to achieve the same blood levels as non-smokers. A study by Sargeant and Jaeckel found that vitamin C may modify the adverse effects of smoking and the risk of COPD in the European population.³⁵ Additionally, Shin *et al* reported that Korean smokers with adequate vitamin C intake had acceptable pulmonary functions³⁶ and Park *et al* showed that dietary vitamin C provides protection against COPD.³⁷ Additionally, Morabia *et al* identified that airway obstruction was reduced by vitamin A in smokers.¹¹

One notable finding in the current study was that the effects of antioxidant vitamin intake on lung function were stronger among male smokers. Additionally, the association between the risk of COPD and antioxidant vitamin intake was clear for male but not female smokers. Male smokers with lower antioxidant vitamin intakes had increased ORs of COPD compared with female smokers. Although the dose-dependent effect on COPD risk was not obvious in vitamin A and carotene, contrary to vitamin C (figure 2), male smokers with Q5 intake showed a clearly reduced risk to develop COPD than male smokers with Q1 intake in all three antioxidant vitamins.

After limiting the analysis to subjects with COPD, a significant association between antioxidant vitamin intake and FEV₁ was observed in male smokers but not in other groups. This finding was similar to that of Joshi *et al*, where changes in COPD risk and dietary vitamin C and vitamin E intake differed between men and women.³⁸

It is not known how gender differences impact pulmonary functions based on antioxidant vitamin intake; however, animal studies have revealed gender differences in antioxidant vitamin requirements. Al-Rejaie *et al* reported gender-related differences in the protective roles of ascorbic acid against oxidative stress,³⁹ whereas Jiao *et al* revealed gender differences in the regulation and expression of oxidative genes in mice.⁴⁰

Studies detailing the effects of antioxidant vitamins on lung function in smokers and non-smokers are lacking.^{5,23} In the US population, Britton *et al* revealed that the relationship between vitamin C intake and FEV₁ was stronger in ex-smokers than non-smokers or current smokers.⁵ Shahar *et al* reported a relationship between individuals

in Q1 of vitamin A intake and airway obstruction among individuals who smoked >41 pack-years.²³

Among male patients with COPD, those smoking ≥ 20 pack-years had improved lung functions as antioxidant vitamin intake increased. These results support that associations between antioxidants and lung function may differ according to smoking status in patients with COPD patients. However, it is unknown what causes such differences. One hypothesis is that the efficacy of antioxidant vitamins is proportional to the level of oxidant burden in COPD. Additional studies are required to determine whether the benefits of antioxidant vitamins depend on the smoking duration or dose in patients with COPD.

This study has several limitations that should be noted. As we used a cross-sectional design, the data cannot be used to answer questions regarding causation. Additionally, because data on nutritional intake were obtained by 24 hours recall, inaccurate responses may have been offered. This study used the prebronchodilator FEV₁ for determining COPD; however, the definition of COPD is based on postbronchodilator FEV₁.¹⁵ This study failed to obtain data regarding air pollution or occupational exposure and, therefore, could not associate these variables with lung function; however, the strength of this study is that these data represent the Korean population.

CONCLUSION

This study supports that antioxidant vitamins have beneficial effects on pulmonary function in the Korean population. The data indicate that there is a stronger association between antioxidant vitamin intake and the risk of COPD in male smokers. The beneficial effects of antioxidant vitamins in patients with COPD differed by gender and smoking status, and future investigations should determine the roles of dietary antioxidant vitamins in specific groups.

Contributors JYH and YSK equally contributed to the conception and design of the research; YSK contributed to the design of the research; CYL contributed to the acquisition and analysis of the data; MGL and YSK contributed to the interpretation of the data and JYH drafted the manuscript. All authors critically revised the manuscript, agreed to be fully accountable for ensuring the integrity and accuracy of the work and read and approved the final manuscript.

Funding This research was supported by the Bio and Medical Technology Development Program of the National Research Foundation (NRF) and funded by the Korean Government (MSIT) (NRF-2017M3A9E8033225). This work was supported by the National Research Foundation of Korea Grant funded by the Korean Government (NRF 2017R1C1B5017879).

Competing interests None declared.

Patient consent Obtained.

Ethics approval Institutional Review Board of the Korean Centers for Disease Control and Prevention.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement No additional data are available.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is

properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- Viegi G, Pistelli F, Sherrill DL, *et al.* Definition, epidemiology and natural history of COPD. *Eur Respir J* 2007;30:993–1013.
- Capocaccia R, Verdecchia A, Micheli A, *et al.* Breast cancer incidence and prevalence estimated from survival and mortality. *Cancer Causes Control* 1990;1:23–9.
- Northrop-Clewes CA, Thurnham DI. Monitoring micronutrients in cigarette smokers. *Clin Chim Acta* 2007;377:14–38.
- Kan H, Stevens J, Heiss G, *et al.* Dietary fiber, lung function, and chronic obstructive pulmonary disease in the atherosclerosis risk in communities study. *Am J Epidemiol* 2008;167:570–8.
- Britton JR, Pavord ID, Richards KA, *et al.* Dietary antioxidant vitamin intake and lung function in the general population. *Am J Respir Crit Care Med* 1995;151:1383–7.
- Schünemann HJ, Freudenheim JL, Grant BJ. Epidemiologic evidence linking antioxidant vitamins to pulmonary function and airway obstruction. *Epidemiol Rev* 2001;23:248–67.
- Ford ES, Li C, Cunningham TJ, *et al.* Associations between antioxidants and all-cause mortality among US adults with obstructive lung function. *Br J Nutr* 2014;112:1662–73.
- Rodríguez-Rodríguez E, Ortega RM, Andrés P, *et al.* Antioxidant status in a group of institutionalised elderly people with chronic obstructive pulmonary disease. *Br J Nutr* 2016;115:1740–7.
- Wu TC, Huang YC, Hsu SY, *et al.* Vitamin E and vitamin C supplementation in patients with chronic obstructive pulmonary disease. *Int J Vitam Nutr Res* 2007;77:272–9.
- Heart Protection Study Collaborative Group. MRC/BHF Heart Protection Study of antioxidant vitamin supplementation in 20 536 high-risk individuals: a randomised placebo-controlled trial. *Lancet* 2002;360:23–33.
- Morabia A, Sorenson A, Kumanyika SK, *et al.* Vitamin A, cigarette smoking, and airway obstruction. *Am Rev Respir Dis* 1989;140:1312–6.
- Hu G, Cassano PA. Antioxidant nutrients and pulmonary function: the Third National Health and Nutrition Examination Survey (NHANES III). *Am J Epidemiol* 2000;151:975–81.
- Miller MR, Hankinson J, Brusasco V, *et al.* Standardisation of spirometry. *Eur Respir J* 2005;26:319–38.
- Choi JK, Paek D, Lee JO. Normal predictive values of spirometry in Korean population. *Tuberc Respir Dis* 2005;58:230–42.
- Rabe KF, Hurd S, Anzueto A, *et al.* Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease: GOLD executive summary. *Am J Respir Crit Care Med* 2007;176:532–55.
- RRD I. *Food composition table*. 7th ed. Suwon, Korea: Rural Resource Development Institute, 2006.
- Yoo KH, Kim YS, Sheen SS, *et al.* Prevalence of chronic obstructive pulmonary disease in Korea: the fourth Korean National Health and Nutrition Examination Survey, 2008. *Respirology* 2011;16:659–65.
- Hong JY, Kim SY, Chung KS, *et al.* Factors associated with the quality of life of Korean COPD patients as measured by the EQ-5D. *Qual Life Res* 2015;24:2549–58.
- Kelly Y, Sacker A, Marmot M. Nutrition and respiratory health in adults: findings from the health survey for Scotland. *Eur Respir J* 2003;21:664–71.
- McKeever TM, Lewis SA, Smit HA, *et al.* A multivariate analysis of serum nutrient levels and lung function. *Respir Res* 2008;9:67.
- Miedema I, Feskens EJ, Heederik D, *et al.* Dietary determinants of long-term incidence of chronic nonspecific lung diseases. The Zutphen Study. *Am J Epidemiol* 1993;138:37–45.
- McKeever TM, Scrivener S, Broadfield E, *et al.* Prospective study of diet and decline in lung function in a general population. *Am J Respir Crit Care Med* 2002;165:1299–303.
- Shahar E, Folsom AR, Melnick SL, *et al.* Does dietary vitamin A protect against airway obstruction? The Atherosclerosis Risk in Communities (ARIC) Study Investigators. *Am J Respir Crit Care Med* 1994;150:978–82.
- Keranis E, Makris D, Rodopoulou P, *et al.* Impact of dietary shift to higher-antioxidant foods in COPD: a randomised trial. *Eur Respir J* 2010;36:774–80.
- Alberg AJ, Chen JC, Zhao H, *et al.* Household exposure to passive cigarette smoking and serum micronutrient concentrations. *Am J Clin Nutr* 2000;72:1576–82.
- Pryor WA, Stone K. Oxidants in cigarette smoke. Radicals, hydrogen peroxide, peroxyacetaldehyde, and peroxyacetyl nitrate. *Ann N Y Acad Sci* 1993;686:12–27.
- Koike K, Ishigami A, Sato Y, *et al.* Vitamin C prevents cigarette smoke-induced pulmonary emphysema in mice and provides pulmonary restoration. *Am J Respir Cell Mol Biol* 2014;50:347–57.
- Iakovleva OA, Pentiu AA G VI, *et al.* Effect of vitamin A deficiency on surfactants and enzymes of xenobiotic metabolism in the rat lung]. *Vopr Pitan* 1987;4:61–3.
- Nettesheim P, Snyder C, Kim JC. Vitamin A and the susceptibility of respiratory tract tissues to carcinogenic insult. *Environ Health Perspect* 1979;29:89–93.
- Kalner AB, Hartmann D, Hornig DH. On the requirements of ascorbic acid in man: steady-state turnover and body pool in smokers. *Am J Clin Nutr* 1981;34:1347–55.
- Lykkesfeldt J, Loft S, Nielsen JB, *et al.* Ascorbic acid and dehydroascorbic acid as biomarkers of oxidative stress caused by smoking. *Am J Clin Nutr* 1997;65:959–63.
- Cser MA, Majchrzak D, Rust P, *et al.* Serum carotenoid and retinol levels during childhood infections. *Ann Nutr Metab* 2004;48:156–62.
- Thurnham DI, McCabe GP, Northrop-Clewes CA, *et al.* Effects of subclinical infection on plasma retinol concentrations and assessment of prevalence of vitamin A deficiency: meta-analysis. *The Lancet* 2003;362:2052–8.
- Pelletier O. Smoking and vitamin C levels in humans. *Am J Clin Nutr* 1968;21:1259–67.
- Sargeant LA, Jaeckel A, Wareham NJ. Interaction of vitamin C with the relation between smoking and obstructive airways disease in EPIC Norfolk. European Prospective Investigation into Cancer and Nutrition. *Eur Respir J* 2000;16:397–403.
- Shin JY, Shim JY, Lee DC, *et al.* Smokers with adequate vitamin C intake show a preferable pulmonary function test. *J Am Coll Nutr* 2015;34:385–90.
- Park HJ, Byun MK, Kim HJ, *et al.* Dietary vitamin C intake protects against COPD: the Korea National Health and Nutrition Examination Survey in 2012. *Int J Chron Obstruct Pulmon Dis* 2016;11:2721–8.
- Joshi P, Kim WJ, Lee SA. The effect of dietary antioxidant on the COPD risk: the community-based KoGES (Ansan-Anseong) cohort. *Int J Chron Obstruct Pulmon Dis* 2015;10:2159–68.
- Al-Rejaie S, Abuhashish H, Alkhamees O, *et al.* Gender difference following high cholesterol diet induced renal injury and the protective role of rutin and ascorbic acid combination in Wistar albino rats. *Lipids Health Dis* 2012;11:41.
- Jiao Y, Chen H, Yan J, *et al.* Genome-wide gene expression profiles in antioxidant pathways and their potential sex differences and connections to vitamin C in mice. *Int J Mol Sci* 2013;14:10042–62.