


RESEARCH ARTICLE

Big data and analysis of risk factors for gallbladder disease in the young generation of Korea

Hyung Sun Kim¹, Seong Kyung Cho², Chang Soo Kim^{1,2}, Joon Seong Park^{1*}

1 Pancreatobiliary Cancer Clinic, Department of Surgery, Gangnam Severance Hospital, Yonsei University, Seoul, Korea, **2** Department of Preventive Medicine, Yonsei University College of Medicine, Seoul, Korea

* JSPARK330@yuhs.ac



Abstract

Background/Purpose

Few studies have examined the risk factors for gallbladder (GB) disease in young adults. This study aimed to evaluate risk factors for GB disease in young adults based on big data in Korea.

Methods

All participants underwent routine checkup at the Korea Medical Institute from June 2014 to May 2015. After excluding 677 individuals with missing information in records, 724,114 individuals (435,635 men, 288,479 women) were finally included. The definition of abnormal GB finding included stones, sludge, polyps, and adenomyomatosis detected using ultrasonography. All statistical analyses were performed using SAS software version 9.2.

Results

Overall, 27,130 (17.5%) individuals were diagnosed as having abnormal GB finding in the young age group ($N = 154,463$, aged 20–39 years). In men, significant differences in low-density lipoprotein (LDL) and cholesterol levels were observed between the abnormal GB finding group and normal GB group ($p < 0.05$). In women, a significant difference in smoking history was noted between the abnormal GB finding group and normal GB group ($p < 0.05$). The prevalence rate of GB stones was 1.9% (27,979/154,463) in the young age group. High body mass index (BMI), large thigh circumference, and low high-density lipoprotein (HDL) level in women and low HDL level in men were independent risk factors for the presence of GB stones ($p < 0.05$).

Conclusion

In this study, obesity-related factors (BMI, waist size, thigh circumference, and cholesterol, LDL, and HDL levels) correlated with GB disease in the young generation of Korea.

OPEN ACCESS

Citation: Kim HS, Cho SK, Kim CS, Park JS (2019) Big data and analysis of risk factors for gallbladder disease in the young generation of Korea. *PLoS ONE* 14(2): e0211480. <https://doi.org/10.1371/journal.pone.0211480>

Editor: David Meyre, McMaster University, CANADA

Received: August 20, 2018

Accepted: January 15, 2019

Published: February 22, 2019

Copyright: © 2019 Kim et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All data underlying the study are within the paper and its Supporting Information files.

Funding: This thesis was financially supported by Korea Medical Institute (KMI).

Competing interests: This article was presented as an oral presentation with interim findings at the 6th Congress of the Asian-Pacific Hepato-Pancreato-Biliary Association (6th A-PPBA) in Pacifico Yokohama, Japan, held on June 7–10, 2017. The oral presentation's abstract was published as an "oral abstract" in the *Journal of Hepato-Biliary-*

Pancreatic Sciences. This does not alter our adherence to PLOS ONE policies on sharing data and materials.

Introduction

Gallbladder (GB) disease is a common disease of the digestive system known to occur in approximately 20% of healthy adults. In general, the prevalence of GB disease is more than double in women than in men and increases with age in both sexes, reaching approximately 30% at the age of 70 years [1–8]. In 2015, cholecystectomy ranked seventh among operative cases in Korea that were filed with the Health Insurance Review and Assessment Service [9]. Recent studies have reported a 27% growth in the number of cholecystectomies in 2015 compared with that in 2010 [9]. In Asian countries, the prevalence rate of cholelithiasis ranges from 3% to 10%; specifically, according to recent studies, the prevalence rates of cholelithiasis were 3.2% in Japan [1], 10.7% in China [2], 7.1% in North India [3], and 5.0% in Taiwan [4].

Risk factors for GB disease can be categorized into two, namely, (1) immutable factors, such as ethnicity, advanced age, female sex, and pregnancy, and (2) modifiable factors. Obesity is a major risk factor for the development of gallstones. High low-density lipoprotein (LDL), low high-density lipoprotein (HDL), and high triglyceride levels are positively correlated with gallstone formation [10, 11]. Additionally, physical activity appears to be protective, as it reduces one's risk of developing cholelithiasis [12, 13]. Diet and lifestyle, particularly with diet westernization (increased fat), changed the stone composition from pigment to cholesterol gallstones [14–16]. The prevalence of obesity, which accounts for the largest percentage of risk factors, is increasing in Korea. Furthermore, from 1979 to 2015, the average body mass index (BMI) and waist size increased among young male and female adults [9].

Taking these states into account, we need to investigate the prevalence of GB disease in young individuals undergoing health screening owing to the change in dietary habits in recent years. Few studies have examined the risk factors for GB disease in the young population [17, 18]. Therefore, this study aimed to evaluate the risk factors for GB disease in young adults based on health checkup data in Korea.

Materials and methods

Study population and screening program

This study was conducted using data maintained by the Korea Medical Institute (KMI) from September 2014 to August 2015. The included cohort comprised 439,792 men and 295,498 women aged 20–89 years who underwent screening via the KMI between 2014 and 2015. Among them, 677 individuals with missing information in records were excluded, and 724,114 individuals were finally included (435,635 men, 288,479 women) (Fig 1).

In this study, we reviewed young adults aged 20–39 years (217,986 male patients, 157,756 female patients) and collected data from the KMI database, which included information on participants' anthropometric characteristics and laboratory and imaging studies. Anthropometric data included height, weight, and waist and thigh circumferences. A questionnaire that included questions about life habits (history of alcohol consumption and smoking) was used in the examination. Laboratory tests included a blood test (complete blood cell count, liver function, renal function, thyroid function, lipid profile, anemia, and tumor marker and urine analyses). Metabolic syndrome was defined as the presence of three risk factors (waist circumference ≥ 102 cm for men or ≥ 88 cm for women; triglyceride level ≥ 150 mg/dL; HDL level < 40 mg/dL for men or < 50 mg/dL for women; blood pressure $\geq 130/85$ mmHg, and fasting glucose level ≥ 110 mg/dL), according to the National Cholesterol Education Program criteria [19].

The cutoff value for LDL level was 100 mg/dL in concordance with the indication by the Adult Treatment Panel III guidelines that the LDL level for initiating therapeutic lifestyle




Fig 1. Study flow diagram.

<https://doi.org/10.1371/journal.pone.0211480.g001>

changes is ≥ 100 mg/dL. According to these guidelines, the optimal value for LDL level was < 100 mg/dL [19]. The cutoff value for HDL level was 40 mg/dL for men and 50 mg/dL for women.

Based on average values in Koreans aged 20–30 years, the mean waist circumference is 83.08 cm for men and 73.89 cm for women, whereas the mean thigh circumference is 57.85 cm for men and 55.12 cm for women [20]. The cutoff value for BMI was 23 (BMI: overweight, 23–25 kg/m²; obese, 25–30 kg/m²; severely obese, ≥ 30 kg/m²). The cutoff value for cholesterol level was the mean value in our data. We performed further analysis of the value based on the cutoff value. Abdominal ultrasonographic findings were reported as either negative or positive (abnormal finding for the GB, liver, kidney, and pancreas). This study received approval from the institutional review board of Gangnam Severance Hospital (2015-0519-002), and informed consent was obtained from all patients.

Case identification and data selection

The outlier group was defined based on anthropometric and laboratory data. We excluded the outlier group from the total datasets. We defined the outlier group as below 1% and above 99%. Therefore, 69,568 young (20s and 30s) men (n = 41,974) and women (n = 27,594) were included in this analysis. With respect to abnormal abdominal ultrasonographic findings, we defined abnormal GB finding as GB stones, polyps, and adenomyomatosis (Fig 1).

Data are contained within the manuscript.

Statistical analysis

The calculated prevalence of abnormal GB finding was compared with the previously reported prevalence using tests of proportions, for which a p-value < 0.05 was considered statistically significant. Anthropometric indices and laboratory values were compared for any difference, and we evaluated whether the difference was significant using paired t-test.

Continuous variables, including age, height, weight, BMI, waist and thigh circumferences, and laboratory data, are presented as mean ± standard deviation (SD) and were compared using Student's t-test. Categorical variables, including metabolic syndrome and a history of alcohol consumption and smoking, are expressed as counts and percentages and were compared using Fisher's exact test. The results of this model are presented as odds ratio (OR) with 95% confidence intervals (CIs) using multivariate logistic regression analysis. Data were processed using SAS software version 9.2 (SAS Institute, Cary, NC, USA).

Results

Baseline characteristics

Anthropometric indices and laboratory data of the young age group were analyzed (Table 1). In the young age group, the mean age of men and women was 32.9 (SD 4.2) years and 31.1 (SD 4.8) years, respectively. In addition, the mean height and weight were 172.7 (SD 19.5) cm and 74.1 (SD 12.9) kg in men and 159.9 (SD 18.8) cm and 55.4 (SD 10.2) kg in women, respectively. The BMI was 24.2 (SD 3.9) kg/m² in men and 21.1 (SD 3.7) kg/m² in women. A BMI of 24.2 kg/m² in men is defined as overweight in Korea (BMI: overweight, 23–25 kg/m²; obese, 25–30 kg/m²; severely obese, ≥30 kg/m²) [10].

The mean waist and thigh circumference in the young age group were 81.9 (SD 11.5) cm and 53.2 (SD 7.2) cm in men and 69.4 (SD 10.5) cm and 49.3 (SD 7.2) cm in women, respectively. All variables showed statistically significant differences between men and women.

Metabolic syndrome

In the young age group, 1,181 (1.7%) participants had metabolic syndrome. Among participants, 20337 (29.2%) were smokers, 69227 (99.5%) were drinkers, and 289 were diagnosed

Table 1. Anthropometric indices and laboratory data of the young age group (age, 20s to 30s).

	Men (n = 41,974)	Women (n = 27,594)	P-value ^a
Age, yr	32.9±4.2	31.1±4.8	<0.01
Height, cm	172.7±19.5	159.9±18.8	<0.01
Weight, kg	74.1±12.9	55.4±10.2	<0.01
BMI	24.2±3.9	21.1±3.7	<0.01
Waist circumference, cm	81.9±11.5	69.4±10.5	<0.01
Thigh circumference, cm	53.2±7.2	49.3±7.2	<0.01
Hb [12–16g/dl]	15.5±0.8	13.1±0.8	<0.01
Hct [37–47%]	44.9±2.2	39.1±2.2	<0.01
AST [16–37U/L]	23.0±7.6	18.0±5.0	<0.01
ALT [11–46U/L]	28.0±16.8	14.2±7.9	<0.01
Bilirubin [0.3–1.8mg/dl]	0.9±0.3	0.8±0.2	<0.01
Albumin [3.4–5.3g/dl]	4.5±0.2	4.4±0.2	<0.01
r-GTP [8–46U/L]	39.8±27.8	18.1±11.4	<0.01
ALP [35–83U/L]	54.4±11.7	43.8±10.0	<0.01
Cholesterol [<200mg/dl]	190.1±30.2	177.4±26.8	<0.01
HDL [40–99mg/dl]	55.1±12.4	68.7±13.8	<0.01
LDL [<100mg/dl]	109.9±27.5	94.4±24.1	<0.01
Triglyceride [<150mg/dl]	127.1±73.2	72.1±38.8	<0.01

Abbreviations: yr, years; BMI, body mass index; Hb, hemoglobin; Hct, hematocrit; AST, aspartate aminotransferase; ALT, alanine aminotransferase; r-GTP, gamma-glutamyl transpeptidase; ALP, alkaline phosphatase; HDL, high-density lipoprotein; LDL, low-density lipoprotein

^a p values were calculated using the t test.

<https://doi.org/10.1371/journal.pone.0211480.t001>

with diabetes. All variables showed statistically significant differences between men and women (Table 2).

Young age group with abnormal GB finding

Baseline characteristics. In men and women, there were significant differences in age values between the abnormal GB finding group and normal GB group. In men, significant differences in BMI, triglyceride level, and laboratory data (hemoglobin, hematocrit, aspartate aminotransferase, alanine aminotransferase, albumin, gamma-glutamyl transpeptidase, alkaline phosphatase) were observed between the abnormal GB finding group and normal GB group. In women, significant differences in the values of anthropometric indices (weight, BMI, waist and thigh circumferences), LDL and HDL levels, and laboratory data (hemoglobin, hematocrit, alanine aminotransferase, albumin) between the abnormal GB finding group and normal GB group (Table 3).

Risk factors for abnormal GB finding. In the univariate analysis, significant differences in cholesterol level among men were observed between the abnormal GB finding group and normal GB group (cholesterol: OR = 0.928, 95% CI = 0.882–0.976, $p < 0.01$). Among women, there were significant differences in waist circumference (OR = 1.177, 95% CI = 1.063–1.304, $p < 0.01$), thigh circumference (OR = 1.111, 95% CI = 1.021–1.20, $p = 0.01$), BMI (OR = 1.124, 95% CI = 1.027–1.231, $p = 0.01$), LDL level (OR = 1.098, 95% CI = 1.015–1.187, $p = 0.02$), cholesterol level (OR = 1.082, 95% CI = 0.999–1.171, $p = 0.05$), and smoking history (OR = 1.189, 95% CI = 1.013–1.395, $p = 0.03$) between the abnormal GB finding group and normal GB group (Table 4).

In the multivariate analysis, significant differences in LDL level (OR = 1.081, 95% CI = 1.007–1.161, $p = 0.03$) and cholesterol level (OR = 0.88, 95% CI = 0.821–0.943, $p < 0.01$) were noted among men between the abnormal GB finding group and normal GB group (Fig 2A). Furthermore, significant difference in smoking history was observed among women between the abnormal GB finding group and normal GB group (smoking history: OR = 1.196, 95% CI = 1.019–1.405, $p = 0.03$) (Fig 2B).

Young age group with GB stones

We compared the GB stone group and non-GB stone group among young adults. In the univariate analysis, significant differences in waist circumference (OR = 1.192, 95% CI = 1.003–1.417, $p = 0.04$), BMI (OR = 1.204, 95% CI = 1.035–1.401, $p = 0.01$), and HDL level (OR = 0.813, 95% CI = 0.705–0.936, $p < 0.01$) were observed among men between the GB stone

Table 2. Metabolic syndrome and health behaviors in the young age group.

Variable		Men(n = 41974)	Women(n = 27594)	P -value ^a
Metabolic Syndrome	Yes	656	525	<0.01
	No	41318	27069	
Smoking	Non smoker	23664	25567	<0.01
	Smoker	18310	2027	
Drinking	Non drinker	144	197	<0.01
	Drinker	41830	27397	
Diabetes mellitus	Yes	208	81	<0.01
	No	41766	27513	

^a p values were calculated using the t test.

<https://doi.org/10.1371/journal.pone.0211480.t002>

Table 3. Anthropometric indices and laboratory data of the young age group with abnormal GB finding (stones, sludge, polyps, and adenomyomatosis) and normal GB finding.

	Men (n = 7631)			Women (n = 3042)		
	Abnormal GB	Normal GB	p-value ^a	Abnormal GB	Normal GB	p-value ^a
Age, yr	33.5±3.9	32.9±4.2	<0.01	32.7±4.3	31.3±4.8	<0.01
Height, cm	174.8±5.4	174.7±5.6	0.41	162.0±4.9	162.0±5.1	0.85
Weight, kg	74.9±9.6	75.0±10.1	0.11	57.1±8.2	56.5±7.8	<0.01
BMI	24.5±2.8	24.6±2.9	0.03	21.7±2.9	21.5±2.8	<0.01
Waist circumference, cm	82.8±7.1	82.9±7.3	0.08	71.3±7.0	70.7±6.7	<0.01
Thigh circumference, cm	53.9±4.2	53.9±4.4	0.83	50.4±4.4	50.2±4.3	0.01
Hb [12-16g/dl]	15.4±0.8	15.5±0.8	<0.01	13.1±0.8	13.0±13.0	<0.01
Hct [37-47%]	44.8±2.2	44.9±2.2	<0.01	38.9±2.2	39.1±2.2	<0.01
AST [16-37U/L]	22.3±7.3	23.3±7.9	<0.01	17.9±5.1	18.1±5.1	0.14
ALT [11-46U/L]	26.7±15.4	28.6±17.3	<0.01	14.7±8.3	14.3±7.9	0.01
Bilirubin [0.3-1.8mg/dl]	0.9±0.3	0.9±0.3	0.56	0.8±0.2	0.8±0.3	0.15
Albumin [3.4-5.3g/dl]	4.4±0.2	4.5±0.2	<0.01	4.3±0.2	4.4±0.2	<0.01
r-GTP [8-46U/L]	39.0±27.8	40.3±28.3	<0.01	18.3±12.0	18.1±11.4	0.24
ALP [35-83U/L]	53.9±11.8	54.4±11.7	<0.01	43.6±10.4	43.7±9.9	0.72
Cholesterol [<200mg/dl]	189.6±29.9	190.7±30.3	<0.01	178.5±27.3	177.6±26.8	0.08
HDL [40-99mg/dl]	54.7±12.3	54.9±12.4	0.07	67.7±13.7	68.4±13.7	0.02
LDL [<100mg/dl]	110.3±27.1	110.3±27.5	0.94	96.3±24.3	94.7±21.2	<0.01
Triglyceride [<150mg/dl]	124.1±70.8	128.2±72.8	<0.01	72.8±39.3	72.5±38.6	0.65

Abbreviations: GB, gallbladder; yr, years; BMI, body mass index; Hb, hemoglobin; Hct, hematocrit; AST, aspartate aminotransferase; ALT, alanine aminotransferase; r-GTP, gamma-glutamyl transpeptidase; ALP, alkaline phosphatase; HDL, high-density lipoprotein; LDL, low-density lipoprotein

^a p values were calculated using the t test.

<https://doi.org/10.1371/journal.pone.0211480.t003>

group and normal GB group. Moreover, significant differences in waist circumference (OR = 2.021, 95% CI = 1.666–2.452, p < 0.01), thigh circumference (OR = 1.785, 95% CI = 1.502–2.123, p < 0.01), BMI (OR = 2.079, 95% CI = 1.742–2.48, p < 0.01), and HDL level (OR = 0.583, 95% CI = 0.454–0.75, p < 0.01) were noted among women between the GB stone group and normal GB group (Table 4).

With respect to anthropometric indices, BMI, waist circumference, and HDL level were significant risk factors for the presence of GB stones in the male GB stone group (p < 0.01). In women, the LDL level was also a significant risk factor for the presence of GB stones (p < 0.01) (Table 5).

Multivariate analysis of the risk of gallstones by some variables, including anthropometric indices and obesity-related factors, was performed in the young age group. In men, low HDL level (OR = 0.858, 95% CI = 0.741–0.994, p = 0.01) was an independent risk factor for the presence of GB stones (Fig 3A). In women, high BMI (OR = 1.512, 95% CI = 1.17–1.955, p < 0.01), large thigh circumference (OR = 1.288, 95% CI = 1.04–1.596, p = 0.02), and low HDL level (OR = 0.747, 95% CI = 0.573–0.972, p = 0.03) were independent risk factors for the presence of GB stones (Fig 3B).

Discussion

BMI, obesity, abdominal fat, metabolic syndrome, and diabetes mellitus are confirmed risk factors for gallstones [5, 10, 12, 21–24]. However, the prevalence of abnormal GB finding among young adults with a high level of obesity has not been investigated. The present study is the first large-scale study on abnormal GB findings detected by abdominal ultrasonography in

Table 4. Univariate analysis of risk factors for abnormal GB finding and GB stones.

Variables	Men				Women			
	Abnormal GB		GB stone		Abnormal GB		GB stone	
	Odds ratio	P value ^b	Odds ratio	P value ^b	Odds ratio	P value ^b	Odds ratio	P value ^b
	[95%CI]		[95%CI]		[95%CI]		[95%CI]	
Waist circumference	1.013	0.67	1.192	0.04	1.177	<0.01	2.021	<0.01
	[0.955–1.075]		[1.003–1.417]		[1.063–1.304]		[1.666–2.452]	
Thigh circumference	1.012	0.65	1.016	0.82	1.111	0.01	1.785	<0.01
	[0.961–1.066]		[0.88–1.173]		[1.021–1.209]		[1.502–2.123]	
BMI	1.004	0.88	1.204	0.01	1.124	0.01	2.079	<0.01
	[0.952–1.059]		[1.035–1.401]		[1.027–1.231]		[1.742–2.48]	
LDL	0.993	0.78	1.017	0.81	1.098	0.02	1.132	0.16
	[0.943–1.046]		[0.881–1.176]		[1.015–1.187]		[0.953–1.344]	
HDL	0.967	0.21	0.813	<0.01	0.888	0.08	0.583	<0.01
	[0.918–1.019]		[0.705–0.936]		[0.777–1.016]		[0.454–0.75]	
Cholesterol	0.928	<0.01	0.921	0.25	1.082	0.05	0.999	0.99
	[0.882–0.976]		[0.801–1.059]		[0.999–1.171]		[0.837–1.191]	
Smoking	0.989	0.65	0.921	0.25	1.189	0.03	0.854	0.32
	[0.94–1.04]		[0.801–1.06]		[1.013–1.395]		[0.625–1.167]	
Drinking	1.138	0.54	1.909	0.16	0.953	0.83	0.251	0.17
	[0.747–1.736]		[0.779–4.682]		[0.602–1.51]		[0.035–1.798]	

Abbreviations: GB, gallbladder; BMI, body mass index; HDL, high-density lipoprotein; LDL, low-density lipoprotein

^bp values were calculated using the cox-proportional hazard model

<https://doi.org/10.1371/journal.pone.0211480.t004>

healthy individuals. As the prevalence of obesity increases globally [25–27], the prevalence of gallstones associated with obesity is expected to also increase. In particular, the prevalence of obesity is rapidly increasing in Korea owing to the westernization of food and lifestyle [28]. In recent years, obesity has emerged as a major health threat in Korea [29]. The Korean National Health Insurance Service reported that the percentage of obese Koreans (those with a BMI of 30 kg/m² or above) increased from 2.5% of the entire population in 2002 to 4.2% in 2012 and that the prevalence rate of GB stones in the 20s and 30s was 0.95% and 2.26%, respectively [9]. According to another study, the prevalence rate of gallstones in the early 2000s was 0.39% for those aged <30 years and 0.99% for those in their 30s [21]. In the present study, which was performed using data from September 2014 to August 2015, the prevalence rate of abnormal GB finding and GB stones was 17.5% and 1.9%, respectively. This result showed that the increase in obesity in Korea for approximately 10 years was associated with an increase in the prevalence of gallstones.

However, abdominal ultrasonography is performed to identify the cause of persistent abdominal discomfort, and several young individuals have been diagnosed with cholecystitis. Therefore, detecting young adults with asymptomatic GB disease is important because early diagnosis of GB disease can be the first step to finding the cause of indigestion and abdominal pain.

In our data, abnormal ultrasonographic findings of GB disease included cholesterol and pigment stones, polyps, and adenomyomatosis. Traditionally, pigment and cholesterol stones were thought to represent a different pathogenesis. However, recent findings have indicated that these types of stones share the same risk factors [30–32].

Variants of the gene encoding UGT1A1 (uridine 5'-diphosphate (UDP)- glucuronosyl-transferase 1A1) responsible for bilirubin conjugation were correlated with the risk of

a. Odds ratio for abnormal GB finding in young men observed from health checkup data.

Risk factors		Odds Ratio	95% CI	P value
Waist circumference		1.02	0.945-1.1	0.61
Thigh circumference		1.011	0.95-1.075	0.73
BMI		0.992	0.924-1.066	0.83
LDL		1.081	1.007-1.161	0.03
HDL		0.979	0.927-1.033	0.43
Cholesterol		0.88	0.821-0.943	<0.01
Smoking		0.987	0.938-1.039	0.62
Drinking		1.122	0.735-1.711	0.59

Abbreviations: BMI, body mass index; LDL, low density lipoprotein; HDL, high density lipoprotein;

b. Odds ratio for abnormal GB finding in young women observed from health checkup data.

Risk factors		Odds Ratio	95% CI	P value
Waist circumference		1.129	0.987-1.291	0.07
Thigh circumference		1.05	0.949-1.162	0.34
BMI		1.001	0.881-1.137	0.98
LDL		1.046	0.937-1.167	0.42
HDL		0.916	0.797-1.053	0.21
Cholesterol		1.037	0.929-1.158	0.51
Smoking		1.196	1.019-1.405	0.03
Drinking		0.879	0.543-1.423	0.59

Abbreviations: BMI, body mass index; LDL, low density lipoprotein; HDL, high density lipoprotein;

Fig 2. Odds ratios for abnormal GB finding in young men (A) and women (B) observed from health checkup data.

<https://doi.org/10.1371/journal.pone.0211480.g002>

gallstones and stone bilirubin content, suggesting common factors in the pathogenesis of cholesterol and pigment gallstones [31].

As these stones were not classified in our data, this was a limitation in determining the risk factors for GB disease. In this study, the mean values for all obesity-related factors among young men and women were higher in the abnormal GB finding group than in the normal GB group. This finding suggests that obesity-related factors affect GB disease in young adults.

Among men, BMI was higher in the abnormal GB finding group than in the normal GB group, which was contrary to our conclusion. BMI and waist circumference among women were higher in the abnormal GB finding group than in the normal GB group (Table 3). For this reason, metabolic factors in women seem to exert a greater effect on GB disease. Lifestyle

Table 5. Comparison of the GB stone group and non-GB stone group among young adults.

	Men (n = 37301)			Women (n = 22098)		
	GB stone (n = 806)	Non GB stone (n = 36495)	p-value ^a	GB stone (n = 554)	Non GB stone (n = 21544)	p-value ^a
BMI	24.9 ± 3.1	24.6 ± 2.9	<0.01	22.8 ± 3.6	21.5 ± 2.8	<0.01
Waist circumference, cm	83.8 ± 7.8	82.9 ± 7.3	<0.01	73.5 ± 8.2	70.7 ± 6.7	<0.01
Thigh circumference, cm	54.1 ± 4.4	53.9 ± 4.4	0.21	51.6 ± 4.9	50.2 ± 4.3	0.21
Cholesterol [$<200\text{mg/dl}$]	189.3 ± 30.9	190.5 ± 30.2	0.29	178.9 ± 28.2	177.7 ± 26.8	0.30
HDL [40-99mg/dl]	53.2 ± 11.6	55.0 ± 12.4	<0.01	65.1 ± 13.9	68.4 ± 13.7	<0.01
LDL [$<100\text{mg/dl}$]	110.8 ± 27.1	110.3 ± 27.5	0.61	97.9 ± 24.6	94.9 ± 24.2	<0.01

Abbreviations: GB, gallbladder; BMI, body mass index; HDL, high-density lipoprotein; LDL, low-density lipoprotein

^a p values were calculated using the t test.

<https://doi.org/10.1371/journal.pone.0211480.t005>

habits, such as smoking and drinking, seem to have a greater effect on risk factors for GB disease in men.

The risk of abnormal GB finding in young men was 1.5 times higher than that in young women (OR = 1.553, 95% CI = 1.464–1.648, $p < 0.01$). This result is in contrast to a previous finding that indicated generally higher prevalence of GB disease in women than in men [5]. In this study, differences in the results for obesity-related factors in the abnormal GB finding group were larger in men than in women. Compared with young women, young men in Korea are familiar with the culture of having dinner and drinking with company. This culture results in more obesity, especially among young men [33, 34]. Relatively, young women prefer to have a slender appearance [35, 36], and most obesity-related factors in the abnormal GB finding group were within the normal range.

In men, obesity-related factors such as LDL level are proportional to the risk of abnormal GB finding. However, cholesterol level has the opposite effect. There was only a subtle difference in cholesterol level between the abnormal GB finding group and normal GB group; furthermore, cholesterol level was within the normal range. In women, smoking is proportional to the risk of abnormal GB finding. The role of smoking as a risk factor has not yet been clarified. Some studies have reported a correlation between smoking and GB disease [37–39]; nevertheless, the definite mechanism of smoking in GB disease remains unknown. GB disease associated with smoking may be explained by changes in serum cholesterol concentrations [38].

In this study, abdominal ultrasonography was performed in adults over 20 years of age. However, most young adults in their 20s to 40s until now do not undergo ultrasonography because abdominal ultrasonography is not covered by insurance. However, our results would be meaningful because insurance for ultrasonography is implemented in Korea starting from this year.

The present study has limitations, particularly its retrospective design and data from young, healthy workers who underwent health screening. Because of the high proportion of young individuals, the prevalence in young adults in relation to the overall prevalence is not definitely indicative of the actual prevalence. In addition, as we used data from one year in 2014, the study included recent data; nonetheless, the fact that it is not a multiyear study is a disadvantage. Additionally, the participants in the study were not followed up. Thus, we could not evaluate their treatment for GB disease (cholecystectomy or medications) after their health checkup. In our study, we did not classify each ultrasonographic finding for GB disease (cholesterol stone, pigment stone, polyp, or adenomyomatosis). We could not also evaluate each risk factor. However, for GB stones, risk factors were analyzed separately. These results were

a. Odds ratio for the presence of gallstones in young men observed from health checkup data.

Risk factors		Odds Ratio	95% CI	P value
Waist circumference		1.096	0.881-1.365	0.41
Thigh circumference		0.894	0.755-1.059	0.19
BMI		1.195	0.976-1.464	0.08
LDL		1.094	0.901-1.329	0.36
HDL		0.858	0.741-0.994	0.01
Cholesterol		0.845	0.7-1.02	0.08
Smoking		0.931	0.809-1.072	0.32
Drinking		1.838	0.748-4.517	0.18

Abbreviations: BMI, body mass index; LDL, low density lipoprotein; HDL, high density lipoprotein;

b. Odds ratio for the presence of gallstones in young women observed from health checkup data.

Risk factors		Odds Ratio	95% CI	P value
Waist circumference		1.264	0.975-1.638	0.09
Thigh circumference		1.288	1.04-1.596	0.02
BMI		1.512	1.17-1.955	<0.01
LDL		1.066	0.839-1.355	0.6
HDL		0.747	0.573-0.972	0.03
Cholesterol		0.877	0.688-1.119	0.29
Smoking		0.9	0.657-1.232	0.51
Drinking		0.241	0.034-1.73	0.15

Abbreviations: BMI, body mass index; LDL, low density lipoprotein; HDL, high density lipoprotein;

Fig 3. Odds ratios for the presence of gallstones in young men (A) and women (B) observed from health checkup data.

<https://doi.org/10.1371/journal.pone.0211480.g003>

different from those of the overall analysis of GB disease. GB stones are more likely to be associated with obesity-related factors (BMI, HDL level, and thigh circumference). In this study, we could not describe the degree of alcohol consumption according to the amount of alcohol because the questionnaire used for collecting health checkup data was not designed to assess the exact amount of alcohol.

Conclusions

This study is the first attempt to analyze the prevalence of and risk factors for GB disease in young adults based on large-scale health screening data. The use of big data aims to analyze large, heterogeneous datasets to provide in-depth insights into complex processes, and it has the advantages of evaluating risk factors. Based on these results, we might consider the

probability of GB disease in young adults with digestive symptoms, such as persistent abdominal discomfort.

Supporting information

S1 File. Raw data of health checkup.

(XLSX)

Author Contributions

Conceptualization: Joon Seong Park.

Data curation: Hyung Sun Kim, Seong Kyung Cho.

Formal analysis: Chang Soo Kim.

Investigation: Hyung Sun Kim.

Methodology: Seong Kyung Cho.

Resources: Seong Kyung Cho, Chang Soo Kim, Joon Seong Park.

Supervision: Chang Soo Kim.

Writing – original draft: Hyung Sun Kim.

Writing – review & editing: Joon Seong Park.

References

1. Nomura H, Kashiwagi S, Hayashi J, Kajiyama W, Ikematsu H, Noguchi A, et al. Prevalence of gallstone disease in a general population of Okinawa, Japan. *Am J Epidemiol*. 1988; 128(3):598–605. Epub 1988/09/01. PMID: [3046339](https://pubmed.ncbi.nlm.nih.gov/3046339/).
2. Sun H, Tang H, Jiang S, Zeng L, Chen EQ, Zhou TY, et al. Gender and metabolic differences of gallstone diseases. *World J Gastroenterol*. 2009; 15(15):1886–91. Epub 2009/04/17. <https://doi.org/10.3748/wjg.15.1886> PMID: [19370788](https://pubmed.ncbi.nlm.nih.gov/19370788/); PubMed Central PMCID: [PMC2670418](https://pubmed.ncbi.nlm.nih.gov/PMC2670418/).
3. Unisa S, Jagannath P, Dhir V, Khandelwal C, Sarangi L, Roy TK. Population-based study to estimate prevalence and determine risk factors of gallbladder diseases in the rural Gangetic basin of North India. *HPB (Oxford)*. 2011; 13(2):117–25. Epub 2011/01/19. <https://doi.org/10.1111/j.1477-2574.2010.00255.x> PMID: [21241429](https://pubmed.ncbi.nlm.nih.gov/21241429/); PubMed Central PMCID: [PMC3044346](https://pubmed.ncbi.nlm.nih.gov/PMC3044346/).
4. Chen CH, Huang MH, Yang JC, Nien CK, Etheredge GD, Yang CC, et al. Prevalence and risk factors of gallstone disease in an adult population of Taiwan: an epidemiological survey. *J Gastroenterol Hepatol*. 2006; 21(11):1737–43. Epub 2006/09/21. <https://doi.org/10.1111/j.1440-1746.2006.04381.x> PMID: [16984599](https://pubmed.ncbi.nlm.nih.gov/16984599/).
5. Shaffer EA. Epidemiology and risk factors for gallstone disease: has the paradigm changed in the 21st century? *Curr Gastroenterol Rep*. 2005; 7(2):132–40. PMID: [15802102](https://pubmed.ncbi.nlm.nih.gov/15802102/).
6. European Association for the Study of the Liver. Electronic address eee. EASL Clinical Practice Guidelines on the prevention, diagnosis and treatment of gallstones. *J Hepatol*. 2016; 65(1):146–81. Epub 2016/04/18. <https://doi.org/10.1016/j.jhep.2016.03.005> PMID: [27085810](https://pubmed.ncbi.nlm.nih.gov/27085810/).
7. Lammert F, Gurusamy K, Ko CW, Miquel JF, Mendez-Sanchez N, Portincasa P, et al. Gallstones. *Nat Rev Dis Primers*. 2016; 2:16024. Epub 2016/04/29. <https://doi.org/10.1038/nrdp.2016.24> PMID: [27121416](https://pubmed.ncbi.nlm.nih.gov/27121416/).
8. Portincasa P, Moschetta A, Palasciano G. Cholesterol gallstone disease. *Lancet*. 2006; 368(9531):230–9. Epub 2006/07/18. [https://doi.org/10.1016/S0140-6736\(06\)69044-2](https://doi.org/10.1016/S0140-6736(06)69044-2) PMID: [16844493](https://pubmed.ncbi.nlm.nih.gov/16844493/).
9. Service NHI. Korea Health Industry Statistic System 2015.
10. Aune D, Norat T, Vatten LJ. Body mass index, abdominal fatness and the risk of gallbladder disease. *Eur J Epidemiol*. 2015; 30(9):1009–19. <https://doi.org/10.1007/s10654-015-0081-y> PMID: [26374741](https://pubmed.ncbi.nlm.nih.gov/26374741/).
11. Tsai CJ, Leitzmann MF, Willett WC, Giovannucci EL. Long-chain saturated fatty acids consumption and risk of gallstone disease among men. *Ann Surg*. 2008; 247(1):95–103. <https://doi.org/10.1097/SLA.0b013e31815792c2> PMID: [18156928](https://pubmed.ncbi.nlm.nih.gov/18156928/).

12. Leitzmann MF, Rimm EB, Willett WC, Spiegelman D, Grodstein F, Stampfer MJ, et al. Recreational physical activity and the risk of cholecystectomy in women. *New Engl J Med*. 1999; 341(11):777–84. <https://doi.org/10.1056/NEJM199909093411101> WOS:000082534000001. PMID: 10477775
13. Molina-Molina E, Lunardi Baccetto R, Wang DQ, de Bari O, Krawczyk M, Portincasa P. Exercising the hepatobiliary-gut axis. The impact of physical activity performance. *Eur J Clin Invest*. 2018; 48(8): e12958. Epub 2018/05/26. <https://doi.org/10.1111/eci.12958> PMID: 29797516.
14. Larsson SC, Hakansson N, Wolk A. Healthy dietary patterns and incidence of biliary tract and gallbladder cancer in a prospective study of women and men. *Eur J Cancer*. 2017; 70:42–7. <https://doi.org/10.1016/j.ejca.2016.10.012> PMID: 27870981.
15. McConnell TJ, Appleby PN, Key TJ. Vegetarian diet as a risk factor for symptomatic gallstone disease. *Eur J Clin Nutr*. 2017; 71(6):731–5. <https://doi.org/10.1038/ejcn.2016.252> WOS:000403057500006. PMID: 28272400
16. Park Y, Kim D, Lee JS, Kim YN, Jeong YK, Lee KG, et al. Association between diet and gallstones of cholesterol and pigment among patients with cholecystectomy: a case-control study in Korea. *J Health Popul Nutr*. 2017; 36. ARTN 39 <https://doi.org/10.1186/s41043-017-0116-y> WOS:000416123500001. PMID: 29169372
17. Constantinescu T, Huwood Al Jabouri AK, Bratucu E, Olteanu C, Toma M, Stoiculescu A. Gallstone disease in young population: incidence, complications, therapeutic approach. *Chirurgia (Bucur)*. 2012; 107(5):579–82. PMID: 23116830.
18. Chilimuri S, Gaduputi V, Tariq H, Nayudu S, Vakde T, Glandt M, et al. Symptomatic Gallstones in the Young: Changing Trends of the Gallstone Disease-Related Hospitalization in the State of New York: 1996–2010. *J Clin Med Res*. 2017; 9(2):117–23. <https://doi.org/10.14740/jocmr2847w> PMID: 28090227; PubMed Central PMCID: PMC5215015.
19. Program TNCE. Third Report of the Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III, or ATP III). 2002.
20. <https://sizekorea.kr>.
21. Yun-Jin Chung MD, Young-Dae Park M.D., Ho-Chul Lee M.D., Han-Jin Cho M.D., Kwan-Sik Park MD, Eun-Hee Seo M.D., et al. Prevalence and risk factors of gallstones in a general health screened population. *Korean Journal of Medicine* 2007; 72(5).
22. Banim PJ, Luben RN, Bulluck H, Sharp SJ, Wareham NJ, Khaw KT, et al. The aetiology of symptomatic gallstones quantification of the effects of obesity, alcohol and serum lipids on risk. Epidemiological and biomarker data from a UK prospective cohort study (EPIC-Norfolk). *Eur J Gastroenterol Hepatol*. 2011; 23(8):733–40. Epub 2011/05/31. <https://doi.org/10.1097/MEG.0b013e3283477cc9> PMID: 21623190.
23. Chen Y, Kong J, Wu S. Cholesterol gallstone disease: focusing on the role of gallbladder. *Lab Invest*. 2015; 95(2):124–31. <https://doi.org/10.1038/labinvest.2014.140> PMID: 25502177.
24. Cai JS, Qiang S, Bao-Bing Y. Advances of recurrent risk factors and management of choledocholithiasis. *Scand J Gastroenterol*. 2017; 52(1):34–43. <https://doi.org/10.1080/00365521.2016.1224382> PMID: 27610642.
25. James PT. Obesity: the worldwide epidemic. *Clin Dermatol*. 2004; 22(4):276–80. Epub 2004/10/12. <https://doi.org/10.1016/j.clindermatol.2004.01.010> PMID: 15475226.
26. Ezzati M, Bentham J, Di Cesare M, Bilano V, Bixby H, Zhou B, et al. Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128.9 million children, adolescents, and adults. *Lancet*. 2017; 390(10113):2627–42. [https://doi.org/10.1016/S0140-6736\(17\)32129-3](https://doi.org/10.1016/S0140-6736(17)32129-3) WOS:000418101000024. PMID: 29029897
27. Ogden CL, Carroll MD, Lawman HG, Fryar CD, Kruszon-Moran D, Kit BK, et al. Trends in Obesity Prevalence Among Children and Adolescents in the United States, 1988–1994 Through 2013–2014. *JAMA*. 2016; 315(21):2292–9. <https://doi.org/10.1001/jama.2016.6361> WOS:000377183100018. PMID: 27272581
28. Ramachandran A, Snehalatha C. Rising burden of obesity in Asia. *J Obes*. 2010; 2010. Epub 2010/09/28. <https://doi.org/10.1155/2010/868573> PMID: 20871654; PubMed Central PMCID: PMC2939400.
29. Nam-Kyou Bae K-HK, In-Sun Kwon, Young-Chae Cho. Changes in Prevalence of Obesity for 10 Years (1997–2007) and its related factors in Health Checkup Examinees. *Journal of the Korea Academia-Industrial cooperation Society*. 2009; 10(5):1091–9.
30. Weerakoon HT, Ranasinghe JG, Navaratna A, Sivakanesan R, Galketiya KB, Rosairo S. Can the type of gallstones be predicted with known possible risk factors?: A comparison between mixed cholesterol and black pigment stones. *BMC Gastroenterol*. 2014; 14:88. <https://doi.org/10.1186/1471-230X-14-88> PMID: 24884475; PubMed Central PMCID: PMC4017087.

31. Van Erpecum KJ. Pathogenesis of cholesterol and pigment gallstones: An update. *Clin Res Hepatol Gas*. 2011; 35(4):281–7. <https://doi.org/10.1016/j.clinre.2011.01.009> WOS:000289839700006. PMID: [21353662](https://pubmed.ncbi.nlm.nih.gov/21353662/)
32. Portincasa P, Moschetta A, Berardino M, Di-Ciula A, Vacca M, Baldassarre G, et al. Impaired gallbladder motility and delayed orocecal transit contribute to pigment gallstone and biliary sludge formation in beta-thalassemia major adults. *World J Gastroenterol*. 2004; 10(16):2383–90. Epub 2004/07/31. <https://doi.org/10.3748/wjg.v10.i16.2383> PMID: [15285024](https://pubmed.ncbi.nlm.nih.gov/15285024/); PubMed Central PMCID: [PMCPMC4576293](https://pubmed.ncbi.nlm.nih.gov/PMC4576293/).
33. Amudha S. Poobalana LSA, Amanda Clarke and William Cairns S. Smitha. Diet behaviour among young people in transition to adulthood (18–25 year olds): a mixed method study. *Health Psychology & Behavioural Medicine*. 2014; 2(1):909–28.
34. Martinez-Gonzalez MA, Martin-Calvo N. The major European dietary patterns and metabolic syndrome. *Rev Endocr Metab Dis*. 2013; 14(3):265–71. <https://doi.org/10.1007/s11154-013-9264-6> WOS:000324881200006. PMID: [23979531](https://pubmed.ncbi.nlm.nih.gov/23979531/)
35. Kim CS, Ko SH, Kwon HS, Kim NH, Kim JH, Lim S, et al. Prevalence, awareness, and management of obesity in Korea: data from the Korea national health and nutrition examination survey (1998–2011). *Diabetes Metab J*. 2014; 38(1):35–43. Epub 2014/03/15. <https://doi.org/10.4093/dmj.2014.38.1.35> PMID: [24627826](https://pubmed.ncbi.nlm.nih.gov/24627826/); PubMed Central PMCID: [PMCPMC3950193](https://pubmed.ncbi.nlm.nih.gov/PMC3950193/).
36. Mi-Ok Kim J-GE, Un-Jae Chang, Kayoko Sawano, and Tokuko Miyamoto. The Comparison of Korean and Japanese College Women's Self-evaluation on Obesity and Attitude to Weight Control. *J Food Sci Nutr*. 2009; 14:303–9.
37. Degirmenci B, Albayrak R, Haktanir A, Acar M, Yucel A. Acute effect of smoking on gallbladder emptying and refilling in chronic smokers and nonsmokers: A sonographic study. *World J Gastroenterol*. 2006; 12(34):5540–3. <https://doi.org/10.3748/wjg.v12.i34.5540> WOS:000240647100019. PMID: [17006996](https://pubmed.ncbi.nlm.nih.gov/17006996/)
38. Craig WY, Palomaki GE, Haddow JE. Cigarette-Smoking and Serum-Lipid and Lipoprotein Concentrations—an Analysis of Published Data. *Brit Med J*. 1989; 298(6676):784–8. WOS:A1989T847900019. PMID: [2496857](https://pubmed.ncbi.nlm.nih.gov/2496857/)
39. Aune D, Vatten LJ, Boffetta P. Tobacco smoking and the risk of gallbladder disease. *Eur J Epidemiol*. 2016; 31(7):643–53. <https://doi.org/10.1007/s10654-016-0124-z> PMID: [26898907](https://pubmed.ncbi.nlm.nih.gov/26898907/); PubMed Central PMCID: [PMCPMC4977331](https://pubmed.ncbi.nlm.nih.gov/PMC4977331/).