University of New Hampshire University of New Hampshire Scholars' Repository

NH Water Resources Research Center Scholarship

NH Water Resources Research Center

6-1-1998

RELEASE OF NITRATE-NITROGEN AND HEAVY METALS FROM LAND-APPLIED BIOSOLIDS IN NORTHERN AREAS

George O. Estes University of New Hampshire

Jiangiang Zhao
University of New Hampshire

Thomas P. Ballestero
University of New Hampshire, tom.ballestero@unh.edu

Follow this and additional works at: https://scholars.unh.edu/nh wrrc scholarship

Recommended Citation

Estes, George O.; Zhao, Jiangiang; and Ballestero, Thomas P., "RELEASE OF NITRATE-NITROGEN AND HEAVY METALS FROM LAND-APPLIED BIOSOLIDS IN NORTHERN AREAS" (1998). NH Water Resources Research Center Scholarship. 93. https://scholars.unh.edu/nh_wrrc_scholarship/93

This Report is brought to you for free and open access by the NH Water Resources Research Center at University of New Hampshire Scholars' Repository. It has been accepted for inclusion in NH Water Resources Research Center Scholarship by an authorized administrator of University of New Hampshire Scholars' Repository. For more information, please contact nicole.hentz@unh.edu.

RELEASE OF NITRATE-NITROGEN AND HEAVY METALS FROM LAND-APPLIED BIOSOLIDS IN NORTHERN AREAS

Principal Investigators: Dr. George O. Estes, Jiangiang Zhao, and Dr. Thomas P. Ballestero, University of New Hampshire

Descriptors: Soil water relationships, nitrogen, land disposal, soil contamination, sludge, agriculture, soil profile

Research Objectives:

Field studies at two New Hampshire sites were conducted from 1993-1995 to evaluate the effects of various rates of broadcast, soil-incorporated wastewater treatment facility (WWTF) biosolids and dairy manure on water, crops and soils. The principal focus was on N release and its impact on NO3-N concentrations in soil and water.

Principal Findings and Significance:

Chemical composition of lime- stabilized, dewatered biosolids from the Concord, NH Waste Water Treatment Facility (WWTF) was highly variable ranging from 0.6-5.2% total N over a four-year period. The N content of anaerobically digested biosolids from the Hanover, NH WWTF ranged from 5.0-5.4% over the 1993-95 period; the N range in manure was 0.8-3.4%. Nitrate-N measurements made on soil with the pre-sidedress N test (PSNT) and in water collected via suction lysimeters and from monitoring wells show a slower N release from biosolids. Elevated NO3-N concentrations suggest the need for improved management of both manure and biosolids. Acceptance of higher values for mineralizable N would justify lower application rates of soil amendments rich in organic-N. Copper and zinc were the principal metals in biosolids; quantities applied were low compared to naturally occurring levels in soil.