

**HHS PUBLIC ACCESS**

Author manuscript

Transl Anim Sci. Author manuscript; available in PMC 2019 August 01.

Published in final edited form as:

Transl Anim Sci. 2017 February ; 1(1): 69–76. doi:10.2527/tas2016.0008.**Weaned pigs experimentally infected with *Salmonella* display sexually dimorphic innate immune responses without affecting pathogen colonization patterns^{1,2}****Nicole C. Burdick Sanchez^{*}, Paul R. Broadway^{*}, Jeffery A. Carroll^{*,4}, Elena V. Gart[†], Laura K. Bryan[†], Sara D. Lawhon[†]**^{*}Livestock Issues Research Unit, USDA-ARS, Lubbock, TX 79403[†]Department of Veterinary Pathology, College of Veterinary Medicine and Biomedical Sciences, Texas A&M University, College Station 77843**Abstract**

Sexually dimorphic innate immune responses have been observed in several species, but have not been studied in response to a live pathogen challenge in pigs. This study aimed to elucidate sexually dimorphic innate immune responses along with *Salmonella* translocation patterns in newly weaned pigs orally inoculated with *Salmonella*. Newly weaned pigs ($n = 8$ gilts and 12 barrows; 6.2 ± 0.2 kg BW) were obtained from a commercial swine facility and were maintained in an environmentally-controlled facility in individual pens equipped with feeders and nipple waterers. Pigs were allowed ad libitum access to a commercial non-medicated starter ration and water throughout the study. On d 12 post-weaning, pigs were anesthetized to allow placement of a temperature measuring device in the abdominal cavity for measurement of intraperitoneal temperature (TEMP). On d 17, pigs were anesthetized and fitted with indwelling jugular vein catheters. On the following day (d 18), pigs were orally inoculated with 4.7×10^9 *Salmonella typhimurium*. Blood samples were collected at 0.5-h intervals from -2 to 8 h, and at 8-h intervals from 8 to 72 h post-challenge. Whole blood was analyzed for complete blood cell counts. Serum was isolated for measurement of cortisol. Following collection of the 72 h sample, pigs were humanely euthanized and tissues were collected for *Salmonella* isolation. There was a sex \times time interaction ($P < 0.001$) for TEMP such that gilts had a greater TEMP response to the *Salmonella* challenge compared to barrows. There was also a sex \times time interaction ($P = 0.03$) for serum cortisol with gilts having decreased cortisol at 16 h yet greater cortisol at 32 h than barrows. Barrows had greater total white blood cells (17.8 vs. $16.2 \pm 0.4 \times 10^3$ cells/ μ L; $P < 0.01$;

¹Mention of trade names or commercial products in this article is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the U.S. Department of Agriculture.

²The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotope, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, DC, 20250-9410, or call (800) 795-3272 (voice) or (202) 720-6382 (TDD).

This is an open access article distributed under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

⁴Corresponding author: Jeff.Carroll@ars.usda.gov.

respectively) and neutrophils (7.8 vs. $6.1 \pm 0.4 \times 10^3$ cells/ μL ; $P < 0.01$; respectively) than gilts. However, gilts had greater lymphocytes (9.6 vs. $9.0 \pm 0.2 \times 10^3$ cells/ μL ; $P = 0.05$; respectively) than barrows. While immune parameters were influenced by sex, there was no effect of sex ($P > 0.05$) on *Salmonella* concentrations from fecal shedding 3 d post-inoculation in the cecum, mesenteric and subiliac lymph nodes, liver, spleen, gallbladder, or kidney tissues. These data demonstrate that weaned gilts appear to produce a stronger acute phase response to a *Salmonella* challenge compared to barrows, without affecting the tissue translocation or shedding of *Salmonella*.

Keywords

cortisol; innate immunity; pigs; *Salmonella*; sexual dimorphism

INTRODUCTION

The responsiveness of the immune system to a perceived challenge can be modulated by various factors, including internal factors such as the hypothalamic-pituitary-adrenal (HPA) axis and energy availability/storage, as well as naturally-occurring factors such as behavior, breed, and sex (Burdick et al., 2009; Carroll et al., 2011; Carroll et al., 2015). Specifically regarding sex, studies conducted by our lab have demonstrated sex-specific innate immune responses to a lipopolysaccharide (LPS), and to a dual corticotropin-releasing hormone (CRH) and vasopressin challenge in cattle (Hulbert et al., 2013; Carroll et al., 2015). Additionally, Williams et al. (2009) observed greater baseline and peak serum TNF- α and IL-6, but decreased IL-1 β concentrations in weaned gilts in response to an LPS challenge compared to barrows. Thus, sexual dimorphic effects are found in livestock species, and particularly in pigs utilizing exogenous challenges.

Utilizing substances such as LPS and CRH to exogenously stimulate immune and stress responses in the body have proven to be useful models for elucidating these complex biological systems. However, the use of live pathogen challenge models provides a further means of studying immune and stress responses that are more reflective of real-world scenarios. Live pathogen challenge studies are more indicative and representative of the immunological challenges pigs encounter in the commercial production setting; however increased variation is introduced due to endogenous factors associated with each animal and the specific microorganism. The use of a live *Salmonella* challenge in swine is of particular relevance due to the health issues associated with *Salmonella* outbreaks in swine herds, and due to an increasing concern within the livestock industry associated with post-harvest contamination, food safety and human illness (Ojha and Kostrzynska, 2007). Aligned with the observed effects of sex on the innate immune response that have been observed in response to various immune challenges in swine and other species, this study sought to determine whether newly weaned pigs exhibit sexually dimorphic innate immune responses to an oral *Salmonella typhimurium* challenge.

MATERIALS AND METHODS

All experimental procedures were in compliance with the *Guide for the Care and Use of Agricultural Animals in Research and Teaching* and approved by the Institutional Animal

Care and Use Committee at the Livestock Issues Research Unit (Protocol # 2014–04-JAC18).

Experimental Design

Newly weaned pigs ($n = 8$ gilts and 12 barrows; 6.2 ± 0.2 kg body weight) were obtained from a commercial swine facility and transported to the Livestock Issues Research Unit's Liberty Farm facility. Pigs were housed in an enclosed, environmentally-controlled facility in individual pens equipped with stainless steel feeders and nipple waterers. Pigs were allowed ad libitum access to a commercial non-medicated starter ration and water throughout the study. On d 12 post-weaning, pigs were anesthetized and a small incision (2 to 2.5 cm) was made in the lower abdominal region for the placement of an indwelling temperature recording device (25.4 mm in length, 8.3 mm in diameter, 3.3 g; Star Oddi DST micro-T; MeterMall USA, Marysville, OH) into the peritoneal cavity. Intraperitoneal temperature was measured at 5-min intervals from the time of the placement of the temperature device until the end of the study. On d 17, pigs were anesthetized and non-surgically fitted with indwelling jugular vein catheters (Carroll et al., 1999). On d 18, pigs were orally inoculated with 4.7×10^9 cfu/pig of a nalidixic acid-resistant *Salmonella typhimurium* in 10 mL of culture media. One 4.5-mL whole blood sample was collected in Sarstedt tubes containing no additive (Sarstedt Inc., Newton, NC) at 0.5 h intervals from –2 to 8 h, and at 8-h intervals from 8 to 72 h post-challenge. Samples were allowed to clot at room temperature for 30 min prior to centrifugation at 1,500 g for 20 min at 4°C. Isolated serum was stored at –80°C until analyzed for cortisol concentrations. A second 4-mL sample was collected in a vacutainer containing EDTA for determination of complete blood counts (CBC) using a ProCyt Dx Hematology Analyzer (IDEXX, Westbrook, ME). Sickness behavior scores were recorded for each pig following the collection of each blood sample. Additionally, individual pig feeders were weighed on d 0, 12, 18, and 22 to measure average daily feed disappearance. Fecal samples were collected on d 19 (0 h), 20 (24 h), 21 (48 h) and 22 (72 h) after the challenge to determine fecal shedding of the inoculated *Salmonella*. Following collection of the 72 h sample, pigs were humanely euthanized and samples of the cecum, mesenteric lymph node, subiliac lymph node, liver, spleen, gallbladder and kidney were collected to detect any *Salmonella* translocation into these tissues.

Sickness Behavior Scores

Sickness behavior scores were assigned to each pig by a single observer skilled at assessing sickness behavior in cattle and pigs. Pigs assigned a score of 1 exhibited normal maintenance behaviors. Pigs receiving a score of 2 were calm, but were lying down with increased respiration. A score of 3 was given to pigs exhibiting clinical signs of sickness including increased respiration with drool, scours, and(or) coughing. If a pig was observed to be lying on its side with labored breathing and frothing at the mouth it was given a score of 4, and then was humanely euthanized.

Cortisol Analysis

Serum cortisol concentrations were determined in duplicate using a commercially available, porcinespecific enzyme immunoassay kit (Abnova, Walnut, CA) according to the

manufacturer's directions by comparison of unknowns to standard curves generated with known concentrations of cortisol. The minimum detectable cortisol concentration was 0.2 ng/mL, and the intra- and inter-assay coefficients of variation were 10.1% and 8.2%, respectively.

Salmonella Isolation

Prior to experimental infection, fecal samples from all pigs were collected to confirm the absence of *Salmonella* or the presence of bacteria with similar morphology resistant to nalidixic acid. All fecal and tissue samples were aseptically collected, weighed (10 g), homogenized, and placed in a 1:10 dilution of phosphate buffered saline (PBS). Overnight cultures were serially diluted in PBS and plated on Luria-Bertani (LB) agar with 50µg/mL nalidixic acid and manually counted the following day after incubation for 18 to 24 h at 37°C. Isolates were confirmed as *Salmonella* via biochemical reactions on 4 bacterial test media: Triple Sugar Iron Slants, Indole Test, Lysine Iron Agar, and Urease Test. Negative cultures were enriched overnight in tetrathionate broth and re-plated.

Statistical Analysis

Prior to analysis, intraperitoneal temperature (TEMP) was averaged into 1-h intervals. Intraperitoneal temperature, serum cortisol, and CBC data were analyzed using the MIXED procedure of SAS specific for repeated measures (SAS Inst. Inc., Cary, NC). Sex, time and their interaction were included as fixed effects with pig within sex as the experimental unit. Specific treatment comparisons were made using the PDIF option in SAS, with $P < 0.05$ considered significant and $P < 0.10$ considered a tendency. *Salmonella* tissue and fecal count data were analyzed in SAS using Proc Glimmix of SAS at an $\alpha = 0.05$ utilizing the Tukey option for mean separation. Positive/negative binomial data to indicate the presence or absence of *Salmonella* in a tissue were analyzed using logistics regression in Proc Glimmix. All data are presented as the LSM \pm SEM.

RESULTS

There was a sex \times time interaction ($P < 0.001$) for TEMP. Specifically, gilts had greater TEMP than barrows from 36 to 63 h and at 72 h post-*Salmonella* challenge (Fig. 1). Additionally, there were also sex ($P < 0.001$) and time ($P < 0.001$) effects for TEMP. Intraperitoneal temperature was greater in gilts ($39.72 \pm 0.01^\circ\text{C}$) than barrows ($39.61 \pm 0.01^\circ\text{C}$). Intraperitoneal temperature generally increased from -144 h (insertion of temperature probes) to 0 h (immediately prior to oral *Salmonella* challenge; $P < 0.001$; -144 h vs. 0 h), and subsequently increased in response to the *Salmonella* challenge beginning at 12 h which persisted through the end of the study ($P < 0.02$).

Sickness behavior score was measured on pigs throughout the study. Interestingly, all pigs exhibited normal maintenance behaviors (i.e., a score of 1) throughout the entire study. Therefore, no differences in sickness behaviors were observed. However, differences in feed disappearance were observed during the study. Specifically, average daily feed disappearance increased from d 0 to 12 (0.21 ± 0.02 kg/d) to d 12 to 18 (0.51 ± 0.02 kg/d; $P < 0.001$), but decreased following *Salmonella* challenge to the end of the study (d 18 to 22;

0.38 ± 0.02; $P < 0.001$). There was no effect of sex ($P = 0.38$) or sex × time ($P = 0.75$) on average daily feed disappearance.

There was a sex × time interaction ($P = 0.03$) for serum cortisol concentrations. Specifically, cortisol concentrations were greater in barrows than gilts at 16 h (5.6 ± 0.4 vs. 3.8 ± 0.5 ng/mL; $P = 0.009$), yet cortisol was greater in gilts than barrows 32 h post *Salmonella* challenge (6.4 ± 0.5 vs. 2.9 ± 0.4 ng/mL; $P < 0.001$; Fig. 2). Overall, there was no effect of sex ($P = 0.16$) but there was an effect of time ($P < 0.01$) such that cortisol concentrations decreased below -2 h values at 3.5 h post-challenge ($P = 0.01$) and remained there before increasing above -2 h values at 16 h postchallenge ($P < 0.001$) and returning to baseline values by 64 h post-challenge ($P = 0.87$; -2 h vs. 64 h).

There was no effect of sex on red blood cells, hemoglobin, monocytes, or eosinophils (Table 1). There were sex ($P = 0.02$) and time ($P < 0.001$) effects for hematocrit such that barrows had greater hematocrit than gilts, and hematocrit decreased over time (Table 1). Also, similar results were found for platelet counts, such that barrows had greater platelets than gilts ($P < 0.001$), with counts fluctuating over time ($P = 0.003$; Table 1). Total white blood cells (WBC) were affected by sex ($P = 0.003$; Fig. 3). Specifically, barrows had greater total WBC counts than gilts. There was a tendency ($P = 0.06$) for a decrease in WBC following *Salmonella* challenge, and there was no sex × time interaction ($P = 0.49$). There was also a sex effect for neutrophils ($P < 0.001$) such that barrows had greater neutrophils than gilts (Fig. 4). Neutrophil counts decreased following *Salmonella* challenge ($P < 0.001$), but there was no sex × time interaction ($P = 0.55$). In contrast to total WBC and neutrophils, lymphocytes were greater in gilts than barrows ($P = 0.05$), yet there was no effect of time ($P = 0.27$) or sex × time interaction ($P = 0.71$; Fig. 5). There was a sex × time interaction ($P = 0.002$) for basophil counts, with gilts having greater basophils from 56 to 72 h ($P = 0.05$; Fig. 6). There was also an effect of time, with basophil counts increasing from 56 to 72 h post-*Salmonella* challenge, and there was a tendency ($P = 0.10$) for gilts to have greater basophil counts than barrows.

While many immune parameters were influenced by sex, there were no effects of sex ($P > 0.05$) on *Salmonella* concentrations from fecal shedding 4 d post-inoculation, or within the tissues of the cecum, mesenteric and subiliac lymph nodes, liver, spleen, gallbladder, or kidney tissues (Table 2). The only exception was fecal shedding at 48 h post-inoculation, which was greater in barrows than gilts ($P = 0.05$).

DISCUSSION

Sexual dimorphism has been reported in many animal species, including humans, cattle, dogs and pigs (Glucksmann, 1974; Williams et al., 2009; Carroll et al., 2015). Understanding the sexually dimorphic responses exhibited in livestock, such as pigs, is important as it may lead to sex-specific treatment or management to address the potential impacts of sexual dimorphism on health and subsequently growth. After thorough review of the literature, we believe this is the first study to demonstrate sexually dimorphic innate immune responses to an oral *Salmonella typhimurium* challenge in pigs.

Body temperature is an important physiological variable that can be used to detect an inflammatory response, and has been used by industry as an indicator of illness along with behavioral observations. Use of an intraperitoneal temperature device in the current study allowed for the measurement of body temperature at a constant interval throughout the study without having to handle the pigs, which can be stressful and further alter body temperature. Intraperitoneal temperature increased in response to the oral *Salmonella* challenge, but this increase was not observed until 12 h following the challenge. The temporal response to *Salmonella* is similar to what has been observed previously in pigs, with an increase in body temperature occurring approximately 12 to 24 h following inoculation (Balaji et al., 2000; Jenkins et al., 2004; Price et al., 2010). In contrast, there was no change in body temperature in finishing barrows nasally inoculated with *Salmonella* compared to non-challenged controls (Rostagno et al., 2011), which may be a result of route of administration or the sampling timeline. This appears to be the first study to report a sexually dimorphic temperature response to *Salmonella* challenge in pigs, with gilts producing a greater temperature response compared to barrows. Studies in cattle in response to LPS and CRH challenges have reported differences in rectal temperature responses between heifers and bulls, such that rectal temperature was greater in heifers than bulls (Hulbert et al., 2013; Carroll et al., 2015). Additionally, greater corticosterone responses to LPS have been reported in female mice (Spinedi et al., 1992). Changes in body temperature in response to an infectious agent are regulated by cytokines, including TNF- α , IL-6 and IL-1 β . The current study did not measure serum cytokine concentrations in response to *Salmonella* challenge. Although, a study by Williams et al. (2009) utilizing an LPS challenge in weaned pigs found greater peak TNF- α and IL-6 concentrations in gilts compared to barrows. Perhaps the greater peak concentrations of these 2 thermogenic cytokines observed in response to LPS partially explains the greater intraperitoneal temperature response to *Salmonella* challenge in the current study. However, further research is warranted to determine if sex alters *Salmonella*-induced cytokine concentrations in weaned pigs.

It is interesting to note that no differences were observed in sickness behavior. In fact, there were no changes in behavior outside of normal maintenance behaviors observed in any of the pigs throughout the post-challenge period. Typically, animal behavior is used by producers as an objective measurement when selecting animals for treatment of an illness or injury. These behaviors include decreased movement in the pen, decreased feeding and drinking behavior, and lethargy (Ahmed et al., 2015). Ahmed et al. (2015) observed decreases in pig movement and feeding and drinking behavior in response to a *Salmonella* challenge in growing pigs observed over a 4 wk period using video monitoring. While the current study observed a measureable decrease in average daily feed disappearance following the challenge, the decrease from pre-challenge to post-challenge was minimal (0.13 kg/d), which may not be significant enough to be detected by an observer. Yet, feed disappearance was only measured for a short period following *Salmonella* challenge, as pigs were euthanized at 72 h post-challenge, and sickness behavior was measured at specific time points throughout the study and did not utilize video surveillance. However, in the current study it could be suggested, based on the lack of sickness behaviors, that these pigs would not be selected for treatment based on periodically-observed behaviors alone, similar to that

performed in industry, regardless of the variables that indeed indicate an active infection (i.e., elevated body temperature and changes in blood cell parameters).

Cortisol concentrations varied between barrows and gilts at 2 time points in response to *Salmonella*, suggesting that barrows had an earlier peak in cortisol compared to gilts. However, overall there was no sex effect on serum cortisol concentrations. This is similar to what was observed by Williams et al. (2009) in response to an LPS challenge in pigs. A similar temporal response was observed in barrows challenged orally with *Salmonella* (Balaji et al., 2000). Studies in cattle did not find a difference in serum cortisol between heifers and bulls in response to an LPS challenge (Carroll et al., 2015), but a greater cortisol response was observed in response to a CRH challenge (Hulbert et al., 2013). The temporal pattern relative to the increase in cortisol concentration follows that of intraperitoneal temperature such that cortisol concentrations increased at 16 h post-challenge. Larzul et al. (2015) observed differences in cortisol binding globulin (CBG) capacity in pigs, with a correlation between basal cortisol concentrations and CBG in males but not in females. Additionally, female mammals have been noted to have greater adrenal gland weights (Glucksmann, 1974). However, the differences observed in the cortisol response to *Salmonella* challenge in the current study were minimal.

White blood cells are the cellular component of the immune response, and thus changes in these cell populations within the blood can suggest changes in the response to pathogens. In the current study, various differences in blood cell parameters were observed between gilts and barrows. This is in contrast to Williams et al. (2009) who found no differences in total white blood cell or differential counts in response to LPS between barrows and gilts. Perhaps the nature of the live challenge compared to a more provocative LPS challenge resulted in the differences observed between the 2 studies. Similar results have been observed in cattle, such that Brahman bulls had greater total leukocytes and neutrophils compared to heifers (Hulbert et al., 2013; Carroll et al., 2015). Females have been reported to have a heavier spleen and thymus (Glucksmann, 1974); however, this does not correspond with the decreased total white blood cells and neutrophils observed in gilts compared to barrows. It is possible that the decrease in neutrophils is a result of a greater number of neutrophils leaving the periphery and entering the gut. A study of neutrophil activity and adhesion molecules would help elucidate if this is indeed happening. A study in rats and mice found a greater neutrophil accumulation in tissues following ischemia/reperfusion, and was believed to be regulated by the chemokine Cxcl5, which was able to increase systemic neutrophils in the absence of ischemia/reperfusion (Madalli et al., 2015). Thus, perhaps differences in cytokine and chemokine concentrations are also driving the differences in various immune cell parameters.

Observing almost no differences in fecal shedding and tissue *Salmonella* content was surprising due to the differences observed in intraperitoneal temperature, serum cortisol, and the complete blood counts. It is possible that although there were differences in blood parameters, this did not influence colonization within the gastrointestinal system and did not influence migration of *Salmonella* from the gut to peripheral tissues. It is unclear why fecal shedding was influenced by sex 48 h following the challenge, but not at any other time point measured. Further, evidence of sexual dimorphism regarding fecal shedding of bacteria is

limited. Reports have indicated a greater incidence of bacterial infections in males compared to females (Green, 1992; Strachan et al., 2008). Yet, when experimentally introduced, it appears there is no difference between weaned gilts and barrows in the tissue *Salmonella* content.

There have been many studies in several species aimed at elucidating sexually dimorphic responses. From these studies it appears that females are more resilient to infection, yet are more prone to autoimmune diseases (Homodelarche et al., 1991; Spitzer, 1999). For example, female mice of reproductive age appear to have more active polymorphonuclear leukocytes than male counterparts (Spitzer, 1999). Sex hormones, estrogens and androgens, have been implicated in these effects. For example, Kahl and Elsasser (2006) reported greater serum TNF- α and serum amyloid A concentrations in steers administered testosterone 2 d prior to an LPS challenge. Additionally, heifers in estrus (i.e., low progesterone concentrations) had greater mean TNF- α concentrations after LPS administration compared to heifers in diestrus (i.e., high progesterone concentrations). In humans, Dosiou et al. (2004) reported a decrease in the Th1 response during the luteal phase of the menstrual cycle. However it is interesting to note that there are several studies in which sexually dimorphic effects were observed in pre-pubertal animals. Thus, the observed sexually dimorphic effects are likely initiated earlier in life, and likely *in utero* and/or in the early perinatal period. This is supported by research that reports a change in thymus and T lymphocyte development when female mice were administered testosterone early in life (Leposavic et al., 2009). Based on the findings within our current study, and that of prior reports, it is clear that pigs exhibit a sexually dimorphic immune response which warrants further study.

Conclusion

These data provide further proof that sex can influence the acute phase immune response to a provocative immune challenge. Specifically, gilts had a greater intraperitoneal temperature response, greater lymphocyte and basophil counts, yet decreased hematocrit, platelets, total white blood cell, and neutrophil counts than barrows. These data demonstrate that weaned gilts appear to produce a stronger acute phase response to a *Salmonella* challenge compared to barrows, without affecting sickness behavior or the tissue translocation or shedding of *Salmonella*. Based on the differences in cellular immune and serum variables, but the lack of differences in sickness behavior, managing gilts and barrows separately may have significant impacts on herd health management during the weaning/early post-weaning period.

Acknowledgments

The authors would like to acknowledge the technical assistance of J. W. Dailey and J. R. Carroll (USDA-ARS). Elena Gart was supported by a Texas A&M University Association of Former Students Diversity Fellowship. Laura Bryan was supported by NIH T32 grant 5T320D011083.

LITERATURE CITED

Ahmed ST, Mun HS, Yoe H, and Yang CJ. 2015 Monitoring of behavior using a video-recording system for recognition of *Salmonella* infection in experimentally infected growing pigs. *Animal* 9:115–121. doi:10.1017/S1751731114002213 [PubMed: 25166898]

- Balaji R, Wright KJ, Hill CM, Dritz SS, Knoppel EL, and Minton JE. 2000 Acute phase responses of pigs challenged orally with *Salmonella typhimurium*. *J. Anim. Sci* 78:1885–1891. doi: 10.2527/2000.7871885x [PubMed: 10907831]
- Burdick NC, Banta JP, Neuendorff DA, White JC, Vann RC, Laurenz JC, Welsh TH, and Randel RD. 2009 Interrelationships among growth, endocrine, immune, and temperament variables in neonatal Brahman calves. *J. Anim. Sci* 87:3202–3210. doi:10.2527/jas.2009-1931 [PubMed: 19542503]
- Carroll JA, Burdick NC, Reuter RR, Chase CC Jr., Spiers DE, Arthington JD, and Coleman SW. 2011 Differential acute phase immune responses by Angus and Romosinuano steers following an endotoxin challenge. *Domest. Anim. Endocrinol* 41:163–173. doi:10.1016/j.domaniend.2011.06.002 [PubMed: 21865005]
- Carroll JA, Burdick Sanchez NC, Hulbert LE, Ballou MA, Dailey JW, Caldwell LC, Vann RC, Welsh TH Jr., and Randel RD. 2015 Sexually dimorphic innate immunological responses of pre-pubertal Brahman cattle following an intravenous lipopolysaccharide challenge. *Vet. Immunol. Immunopathol* 166:108–115. doi:10.1016/j.vetimm.2015.06.009 [PubMed: 26144890]
- Carroll JA, Daniel JA, Keisler DH, and Matteri RL. 1999 Non-surgical catheterization of the jugular vein in young pigs. *Lab. Anim* 33:129–134. doi:10.1258/002367799780578345 [PubMed: 10780815]
- Dosiou C, Lathi RB, Tulac S, Huang STJ, and Giudice LC. 2004 Interferon-related and other immune genes are downregulated in peripheral blood leukocytes in the luteal phase of the menstrual cycle. *J. Clin. Endocrinol. Metab* 89:2501–2504. doi:10.1210/jc.2003-031647 [PubMed: 15126584]
- Glucksmann A 1974 Sexual Dimorphism in Mammals. *Biol. Rev. Camb. Philos. Soc* 49:423–475. doi: 10.1111/j.1469-185X.1974.tb01171.x [PubMed: 4616731]
- Green NS 1992 The Male Predominance in the Incidence of Infectious-Disease in Children: A Postulated Explanation for Disparities in the Literature. *Int. J. Epidemiol* 21:381–386. doi: 10.1093/ije/21.2.381 [PubMed: 1428496]
- Homo-Delarche F, Fitzpatrick F, Christeff N, Nunez EA, Bach JF, and Dardenne M. 1991 Sex Steroids, Glucocorticoids, Stress and Autoimmunity. *J. Steroid Biochem. Mol. Biol* 40:619–637. doi: 10.1016/0960-0760(91)90285-D [PubMed: 1958562]
- Hulbert LE, Carroll JA, Ballou MA, Burdick NC, Dailey JW, Caldwell LC, Loyd AN, Vann RC, Welsh TH Jr., and Randel RD. 2013 Sexually dimorphic stress and pro-inflammatory cytokine responses to an intravenous corticotropin-releasing hormone challenge of Brahman cattle following transportation. *Innate Immun.* 19:378–387. doi:10.1177/1753425912462752 [PubMed: 23112011]
- Jenkins NL, Turner JL, Dritz SS, Durham SK, and Minton JE. 2004 Changes in circulating insulin-like growth factor-I, insulin-like growth factor binding proteins, and leptin in weaned pigs infected with *Salmonella enterica* serovar Typhimurium. *Domest. Anim. Endocrinol* 26:49–60. doi: 10.1016/j.domaniend.2003.09.001 [PubMed: 14732452]
- Kahl S, and Elsasser TH. 2006 Exogenous testosterone modulates tumor necrosis factor-alpha and acute phase protein responses to repeated endotoxin challenge in steers. *Domest. Anim. Endocrinol* 31:301–311. doi:10.1016/j.domaniend.2005.11.005 [PubMed: 16386401]
- Larzul C, Terenina E, Foury A, Billon Y, Louveau I, Merlot E, and Mormede P. 2015 The cortisol response to ACTH in pigs, heritability and influence of corticosteroid-binding globulin. *Animal* 9:1929–1934. doi:10.1017/S1751731115001767 [PubMed: 26302113]
- Leposavi G, Perisic M, Kosec D, Arsenovic-Ranin N, Radojevic K, Stojic-Vukanic Z, and Pilipovic I. 2009 Neonatal testosterone imprinting affects thymus development and leads to phenotypic rejuvenation and masculinization of the peripheral blood T-cell compartment in adult female rats. *Brain Behav. Immun* 23:294–304. doi:10.1016/j.bbi.2008.11.002 [PubMed: 19028560]
- Madalli S, Beyrau M, Whiteford J, Duchene J, Nandhra IS, Patel NSA, Motwani MP, Gilroy DW, Thiernemann C, Nourshargh S, and Scotland RS. 2015 Sex-specific regulation of chemokine Cxcl5/6 controls neutrophil recruitment and tissue injury in acute inflammatory states. *Biol. Sex Differ* 6:27. doi:10.1186/s13293-015-0047-5 [PubMed: 26617980]
- Ojha S, and Kostrzynska M. 2007 Approaches for reducing *Salmonella* in pork production. *J. Food Prot* 70:2676–2694. [PubMed: 18044456]
- Price KL, Totty HR, Lee HB, Utt MD, Fitzner GE, Yoon I, Ponder MA, and Escobar J. 2010 Use of *Saccharomyces cerevisiae* fermentation product on growth performance and microbiota of weaned

pigs during Salmonella infection. *J. Anim. Sci* 88:3896–3908. doi:10.2527/jas.2009-2728 [PubMed: 20656973]

Rostagno MH, Eicher SD, and Lay DC. 2011 Immunological, Physiological, and Behavioral Effects of Salmonella enterica Carriage and Shedding in Experimentally Infected Finishing Pigs. *Food. Pathog. Dis* 8:623–630. doi:10.1089/fpd.2010.0735

Spinedi E, Suescun MO, Hadid R, Daneva T, and Gaillard RC. 1992 Effects of Gonadectomy and Sex-Hormone Therapy on the Endotoxin-Stimulated Hypothalamo-Pituitary-Adrenal Axis- Evidence for a Neuroendocrine-Immunological Sexual Dimorphism. *Endocrinology* 131:2430–2436. doi: 10.1210/endo.131.5.1330501 [PubMed: 1330501]

Spitzer JA 1999 Gender differences in some host defense mechanisms. *Lupus* 8:380–383. doi: 10.1177/096120339900800510 [PubMed: 10455517]

Strachan NJC, Watson RO, Novik V, Hofreuter D, Ogden ID, and Galan JE. 2008 Sexual dimorphism in campylobacteriosis. *Epidemiol. Infect* 136:1492–1495. doi:10.1017/S0950268807009934 [PubMed: 18062834]

Williams PN, Collier CT, Carroll JA, Welsh TH, and Laurenz JC. 2009 Temporal pattern and effect of sex on lipopolysaccharide-induced stress hormone and cytokine response in pigs. *Domest. Anim. Endocrinol* 37:139–147. doi:10.1016/j.domaniend.2009.04.004 [PubMed: 19523782]




Figure 1.

Effect of sex on the intraperitoneal temperature (TEMP) response to an oral *Salmonella* challenge (4.7×10^9 cfu/pig *Salmonella typhimurium*) in weaned pigs. Gilts $n = 8$, Barrows $n = 12$. Data presented as the LSM. SEM is ± 0.13 for Gilts and ± 0.10 for Barrows. There was a sex \times time interaction ($P < 0.001$) with gilts having greater TEMP response than barrows from 36 to 63 and at 72 h post-*Salmonella* challenge.




Figure 2. Effect of sex on the serum cortisol response to an oral *Salmonella* challenge (4.7×10^9 cfu/pig *Salmonella typhimurium*) in weaned pigs. Gilts $n = 8$, Barrows $n = 12$. Data presented as the LSM \pm SEM. Sex \times time interaction ($P = 0.03$). *Gilts and Barrows differ $P = 0.009$.




Figure 3. Effect of sex on the total white blood cell (WBC) response to an oral *Salmonella* challenge (4.7×10^9 cfu/pig *Salmonella typhimurium*) in weaned pigs. Gilts $n = 8$, Barrows $n = 12$. Data presented as the LSM \pm SEM. Barrows had greater WBC counts than gilts (sex: $P = 0.003$).




Figure 4. Effect of sex on the neutrophil response to an oral *Salmonella* challenge (4.7×10^9 cfu/pig *Salmonella typhimurium*) in weaned pigs. Gilts $n = 8$, Barrows $n = 12$. Data presented as the LSM \pm SEM. Barrows had greater neutrophils than gilts (sex: $P < 0.001$).




Figure 5. Effect of sex on the lymphocyte response to an oral *Salmonella* challenge (4.7×10^9 cfu/pig *Salmonella typhimurium*) in weaned pigs. Gilts $n = 8$, Barrows $n = 12$. Data presented as the LSM \pm SEM. Lymphocytes were greater in gilts than barrows (sex: $P = 0.05$).




Figure 6.

Effect of sex on the basophil response to an oral *Salmonella* challenge (4.7×10^9 cfu/pig *Salmonella typhimurium*) in weaned pigs. Gilts $n = 8$, Barrows $n = 12$. Data presented as the LSM \pm SEM. Sex \times time interaction ($P = 0.002$). *Gilts and Barrows differ $P < 0.05$.

Table 1. Summary of hematology variables measured in gilts and barrows in response to an oral *Salmonella typhimurium* challenge

Variable	Sex ¹				P-value		
	Gilts	Barrows	SEM	Sex	Time	Sex × time	
Red blood cells, 10 ⁶ /μL	4.83	4.82	0.07	0.95	< 0.001	1.00	
Hemoglobin, g/dL	8.22	8.35	0.06	0.11	< 0.001	0.99	
Hematocrit, %	26.91	27.58	0.23	0.02	< 0.001	0.99	
Platelets, 10 ³ /μL	402.44	446.11	6.41	< 0.001	0.003	0.99	
White blood cells, 10 ³ /μL	16.20	17.78	0.40	0.003	0.06	0.49	
Neutrophils, 10 ³ /μL	6.05	7.80	0.40	< 0.001	< 0.001	0.55	
Lymphocytes, 10 ³ /μL	9.62	9.03	0.24	0.05	0.27	0.71	
Monocytes, 10 ³ /μL	1.10	1.09	0.05	0.80	< 0.001	0.98	
Eosinophils, 10 ³ /μL	0.23	0.25	0.01	0.22	< 0.001	0.77	
Basophils, 10 ³ /μL	0.02	0.01	0.00	0.10	< 0.001	0.002	

¹Values for sex represent average values for each sex pooled over time.

Table 2.

Summary of fecal shedding and tissue *Salmonella* counts (\log_{10}) in response to an oral *Salmonella typhimurium* challenge in weaned pigs

Tissue	Sex ¹		SEM	Sex effect
	Gilts	Barrows		
Fecal Shedding 24 h	3.9	4.9	0.64	0.34
Fecal Shedding 48 h	4.8	6.0	0.35	0.05
Fecal Shedding 72 h	4.1	5.2	0.45	0.12
Cecum	4.7	4.2	0.26	0.33
Mesenteric lymph node	3.5	3.5	0.11	0.45
Subiliac lymph node	0.5	0.4	0.29	0.88
Liver	0.5	0.2	0.22	0.49
Spleen	0.6	0.4	0.25	0.60
Gallbladder	0.8	0.4	0.16	0.12
Kidney	0.3	0.2	0.18	0.78

¹Values for sex represent average *Salmonella* counts (\log_{10} ; log/g) for each sex.