



# NDH-1 Is Important for Photosystem I Function of *Synechocystis* sp. Strain PCC 6803 under Environmental Stress Conditions

Jiaohong Zhao<sup>1</sup>, Fudan Gao<sup>1</sup>, Da-Yong Fan<sup>2,3</sup>, Wah Soon Chow<sup>3</sup> and Weimin Ma<sup>1\*</sup>

<sup>1</sup> Department of Biology, College of Life and Environment Sciences, Shanghai Normal University, Shanghai, China, <sup>2</sup> State Key Laboratory of Vegetation and Environmental Change, Institute of Botany, Chinese Academy of Sciences, Beijing, China,

<sup>3</sup> Division of Plant Sciences, Research School of Biology, Australian National University, Canberra, ACT, Australia

## OPEN ACCESS

### Edited by:

Hongbo Gao,  
Beijing Forestry University, China

### Reviewed by:

Hualing Mi,  
Shanghai Institutes for Biological  
Sciences, Chinese Academy of  
Science, China  
Fang Huang,  
Key Laboratory of Photosynthesis and  
Environmental Molecular Physiology,  
Institute of Botany, (CAS), China  
Qiang Wang,  
Institute of Hydrobiology (CAS), China  
Stefano Santabarbara,  
Consiglio Nazionale Delle Ricerche  
(CNR), Italy

### \*Correspondence:

Weimin Ma  
wma@shnu.edu.cn

### Specialty section:

This article was submitted to  
Plant Physiology,  
a section of the journal  
Frontiers in Plant Science

**Received:** 12 May 2017

**Accepted:** 12 December 2017

**Published:** 17 January 2018

### Citation:

Zhao J, Gao F, Fan D-Y, Chow WS  
and Ma W (2018) NDH-1 Is Important  
for Photosystem I Function of  
*Synechocystis* sp. Strain PCC 6803  
under Environmental Stress  
Conditions. *Front. Plant Sci.* 8:2183.  
doi: 10.3389/fpls.2017.02183

Cyanobacterial NDH-1 interacts with photosystem I (PSI) to form an NDH-1-PSI supercomplex. Here, we observed that absence of NDH-1 had little, if any, effect on the functional fractions of PSI under growth conditions, but significantly reduced the functional fractions of PSI when cells of *Synechocystis* sp. strain PCC 6803 were moved to conditions of multiple stresses. The significant reduction in NDH-1-dependent functional fraction of PSI was initiated after PSII activity was impaired. This finding is consistent with our observation that the functional fraction of PSI under growth conditions was rapidly and significantly decreased with increasing concentrations of DCMU, which rapidly and significantly suppressed PSII activity by blocking the transfer of electrons from Q<sub>A</sub> to Q<sub>B</sub> in the PSII reaction center. Furthermore, absence of NDH-1 resulted in the PSI limitation at the functionality of PSI itself but not its donor-side and acceptor-side under conditions of multiple stresses. This was supported by the result of a significant destabilization of the PSI complex in the absence of NDH-1 but the presence of multiple stresses. Based on the above results, we propose that NDH-1 is important for PSI function of *Synechocystis* sp. strain PCC 6803 mainly via maintaining stabilization of PSI under conditions of environmental stresses.

**Keywords:** NDH-1, NDH-1-PSI, PSI function, environmental stress, *Synechocystis* sp. strain PCC 6803

## INTRODUCTION

Cyanobacterial NDH-1 is predominantly, if not totally, located in the thylakoid membrane (Ohkawa et al., 2001, 2002; Zhang et al., 2004; Xu et al., 2008; Battchikova et al., 2011), and participates in a variety of bioenergetic reactions, including respiration, cyclic electron transport around PSI, and CO<sub>2</sub> acquisition (Ogawa, 1991; Mi et al., 1992; Ohkawa et al., 2000). The function of NDH-1 is usually minor under normal growth conditions and becomes important under conditions of environmental stresses. In line with this, absence of NDH-1 has little, if any, effect on cell growth under normal conditions but retards cell growth and even causes cell death under conditions of environmental stresses, such as high light (Battchikova et al., 2011; Dai et al., 2013; Zhang et al., 2014; Zhao et al., 2014b, 2015; Gao et al., 2016a; Wang et al., 2016) and high temperature (Zhao et al., 2014a; Gao et al., 2016b). Therefore, cyanobacterial NDH-1 plays an important role in coping with various environmental stresses.

When cyanobacterial cells are transferred from normal growth conditions to multiple stressful environments, the amount of NDH-1 and its activity, for example, NDH-1-dependent cyclic electron transport around PSI (NDH-CET), are significantly increased under conditions of high temperature (Rowland et al., 2010; Zhao et al., 2014a), high salt (Hibino et al., 1996; Tanaka et al., 1997), high light (Mi et al., 2001), and low CO<sub>2</sub> (Deng et al., 2003). Such an increase is assumed to be important in repairing the photodamaged PSII, optimizing photosynthesis by increasing the proton gradient across the thylakoid membrane and supplying additional ATP (Allakhverdiev et al., 2005). However, the effect of cyanobacterial NDH-1 on PSI remains elusive.

Recently, cyanobacterial NDH-1 was found to interact with PSI to form an NDH-1-PSI supercomplex via CpcG2-phycobilisome, a PSI-specific antenna (Kondo et al., 2007; Gao et al., 2016a). Further, the supercomplex was found to be involved in NDH-CET but not respiration and CO<sub>2</sub> uptake (Gao et al., 2016a). Here, our results demonstrate that NDH-1 is important for PSI function of *Synechocystis* sp. strain PCC 6803 (hereafter *Synechocystis* 6803) under conditions of multiple stresses. We further found that NDH-1-dependent PSI function was initiated after PSII was impaired and its impairment is mostly the result of destabilization of PSI. The contribution of NDH-1-PSI supercomplex to PSI function is discussed.

## MATERIALS AND METHODS

### Culture Conditions

A glucose tolerant strain of wild-type (WT) *Synechocystis* 6803 and its mutants,  $\Delta ndhB$  (M55) (Ogawa, 1991), M55+NdhB and  $\Delta pgr5$  were cultured at 30°C in BG-11 medium (Allen, 1968) buffered with Tris-HCl (5 mM, pH 8.0) by bubbling with 2% (v/v) CO<sub>2</sub> in air. Continuous illumination was provided by fluorescence lamps at 40  $\mu\text{mol photons m}^{-2}\text{s}^{-1}$ . The mutant strains were grown in the presence of appropriate antibiotics.

### Construction of $\Delta pgr5$ Mutant

The  $\Delta pgr5$  mutant was constructed as follows. The upstream and downstream regions of *ssr2016* (*pgr5*) were amplified by PCR, creating appropriate restriction sites. A DNA fragment encoding a kanamycin resistance (Kam<sup>R</sup>) cassette was also amplified by PCR, creating *SalI* and *XbaI* sites using appropriate PCR primers, *pgr5-C* and *pgr5-D* (Supplementary Table S1). These three PCR products were ligated into the multiple cloning site of pUC19 (Figure 5A) and used to transform the WT cells of *Synechocystis* 6803 to generate the  $\Delta pgr5$  mutant. The transformants were spread on agar plates containing BG-11 medium and kanamycin (10  $\mu\text{g mL}^{-1}$ ) buffered at pH 8.0, and the plates were incubated in 2% (v/v) CO<sub>2</sub> in air under illumination by fluorescent lamps

**Abbreviations:** BN-PAGE, blue-native-PAGE; CBB, Coomassie Brilliant Blue; Chl, chlorophyll; DCMU, 3-(3,4-dichlorophenyl)-1,1-dimethylurea;  $F_v/F_m$ , maximal quantum yield of PSII; HL, high light; HS, high salt; M55,  $\Delta ndhB$ ; NDH-CET, NDH-1-dependent cyclic electron transport around PSI; MV, methyl viologen; PRG5, PROTON GRADIENT REGULATION 5;  $P_m$ , the maximal P700 change;  $qP$ , the ratio of the oxidized PQ pool; *Synechocystis* 6803, *Synechocystis* sp. strain PCC 6803; WT, wild-type.

at 40  $\mu\text{mol photons m}^{-2}\text{s}^{-1}$  as described previously (Williams and Szalay, 1983; Long et al., 2011). The mutated *pgr5* in the transformants was segregated to homogeneity (by successive streak purification) as determined by PCR amplification and reverse transcription (RT)-PCR analysis (Figures 5B,C).

### Construction of M55 Complementation Strain

The M55 complementation strain was constructed as follows. A DNA fragment containing the *ndhB* gene was amplified by PCR and then inserted into *NdeI* site of *PpsbAII* expression vector (Jiang et al., 2012) to form the *PpsbAII-ndhB* expression plasmid (Supplementary Figure S1A; primers are shown in Supplementary Table S1), which was used to transform the M55 mutant of *Synechocystis* 6803 to generate the M55 complementation strain. The transformants were spread on agar plates containing BG-11 medium, spectinomycin (10  $\mu\text{g mL}^{-1}$ ) and kanamycin (10  $\mu\text{g mL}^{-1}$ ) buffered at pH 8.0, and the plates were incubated in 2% (v/v) CO<sub>2</sub> in air under illumination by fluorescent lamps at 40  $\mu\text{mol photons m}^{-2}\text{s}^{-1}$  as described previously (Williams and Szalay, 1983; Long et al., 2011). Complete segregation of the transformants was confirmed by PCR (Supplementary Figure S1B) and the expression level of *ndhB* in the transformants was estimated by RT-PCR (Supplementary Figure S1C).

### RNA Extraction and RT-PCR Analysis

Total RNA was isolated and analyzed as described previously (McGinn et al., 2003). RT-PCR was performed using the Access RT-PCR system (Promega) to generate products corresponding to *pgr5* and *16S rRNA*, with 0.5  $\mu\text{g}$  of DNase-treated total RNA as starting material. RT-PCR conditions were 95°C for 5 min followed by cycles of 95, 62, and 72°C for 30 s each. The reactions were stopped after 25 cycles for *16S rRNA* and after 35 cycles for *ndhB* and *pgr5*. The primers used are summarized in Supplementary Table S1.

### Chlorophyll Fluorescence

The yield of chlorophyll (Chl) fluorescence at the steady state of electron transport was measured using a Dual-PAM-100 monitoring system (Walz, Effeltrich, Germany) equipped with an ED-101US/MD unit (Schreiber et al., 1986; Ma et al., 2008), as shown in Supplementary Figure S2. Minimal fluorescence corresponding to open PSII centers in the dark-adapted state ( $F_o$ ) and in the far-red (FR) light-adapted state ( $F_o'$ ) was excited by a weak measuring light (650 nm) at a PFD of 0.05–0.15  $\mu\text{mol photons m}^{-2}\text{s}^{-1}$ . A saturation pulse of red light (100 ms, 10,000  $\mu\text{mol photons m}^{-2}\text{s}^{-1}$ ) was applied to determine the maximal fluorescence at closed PSII centers in the dark-adapted state ( $F_m$ ) and in the red actinic light (AL)-adapted state ( $F_m'$ ) as described previously (Klughammer and Schreiber, 2008a). Subsequently, red AL was applied to monitor fluorescence under the steady-state condition ( $F_s$ ).  $F_v/F_m$  and  $qP$  were calculated as  $(F_m - F_o)/F_m$  (Kitajima and Butler, 1975) and  $(F_m' - F_s)/(F_m' - F_o')$  (van Kooten and Snel, 1990), respectively.

## The P700<sup>+</sup> Signal Monitored as an 830 nm Absorbance Change

With the Dual-PAM-100, P700<sup>+</sup> was monitored as the absorption difference between 830 and 875 nm in transmission mode. The quantum yields of PSI were determined using the saturation pulse method as described previously (Klughammer and Schreiber, 1994, 2008b; Supplementary Figure S3). The level of maximal P700<sup>+</sup> signal observed upon full oxidation,  $P_m$ , was determined by application of a saturation pulse of red light (100 ms; 10,000  $\mu\text{mol photons m}^{-2}\text{s}^{-1}$ ) in the presence of far-red light (about 0.3  $\mu\text{mol photons m}^{-2}\text{s}^{-1}$ ) at 720 nm.


## Photosynthetic Oxygen Evolution

Oxygen evolution by photosynthesis was measured at 30°C in the presence of 10 mM NaHCO<sub>3</sub> with a Clark-type oxygen electrode (Hansatech, Hertfordshire, UK)


according to the method described by Ma and Mi (2005). Cells were suspended in fresh BG-11 medium at a Chl *a* concentration of 5  $\mu\text{g mL}^{-1}$  and, different concentrations of 3-(3,4-dichlorophenyl)-1,1-dimethylurea (DCMU) were added to the cell suspension cultures prior to measurement. The intensity of AL used for the measurement of photosynthetic oxygen evolution was 800  $\mu\text{mol photons m}^{-2}\text{s}^{-1}$ .

## Isolation of Crude Thylakoid Membranes

Cell in cultured medium (800 mL) were harvested at the logarithmic phase of growth and washed twice by suspending in 50 mL of fresh BG-11 medium, and the thylakoid membranes were isolated according to Gombos et al. (1994) with some modifications as follows. The cells suspended in 5 mL of disruption buffer (10 mM HEPES-NaOH, 5 mM sodium phosphate, pH 7.5, 10 mM MgCl<sub>2</sub>, 10 mM NaCl, and 25% [*v/v*]



**FIGURE 1** | PSII activity and PSI functional fraction of WT (A,B) and M55 (C,D) under high temperature conditions. Cells were grown at 30°C for 24 h and were transferred to 45°C for different periods. Prior to the measurements, the Chl *a* concentration was adjusted to 20  $\mu\text{g mL}^{-1}$ . PSII activity and PSI functionality were determined by the  $F_v/F_m$  and  $P_m$  parameters, respectively, expressed as percentage of the WT (100%). The  $F_v/F_m$  and  $P_m$  values that correspond to 100% are 0.46  $\pm$  0.02 and 0.58  $\pm$  0.02, respectively. Values are means  $\pm$  SD ( $n = 5$ ).



**FIGURE 2** | PSII activity and PSI functional fraction of WT (A,B) and M55 (C,D) under conditions of high light and high salt. Cells were grown under normal salt conditions at 40  $\mu\text{mol photons m}^{-2}\text{s}^{-1}$  for 24 h and were transferred to 300  $\mu\text{mol photons m}^{-2}\text{s}^{-1}$  for 36 h or to 0.8 M NaCl for 12 h. Prior to the measurements, the Chl *a* concentration was adjusted to 20  $\mu\text{g mL}^{-1}$ . PSII activity and PSI functionality were determined by the  $F_v/F_m$  and  $P_m$  parameters, respectively, expressed as percentage of the WT (100%). The  $F_v/F_m$  and  $P_m$  values that correspond to 100% are shown in the legend of **Figure 1**. Values are means  $\pm$  SD ( $n = 5$ ). CK, control check; HL, high light; HS, high salt.

glycerol) were supplemented by zirconia/silica beads and broken by vortexing 20 times at the highest speed for 30 s at 4°C with 5 min of cooling on ice between the runs. The crude extract was centrifuged at 5,000×g for 5 min to remove the glass beads and unbroken cells. By further centrifugation at 20,000×g for 30 min, we obtained crude thylakoid membranes as precipitates.

## Electrophoresis

Blue-native (BN)-PAGE of *Synechocystis* 6803 membranes was performed as described previously (Kügler et al., 1997) with slight modifications (Battchikova et al., 2011; Dai et al., 2013; Zhang et al., 2014; Zhao et al., 2014a, 2015; Gao et al., 2016a,b; Wang et al., 2016). Isolated membranes were prepared for BN-PAGE as follows. Membranes were washed with 330 mM sorbitol, 50 mM Bis-Tris (pH 7.0), and 0.5 mM phenylmethylsulfonyl fluoride (PMSF; Sigma) and resuspended in 20% (w/v) glycerol, 25 mM Bis-Tris (pH 7.0), 10 mM MgCl<sub>2</sub>, 0.1 units of RNase-free DNase RQ1 (Promega) at a Chl *a* concentration of 0.25 mg mL<sup>-1</sup>, and 0.5 mM PMSF. The samples were incubated on ice for 10 min, and an equal volume of 3% DM was added. Solubilization was performed for 40 min on ice. Insoluble components were removed by centrifugation at 18,000×g for 15 min. The collected supernatant was mixed with one-tenth volume of sample buffer, 5% Serva Blue G, 100 mM Bis-Tris (pH 7.0), 30% (w/v) sucrose, 500 mM ε-amino-*n*-caproic acid, and 10 mM EDTA. Solubilized membranes were then applied to a 0.75-mm-thick, 5–12.5% acrylamide gradient gel (Hofer Mighty Small mini-vertical unit). Samples were loaded on an equal Chl *a* basis per lane. Electrophoresis was performed at 4°C by increasing the voltage gradually from 50 up to 200 V during the 5.5-h run.

## RESULTS

### NDH-1 Is Important for PSI Function under Conditions of Multiple Stresses

In *Synechocystis* 6803, NDH-1 interacts with PSI to form an NDH-1-PSI supercomplex (Gao et al., 2016a). To test whether the absence of NDH-1 affected the function of PSI under conditions of growth and environmental stress, we monitored the fractions of functional PSI when cells of WT and  $\Delta ndhB$  (M55), which lacks all functional NDH-1 complexes (Zhang et al., 2004), were incubated under high temperature conditions for different periods. The fractions of functional PSI of WT and M55 were similar under conditions of growth temperature (30°C), as deduced from the maximal P700 change ( $P_m$ ) level (0 h in **Figures 1B,D**). When cells were moved to conditions of high temperature (45°C) for different periods, however, we observed some expected and unexpected results. As expected, the activity of PSII in WT and M55 was gradually decreased (**Figures 1A,C**) whereas the fraction of functional PSI in WT was always maintained at a high level (**Figure 1B**) with the extension of heating time. Unexpectedly, the fraction of functional PSI in M55 was gradually and drastically decreased with the increase in heat incubation (**Figure 1D**).


We also measured the fractions of functional PSI upon exposure of WT and M55 cells to conditions of high light and high salt. As expected, the remaining functional fraction of PSI in

M55 was about 20% of that in the WT under high light and was about 40% of that in the WT under high salt (**Figures 2B,D**). It is worthy of note that these decreased fractions of functional PSI in M55 under conditions of heat, high light, and high salt could be recovered in its complementation strain M55+NdhB to the levels of WT (Supplementary Figures S1D–F). Taking all these results together, we can conclude that NDH-1 is important for the function of PSI under conditions of environmental stresses but not under growth conditions.



### NDH-1-Dependent PSI Function Is Initiated after PSII Activity Is Impaired

Under conditions of normal growth, WT and M55 cells had a similar activity of PSII (measured as the maximal quantum yield of PSII,  $F_v/F_m$ ) and had also a similar functional activity of PSI (0 h in **Figures 1A–D**). When cells of WT and M55 were transferred to conditions of multiple stresses and the activity of PSII was significantly suppressed in both strains, the functional fraction of PSI was significantly decreased in M55 mutant but not (**Figure 1B**) or to a lesser extent (**Figure 2B**) in the WT. It appears plausible that the decrease in PSII activity is a prerequisite for initiating the NDH-1-dependent PSI function.

To confirm this possibility, we measured the fractions of functional PSI in WT and M55 cells without or with different concentrations of DCMU, which blocks the transfer of electrons from  $Q_A$  to  $Q_B$  in the PSII reaction center (Trebst, 1980). As expected, with the increase in DCMU concentration, the activity of PSII (measured as oxygen evolution) was rapidly and similarly suppressed in WT and M55 (**Figure 3**). By contrast, the fractions of functional PSI were rapidly and significantly decreased in M55 but much less in the WT with the increase in DCMU concentration (**Figure 4A**), although both strains had a similar




**FIGURE 3** | Effects of DCMU treatment at different concentrations on the PSII activity in WT and M55 strains. PSII activity was assessed by the light-saturated photosynthetic oxygen evolution rate in the presence of DCMU and NaHCO<sub>3</sub>, expressed as percentage of the WT (100%). The rate of photosynthetic oxygen evolution that corresponds to 100% is 128.2 ± 8.6 µmol O<sub>2</sub> mg<sup>-1</sup> Chl *a* h<sup>-1</sup>. Values are means ± SD (*n* = 5).



function of PSI in the absence of DCMU ( $0\ \mu\text{M}$  in Figure 4A). Under conditions of DCMU addition, however, these changes in PSI function in the absence of NDH-1 were not observed in the absence of PROTON GRADIENT REGULATION 5 (PGR5) (Figures 5A–D), an alternative to NDH-1 in cyclic electron transport around PSI (Yeremenko et al., 2005). Based on the above results, we can conclude that NDH-1-dependent PSI function is specifically initiated after PSII activity is impaired but PGR5 is not involved in this process.

## Functionality of PSI Itself Is Impaired in the Background of NDH-1 Absence and Multiple Stresses

To test how NDH-1 affected PSI function under conditions of multiple stresses, we measured the PSI limitation at donor-side and acceptor-side as well as the functionality of PSI itself in WT and M55 strains. The reduced fractions of functional PSI under conditions of high temperature, high light and high salt were





**FIGURE 6** | PSI functional fractions of WT and M55 under multiple stresses with PSI electron donor and/or acceptor. Cells were grown under normal salt conditions at  $30^{\circ}\text{C}$  and  $40\ \mu\text{mol photons m}^{-2}\ \text{s}^{-1}$  for 24 h and were transferred to  $45^{\circ}\text{C}$  for 12 h,  $300\ \mu\text{mol photons m}^{-2}\ \text{s}^{-1}$  for 36 h, or to  $0.8\ \text{M NaCl}$  for 12 h. The Chl a concentration of WT and M55 cells was adjusted to  $20\ \mu\text{g mL}^{-1}$ . Prior to the  $P_m$  measurements,  $200\ \mu\text{M}$  2,6-dichlorophenol-indophenol and  $5\ \text{mM}$  sodium ascorbate and/or  $200\ \mu\text{M}$  methyl viologen were added to the cell suspension, as shown in (A–I). The functionality of PSI reaction centers was determined by the  $P_m$  parameter, expressed as a percentage of the WT (100%). The  $P_m$  value that corresponds to 100% is shown in the legend of Figure 1. Values are means  $\pm$  SD ( $n = 5$ ).

not recovered in M55 and were even slightly decreased by the addition of an exogenous PSI electron donor and/or acceptor to the cell suspension cultures of M55 (Figures 6A–I). Here, 2,6-dichlorophenolindophenol together with ascorbate (DCPIPH<sub>2</sub>) and methyl viologen (MV) were used as an exogenous electron donor (Mi et al., 1995) and acceptor (Takahashi and Katoh, 1984) for the PSI complex, respectively. Taking all these results together, we suggest that the impairment of PSI function in the absence of NDH-1 but the presence of multiple stresses is a result of impaired functionality of PSI itself.

DCMU addition and multiple stress treatments have a similar damage effect on PSI function in the absence of NDH-1 (see Figures 1, 2, 4) possibly because of suppressed

PSII activity (Figures 1–3). To see whether the impairment of PSI function in DCMU-treated M55 cells also resulted from impaired functionality of PSI itself, we monitored the fractions of functional PSI in DCMU-treated cells of WT and M55 with or without DCPIPH<sub>2</sub> and/or MV. Unexpectedly, the reduced fractions of functional PSI were completely recovered in M55 by the addition of DCPIPH<sub>2</sub> with or without MV to the cell suspension cultures of M55 (Figures 4B,D), although they were still not recovered in M55 by the addition of only MV to the cell suspension cultures of M55 (Figure 4C). Therefore, the impairment of PSI function in the absence of NDH-1 but the presence of DCMU is a result of donor-side limitation of PSI. Based on the above results, we suggest that DCMU addition and multiple stress treatments have a similar damage effect on PSI function in the absence of NDH-1 but their impaired targets are different.

To obtain insights into the reason why DCMU addition and multiple stress treatments have different impaired targets in PSI, we measured the redox state of plastoquinone (PQ) pool in WT and M55 grown under conditions of multiple stresses. The fluorescence parameter *qP* can reflect the redox state of the PQ pool and its decrease is closely linked with reduction of the PQ pool (Misumi et al., 2016). Under conditions of high temperature, high light or high salt, the reduction level of the PQ pool in M55 was higher than that in WT as deduced from the *qP* values (Figure 7, right), regardless of similar redox state of the PQ pool under conditions of growth temperature (Figure 7, left). In addition, it is known that DCMU addition results in oxidation of the PQ pool via blocking the transfer of electrons from Q<sub>A</sub> to Q<sub>B</sub> in PSII (Trebst, 1980) and in the presence of DCMU, the oxidation level of the PQ pool in M55 might be higher than that in WT, because of absence of NDH-1-dependent cyclic and respiratory flows in M55 (Mi et al., 1992). Taking all these results together, we propose that multiple stresses and DCMU addition



cause an opposite change in the redox state of the PQ pool, which may give a clue to the reason why they have different impaired targets in PSI.

## Absence of NDH-1 Destabilizes the PSI Complex under Multiple Stresses

To understand why the functionality of PSI itself is impaired under conditions of high temperature, high light and high salt, we examined various types of PSI complexes in WT and M55 cells grown under these stress conditions. Under normal growth conditions, the absence of NDH-1 in M55 resulted in a slight destabilization of the PSI complexes compared to the WT (see red box in **Figure 8**; left), just like the results of previous studies (Gao et al., 2016a). Such destabilization became more evident under conditions of multiple stresses, as reflected by the relative contents of the PSI trimer, dimer and monomer (**Figure 8**; left); these differences between WT and M55 were clearer after staining with Coomassie Brilliant Blue (**Figure 8**; right). These results strongly suggest that absence of NDH-1 destabilized the PSI complex, specifically under multiple stresses, and consequently impaired the functionality of PSI itself.

## DISCUSSION

Over the past decades, a significant achievement has been made in identifying the important role of cyanobacterial NDH-1 in protecting photosynthesis against environmental stresses, including protecting PSII and optimizing carbon assimilation via producing additional ATP and improving the ATP/NADPH ratio (Battchikova et al., 2011; Dai et al., 2013; Zhang et al., 2014; Zhao et al., 2014a,b, 2015; Gao et al., 2016a,b; Wang et al., 2016). The results of this study further indicate that cyanobacterial NDH-1 is important in maintaining the functionality of PSI itself (**Figure 6**), thereby guaranteeing a high level of PSI function under multiple stress conditions (**Figures 1, 2**). To our knowledge, this is the first study that reveals the contribution of NDH-1 to maintaining PSI functionality in cyanobacteria.

Recently, an NDH-1-PSI supercomplex was identified in the cyanobacterium *Synechocystis* 6803 (Gao et al., 2016a). We further found that absence of NDH-1 collapsed the supercomplex and resulted in a slight destabilization of the PSI complex, as deduced from the results of this study (**Figure 8**) and a previous study (Gao et al., 2016a) and that such destabilization became more evident under conditions of environmental stresses, as deduced from the results of this study (**Figure 8**) and a previous study (Zhao et al., 2014a). This finding may (1) explain why multiple stress treatments impairs the functionality of PSI itself, and (2) indicate that formation of the NDH-1-PSI supercomplex is important to keep a high level of PSI function under various stressful conditions.

Although the NDH-1-PSI supercomplex was also identified in angiosperms (Peng et al., 2008, 2009; Kouřil et al., 2014), the linker protein between NDH-1 and PSI as well as the component and function of NDH-1 were altered during evolution from cyanobacteria to higher plants. Stable formation of the supercomplex in higher plants needs two light-harvesting complex I (LHCI) proteins, Lhca5 and Lhca6, and in

cyanobacteria needs a PSI-specific peripheral antenna, CpcG2-phycobilisome, but their homologs lack in each other (Peng et al., 2009; Gao et al., 2016a). In addition, NDH-1 included in the NDH-1-PSI supercomplex of cyanobacteria and higher plants has a similar L-shaped skeleton. Despite their similarity, a large number of NDH-1 subunits in higher plants, including ferredoxin-binding subcomplex subunits NdhT and NdhU and all the subunits of subcomplex B and lumen subcomplex, are absent in the cyanobacterial NDH-1 (for review, see Ma and Ogawa, 2015).

It is known that the NDH-1-PSI supercomplex mainly participates in NDH-CET (Peng et al., 2008, 2012; Gao et al., 2016a). NDH-CET can produce ATP via building a proton gradient across the thylakoid membrane, which is important for running the Calvin-Benson cycle under conditions of environmental stress; consequently, the presence of NDH-1 keeps a high PSI functional activity. In Arabidopsis, NDH-CET had no effect on PSI function even under conditions of fluctuating light, regardless of an important protecting role of PGR5-CET in PSI function under fluctuating light (Suorsa et al., 2012; Kono et al., 2014). Unexpectedly, in rice, NDH-CET plays an important contribution on PSI function, although the contribution is still minor compared with the PGR5-CET (Yamori et al., 2016). What's even more amazing is that NDH-1 seems to aid the antimycin A-sensitive PGR5-CET in Arabidopsis (Kou et al., 2015). By contrast, in *Synechocystis* 6803, NDH-CET is found to be important for PSI function under conditions of multiple stresses (**Figures 1, 2**) but PGR5-CET had little, if any, effect on PSI function even in the presence of DCMU (**Figure 5**). Based on the above results, we propose that the role of NDH-1 on PSI function has been altered during evolution from cyanobacteria to higher plants.

## AUTHOR CONTRIBUTIONS

WM: designed and supervised the experiments; JZ and FG: performed the experiments and analyzed the data; D-YF, WC, and WM: analyzed and interpreted the data and wrote the article.

## ACKNOWLEDGMENTS

We thank Professors Teruo Ogawa (Nagoya University) and Bao-Sheng Qiu (Central China Normal University) for kindly providing the M55 mutant and the *PpsbAII* expression vector, respectively. This work was supported by the National Natural Science Foundation of China (grant nos. 31370270, 31570235, 31770259, and 31700205), China Postdoctoral Science Foundation (grant nos. 2015M581643 and 2017T100304), Shanghai Science and Technology Committee (grant no. 17070502900) and Project of Shanghai Normal University (grant no. SK201705).

## SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fpls.2017.02183/full#supplementary-material>



## REFERENCES

- Allakhverdiev, S. I., Nishiyama, Y., Takahashi, S., Miyairi, S., Suzuki, I., and Murata, N. (2005). Systematic analysis of the relation of electron transport and ATP synthesis to the photodamage and repair of photosystem II in *Synechocystis*. *Plant Physiol.* 137, 263–273. doi: 10.1104/pp.104.054478
- Allen, M. M. (1968). Simple conditions for growth of unicellular blue-green algae on plates. *J. Phycol.* 4, 1–4. doi: 10.1111/j.1529-8817.1968.tb04667.x
- Battchikova, N., Wei, L., Du, L., Bersanini, L., Aro, E. M., and Ma, W. (2011). Identification of novel Ssl0352 protein (NdhS), essential for efficient operation of cyclic electron transport around photosystem I, in NADPH: plastoquinone oxidoreductase (NDH-1) complexes of *Synechocystis* sp. PCC 6803. *J. Biol. Chem.* 286, 36992–37001. doi: 10.1074/jbc.M111.263780
- Dai, H., Zhang, L., Zhang, J., Mi, H., Ogawa, T., and Ma, W. (2013). Identification of a cyanobacterial CRR6 protein, Slr1097, required for efficient assembly of NDH-1 complexes in *Synechocystis* sp. PCC 6803. *Plant J.* 75, 858–866. doi: 10.1111/tj.12251
- Deng, Y., Ye, J., and Mi, H. (2003). Effects of low CO<sub>2</sub> on NAD(P)H dehydrogenase, a mediator of cyclic electron transport around photosystem I in the cyanobacterium *Synechocystis* PCC6803. *Plant Cell Physiol.* 44, 534–540. doi: 10.1093/pcp/pcg067
- Gao, F., Zhao, J., Chen, L., Battchikova, N., Ran, Z., Aro, E. M., et al. (2016a). The NDH-1L-PSI supercomplex is important for efficient cyclic electron transport in cyanobacteria. *Plant Physiol.* 172, 1451–1464. doi: 10.1104/pp.16.00585
- Gao, F., Zhao, J., Wang, X., Qin, S., Wei, L., and Ma, W. (2016b). NdhV is a subunit of NADPH dehydrogenase essential for cyclic electron transport in *Synechocystis* sp. strain PCC 6803. *Plant Physiol.* 170, 752–760. doi: 10.1104/pp.15.01430
- Gombos, Z., Wada, H., and Murata, N. (1994). The recovery of photosynthesis from low-temperature photoinhibition is accelerated by the unsaturation of membrane lipids: a mechanism of chilling tolerance. *Proc. Natl. Acad. Sci. U.S.A.* 91, 8787–8791. doi: 10.1073/pnas.91.19.8787
- Hibino, T., Lee, B. H., Rai, A., Ishikawa, H., Kojima, H., Tawada, M., et al. (1996). Salt enhances photosystem I content and cyclic electron flow via NAD(P)H dehydrogenase in the halotolerant cyanobacterium *Aphanothece halophytica*. *Aust. J. Plant Physiol.* 23, 321–330. doi: 10.1071/PP9960321
- Jiang, H. B., Lou, W. J., Du, H. Y., Price, N. M., and Qiu, B. S. (2012). Sll1263, a unique cation diffusion facilitator protein that promotes iron uptake in the cyanobacterium *Synechocystis* sp. strain PCC 6803. *Plant Cell Physiol.* 53, 1404–1417. doi: 10.1093/pcp/pcs086
- Kitajima, M., and Butler, W. (1975). Quenching of chlorophyll fluorescence and primary photochemistry in chloroplasts by dibromothymoquinone. *Biochim. Biophys. Acta* 376, 105–115. doi: 10.1016/0005-2728(75)90209-1
- Klughammer, C., and Schreiber, U. (1994). An improved method, using saturating light pulses, for the determination of photosystem I quantum yield via P700<sup>+</sup>-absorbance changes at 830 nm. *Planta* 192, 261–268. doi: 10.1007/BF01089043
- Klughammer, C., and Schreiber, U. (2008a). Complementary PS II quantum yields calculated from simple fluorescence parameters measured by PAM fluorometry and the Saturation Pulse method. *PAM Appl. Notes* 1, 27–35. Available online at: <http://walz.asia/downloads/pan/PAN078007.pdf>
- Klughammer, C., and Schreiber, U. (2008b). Saturation pulse method for assessment of energy conversion in PS I. *PAM Appl. Notes* 1, 11–14. Available online at: <http://www.photosynthesisystem.com/downloads/pan/PAN07002.pdf>
- Kondo, K., Ochiai, Y., Katayama, M., and Ikeuchi, M. (2007). The membrane-associated CpcG2-phycoobilisome in *Synechocystis*: a new photosystem I antenna. *Plant Physiol.* 144, 1200–1210. doi: 10.1104/pp.107.099267
- Kono, M., Noguchi, K., and Terashima, I. (2014). Roles of the cyclic electron flow around PSI (CEF-PSI) and O<sub>2</sub>-dependent alternative pathways in regulation of the photosynthetic electron flow in short-term fluctuating light in *Arabidopsis thaliana*. *Plant Cell Physiol.* 55, 990–1004. doi: 10.1093/pcp/pcu033
- Kou, J., Takahashi, S., Fan, D. Y., Badger, M. R., and Chow, W. S. (2015). Partially dissecting the steady-state electron fluxes in photosystem I in wild-type and *pgr5* and *ndh* mutants of *Arabidopsis*. *Front. Plant Sci.* 6:758. doi: 10.3389/fpls.2015.00758
- Kouřil, R., Strouhal, O., Nosek, L., Lenobel, R., Chamrád, I., Boekema, E. J., et al. (2014). Structural characterization of a plant photosystem I and NAD(P)H dehydrogenase supercomplex. *Plant J.* 77, 568–576. doi: 10.1111/tj.12402
- Kügler, M., Jansch, L., Kruft, V., Schmitz, U. K., and Braun, H.-P. (1997). Analysis of the chloroplast protein complexes by blue-native polyacrylamide gel electrophoresis (BN-PAGE). *Photosyn. Res.* 53, 35–44. doi: 10.1023/A:1005882406718
- Long, Z., Zhao, J., Zhang, J., Wei, L., Wang, Q., and Ma, W. (2011). Effects of different light treatments on the natural transformation of *Synechocystis* sp. strain PCC 6803. *Afr. J. Microbiol. Res.* 5, 3603–3610. doi: 10.5897/AJMR10.123
- Ma, W., and Mi, H. (2005). Expression and activity of type I NAD(P)H dehydrogenase at different growth phases of the cyanobacterium, *Synechocystis* PCC 6803. *Physiol. Plant.* 125, 135–140. doi: 10.1111/j.1399-3054.2005.00555.x
- Ma, W., and Ogawa, T. (2015). Oxygenic photosynthesis-specific subunits of cyanobacterial NADPH dehydrogenases. *IUBMB Life* 67, 3–8. doi: 10.1002/iub.1341
- Ma, W., Wei, L., and Wang, Q. (2008). The response of electron transport mediated by active NADPH dehydrogenase complexes to heat stress in the cyanobacterium *Synechocystis* 6803. *Sci. China C Life Sci.* 51, 1082–1087. doi: 10.1007/s11427-008-0139-0
- McGinn, P. J., Price, G. D., Maleszka, R., and Badger, M. R. (2003). Inorganic carbon limitation and light control the expression of transcripts related to the CO<sub>2</sub>-concentrating mechanism in the cyanobacterium *Synechocystis* sp. strain PCC6803. *Plant Physiol.* 132, 218–229. doi: 10.1104/pp.019349
- Mi, H., Deng, Y., Tanaka, Y., Hibino, T., and Takabe, T. (2001). Photo-induction of an NADPH dehydrogenase which functions as a mediator of electron transport to the intersystem chain in the cyanobacterium *Synechocystis* PCC6803. *Photosyn. Res.* 70, 167–173. doi: 10.1023/A:1017946524199
- Mi, H., Endo, T., Ogawa, T., and Asada, K. (1995). Thylakoid membrane-bound, NADPH-specific pyridine nucleotide dehydrogenase complex mediated cyclic electron transport in the cyanobacterium *Synechocystis* sp. PCC 6803. *Plant Cell Physiol.* 36, 661–668. doi: 10.1093/oxfordjournals.pcp.a078807
- Mi, H., Endo, T., Schreiber, U., Ogawa, T., and Asada, K. (1992). Electron donation from cyclic and respiratory flows to the photosynthetic intersystem chain is mediated by pyridine nucleotide dehydrogenase in the cyanobacterium *Synechocystis* PCC 6803. *Plant Cell Physiol.* 33, 1233–1237. doi: 10.1093/oxfordjournals.pcp.a078378
- Misumi, M., Katoh, H., Tomo, T., and Sonoike, K. (2016). Relationship between photochemical quenching and non-photochemical quenching in six species of cyanobacteria reveals species difference in redox state and species commonality in energy dissipation. *Plant Cell Physiol.* 57, 1510–1517. doi: 10.1093/pcp/pcv185
- Ogawa, T. (1991). A gene homologous to the subunit-2 gene of NADH dehydrogenase is essential to inorganic carbon transport of *Synechocystis* PCC6803. *Proc. Natl. Acad. Sci. U.S.A.* 88, 4275–4279. doi: 10.1073/pnas.88.10.4275
- Ohkawa, H., Pakrasi, H. B., and Ogawa, T. (2000). Two types of functionally distinct NAD(P)H dehydrogenases in *Synechocystis* sp. strain PCC6803. *J. Biol. Chem.* 275, 31630–31634. doi: 10.1074/jbc.M003706200
- Ohkawa, H., Sonoda, M., Hagino, N., Shibata, M., Pakrasi, H. B., and Ogawa, T. (2002). Functionally distinct NAD(P)H dehydrogenases and their membrane localization in *Synechocystis* sp. PCC6803. *Funct. Plant Biol.* 29, 195–200. doi: 10.1071/PP01180
- Ohkawa, H., Sonoda, M., Shibata, M., and Ogawa, T. (2001). Localization of NAD(P)H dehydrogenase in the cyanobacterium *Synechocystis* sp. strain PCC 6803. *J. Bacteriol.* 183, 4938–4939. doi: 10.1128/JB.183.16.4938-4939.2001
- Peng, L., Fukao, Y., Fujiwara, M., and Shikanai, T. (2012). Multistep assembly of chloroplast NADH dehydrogenase-like subcomplex A requires several nucleus-encoded proteins, including CRR41 and CRR42, in *Arabidopsis*. *Plant Cell* 24, 202–214. doi: 10.1105/tpc.111.090597
- Peng, L., Fukao, Y., Fujiwara, M., Takami, T., and Shikanai, T. (2009). Efficient operation of NAD(P)H dehydrogenase requires supercomplex formation with photosystem I via minor LHCl in *Arabidopsis*. *Plant Cell* 21, 3623–3640. doi: 10.1105/tpc.109.068791
- Peng, L., Shimizu, H., and Shikanai, T. (2008). The chloroplast NAD(P)H dehydrogenase complex interacts with photosystem I in *Arabidopsis*. *J. Biol. Chem.* 283, 34873–34879. doi: 10.1074/jbc.M803207200
- Rowland, J. G., Simon, W. J., Nishiyama, Y., and Slabas, A. R. (2010). Differential proteomic analysis using iTRAQ reveals changes in thylakoids associated with Photosystem II-acquired thermotolerance in *Synechocystis* sp. PCC 6803. *Proteomics* 10, 1917–1929. doi: 10.1002/pmic.200900337

- Schreiber, U., Schliwa, U., and Bilger, W. (1986). Continuous recording of photochemical and non-photochemical chlorophyll fluorescence quenching with a new type of modulation fluorometer. *Photosyn. Res.* 10, 51–62. doi: 10.1007/BF00024185
- Suorsa, M., Järvi, S., Grieco, M., Nurmi, M., Pietrzykowska, M., Rantala, M., et al. (2012). PROTON GRADIENT REGULATION5 is essential for proper acclimation of Arabidopsis photosystem I to naturally and artificially fluctuating light conditions. *Plant Cell* 24, 2934–2948. doi: 10.1105/tpc.112.097162
- Takahashi, Y., and Katoh, S. (1984). Triplet states in a photosystem I reaction center complex. Inhibition of radical pair recombination by bipyridinium dyes and naphthoquinones. *Plant Cell Physiol.* 25, 785–794. doi: 10.1093/oxfordjournals.pcp.a076773
- Tanaka, Y., Katada, S., Ishikawa, H., Ogawa, T., and Takabe, T. (1997). Electron flow from NAD(P)H dehydrogenase to photosystem I is required for adaptation to salt shock in the cyanobacterium *Synechocystis* sp. PCC 6803. *Plant Cell Physiol.* 38, 1311–1318. doi: 10.1093/oxfordjournals.pcp.a029123
- Trebst, A. (1980). Inhibitors in electron flow: tools for the functional and structural localization of carriers and energy conservation sites. *Methods Enzymol.* 69, 675–715. doi: 10.1016/S0076-6879(80)69067-3
- van Kooten, O., and Snel, J. F. (1990). The use of chlorophyll fluorescence nomenclature in plant stress physiology. *Photosyn. Res.* 25, 147–150. doi: 10.1007/BF00033156
- Wang, X., Gao, F., Zhang, J., Zhao, J., Ogawa, T., and Ma, W. (2016). A cytoplasmic protein Ssl3829 is important for NDH-1 hydrophilic arm assembly in *Synechocystis* sp. strain PCC 6803. *Plant Physiol.* 171, 864–877. doi: 10.1104/pp.15.01796
- Williams, J. G., and Szalay, A. A. (1983). Stable integration of foreign DNA into the chromosome of the cyanobacterium *Synechococcus* R2. *Gene* 24, 37–51. doi: 10.1016/0378-1119(83)90129-4
- Xu, M., Ogawa, T., Pakrasi, H. B., and Mi, H. (2008). Identification and localization of the CupB protein involved in constitutive CO<sub>2</sub> uptake in the cyanobacterium, *Synechocystis* sp. strain PCC 6803. *Plant Cell Physiol.* 49, 994–997. doi: 10.1093/pcp/pcn074
- Yamori, W., Makino, A., and Shikanai, T. (2016). A physiological role of cyclic electron transport around photosystem I in sustaining photosynthesis under fluctuating light in rice. *Sci. Rep.* 6:20147. doi: 10.1038/srep20147
- Yeremenko, N., Jeanjean, R., Prommeenate, P., Krasikov, V., Nixon, P. J., Vermaas, W. F., et al. (2005). Open reading frame *ssr2016* is required for antimycin A-sensitive photosystem I-driven cyclic electron flow in the cyanobacterium *Synechocystis* sp. PCC 6803. *Plant Cell Physiol.* 46, 1433–1436. doi: 10.1093/pcp/pci147
- Zhang, J., Gao, F., Zhao, J., Ogawa, T., Wang, Q., and Ma, W. (2014). NdhP is an exclusive subunit of large complex of NADPH dehydrogenase essential to stabilize the complex in *Synechocystis* sp. strain PCC 6803. *J. Biol. Chem.* 289, 18770–18781. doi: 10.1074/jbc.M114.553404
- Zhang, P., Battchikova, N., Jansen, T., Appel, J., Ogawa, T., and Aro, E.-M. (2004). Expression and functional roles of the two distinct NDH-1 complexes and the carbon acquisition complex NdhD3/NdhF3/CupA/Sll1735 in *Synechocystis* sp. PCC 6803. *Plant Cell* 16, 3326–3340. doi: 10.1105/tpc.104.026526
- Zhao, J., Gao, F., Qiu, Z., Wang, Q., and Ma, W. (2014a). Deletion of an electron donor-binding subunit of the NDH-1 complex, NdhS, results in a heat-sensitive growth phenotype in *Synechocystis* sp. PCC 6803. *Chin. Sci. Bull.* 59, 4484–4490. doi: 10.1007/s11434-014-0596-8
- Zhao, J., Gao, F., Zhang, J., Ogawa, T., and Ma, W. (2014b). NdhO, a subunit of NADPH dehydrogenase, destabilizes medium size complex of the enzyme in *Synechocystis* sp. Strain PCC 6803. *J. Biol. Chem.* 289, 26669–26676. doi: 10.1074/jbc.M114.553925
- Zhao, J., Rong, W., Gao, F., Ogawa, T., and Ma, W. (2015). Subunit Q is required to stabilize the large complex of NADPH dehydrogenase in *Synechocystis* sp. strain PCC 6803. *Plant Physiol.* 168, 443–451. doi: 10.1104/pp.15.00503

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Zhao, Gao, Fan, Chow and Ma. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.