

Characterizing circumgalactic gas around massive ellipticals at $z \sim 0.4$ – II. Physical properties and elemental abundances

Fakhri S. Zahedy¹,^{1*} Hsiao-Wen Chen¹,¹ Sean D. Johnson^{2,3},^{3†} Rebecca M. Pierce^{1,4}, Michael Rauch³, Yun-Hsin Huang⁵, Benjamin J. Weiner⁵ and Jean-René Gauthier⁶

¹Department of Astronomy and Astrophysics, The University of Chicago, Chicago, IL 60637, USA

²Department of Astrophysical Sciences, Princeton University, Princeton, NJ 08544, USA

³The Observatories of the Carnegie Institution for Science, 813 Santa Barbara Street, Pasadena, CA 91101, USA

⁴Department of Aerospace Engineering, University of Maryland, MD 20742, USA

⁵Steward Observatory, University of Arizona, Tucson, AZ 85721, USA

⁶Oracle Corporation, Culver City, CA 90230, USA

Accepted 2018 December 19. Received 2018 December 17; in original form 2018 September 6

ABSTRACT

We present a systematic investigation of the circumgalactic medium (CGM) within projected distances $d < 160$ kpc of luminous red galaxies (LRGs). The sample comprises 16 intermediate-redshift ($z = 0.21$ – 0.55) LRGs of stellar mass $M_{\text{star}} > 10^{11} M_{\odot}$. Combining far-ultraviolet Cosmic Origin Spectrograph spectra from the *Hubble Space Telescope* and optical echelle spectra from the ground enables a detailed ionization analysis based on resolved component structures of a suite of absorption transitions, including the full H I Lyman series and various ionic metal transitions. By comparing the relative abundances of different ions in individually matched components, we show that cool gas ($T \sim 10^4$ K) density and metallicity can vary by more than a factor of 10 in an LRG halo. Specifically, metal-poor absorbing components with $< 1/10$ solar metallicity are seen in 50 per cent of the LRG haloes, while gas with solar and super-solar metallicity is also common. These results indicate a complex multiphase structure and poor chemical mixing in these quiescent haloes. We calculate the total surface mass density of cool gas, Σ_{cool} , by applying the estimated ionization fraction corrections to the observed H I column densities. The radial profile of Σ_{cool} is best described by a projected Einasto profile of slope $\alpha = 1$ and scale radius $r_s = 48$ kpc. We find that typical LRGs at $z \sim 0.4$ contain cool gas mass of $M_{\text{cool}} = (1 - 2) \times 10^{10} M_{\odot}$ at $d < 160$ kpc (or as much as $M_{\text{cool}} \approx 4 \times 10^{10} M_{\odot}$ at $d < 500$ kpc), comparable to the cool CGM mass of star-forming galaxies. Furthermore, we show that high-ionization O VI and low-ionization absorption species exhibit distinct velocity profiles, highlighting their different physical origins. We discuss the implications of our findings for the origin and fate of cool gas in LRG haloes.

Key words: surveys – galaxies: elliptical and lenticular, cD – galaxies: formation – galaxies: haloes – intergalactic medium – quasars: absorption lines.

1 INTRODUCTION

Substantial efforts have been made in the last two decades to identify and characterize the physical processes which are at play in the gaseous halo surrounding galaxies, known as the circumgalactic medium (CGM; see recent reviews by Chen 2017; Tumlinson, Peebles & Werk 2017, and references therein). The CGM is situated

between the intergalactic medium (IGM), where most baryons in the Universe reside, and galaxies, where star formation occurs and heavy metals are synthesized. This unique characteristic makes the CGM a prime location to investigate the intricate interplay between gas accretion from the IGM and feedback processes originating in galaxies, in order to understand the baryon cycles that regulate galaxy evolution over cosmic time.

Some of the major unanswered questions in the study of galaxy evolution concern the origin and nature of cool ($T \sim 10^{4-5}$ K) gas in and around massive quiescent galaxies. Among the most massive galaxies in the Universe, they consist of predominantly

* E-mail: fsz@uchicago.edu

† Hubble & Carnegie-Princeton Fellow

old ($\gtrsim 1$ Gyr) stars and do not show any recent star formation (e.g. Eisenstein et al. 2003; Roseboom et al. 2006; Gauthier & Chen 2011). While it is tempting to attribute the ‘red and dead’ nature of quiescent galaxies as due to the absence of cool gas needed to fuel star formation, successive QSO absorption-line studies probing the CGM of luminous red galaxies (LRGs) have established that a significant fraction of these $z \sim 0.5$ massive elliptical galaxies host chemically enriched cool gas (e.g. Gauthier, Chen & Tinker 2009, 2010; Lundgren et al. 2009; Bowen & Chelouche 2011; Gauthier & Chen 2011; Thom et al. 2012; Zhu et al. 2014; Huang et al. 2016; Chen et al. 2018a). These findings at intermediate redshifts are consistent with observations in the local Universe, where HI and CO surveys found that at least a third of nearby ellipticals contain abundant atomic or even molecular gas (e.g. Serra et al. 2012; Young et al. 2014; 2018). The high incidence of cool gas in massive quiescent haloes is puzzling, and it presents a challenge to our current understanding of galaxy formation.

First, how does cool gas survive in massive haloes? The strong clustering of LRGs indicates that these galaxies reside inside massive dark-matter haloes with $M_h \gtrsim 10^{13} M_\odot$, where gas accreted from the IGM is expected to be shock heated to the virial temperature of the halo, $T \sim 10^{6.5-7}$ K (see Faucher-Giguère 2017 for a recent review). Recent cosmological simulations predict that massive galaxies at high redshifts can still acquire cool gas via dense and narrow filaments that penetrate deep into the halo (e.g. Dekel, Sari & Ceverino 2009; Kereš et al. 2009; van de Voort et al. 2012; Nelson et al. 2013; Shen et al. 2013), but they also show that this mechanism may not be effective for massive dark-matter haloes hosting LRGs at $z < 1$ (e.g. Kereš et al. 2009). Alternatively, thermal instabilities may cause cool clumps to condense from the hot halo and fall towards the galaxy (e.g. Mo & Miralda-Escudé 1996; Maller & Bullock 2004; Sharma et al. 2012; Voit et al. 2015). Although some observational results suggest this mechanism as a promising explanation (Huang et al. 2016), infalling cool clumps of gas are subject to disruption from ram pressure drag and thermal conduction with the hot medium. For that reason, it is still unclear whether cool clumps in the gaseous halo of LRGs will survive their journey to the centre of the halo. The detection of high-column density cool gas within projected distances $d < 10$ kpc from $z \sim 0.5$ massive quiescent galaxies (Zahedy et al. 2016; 2017b) indicates that some cool gas may survive, but to address this question quantitatively requires knowledge of the density and size distributions of cool clumps in LRG haloes.

Secondly, what are the dominant feedback mechanisms in massive quiescent haloes? The quiescent nature of both local and intermediate-redshift massive ellipticals indicates that some form of energetic feedback is effective at preventing the cooling of the hot halo over cosmic time, which would otherwise trigger continuing star formation. At the same time, the absence of young stellar populations and strong active galactic nuclei (AGNs) in typical LRGs (e.g. Roseboom et al. 2006; Sadler et al. 2007; Hodge et al. 2009; Gauthier & Chen 2011; Huang et al. 2016) makes it difficult to invoke starburst-driven outflows or AGN feedback to explain the high incidence of chemically enriched cool gas in and around massive quiescent galaxies. On the other hand, recent observational and theoretical studies have emphasized the importance of the old stellar population themselves in providing the necessary heating, through energy injection from Type Ia supernovae (SNe Ia) and/or winds from asymptotic giant branch (AGB) stars (e.g. Conroy, van Dokkum & Kravtsov 2015; Zahedy et al. 2016, 2017a; Li et al. 2018, and references therein). Further insights into the dom-

inant feedback mechanisms in LRGs can be obtained by directly comparing observations in the CGM with theoretical predictions for different feedback prescriptions. Doing so requires knowledge of the ionization states and chemical abundances in the CGM of LRGs.

A systematic study is necessary to characterize the physical properties and chemical abundances in the CGM of LRGs. This is a primary motivation behind our COS-LRG survey, a comprehensive survey of the gaseous haloes of 16 LRGs at $z \sim 0.4$ using a combination of far-ultraviolet (FUV) spectra from the Cosmic Origins Spectrograph (COS) on board the *Hubble Space Telescope* (*HST*) and ground-based optical echelle spectra. The COS-LRG sample was selected without any prior knowledge of the absorption properties of the LRGs. In Chen et al. (2018a, hereafter Paper I), we presented the initial results of our study, which we summarize here. First, high HI column density gas is common in the CGM of LRGs, with a median of $\langle \log N(\text{HI})/\text{cm}^{-2} \rangle = 16.6$ at $d < 160$ kpc. Secondly, we measured a high covering fraction of optically thick gas ($\log N(\text{HI})/\text{cm}^{-2} \gtrsim 17.2$) of $\langle \kappa \rangle_{\text{LLS}} = 0.44_{-0.11}^{+0.12}$ at $d < 160$ kpc, which increases to $\langle \kappa \rangle_{\text{LLS}} = 0.71_{-0.20}^{+0.11}$ at $d < 100$ kpc. Moreover, the CGM of LRGs contains widespread chemically enriched gas traced by low-, intermediate-, and high-ionization metals. The most prominent metal transitions in LRG haloes are those of intermediate-ionization species such as C III and Si III, with a high covering fraction of $\langle \kappa(\text{C III}) \rangle = 0.75_{-0.13}^{+0.08}$ within $d < 160$ kpc, comparable to what have been observed in the CGM of star-forming galaxies (e.g. Werk et al. 2013). In this paper, we expand our investigation with absorption-line and ionization analyses of both metal and HI absorption in LRG haloes, in order to characterize the physical properties and elemental abundances in the CGM of LRGs.

The paper is organized as follows. In Section 2, we discuss the COS-LRG sample and the spectroscopic observations and data reduction of the background QSOs. We describe the absorption-line and ionization analyses in Section 3. In Section 4, we characterize the physical properties and elemental abundances in the gaseous haloes of LRGs. Finally, we discuss the implications of our findings in Section 5 and present a summary of our results/conclusions in Section 6. In addition, we discuss the results of the analysis for each individual LRG halo in Appendix A. A standard Λ cosmology is adopted throughout the paper, with $\Omega_M = 0.3$, $\Omega_\Lambda = 0.7$, and a Hubble constant of $H_0 = 70 \text{ km. s}^{-1} \text{ Mpc}^{-1}$.

2 SAMPLE AND DATA

In this section, we summarize the COS-LRG sample and the observations of background QSOs. We refer the readers to Paper I for a more detailed discussion on the program design, sample selection, and data reduction of the FUV COS spectra and optical echelle spectra of background QSOs in our sample.

The COS-LRG sample was established by cross-correlating spectroscopically identified LRGs in the Sloan Digital Sky Survey (SDSS; York et al. 2000) archive and the literature with all known UV-bright QSOs with FUV $\lesssim 18.5$ mag.¹ No prior knowledge of the absorption properties of the LRGs was used in selecting all the LRG-QSO pairs that make up our sample. The UV magnitude cut was

¹Because the FUV bandpass of GALEX has a minimum wavelength of $\approx 1350 \text{ \AA}$, our FUV-bright selection for the background QSOs does not bias against optically thick Lyman-limit systems at $z \lesssim 0.5$, which is coincident with the redshift range of COS-LRG galaxies.

chosen to ensure that high-quality and high-resolution spectra of the background QSOs could be obtained with the COS (Green et al. 2012) onboard the *HST*. Furthermore, we imposed a lower limit on the LRG stellar mass of $\log M_{\text{star}}/M_{\odot} > 11$, and a maximum projected distance of $d = 160$ kpc from the QSO. Both choices were informed by the well-known finding of a significant incidence (>10 per cent) of Mg II absorbers at $d < 120$ kpc from massive LRGs (e.g. Gauthier et al. 2010; Huang et al. 2016). These selection criteria resulted in a mass-limited sample of 16 quiescent galaxies at $0.21 < z < 0.56$, each probed by a background QSO at $d < 160$ kpc (which corresponds to roughly $1/3$ of the virial radius, R_h , of a $10^{13} M_{\odot}$ dark-matter halo).

All 16 QSOs in the COS-LRG sample were observed with COS, either during our own observing program (PID: 14145) or previously available from the *HST* data archive. *HST*/COS with the G130M and G160M gratings provide high-resolution (FWHM ≈ 17 km s $^{-1}$) FUV spectra of the QSOs over a nearly contiguous wavelength coverage between $\lambda \approx 1150$ and ≈ 1780 Å, allowing us to probe halo gas using observations of the full H I Lyman series and corresponding low-, intermediate-, and high-ionization metal absorption features at the LRG redshift, including C III $\lambda 977$, the O VI $\lambda\lambda 1031, 1037$ doublet, Si III $\lambda 1206$, and Si II $\lambda 1260$. The COS data were downloaded from the *HST* archive and processed using our custom software. The data reduction steps were previously described in detail in Paper I. To summarize, an important aspect of our custom data reduction software is a recalibration of the COS wavelength solution, which was done in two steps. First, relative wavelength offsets between different exposures of the same QSO were corrected using a low-order polynomial that best describes the offsets of common narrow absorption features found in different exposures. Next, different exposures were co-added and an absolute wavelength correction was performed on the combined spectrum by registering non-saturated, low-ionization Galactic absorption lines to their known vacuum wavelengths. The final wavelength solution for our FUV COS spectra is accurate to within ± 3 km s $^{-1}$, based on a comparison with low-ionization absorption features seen in the ground-based optical echelle spectra.

Optical echelle spectra of COS-LRG QSOs are available for 11 out of the 16 QSOs in the sample. The echelle observations were obtained using two high-resolution spectrographs, MIKE (Bernstein et al. 2003) on the Magellan Clay telescope and HIRES (Vogt et al. 1994) on the Keck I telescope. The MIKE observations were obtained during our own observing program, whereas the HIRES data were retrieved from the Keck Observatory Archive (KOA). The instrumental configuration chosen for our MIKE observations provides a spectral resolution of FWHM ≈ 10 km s $^{-1}$ at wavelength $\lambda < 5100$ Å. The archival HIRES observations are characterized by a spectral resolution of FWHM ≈ 6.5 km s $^{-1}$ at $\lambda < 5900$ Å. By extending the spectral coverage of the COS-LRG QSOs to optical wavelengths (from $\lambda \sim 3100$ Å to well over ~ 5000 Å), the echelle spectra of the QSOs allow access to additional prominent absorption features arising in low-ionization gas in LRG haloes, especially the Mg II $\lambda\lambda 2796, 2803$ doublet, the Mg I $\lambda 2852$ transition, and a series of Fe II transitions including Fe II $\lambda 2586$ and Fe II $\lambda 2600$. A detailed description of the data reduction for the MIKE and HIRES spectra can be found in Paper I.

A summary of FUV and optical echelle spectroscopic observations is presented in Table 1, where we list for each background QSO the instrument used for the observations, the spectral coverage of the data, and the mean signal-to-noise ratio (S/N) per resolution element in final reduced spectrum.

Table 1. Summary of spectroscopy of background QSOs.

QSO	Instrument	Spectral window (Å)	S/N	Notes
SDSS J0246–0059	COS	1140–1790	10	PID: 14145
	MIKE	3350–9400	41	
SDSS J0803+4332	COS	1160–1800	9	PID: 11598
	HIRES	3150–5870	24	
SDSS J0910+1014	COS	1140–1790	7	PID: 11598
	HIRES	3150–5870	15	
SDSS J0925+4004	COS	1160–1800	6	PID: 11598
	HIRES	3240–5870	16	
SDSS J0946+5123	COS	1140–1780	7	PID: 14145
SDSS J0950+4831	COS	1070–1800	10	PID: 11598 & 13033
	HIRES	3100–5870	30	
SDSS J1111+5547	COS	1140–1800	15	PID: 12025
SDSS J1127+1154	COS	1140–1780	8	PID: 14145
	MIKE	3350–9400	17	
SDSS J1243+3539	COS	1140–1780	14	PID: 14145
SDSS J1244+1721	COS	1420–1780	7	PID: 12466
	MIKE	3350–9400	33	
SDSS J1259+4130	COS	1120–1790	13	PID: 13833
SDSS J1357+0435	COS	1130–1800	13	PID: 12264
	MIKE	3350–9400	25	
SDSS J1406+2509	COS	1140–1780	6	PID: 14145
	MIKE	3350–9400	10	
SDSS J1413+0920	COS	1130–1750	17	PID: 13833
SDSS J1550+4001	COS	1140–1790	8	PID: 11598
	HIRES	3100–5870	31	
SDSS J1553+3548	COS	1140–1790	8	PID: 11598
	HIRES	3100–5870	36	

3 ANALYSIS

To promote a deeper understanding of the circumgalactic environment of massive haloes, we assembled a mass-limited sample of 16 LRGs with $\log M_{\text{star}}/M_{\odot} > 11$. The LRGs were selected without prior knowledge of the presence or absence of CGM absorption features. This uniform sample of galaxies allows an unbiased and accurate characterization of the gaseous halo of intermediate-redshift, massive elliptical galaxies. The two main objectives of the COS-LRG program are: (1) to probe the bulk of cool gas in LRG haloes by obtaining accurate measurements of $N(\text{H I})$; and (2) to constrain the physical properties and elemental abundances in massive quiescent haloes by observing different ionic metal transitions that probe a wide range of ionization states.

In Paper I, we presented $N(\text{H I})$ measurements for the sample and reported significant incidences (>40 per cent) of low-, intermediate-, and high-ionization metal absorptions at $d < 160$ kpc in massive quiescent haloes. To investigate the physical properties of the CGM of LRGs and constrain the chemical abundance of the gas requires (1) accurate column density measurements for the observed metal absorption features, and (2) a detailed ionization modelling of the gas under different physical conditions (e.g. density and metallicity) to explain the observations. Here, we describe the analysis to first measure the ionic column densities and subsequently constrain the physical properties and metallicities of the gas.

3.1 Voigt profile analysis

The available high-resolution FUV and optical echelle spectra of the QSOs enable us to resolve the component structures of different absorption transitions and measure the column densities of metal ions accurately. Utilizing a custom software previously

developed by and described in Zahedy et al. (2016), we performed a forward modelling of Voigt profiles to constrain the ionic column densities of individual absorbing components in each LRG halo. The software was designed to analyse both well-sampled and undersampled absorption spectra with known line-spread function (LSF), and to properly assess the confidence intervals of derived model parameters via a Markov Chain Monte Carlo (MCMC) analysis.

In summary, the Voigt profile of each absorption component is uniquely defined by three parameters: the column density N_c , the Doppler parameter b_c , and the velocity centroid dv_c relative to the redshift of the strongest H I component in the absorption system. To perform the fit, the program first generated a theoretical spectrum using the minimum number of components necessary to explain the observed absorption profile. Then, this model spectrum was convolved by the appropriate instrumental LSF of the spectrograph used to collect the data, and binned to match the spectral pixel width of the data. Finally, the simulated absorption profile was compared to the observed absorption profile, and the best-fitting model was found by minimizing the χ^2 value.

To assess uncertainties in the model parameters, we performed an MCMC analysis using the EMCEE package (Foreman-Mackey et al. 2013). The MCMC analysis allows us to construct the marginalized posterior probability distribution for each model parameter. Each MCMC run consisted of 500 steps performed by an ensemble of 250 walkers. To speed up convergence, the walkers were seeded in a tiny region within the parameter space which is centred at the minimum χ^2 solution.

The absorption transitions which were analysed in a given absorption system include all observed transitions from the following list, ordered by increasing rest wavelength: O I λ 971, C III λ 977, O I λ 988, N III λ 989, the O VI $\lambda\lambda$ 1031, 1037 doublet, C II λ 1036, N II λ 1083, Fe III λ 1122, Fe II λ 1144, Si II λ 1190, Si II λ 1193, Si III λ 1206, Si II λ 1260, O I λ 1302, C II λ 1334, Si IV $\lambda\lambda$ 1393, 1402, Fe II λ 2382, Fe II λ 2586, Fe II λ 2600, the Mg II $\lambda\lambda$ 2796, 2803 doublet, and Mg I λ 2852. In our analysis, we required different transitions of the same species (e.g. Si II λ 1190 and Si II λ 1193) to have the same Voigt profile parameters. Furthermore, we imposed the same kinematic structure (i.e. number of components and velocity structure) among H I, low-, and intermediate-ionization species. This choice was justified by the excellent kinematic agreement among the observed absorption profiles of various low- and intermediate-ionization species, including Mg II, Si II, Si III, and C III (Paper I). Excepted from this requirement was high-ionization O VI absorption, which is known to often exhibit distinct velocity profiles compared to lower ionization gas (e.g. Savage et al. 2010; Werk et al. 2016). For that reason, we performed the Voigt profile analysis for the O VI doublet independently from the analysis for H I and lower ionization metals.

The results of our Voigt profile analysis are presented in Appendix A for each LRG. In Figs A1a–A16a, we present the continuum-normalized absorption profiles of different transitions, the best-fitting Voigt profiles for individual components, and the integrated Voigt profile summed over all components. These figures show the excellent agreement in velocity centroids among individual components of different ionic species, including the H I Lyman series, C II, Mg II, Si II, and C III, which demonstrates the high accuracy of our wavelength calibration. In Tables A1a–A16a, we report the best-fitting Voigt profile parameters and the associated 68 per cent confidence intervals for each component identified in the Voigt profile analysis. For saturated components, we report the 95 per cent lower limits on the column density N_c and the

corresponding 95 per cent upper limits on Doppler parameter b_c . For non-detections, we report the 95 per cent upper limits on N_c based on the error spectrum, calculated over a spectral window that is twice the FWHM (full width at half-maximum) of a line with $b_c = 10 \text{ km s}^{-1}$ for low- and intermediate-ionization species and $b_c = 30 \text{ km s}^{-1}$ for highly ionized O VI. Finally, we also report in Table A1a–Table A16a, the total $N(\text{H I})$ and ionic metal column densities summed over all components in each system.

3.2 Ionization analysis

To constrain the metallicity and other physical quantities of the CGM, it is necessary to determine the ionization state of the gas. The inferred cool temperature of the gas ($T \lesssim$ a few $\times 10^4$ K, see Section 4.2 and Appendix A) is consistent with a photoionized gas. The ionization state of the gas can be determined by comparing the observed column densities of different ionic species to predictions from photoionization calculations (e.g. Chen et al. 2017). An important physical quantity in photoionized gaseous environment is the ionization parameter U , defined as the number of incident ionizing photons per hydrogen atom. For a fixed radiation field characterized by a total flux of hydrogen-ionizing (≥ 1 Ryd) photons Φ , the U parameter is inversely proportional to the hydrogen number density n_{H} , according to $U \equiv \Phi/c n_{\text{H}}$. Higher gas density results in lower U , which leads to a more neutral gas, and vice versa. Another physical quantity which affects the observed ionic column densities is the metallicity of the gas $[M/H]$. High-metallicity gas cools more efficiently than low-metallicity gas, shifting the photoionization equilibrium towards lower ionization (i.e. more neutral) states.

We performed a series of photoionization calculations using CLOUDY v.13.03 (Ferland et al. 2013) package. We considered a plane-parallel column of gas with uniform volume density n_{H} , which was irradiated by an ultraviolet background (UVB) radiation field. To investigate how uncertainty on the UVB affects the derived gas density and metallicity (see Chen et al. 2017 for an extensive discussion), we performed two sets of calculations using two different UVBs: (1) the updated Haardt & Madau (2001) UVB, known as HM05 in CLOUDY; and (2) the Haardt & Madau (2012) UVB, known as HM12 in CLOUDY. The two radiation fields differ in both their spectral slopes and overall intensities between 1 and 10 Ryd. While the HM05 spectrum is softer than HM12 within this energy regime, the HM05 UVB has more 1–3 Ryd photons which have large photoionization cross-sections for neutral hydrogen atoms as well as low- to intermediate-ionization metals. Furthermore, the HM05 UVB has about 2.5 times (0.4 dex) the total number of hydrogen-ionizing photons of the HM12 UVB. In our ionization calculations, both UVBs were adopted at $z = 0.4$, which is roughly the median redshift of the COS-LRG galaxies. At this fiducial redshift, the relationship between U and n_{H} is $\log U = -5.42 - \log n_{\text{H}}$ for HM05, and $\log U = -5.83 - \log n_{\text{H}}$ for HM12. For example, a typical CGM gas density of $n_{\text{H}} = 0.01 \text{ cm}^{-3}$ corresponds to $\log U \approx -3.4$ and ≈ -3.8 for the HM05 and HM12 UVBs, respectively.

For each UVB, we constructed a grid of CLOUDY models spanning a wide range of H I column densities ($14 \leq \log N(\text{H I})/\text{cm}^{-2} \leq 20$ in 0.25 dex steps), gas densities ($-5 \leq \log n_{\text{H}}/\text{cm}^{-3} \leq 1$ in 0.25 dex steps), and metallicities ($-3 \leq [M/H] \leq 1$ in 0.25 dex steps). For each point in the grid, CLOUDY calculated the expected column densities and ionization fractions of different ionic species assuming photoionization equilibrium. We assumed a solar abundance pattern for the gas, although when the predictions were compared to observations, we relaxed this assumption whenever necessary and allowed by the data (see Appendix A).




Figure 1. Example predictions of ionic column densities as a function of ionization parameter U from CLOUDY photoionization calculations. From top to bottom, the panels are ordered by increasing ionization state, shown here for common ionic species O I, Mg I, C II, Mg II, Si II, C III, Si III, and O VI. The prediction curves are shown for a gas with $\log N(\text{H I}) = 16.5$ and a metallicity of $[\text{M}/\text{H}] = -0.5$, which are typical values in our sample. In solid lines, we show the predicted column densities for a gas irradiated by an updated Haardt & Madau (2001) ionizing background radiation field (HM05 in CLOUDY) at $z = 0.4$. In dashed–dotted lines, we show the corresponding predictions under the Haardt & Madau (2012) background radiation field (HM12 in CLOUDY) at the same redshift. A solar abundance pattern is assumed for the model calculations shown here.

An example of CLOUDY calculations is presented in Fig. 1, where the predicted column densities of different ions are plotted as a function of ionization parameter U . The column density curves are shown for a gas with $\log N(\text{H I})/\text{cm}^{-2} = 16.5$ and $[\text{M}/\text{H}] = -0.5$, which are typical values for individual components in our sample. The model predictions for a gas irradiated by the HM05 UVB are shown in solid lines, whereas predictions for a gas irradiated by the HM12 UVB are shown in dashed–dotted lines. Comparing the model expectations under the two different UVBs, it is clear that at fixed ionization parameter, the predicted ionic abundances for neutral and singly ionized species (e.g. Mg I, Si II, and Mg II) are systematically lower under HM12 UVB than HM05 UVB. In addition, the decrements in HM12-predicted column densities relative

to HM05 grow larger for higher U parameter (or equivalently, lower n_{H}). Similar, albeit more modest, trends are also predicted for doubly ionized species such as C III, and Si III. These trends result from of the harder HM12 UVB spectrum, which has a higher fraction of >3 Ryd photons that are needed to produce highly ionized (triply ionized or more) metal species compared to the HM05 UVB. As U increases, both low- and intermediate-ionization species are preferentially lost to higher ionization states under HM12 UVB than HM05 UVB. As a consequence of these intrinsic differences between HM05 and HM12 UVBs, HM05 models require a higher gas metallicity than HM12 models to reproduce the observed ionic abundances (see also Wotta et al. 2016).

To estimate the metallicity and density of the gas, we compared the resulting CLOUDY grid of predictions to the data and performed a statistical analysis which took into account measurements as well as upper limits (non-detections) and lower limits (saturation) in the data. Given a suite of observed ionic transitions $\{y_i\}$ for a kinematically matched absorbing component with n number of measurements, m upper limits, and l lower limits, the probability that the gas has a given density and metallicity is defined to be

$$\mathcal{P}(n_{\text{H}}, [\text{M}/\text{H}] | \{y_i\}) \propto \left(\prod_{i=1}^n \exp \left\{ -\frac{1}{2} \left[\frac{y_i - \bar{y}_i(n_{\text{H}}, [\text{M}/\text{H}])}{\sigma_i} \right]^2 \right\} \right) \times \left(\prod_{i=1}^m \int_{-\infty}^{y_i} dy' \exp \left\{ -\frac{1}{2} \left[\frac{y' - \bar{y}_i(n_{\text{H}}, [\text{M}/\text{H}])}{\sigma_i} \right]^2 \right\} \right) \times \left(\prod_{i=1}^l \int_{y_i}^{+\infty} dy' \exp \left\{ -\frac{1}{2} \left[\frac{y' - \bar{y}_i(n_{\text{H}}, [\text{M}/\text{H}])}{\sigma_i} \right]^2 \right\} \right), \quad (1)$$

where $y_i = \log N_i$ is the observed column density of the i th ionic species, σ_i is the measurement uncertainty of y_i , and $\bar{y}_i = \log \bar{N}_i$ is the corresponding model prediction. Note that in equation (1), the first product is equivalent to calculating $e^{-\frac{1}{2}\chi^2}$ for the n ionic column density measurements, whereas the second and third products extend the calculation over the m upper limits and l lower limits, respectively (see also Chen et al. 2010; Crighton et al. 2015; Stern et al. 2016).

The statistical analysis described above was performed for each absorbing component identified in our Voigt profile analysis (Section 3.1). For each component, all available column density measurements, upper limits, and lower limits for low- and intermediate-ionization species were compared to an interpolated grid of CLOUDY models evaluated at the observed $N(\text{H I})$ of the data. We note that O VI measurements were excluded from this analysis, not only because of the well-known uncertainty in the ionization mechanism of O VI absorbers, but also because of the observed kinematic misalignments between the absorption profiles of O VI and lower ionization gas (H I and metal ions; see Section 4.2 and Appendix A). As discussed in Section 5.3, our observations indicate that contributions from higher ionization gas phase to the observed column densities of lower ionization species are negligible, so the exclusion of higher ionization gas from our ionization analysis should not bias the inferred ionization parameter of cool and lower ionization gas phase considered here. We discuss the possible origins of the high-ionization gas traced by O VI absorbers in LRG haloes in Section 5.3.

The results of the ionization analysis are presented in Tables A1b–A16b in Appendix A, where for each individual component we report the number of detected metal species which are used to constrain the model, N_{metal} , the most probable gas metallicity $[\text{M}/\text{H}]$




Figure 2. Radial profile of integrated absorption column densities in the CGM of quiescent galaxies, shown for H I, Mg II, C III, Si III, and O VI. COS-LRG measurements (black circles) are plotted versus projected distance d . For comparison, absorption measurements from passive galaxies in COS-Halos (Werk et al. 2013; orange squares) and Johnson et al. (2015; red diamonds) are plotted versus normalized projected distance d/R_h . We have excluded five COS-Halos red galaxies which overlap with our LRG sample. Non-detections are shown as downward arrows which represent the 2σ upper limits on ionic column density. Meanwhile, the allowed column density range for saturated absorbers in COS-LRG are shown in empty vertical error bars. For COS-Halos red galaxies, saturated absorbers are represented by upward arrows, which show the lower limits on the absorption column density.

and density n_H under both the HM05 and HM12 UVBs, as well as the estimated 68 per cent confidence intervals for $[M/H]$ and n_H .

For components with $N_{\text{metal}} < 2$, we find that the inferred $[M/H]$ and $\log n_H$ are subject to large uncertainties of >0.5 dex, and in a number of cases, there is no clear point of maximum probability within the parameter space of the models. For these components, we report in Tables A1b–A16b the estimated 95 per cent upper or lower limits on the parameter values. In addition, we report in Tables A1b–A16b the inferred gas metallicity and density considering each absorber as a single clump (SC). For the SC model, the aforementioned ionization analysis was performed using the integrated $N(\text{H I})$ and ionic column densities summed over all components in each system, to facilitate comparisons with existing CGM/IGM ionization studies in the literature (e.g. Werk et al. 2014; Prochaska et al. 2017; Muzahid et al. 2018). Finally, we present in Figs A1b–A16b, the 2D joint probability distribution of $[M/H]$ and n_H for components with $N_{\text{metal}} \geq 2$, under both the HM05 (black contours) and HM12 (blue contours) UVBs. The contours indicate the estimated 68 per cent and 95 per cent confidence levels for the model parameters.

4 PHYSICAL PROPERTIES AND METALLICITIES IN LRG HALOES

Our analysis of CGM absorption in the COS-LRG sample reveals a diversity of gas properties in massive quiescent haloes $z \sim 0.4$. A detailed discussion on the absorption and gas properties in individual LRG haloes is presented in Appendix A, which we summarize as follows. First, a combined Voigt profile analysis on H I and metal absorption lines shows that absorbers in LRG haloes exhibit a multicomponent structure that is distributed over up to $\pm a$ few $\times 100 \text{ km s}^{-1}$ in line-of-sight velocity relative to the LRGs. Furthermore, the excellent kinematic alignments between H I, low ions (e.g. Mg II), and intermediate ions (e.g. C III) indicate a physical connection between these different species. In this section, we characterize the physical properties and chemical abundances in the gaseous haloes of LRGs.

4.1 Column density profiles of H I and heavy ions

In Fig. 2, we present the spatial distribution of absorption column densities for various ions observed in the gaseous haloes of COS-LRG galaxies. From top to bottom panels, we show the integrated column densities versus projected distance d for neutral H I, low-ionization Mg II, intermediate-ionization C III and Si III, and high-ionization O VI species. To facilitate comparisons with other surveys, we include a second horizontal axis showing the halo radius-normalized projected distance d/R_h . Recall from Paper I that the COS-LRG sample of massive quiescent galaxies has a median stellar mass and dispersion of $\log \langle M_{\text{star}}/M_{\odot} \rangle = 11.2 \pm 0.2$, which corresponds to a typical halo mass of $M_h \approx 10^{13} M_{\odot}$ according to the Kravtsov, Vikhlinin & Meshcheryakov (2018) stellar-to-halo-mass relation, and a halo radius of $R_h \approx 500 \text{ kpc}$ at $z = 0.4$.² Given the narrow range in M_{star} , we adopt this R_h for all COS-LRG galaxies plotted in Fig. 2.

For comparison, Fig. 2 shows absorption measurements for the COS-Halos red galaxy subsample (Werk et al. 2013; Prochaska et al. 2017) for H I, Mg II, C III, and Si III. For O VI, we also show measurements from the passive galaxy subsample of Johnson, Chen & Mulchaey (2015), which includes all COS-Halos red

²We approximate R_h as the region with average density of 200 times above the mean matter density of the Universe at a given epoch.

galaxies. Note that column density measurements from these studies are plotted versus the normalized projected distance of the galaxies, d/R_h . Compared to COS-LRG, the COS-Haloes subsample of red galaxies comprises predominantly lower mass galaxies, with a mass range of from $\log M_{\text{star}}/M_{\odot} = 10.3$ to 11.3, and a median of $\log (M_{\text{star}}/M_{\odot}) = 10.8$.

Despite considerable scatter in the observed $N(\text{H I})$ radial profile, there is a general trend of declining $N(\text{H I})$ with increasing d in the COS-LRG sample (top panel of Fig. 2). This trend is consistent with what is seen in COS-Haloes red galaxies as well as previous CGM surveys of the general galaxy populations (e.g. Chen et al. 1998; Johnson et al. 2015). To further examine the decline of $N(\text{H I})$ with increasing distance, we divide our sample into two bins at $d = 100$ kpc, which is approximately the median projected distance. At $d < 100$ kpc from LRGs, the majority of H I absorbers are optically thick (Lyman limit systems (LLSs) with $\log N(\text{H I})/\text{cm}^{-2} > 17.2$). In contrast, there is a significantly higher fraction of optically thin absorbers as well as sightlines with non-detections at $d > 100$ kpc. The mean covering fraction of optically thick H I gas is $\langle \kappa \rangle_{\text{LLS}} = 0.71^{+0.19}_{-0.26}$ at $d < 100$ kpc, which declines to $\langle \kappa \rangle_{\text{LLS}} = 0.22^{+0.22}_{-0.14}$ at $d = 100$ –160 kpc.

A trend of declining column density with increasing d is also seen in low-ionization metal species such as Mg II. At $d < 100$ kpc, strong $N(\text{Mg II})$ absorbers with $\log N(\text{Mg II})/\text{cm}^{-2} > 13$ are common in COS-LRG. In contrast, absorbers at $d \gtrsim 100$ kpc exhibit significantly lower $N(\text{Mg II})$, where $\log N(\text{Mg II})/\text{cm}^{-2} < 13$ is seen in all cases. For strong Mg II absorbers in COS-LRG, we estimate a mean covering fraction of $\langle \kappa(\text{Mg II}) \rangle_{13.0} = 0.60^{+0.25}_{-0.30}$ at $d < 100$ kpc. A caveat of this calculation is that two sightlines at $d < 100$ kpc do not have any Mg II constraints and consequently do not contribute to the covering fraction estimation. Including these two sightlines would lead to a mean Mg II covering fraction of $\langle \kappa(\text{Mg II}) \rangle_{13.0} \approx 0.4 - 0.7$ at $d < 100$ kpc, depending on whether these two absorbers satisfy the strong Mg II absorption criterion or not. In contrast, the lack of strong Mg II absorption at $d = 100$ –160 kpc from LRGs in our sample constrains the mean covering fraction to $\langle \kappa(\text{Mg II}) \rangle_{13.0} \approx 0.0 - 0.2$, for $\log N(\text{Mg II})/\text{cm}^{-2} > 13$.

A surprising finding from Paper I is the high incidence of absorption from intermediate-ionization species C III and Si III, comparable to what have been observed around star-forming galaxies. While the high oscillator strength of the C III $\lambda 977$ transition makes C III absorption easily detectable, it also means that C III absorption profiles are often saturated (see Appendix A). For that reason, it is difficult to draw a strong conclusion on possible radial trends in intermediate ionic column densities using C III absorption. For the comparatively weaker Si III absorption, it is clear that strong Si III absorption with $\log N(\text{Si III})/\text{cm}^{-2} > 13.0$ are more prevalent at smaller d . At $d < 100$ kpc from COS-LRG galaxies, absorbers with $\log N(\text{Si III})/\text{cm}^{-2} > 13.0$ are present in 5 out of 7 cases, which constrains the mean Si III covering fraction to $\langle \kappa(\text{Si III}) \rangle_{13.0} = 0.71^{+0.19}_{-0.26}$. In contrast, Si III absorption are generally weaker at $d = 100$ –160 kpc, with a high fraction (50 per cent) of non-detections. The estimated mean Si III covering fraction absorption at $d = 100$ –160 kpc is $\langle \kappa(\text{Si III}) \rangle_{13.0} = 0.37^{+0.24}_{-0.19}$, for $\log N(\text{Si III})/\text{cm}^{-2} > 13.0$.

For high-ionization gas, measurements of O VI column density in LRG haloes are available for 12 out of 16 COS-LRG galaxies. We detect O VI absorption in seven sightlines at a detection threshold of $\log N(\text{O VI})/\text{cm}^{-2} > 13.5$, which translates to an estimated mean covering fraction of $\langle \kappa(\text{O VI}) \rangle_{13.5} = 0.58^{+0.17}_{-0.18}$ at $d < 160$ kpc ($\sim 0.3 R_h$). The mean O VI covering fraction for LRGs is comparable to what Johnson et al. (2015) found at $d \lesssim 0.3 R_h$ for their passive galaxy subsample, $\langle \kappa(\text{O VI}) \rangle_{11.5} = 0.62^{+0.13}_{-0.17}$ for

$\log N(\text{O VI})/\text{cm}^{-2} > 13.5$. Note that the red galaxies in Johnson et al. (2015) are predominantly less massive than COS-LRG galaxies, with a median stellar mass and dispersion of $\log \langle M_{\text{star}}/M_{\odot} \rangle = 10.7 \pm 0.5$. The comparable O VI covering fractions in massive quiescent haloes spanning over an order of magnitude in halo mass (from $M_h \sim 10^{12}$ to $\gtrsim 10^{13} M_{\odot}$) suggest that O VI absorbers in all quiescent haloes may share a similar physical origin. A two-sample Kolmogorov–Smirnov (K-S) test demonstrates that we cannot rule out at high statistical significance (> 99 per cent) that the lower mass (Johnson et al. 2015) and massive (COS-LRG) quiescent halo samples of O VI absorbers are drawn from the same parent population.

4.2 Kinematic and thermal properties

The line-of-sight kinematics of absorbing gas relay crucial information about the underlying motion of cool clumps within LRG haloes. Our discussion of individual LRG haloes in Appendix A highlights the fact that cool gas absorption profiles in the CGM of LRGs consist of multiple components that are distributed within \pm a few hundred km s^{-1} in line-of-sight velocity relative to the systemic redshift of the galaxy. The distribution of line-of-sight velocities of individual H I components relative to the LRGs is shown in the left-hand panel of Fig. 3. The velocity distribution can be characterized by a mean and dispersion of $\langle \Delta v_{\text{gas-galaxy}} \rangle = 17 \text{ km s}^{-1}$ and $\sigma_{\Delta v_{\text{gas-galaxy}}} = 147 \text{ km s}^{-1}$. The observed velocity dispersion is consistent with what have been reported for large samples of Mg II absorbers around LRGs using low-resolution data (e.g. Zhu et al. 2014; Huang et al. 2016; Lan & Mo 2018). To provide a physical context, the inferred mean mass of LRG haloes is $M_h \approx 10^{13.4} M_{\odot}$ (e.g. Mandelbaum, Seljak & Hirata 2008; Gauthier et al. 2009), and the expected line-of-sight velocity dispersion for virialized motion in LRG haloes is $\sigma_h \approx 260 \text{ km s}^{-1}$. The observed line-of-sight velocity dispersion of the gas, $\sigma_{\Delta v_{\text{gas-galaxy}}} = 147 \text{ km s}^{-1}$, is merely ~ 60 per cent of the expectation from virial motion. The narrow distribution of line-of-sight velocities indicates that an effective dissipative mechanism is at play to slow down the motion of cool gas in the halo (e.g. Huang et al. 2016).

To evaluate whether the observed velocity dispersion varies with projected distance, we divide the absorbing components into two subsamples on d , one for components at $d < 100$ kpc and another for those at $d > 100$ kpc. The resulting velocity distributions of the two subsamples are shown in the middle panel of Fig. 3. While the velocity histograms are understandably noisy due to the smaller size of the two subsamples, no significant trend is detected between the line-of-sight velocity distributions at small and large d . Using a two-sided K-S test, we cannot rule out that the two d subsamples come from the same parent distribution in $\Delta v_{\text{gas-galaxy}}$ at > 50 per cent confidence. We also bisect the sample of individual components into two groups based on their metallicities, a low-metallicity subsample with $[\text{M}/\text{H}] < -0.5$ and a high-metallicity subsample with $[\text{M}/\text{H}] > -0.5$. Again, no statistically significant distinction can be made between the low- and high-metallicity subsamples, with components from each subsample occupying the full range of velocities with respect to the LRGs (Fig. 3, right-hand panel). A K-S test cannot rule out that the two $[\text{M}/\text{H}]$ subsamples come from the same parent distribution in $\Delta v_{\text{gas-galaxy}}$ at > 68 per cent confidence.

Our Voigt profile analysis also allows us to examine the thermal properties of cool clumps in LRG haloes. In the left-hand panel of Fig. 4, we plot the Doppler linewidths of Mg II components as a function of d . With the exception of a few broad components,




Figure 3. Line-of-sight velocity distributions of individual absorption components relative to the LRG systemic redshifts. For the full COS-LRG sample (left-hand panel), we find a mean and dispersion of $\langle \Delta v_{\text{gas-galaxy}} \rangle = 17 \text{ km s}^{-1}$ and $\sigma_{\Delta v_{\text{gas-galaxy}}} = 147 \text{ km s}^{-1}$. In the middle panel, the sample is biseected by projected distance d , whereas in the right-hand panel, the sample is divided by metallicity. We find no statistically significant distinction between the subsamples in either case.




Figure 4. Left: Doppler parameter b_c plotted versus d for individual Mg II absorption components. Saturated components are represented by greyed out data points. Right: distribution of Doppler linewidths for matched H I and Mg II components. The dashed–dotted line shows the expectation for a pure thermal broadening case, $b_c(\text{Mg II}) \approx 0.2 b_c(\text{H I})$, whereas the dotted line shows the expected relationship when Mg II and H I linewidths are dominated by non-thermal broadening, $b_c(\text{Mg II}) = b_c(\text{H I})$. We find that cool CGM gas around LRGs has a mean temperature and dispersion of $\langle T \rangle = 2.0 \times 10^4 \text{ K}$ and $\sigma_T = 1.4 \times 10^4 \text{ K}$, with a modest non-thermal broadening of $\langle b_{\text{nt}} \rangle = 7 \pm 5 \text{ km s}^{-1}$.

most Mg II components in LRG haloes are narrow with $b_c(\text{Mg II}) < 10 \text{ km s}^{-1}$. The narrow linewidths imply that the gas is both cool and kinematically quiescent. Furthermore, no trend in $b_c(\text{Mg II})$ in seen versus d , indicating that cool gas at small- and large-projected distances from LRG have similar thermal properties.

Next, we show the distribution of Doppler linewidths for matched H I and Mg II absorption components in the right-hand panel of Fig. 4. Two straight lines are drawn to indicate two limiting cases. First, the dashed–dotted line in the bottom represents the expectation for a pure thermal-broadening case where the Mg II and H I linewidths are related by the square root of their mass ratio alone, giving $b_c(\text{Mg II}) \approx 0.2 b_c(\text{H I})$. Secondly, the dotted line on top of the

panel shows the expected relation when the Mg II and H I linewidths are dominated by non-thermal broadening, $b_c(\text{Mg II}) \approx b_c(\text{H I})$.³ It is clear from the right-hand panel of Fig. 4 that a large majority data

³While the parameter space outside the region bounded by the two limiting cases is unphysical, two components are found below the thermal-broadening line. One has a $b_c(\text{Mg II})$ that is consistent within 1σ with thermal broadening. The other component has a broad $b_c(\text{H I}) = 33 \text{ km s}^{-1}$, but its Mg II linewidth is only 3 km s^{-1} , which is narrower than expected from thermal broadening. The unphysical relationship between $b_c(\text{H I})$ and $b_c(\text{Mg II})$ for this component implies the presence of unresolved H I components that are not Mg II-bearing.




Figure 5. Cumulative fraction, P , of O VI absorption components with absolute centroid velocity difference less than $|\Delta v|$ from the nearest low-ionization metal and H I components (solid histogram). The mean/median value is $\langle |\Delta v| \rangle = 24 \text{ km s}^{-1}$, with a full range of from $|\Delta v| = 4$ to 71 km s^{-1} . Note that the final wavelength solution for our FUV COS spectra is accurate to within $\pm 3 \text{ km s}^{-1}$, and the expected $P(|\Delta v|)$ for a normal distribution with a width of $\sigma = 3 \text{ km s}^{-1}$ is shown in dotted curve for comparison. The mismatched kinematics between high- and low-ionization gas in COS-LRG suggest different physical origins between the high- and low-ionization gas.

points are situated closer to the thermal-broadening line than to the non-thermal-broadening line. This is consistent with a quiescent gas that is subject to little non-thermal broadening. The ratios of Doppler linewidths for matched H I and Mg II components in the COS-LRG sample show that the gas has a mean temperature and dispersion of $\langle T \rangle = 2.0 \times 10^4 \text{ K}$ and $\sigma_T = 1.4 \times 10^4 \text{ K}$, with a modest mean non-thermal line broadening of $\langle b_{nt} \rangle = 7 \pm 5 \text{ km s}^{-1}$.

Finally, we find that O VI absorption profiles in COS-LRG show distinct kinematic structures from the absorption profiles of lower ionization metal and H I (see Appendix A for a detailed description of individual absorbing systems). The mean/median absolute difference in centroid velocity between O VI absorption components and the nearest H I and low-ionization metal component is $\langle |\Delta v| \rangle = 24 \text{ km s}^{-1}$, with a full range of from $|\Delta v| = 4$ to 71 km s^{-1} (Fig. 5). Recall that the final wavelength solution for our FUV COS spectra is accurate to within $\pm 3 \text{ km s}^{-1}$. The kinematic misalignments between high- and low-ionization gas in the COS-LRG sample suggest that different phases of the CGM gas of LRGs have different physical origins (a more in-depth discussion is presented in Sections 5.2 and 5.3).

4.3 Metallicities and densities

Our ionization analysis on matched absorption components reveals significant variations in gas metallicities and densities in the cool CGM of LRGs, both within individual haloes and among different haloes in the COS-LRG sample. We now discuss and investigate for trends in gas metallicities and densities in the COS-LRG ensemble of galaxies. We begin with a discussion on systematic errors in the ionization analysis.

4.3.1 Systematic errors arising from the uncertain UVB

Metallicity and density estimates in CGM studies are based on comparing the absorption column densities of ionic metals and neutral hydrogen. Because the gas is highly ionized in all but the




Figure 6. Top: difference in gas densities derived under HM05 and HM12 UVBs, plotted versus $N_c(\text{H I})$. Compared to the HM12 UVB, the HM05 UVB leads to higher inferred n_{H} values, with a median difference and dispersion of $\langle \Delta \log n_{\text{H}} \rangle = 0.34 \text{ dex}$ and $\sigma_{\Delta \log n_{\text{H}}} = 0.10 \text{ dex}$. Bottom: difference in metallicities derived under HM05 and HM12 UVBs, plotted versus individual component H I column density, $N_c(\text{H I})$. The HM05 UVB leads to lower inferred $[\text{M}/\text{H}]$ than the HM12 UVB, with a metallicity difference that range from $\langle \Delta [\text{M}/\text{H}] \rangle = -0.1$ for LLSs to $\langle \Delta [\text{M}/\text{H}] \rangle = -0.7$ for optically thin gas.

highest column density absorbers, substantial ionization fraction corrections are necessary to convert the observed ionic column density ratios to the desired elemental abundances. A complicating factor in the ionization analysis of CGM gas is the well-known uncertainties in the shape, intensity, and redshift evolution of the extragalactic UVB (e.g. Faucher-Giguère et al. 2008; Haardt & Madau 2012; Kollmeier et al. 2014; Shull et al. 2015), which affect the expected ionization fraction corrections in the gas. Adopting different UVBs for the ionization analysis can propagate to order-of-magnitude discrepancies in the inferred gas metallicity (see Chen 2017, for an extensive discussion on the subject).

To explore how the uncertain UVB spectrum affects the derived gas densities and metallicities, we performed our ionization analysis using two different photoionizing background radiation fields, the HM05 and HM12 UVBs (see the discussion in Section 3.2). In the top panel of Fig. 6, the difference in gas densities derived under HM05 and HM12 UVBs is plotted versus component H I column density $N_c(\text{H I})$. Over almost five decades in $N_c(\text{H I})$, the gas densities inferred using HM05 UVB are systematically higher than gas densities inferred using the HM12 UVB, with a median difference and dispersion of $\langle \Delta \log n_{\text{H}} \rangle = 0.34 \text{ dex}$ and $\sigma_{\Delta \log n_{\text{H}}} = 0.10 \text{ dex}$. The higher inferred gas density under HM05 can be understood as due to the higher intensity of HM05 UVB compared to the HM12 UVB. Recall from our discussion in Section 3.2 that the total flux of hydrogen-ionizing photons in HM05 UVB is $\sim 0.4 \text{ dex}$ higher than that of the HM12 UVB. Consequently, the higher intensity HM05 UVB requires a higher underlying gas density than the HM12 UVB

for fixed ionization parameter U , which describes the ionization state of the gas.

In the bottom panel of Fig. 6, we plot the difference in metallicities derived under HM05 and HM12 UVBs versus $N_c(\text{H I})$. Over more than five decades in H I column density, not only is the $[\text{M}/\text{H}]$ inferred under HM05 UVB systematically lower than inferred under HM12 UVB, but also the difference in metallicities depends on $N_c(\text{H I})$. For optically thick gas with $\log N_c(\text{H I})/\text{cm}^{-2} \gtrsim 17$, the typical metallicity difference between HM05 and HM12 is modest, $\langle \Delta[\text{M}/\text{H}] \rangle = -0.1 \pm 0.1$ dex. The median metallicity difference is larger for lower $N_c(\text{H I})$ gas, ranging from $\langle \Delta[\text{M}/\text{H}] \rangle \sim -0.3$ dex for gas with $\log N_c(\text{H I})/\text{cm}^{-2} \sim 16$ to $\langle \Delta[\text{M}/\text{H}] \rangle \sim -0.7$ dex for optically thin gas with $\log N_c(\text{H I})/\text{cm}^{-2} < 15$. To understand the origin of this trend, recall from Section 3.2 that not only does the HM12 UVB have a harder spectrum than the HM05 UVB, but also it has a higher fraction of >3 Ryd photons which are required to produce high-ionization (triply ionized or more) metal species. As $N_c(\text{H I})$ decreases and the gas becomes more highly ionized, more low- and intermediate-ionization metals are preferentially lost to higher ionization states under HM12 UVB than under HM05 UVB. Because metallicity estimates of cool CGM gas often rely on suite of low- and intermediate-ionization metal species, the difference in metallicities inferred under HM05 and HM12 UVBs naturally increases with decreasing $N(\text{H I})$.

Finally, we note that in all ionization calculations performed in this work, both HM05 and HM12 UVBs were adopted at redshift $z = 0.4$, which is roughly the median redshift of COS-LRG galaxies. Changing the adopted UVB redshift to $z = 0.2$ or 0.6 would change the intensity of each UVB by no more than ± 0.2 dex. As a result, the inferred gas density n_{H} would change by less than ± 0.2 dex by changing the adopted UVB redshift, which is smaller than the median difference in n_{H} derived under HM05 and HM12 UVBs.

Using two different UVBs that are frequently utilized in CGM/IGM studies, we have quantified the systematic errors resulting from the uncertain shape and intensity of the extragalactic UVB radiation field. It must also be noted that a known issue with the HM12 UVB is that it overpredicts the amplitude of H I column density distribution function in low-redshift ($z < 1$) Ly α forest by a factor of 2–4 (e.g. Kollmeier et al. 2014; Shull et al. 2015; Viel et al. 2017). The HM12 UVB does not match low-redshift IGM observations because of its low hydrogen photoionization rate (Γ_{H}), which is a result of the adopted negligible escape fraction of Ly-continuum photons from low-redshift galaxies (e.g. Shull et al. 2015). In contrast, the HM05 UVB assumes a higher escape fraction of ionizing photons from galaxies, which has been shown to provide better agreement with observations (e.g. Kollmeier et al. 2014; Khaire & Srianand 2015; Viel et al. 2017). For that reason, we adopt the gas metallicities and densities inferred using the HM05 UVB for subsequent analyses and discussions in this work.

4.3.2 Trends in gas metallicities and densities

To illustrate the diversity of inferred gas metallicities and densities in the cool CGM of LRGs, we plot component $[\text{M}/\text{H}]$ versus n_{H} in Fig. 7. While no evidence is seen for any correlation between $[\text{M}/\text{H}]$ and n_{H} , Fig. 7 shows that cool gas in LRG haloes occupy a wide range of metallicities (from less than 0.01 solar to solar and super-solar metallicities) and densities (from $n_{\text{H}} \lesssim 0.001$ to $\sim 0.1 \text{ cm}^{-3}$).




Figure 7. Inferred gas metallicity $[\text{M}/\text{H}]$ versus hydrogen density n_{H} for individual absorbing components in the COS-LRG sample. The vertical and horizontal error bars associated with each data point show the 68 per cent confidence intervals for $[\text{M}/\text{H}]$ and n_{H} , respectively. Greyed out data points show components for which only upper/lower limits on $[\text{M}/\text{H}]$ and/or n_{H} are available, with arrows indicating the 95 per cent upper/lower limits. We find no statistically significant correlation between $[\text{M}/\text{H}]$ and n_{H} .

We present the spatial distribution of component metallicity as a function of d in the left-hand panel of Fig. 8. Two interesting features are revealed by this plot. First, $[\text{M}/\text{H}]$ exhibits large variations among different components detected in the gaseous halo of an LRG, at small and large d alike. A majority of LRG haloes (~ 60 per cent) that exhibit multicomponent absorption profiles show over a factor of 10 difference in $[\text{M}/\text{H}]$ between the most metal-rich and metal-poor components. Such large variations in $[\text{M}/\text{H}]$ within the gaseous halo indicate poor chemical mixing in the CGM of LRGs and underscore the importance of resolving the component structures of CGM absorbers, which is afforded by our high-resolution absorption spectra (see also e.g. Churchill et al. 2012; Rosenwasser et al. 2018). In contrast, any information on intra-halo variations is lost if one utilizes only the integrated H I and metal column densities along individual sightlines in the ionization analysis.

Furthermore, while high-metallicity ($[\text{M}/\text{H}] \gtrsim -1.0$) components are observed in most LRG haloes, metal-poor ($[\text{M}/\text{H}] \lesssim -1.0$) components are found in half of LRG haloes, with a majority these low-metallicity components occurring at $d \gtrsim 100$ kpc. Over the full sample, the median metallicity of individual components is $\langle [\text{M}/\text{H}] \rangle = -0.7 \pm 0.2$, where the uncertainty is calculated using a combined bootstrap and Monte Carlo resampling. In addition, we estimate the 16–84 percentile range in $[\text{M}/\text{H}]$ to be $[\text{M}/\text{H}] = (-1.6, -0.1)$ for the whole sample. Note that components with poor constraints on $[\text{M}/\text{H}]$ (those with metallicity upper limits which are higher than solar metallicity) are excluded from these estimates.

In the right-hand panel of Fig. 8, we present a plot of $[\text{M}/\text{H}]$ versus component H I column density. We find no significant trend in $[\text{M}/\text{H}]$ versus $N_c(\text{H I})$. This lack of correlation in our data stands in contrast to the anticorrelation between metallicity and H I column density that was reported in a number of recent studies (e.g. Prochaska et al. 2017; Muzahid et al. 2018). Considering the




Figure 8. Left: gas metallicity $[M/H]$ versus d in the COS-LRG sample. Each vertical coloured line connects different absorption components detected within the same LRG halo. The vertical error bar associated with each data point shows the 68 per cent confidence interval for $[M/H]$. Greyed out data points are absorbing components with no metal ions detected, with downward arrows indicating the 95 per cent upper limits on $[M/H]$. Large ($\gtrsim 1$ dex) variations in gas metallicities within the CGM are seen in a majority of LRGs that exhibit multicomponent absorption profiles. Right: $[M/H]$ versus component H I column density $N_c(\text{H I})$. There is no evidence for any metallicity trend with $N_c(\text{H I})$. The median metallicity of individual components is $\langle [M/H] \rangle = -0.7 \pm 0.2$, with an estimated 16–84 percentile range of $[M/H] = (-1.6, -0.1)$ for the whole sample.

known trend of declining $N(\text{H I})$ with d in the CGM (e.g. Chen et al. 1998; Johnson et al. 2015; see also Section 4.1), the reported anticorrelation implies that metallicity increases with distance from galaxies, which is difficult to explain. This discrepancy can be attributed as due to two systematic effects. First, the HM12 UVB, which was the adopted UVB in Prochaska et al. (2017), predicts progressively higher metallicities (up to 0.7 dex) with decreasing $N(\text{H I})$ compared to the HM05 UVB (see Section 4.3.1). Secondly, the ionization analyses in these studies utilized integrated H I and metal column densities summed over all components in each absorption system. As we approach lower column density regime, the required data quality (S/N) is higher to detect the gas. Given a fixed S/N and a system with multiple components, weaker metal components is more challenging to uncover. Consequently, relatively more metal-poor gas goes undetected more easily. By treating resolved components separately, we find that several low- $N(\text{H I})$ components only have non-constraining metallicity upper limits. The combination of these two systematic effects explain the reported anticorrelation between $[M/H]$ and $N(\text{H I})$.

A surprising finding from our analysis is the significant incidence of low-metallicity LLSs in the COS-LRG. The right-hand panel of Fig. 8 shows that three optically thick components (out of seven overall) with $\log N_c(\text{H I})/\text{cm}^{-2} \gtrsim 17$ have very low metallicities, $[M/H] \lesssim -1.5$ or less than 0.03 solar metallicity. Two of these components (component 2 along SDSS J0946+5123 and component 4 along SDSS J0246–0059, see Appendices A1 and A10, respectively) contain anomalously little ionic metals despite hosting the bulk of the total H I column density in their respective absorbers. The other component, a remarkable metal-free LLS along SDSS J1357+0435, has the lowest metallicity in the COS-LRG sample, with an estimated metallicity upper limit of $[M/H] < -2.3$ or lower than 0.5 per cent of solar metallicity. Such low metallicities in low-redshift LLSs are consistent with recently accreted gas from the IGM (e.g. Hafen et al. 2017). We estimate the rate of

very-low-metallicity LLS (with $[M/H] \lesssim -1.5$) to be $0.43^{+0.25}_{-0.22}$ assuming binomial statistics (Gehrels 1986), which suggests that chemically pristine gas accreted from the IGM contributes to a substantial fraction of LLS population in LRG haloes at $z \lesssim 0.5$. A more in-depth discussion on the possible origins of low-metallicity gas in LRG haloes is presented in a companion paper on the galaxy environment of the chemically pristine LLS observed along SDSS J1357+0435 (Chen et al. 2018b).

Next, we present a plot of gas density n_{H} versus d in the left-hand panel of Fig. 9. Similar to what is seen with gas metallicities, the inferred n_{H} shows substantial variations among different components detected within the gaseous halo of a given LRG. In half of LRG haloes (6/12) that exhibit multicomponent absorption profiles, we find over a factor of 10 difference in n_{H} between the highest and lowest density components. These large intra-halo variations in n_{H} are observed at both $d < 100$ and > 100 kpc. The median gas density of individual components is $\log \langle n_{\text{H}} \rangle / \text{cm}^{-3} = -2.4 \pm 0.1$, where the uncertainty is calculated using a combined bootstrap and Monte Carlo resampling. In addition, the estimated 16–84 percentile range in gas density is $\log n_{\text{H}} / \text{cm}^{-3} = (-3.0, -1.8) \text{ cm}^{-3}$.

To investigate whether gas density varies with H I column density, we plot n_{H} versus $N_c(\text{H I})$ in the right-hand panel of Fig. 9. The corresponding ionization parameter U for a given n_{H} is indicated on the right y-axis. The data points exhibit a clear trend of rising gas density with increasing H I column density. Because the ionizing background radiation is fixed, the observed correlation is consistent with what is expected from a photoionized gas: more optically thick gas has lower ionization parameter U and is therefore less ionized than optically thin gas. The inferred median U for our sample is $\log \langle U \rangle = -3.0 \pm 0.1$. For stronger absorption components with $\log N_c(\text{H I})/\text{cm}^{-2} \gtrsim 16$, the median ionization parameter is lower, $\log \langle U \rangle \approx -3.5$, which is comparable to what have been found in previous surveys of $z < 1$ partial-LLSs/LLSs (e.g. Lehner et al. 2013).




Figure 9. Left: gas density n_{H} versus d in the COS-LRG sample. Each vertical coloured line connects different absorption components detected within the same LRG halo. The vertical error bar associated with each data point shows the 68 per cent confidence interval for n_{H} . Greyed out data points show components for which only upper/lower limits on n_{H} are available, with upward/downward arrows indicating the 95 per cent upper/lower limits on the underlying gas density. Large ($\gtrsim 1$ dex) variations in n_{H} within the CGM are seen in half of COS-LRG galaxies that exhibit multicomponent absorption profiles. The median gas density of individual components is $\log \langle n_{\text{H}} \rangle / \text{cm}^{-3} = -2.4 \pm 0.1$. Right: n_{H} versus component HI column density $N_{\text{c}}(\text{HI})$. The corresponding ionization parameter U for a given n_{H} is indicated on the right y-axis. The trend of rising gas density with increasing HI column density indicates that high $N_{\text{c}}(\text{HI})$ gas has lower ionization parameter U and is therefore less ionized than low $N_{\text{c}}(\text{HI})$ gas. The median U in the COS-LRG sample is $\log \langle U \rangle \approx -3.0$. Finally, each dotted curve shows the expected $n_{\text{H}}-N_{\text{c}}(\text{HI})$ relation for a cool cloud of a given line-of-sight thickness, from $l = 10$ pc to 10 kpc. The distribution of cool clump sizes shows a clear mode at ~ 100 pc, with an estimated median of $\langle l \rangle = 120_{-40}^{+80}$ pc.

The strong correlation between n_{H} and $N_{\text{c}}(\text{HI})$ also suggests that cool clumps in the CGM of LRGs follow a well-defined distribution of clump sizes. In the right-hand panel of Fig. 9, we plot the expected relationship between n_{H} and $N_{\text{c}}(\text{HI})$ for cool clumps of different thicknesses, from $l = 10$ pc to 10 kpc. It is clear that a large majority of clumps are between ~ 10 pc and ~ 1 kpc thick, with a mode of ~ 100 pc. Furthermore, this characteristic clump thickness of $l \sim 100$ pc is shared by both optically thin and thick clumps, covering a range of nearly three orders of magnitude in $N(\text{HI})$. The median clump size estimated for the COS-LRG sample is $\langle l \rangle = 120_{-40}^{+80}$ pc, where the uncertainty is calculated using a combined bootstrap and Monte Carlo resampling. In addition, we estimate that the range of l containing 68 per cent of individual components is $l = (20, 800)$ pc. The range of inferred clump sizes in LRG haloes is in excellent agreement with transverse clump sizes estimated directly from intervening low-ionization absorbers in the spectra of multiply lensed, high-redshift QSOs (e.g. Rauch, Sargent & Barlow 1999; Rauch et al. 2002).

To put the inferred gas densities of cool CGM clumps in a broader context, we compare the inferred n_{H} in the cool CGM with the expected gas densities in the hot CGM ($T \sim 10^6$ K) of LRGs in Fig. 10. First, n_{H} is plotted versus d and shown in circles/triangles for optically thin/thick cool gas. We also show n_{H} derived for a single-clump model in hollow red squares, where different components within a given LRG halo are imposed to have the same density and metallicity.⁴ For comparison, the solid line in Fig. 10 represents $n_{\text{hot}}(r)|_{r=d}$, the mean radial profile of mean hot gas density in LRG




Figure 10. Spatial distribution of gas densities in the cool ($T \sim 10^4$ K) and hot ($T \sim 10^6$ K) CGM of LRGs. First, n_{H} is plotted versus d and shown in circles/triangles for optically thin/thick components. In hollow red squares, we show n_{H} derived for a single-clump model, where a single density is assumed for different components within an individual halo. For comparison, the solid line shows $n_{\text{hot}}(r)|_{r=d}$, the radial profile of mean hot gas density in massive haloes ($M_{\text{h}} \sim 10^{13} M_{\odot}$; Singh et al. 2018) evaluated at $r = d$. The shaded grey area signifies the 68 per cent confidence region of this power-law density profile. Finally, in dashed line, we show the same hot CGM density profile which has been scaled up by a factor of 100.

⁴Note that the single-clump model results in a positive bias on the inferred distribution of n_{H} , because the inferred density in the single-clump model

haloes, evaluated at $r = d$. For $n_{\text{hot}}(r)$, we chose a power-law model that describes the hot CGM of massive haloes ($M_{\text{h}} \sim 10^{13} M_{\odot}$) from Singh et al. (2018), which is based on a combined analysis of X-ray and Sunyaev–Zel’dovich (SZ) signals from a stack of $\sim 10^5$ massive galaxies at $z \sim 0.1$. The dotted line in Fig. 10 represents a boosted Singh et al. (2018) hot gas density profile which has been scaled up by a factor of 100, for visual comparison.

It is clear that the projected radial density profile of optically thick cool gas sits about 100 times higher than $n_{\text{hot}}(r)|_{r=d}$. Considering the two orders of magnitude of temperature difference between cool CGM gas ($T \sim 10^4$ K, see Section 4.2) and X-ray emitting hot gas ($T \sim 10^6$ K), the inferred density contrast of ~ 100 indicates that optically thick cool CGM gas occurs at $r \sim d$ and is close to being in pressure equilibrium with the hot halo (e.g. Mo & Miralda-Escudé 1996; Maller & Bullock 2004; see also a more in-depth discussion in Section 5.2). In contrast, Fig. 10 shows only ~ 40 per cent of optically thin components have densities consistent with being in thermal pressure equilibrium with the hot halo at $r \sim d$, which implies that a majority of optically thin absorbers likely occur at larger radii in the halo, $r > d$.

5 DISCUSSION

The COS-LRG survey consists of a mass-limited sample of 16 LRGs with $\log M_{\text{star}}/M_{\odot} > 11$ and $d < 160$ kpc from a background QSO, chosen without any prior knowledge of the presence or absence of absorption features in the LRG haloes. This mass-limited and absorption-blind sample enables an unbiased and accurate characterization of the physical properties and metallicities in the CGM of these intermediate-redshift massive ellipticals. Our survey demonstrates that despite their quiescent nature, LRGs are surrounded by widespread and chemically enriched cool gas. By carrying out a detailed ionization analysis on the absorbers, we discover large variations in gas metallicities and number densities in the cool gas, both within individual LRG haloes and across the entire sample. When compared with the expected gas densities in the hot halo, the inferred densities of the cool gas imply that cool clumps in the CGM of LRGs are likely supported by thermal pressure. In addition, we find kinematic mismatches between high-ionization O VI gas and lower ionization gas traced by H I and associated metal ions, which suggest different physical origins of the gas. We now discuss the implications of our study.

5.1 Total mass in the cool CGM of LRGs

The relative amounts of gas that reside in different phases of the CGM are governed by the interplay of accretion and feedback, as well as the detailed gas physics. Empirical constraints on the total mass of the gaseous halo around galaxies are therefore critical to test the validity of current theoretical models of galaxy formation. However, previous estimates of the total mass in the cool CGM of quiescent galaxies suffer from large uncertainties of up to two orders of magnitude, due to the unknown ionization state of the gas (e.g. Thom et al. 2012; Zhu et al. 2014).

Here, we leverage the results of our ionization analysis of the COS-LRG data set in order to infer the surface mass density profile and estimate the total gas mass in the cool CGM of LRGs. For each

is driven predominantly by the densest cool absorption component in each halo.




Figure 11. Surface mass density profile of the cool CGM of LRGs. The surface mass density, Σ_{cool} , is estimated by calculating the total hydrogen column density along each sightline, N_{H} , with the estimated ionization fraction corrections applied to the observed H I column densities according to equation (2). Downward arrows represent estimated upper limits on Σ_{cool} for LRG haloes with non-detected H I. Empty vertical error bars show the range of allowed surface mass density for systems with weakly constrained n_{H} , calculated by imposing that the corresponding clump size is $l \lesssim 1$ kpc. Σ_{cool} exhibits a steep decline with increasing d , which cannot be reproduced by a projected NFW or Einasto profile expected for $M_{\text{h}} \approx 10^{13} M_{\odot}$ dark-matter haloes (dashed and dashed-dotted curves). On the other hand, the radial profile of Σ_{cool} is best described by an exponential profile in either 2D (solid line) or 3D (a projected Einasto profile with $\alpha \approx 1$; dotted curve).

absorption system, we first calculate the total hydrogen column density, N_{H} , according to the following equation,

$$\log N_{\text{H}} = \log \sum_i \frac{N_{\text{c}}(\text{H}1)_i}{f_{\text{H}^0_i}}, \quad (2)$$

where $f_{\text{H}^0_i}$ is the hydrogen neutral fraction for component i determined by our ionization analysis, and the sum is evaluated over all components in the absorption system. For components with poorly constrained ionization state, the range of allowed ionization fraction correction is computed by imposing that the corresponding clump size is not larger than 1 kpc. Once N_{H} is calculated, the corresponding cool gas surface mass density can be computed using the relation $\Sigma_{\text{cool}} = 1.4 m_{\text{H}} N_{\text{H}}$, where m_{H} is the mass of the hydrogen atom and a factor of 1.4 is introduced to account for the contribution of helium to the total gas mass.

The spatial profile of cool gas surface mass density in the CGM is shown in Fig. 11. It is clear that Σ_{cool} exhibits a declining trend with d . At $d < 100$ kpc, the mean N_{H} is $\log \langle N_{\text{H}} \rangle / \text{cm}^{-2} = 19.5 \pm 0.2$, which is equivalent to a mean cool gas surface density of $\langle \Sigma_{\text{cool}} \rangle \approx 0.4_{-0.1}^{+0.2} M_{\odot} \text{pc}^{-2}$. The mean Σ_{cool} at $d < 100$ kpc is comparable to inferred surface mass densities in the predominantly neutral ISM of an LRG lensing galaxy at $z = 0.4$ (Zahedy et al. 2017b). In contrast, the mean N_{H} and Σ_{cool} at $d = 100\text{--}160$ kpc are significantly lower, $\log \langle N_{\text{H}} \rangle / \text{cm}^{-2} = 18.7_{-0.3}^{+0.2}$ and $\langle \Sigma_{\text{cool}} \rangle \approx (0.06 \pm 0.03) M_{\odot} \text{pc}^{-2}$.

To gain insights into the observed cool gas surface mass density profile in LRG haloes, we compare the data with different analytic functions to obtain a best-fitting model that characterizes the relationship between Σ_{cool} and d . We first consider a simple power law in d , which has been used to describe the spatial

distributions of HI and metal equivalent widths in the cool CGM (e.g. Chen et al. 2001; 2010), and find that it cannot reproduce the rapid decline of Σ_{cool} with increasing d . In contrast, we find that the steepness of the Σ_{cool} profile is well fitted by an exponential model in 2D, $\Sigma_{\text{cool}} = \Sigma_0 e^{-d/d_s}$, with best-fitting parameters of $d_s = (27 \pm 4)$ kpc and $\Sigma_0 = (4.1 \pm 1.4) M_{\odot} \text{pc}^{-2}$ determined from a likelihood analysis (Fig. 11, solid line).

Next, to investigate whether cool baryons follows the large-scale dark-matter mass distribution in the halo, we compare the Σ_{cool} profile in Fig. 11 to the projected surface density of spherically symmetric functions commonly used to describe dark-matter mass distributions, including the Einasto and NFW profiles. The Einasto profile (Einasto 1965) is defined by a power-law logarithmic slope, $d \ln \rho / d \ln r = -2 (r/r_s)^\alpha$, in which the scale radius r_s and shape parameter α are free parameters governing its shape. Relatively shallow Einasto profiles with $\alpha \lesssim 0.3$ have been found to produce good fits to the mass distribution of simulated dark-matter haloes (e.g. Hayashi & White 2008; Gao et al. 2008; Dutton & Macciò 2014). By fitting a projected Einasto profile to our data, we find that Σ_{cool} requires a steep Einasto profile with $\alpha = 1.0^{+0.6}_{-0.2}$ and $r_s = 48^{+19}_{-8}$ kpc (Fig. 11, dotted curve). Note that α and r_s is degenerate in a way that models with larger r_s would require still higher values of α in order to fit the observations.

Because an Einasto profile with $\alpha = 1.0$ is equivalent to an exponential profile in 3D, this exercise demonstrates that reproducing the observed Σ_{cool} requires an underlying density profile that is exponentially declining with radius. In contrast, neither a projected NFW profile (Navarro, Frenk & White 1997) with $r_s = 80$ kpc (expected for $M_h \approx 10^{13} M_{\odot}$ haloes at $z = 0.5$, e.g. Dutton & Macciò 2014) nor a shallow Einasto profile with $\alpha < 0.3$ produces a good fit to the data. As illustrated in Fig. 11 (dashed and dashed-dotted curves), these dark-matter-like profiles can be ruled out because they cannot reproduce the sharp decline of Σ_{cool} with d . Therefore, it appears that the mass distribution of cool gas in the CGM of LRGs is different from the expected mass distribution of the underlying dark-matter halo.

The estimated total mass in cool CGM gas within $d = 160$ kpc ($\sim 0.3 R_h$) of LRGs is

$$M_{\text{cool}}(< 160 \text{ kpc}) = 1.5^{+0.7}_{-0.3} \times 10^{10} M_{\odot}, \quad (3)$$

which is obtained by first multiplying the estimated $\langle \Sigma_{\text{cool}} \rangle$ at $d < 100$ and 100–160 kpc by their respective surface areas modulo the covering fraction inferred from Fig. 11 (unity at $d < 100$ kpc and ~ 0.7 at $d = 100$ –160 kpc), and then summing them. We obtain a similar estimate of $M_{\text{cool}} = (1 - 2) \times 10^{10} M_{\odot}$ by integrating the best-fitting exponential model for Σ_{cool} from $d = 0$ to 160 kpc. Note that we choose to limit our mass estimate out to only $d = 160$ kpc in the CGM because it is the largest projected distance probed in the COS-LRG data set. Our estimate above should therefore be considered as a lower limit on the total mass of cool, photoionized gas in massive quiescent haloes. However, note that if we naively adopted the estimated $\langle \Sigma_{\text{cool}} \rangle$ and gas covering fraction at $d = 100$ –160 kpc and extrapolated these values out to $d = 500$ kpc, which is the typical virial radius of $z \sim 0.4$ LRGs, the mass estimate in equation (3) would increase by a factor of three, to $M_{\text{cool}} \approx 4 \times 10^{10} M_{\odot}$.

Our mass estimate demonstrates that despite their quiescent nature, LRGs at $z \sim 0.4$ still host a significant reservoir of cool gas in their circumgalactic space. Furthermore, the estimated cool CGM mass of $\sim 10^{10} M_{\odot}$ is comparable to the inferred total mass in the cool CGM of lower mass and predominantly star-forming L^* galaxies (e.g. Chen et al. 2010; Prochaska et al. 2011; Werk et al.




Figure 12. Comparison between the surface mass density profiles of the cool and hot CGM of LRGs. Symbols are the same as those in Fig. 11, with the solid black line showing the best-fitting exponential model describing the relationship of Σ_{cool} with d . In contrast, the dotted curve shows the inferred surface mass density profile in the hot CGM of LRG-sized haloes ($M_h \approx 10^{13} M_{\odot}$), based on a combined X-ray and SZ analysis (Singh et al. 2018), with the 68 percent confidence region shaded in grey. Within $d < 160$ kpc from LRGs, we estimate a total cool gas mass of $M_{\text{cool}} = 1.5^{+0.7}_{-0.3} \times 10^{10} M_{\odot}$, which is ~ 6 –13 per cent of the expected total mass in the hot CGM.

2014; Stern et al. 2016). At the same time, the total baryon mass budget of the typical LRG in our sample is $\approx 1.6 \times 10^{12} M_{\odot}$ within the virial radius, which is estimated for the median halo mass in COS-LRG, $M_h \approx 10^{13} M_{\odot}$, by adopting a baryon-to-dark-matter mass ratio of $\Omega_b/\Omega_{\text{DM}} = 0.16$. Thus, the inferred M_{cool} in the CGM is at most ≈ 3 per cent of the total baryon budget for typical LRGs.

It is also interesting to compare the total mass contained in cool clumps to the expected total mass of the hot CGM. The dotted curve in Fig. 12 represents Σ_{hot} , the inferred hot gas surface mass density profile in massive $M_h \sim 10^{13} M_{\odot}$ haloes. Σ_{hot} is computed from the hot gas density profile shown in Fig. 10 (Singh et al. 2018), assuming a unity volume filling fraction for the hot gas. The expected spatial mass profile of hot gas is more spatially extended than the observed Σ_{cool} profile (see also Liang, Kravtsov & Agertz 2016). Within $d = 160$ kpc from LRGs, we infer a total hot gas mass of $M_{\text{hot}}(< 160 \text{ kpc}) = (1.7 \pm 0.5) \times 10^{11} M_{\odot}$. Comparing our estimate of M_{cool} with the inferred M_{hot} , the cool-to-hot gas mass ratio in the CGM of LRGs is

$$X_{\text{cool}} \equiv M_{\text{cool}}/M_{\text{hot}} \approx 0.06 - 0.13 \quad (4)$$

at $d < 160$ kpc, which is comparable to the inferred X_{cool} in the interstellar medium (ISM) of one of these massive ellipticals (Zahedy et al. 2017b).

Furthermore, our data also hint at a declining X_{cool} with increasing projected distance from LRGs, from $X_{\text{cool}} \sim 0.1$ –0.2 at $d < 100$ kpc, to no more than $X_{\text{cool}} \sim 0.01$ –0.03 at $d = 100$ –160 kpc. The declining X_{cool} with increasing projected distance implies that the volume filling factor of cool gas is significantly lower in the outer CGM, at galactocentric radius $r \gtrsim 100$ kpc, than it is in the inner CGM, at $r \lesssim 100$ kpc. The mean volume filling factor of cool gas can be estimated using line-of-sight observables according to the following

expression (e.g. McCourt et al. 2018),

$$\langle f_V \rangle = \langle N_{\text{cl}} \rangle \times \frac{l}{L}, \quad (5)$$

where $l \equiv N_H/n_H$ is the clump thickness along the line of sight, L is the path length through the CGM, and $\langle N_{\text{cl}} \rangle$ is the mean number of clumps per line of sight. As shown in the right-hand panel of Fig. 9, the inferred clump sizes in COS-LRG range from 10 pc to 10 kpc, with a median value and mode of $l \sim 100$ pc. For the purpose of this calculation, we approximate $\langle N_{\text{cl}} \rangle$ to be the average number of discrete components identified per sightline. Based on our data, there are on average $\langle N_{\text{cl}} \rangle_{<100} = 3.7^{+0.6}_{-0.4}$ discrete components at $d < 100$ kpc, which subsequently declines to $\langle N_{\text{cl}} \rangle_{>100} = 2.2^{+0.9}_{-0.5}$ at $d > 100$ kpc. Given that $\langle N_{\text{cl}} \rangle_{<100}$ has contributions from both the inner and outer parts of the halo, we can solve for $\langle f_V \rangle$ in the inner and outer CGM separately using the following approximations:

$$\langle f_V \rangle_{\text{inner}} \approx (\langle N_{\text{cl}} \rangle_{<100} - \langle N_{\text{cl}} \rangle_{>100}) \times \frac{l}{L_{\text{inner}}} \quad (6)$$

$$\langle f_V \rangle_{\text{outer}} \approx \langle N_{\text{cl}} \rangle_{>100} \times \frac{l}{L_{\text{outer}}}. \quad (7)$$

By plugging the different quantities above to equations (6) and (7) and adopting $L_{\text{inner}} = 100$ and 500 kpc, we estimate that the mean volume filling factor for typical clumps with $l = 100$ pc is $\langle f_V \rangle_{\text{inner}} \sim 2 \times 10^{-3}$ in the inner halo ($r \lesssim 100$ kpc), and $\langle f_V \rangle_{\text{outer}} \sim 4 \times 10^{-4}$ in the outer ($r \gtrsim 100$ kpc) halo of LRGs. This exercise illustrates that while the cool gas covering fraction in the CGM of LRGs is high, the volume filling factor can remain very low (for possible theoretical explanations, see e.g. McCourt et al. 2018; Liang & Remming 2018)

5.2 On the origin and fate of cool gas in LRG haloes

In the previous section, we show that despite their ‘red and dead’ nature, LRGs at $z \sim 0.4$ harbour as much as $\sim 10^{10} M_\odot$ of photoionized $T \sim 10^4$ K gas in their extended haloes. This massive reservoir of cool gas appears to consist of compact clumps with a characteristic size of ~ 100 pc (Section 4.3.2), which are pressure confined by the hot gaseous halo that is expected to be ubiquitous around LRGs.

To gain a better understanding of the nature of the cool gas around LRGs, we now consider our observational results in the larger context of a multiphase gaseous halo around LRGs.

The physical formalism for a two-phase CGM was first explored by Mo & Miralda-Escudé (1996), who argued that QSO absorption systems in the vicinity of galaxies originate in cool clouds which are in thermal pressure equilibrium with the hot halo. To explain the formation of cool clumps within an otherwise hot corona, Maller & Bullock (2004) elaborated on this simple model by incorporating multiphase cooling in the halo. In their analytic model, cool clumps originate from condensation in a hydrostatically stable hot halo, triggered by thermal instabilities which develop locally when the cooling time (τ_{cool}) is comparable to the dynamical timescale (τ_{ff}) of the gas. Building on these earlier works, more recent numerical simulations have shown that a multiphase halo can form as soon as $\tau_{\text{cool}}/\tau_{\text{ff}} \lesssim 10$ (e.g. Sharma et al. 2012; McCourt et al. 2012), which is consistent with observations of multiphase gas in a number of nearby galaxy clusters and elliptical galaxies (e.g. Voit et al. 2015; Voit & Donahue 2015).

Under the multiphase-cooling paradigm, cool clumps form within the cooling radius, R_c inside the halo, where thermal instability is prone to develop. For LRG-sized haloes, R_c is estimated to be

between 100 and 200 kpc (Maller & Bullock 2004, equation 18), which is qualitatively consistent with a number of COS-LRG findings, including the observed decline in H I covering fraction with d (Paper I and Section 4.1) and the steep drop in inferred cool gas surface mass density and volume filling factor at $d \gtrsim 100$ kpc (Section 5.1).

In the absence of vigorous star formation activity capable of driving large-scale outflows, circumgalactic cool gas is likely falling towards the centre of the halo. The infall interpretation is supported by the observed line-of-sight velocity dispersion of individual absorbing components in the COS-LRG sample, $\sigma_{\text{gas}} \approx 150 \text{ km s}^{-1}$, which is merely ~ 60 per cent of what is expected from virial motion (see also Huang et al. 2016; Lan & Mo 2018). The observed narrow distribution of line-of-sight velocities indicates that dissipative processes are effective in slowing down cool clumps as they undergo orbital motions in the halo. By attributing the observed deceleration as due to ram pressure drag exerted by the hot halo, Huang et al. (2016) calculated an upper limit on the cool clump mass of $m_{\text{cl}} \lesssim 10^4 M_\odot$ in LRG haloes. This dissipative interaction with the hot gas would lead to orbital decay, causing cool clumps to fall towards the galaxy. But does the cool gas survive this inward journey?

The survival of cool clumps depends on whether the infall time is sufficiently short compared to the timescales of cloud disruption processes acting on them. Cloud destruction is driven predominantly by thermal conduction between cool clumps and the surrounding hot gas. If cool clumps are not sufficiently massive, they will not only decelerate due to ram-pressure drag, but also evaporate due to thermal conduction before reaching the LRG at the centre of the halo.

We expect cool clumps to eventually reach terminal speed when the ram-pressure drag force exerted by the hot gas is balanced by the gravitational pull of the halo on the clump. By identifying this terminal speed with the observed $\sigma_{\text{gas}} \sim 0.6 \sigma_{\text{vir}}$ in LRG absorbers, we can compute the typical cool clump mass (Maller & Bullock 2004, equation 39),

$$m_{\text{cl}} \approx 7.7 \times 10^2 T_6^{-3/8} (\Lambda_Z t_8)^{1/2} M_\odot, \quad (8)$$

where $T_6 = T/10^6$ K is the temperature of the hot corona, Λ_Z is a cooling parameter which depends on the gas metallicity, and $t_8 = t_f/8$ Gyr is the halo formation timescale. For typical LRG haloes in our sample, $T \sim 6 \times 10^6$ K assuming an isothermal gas, and $t_f \sim 9$ Gyr assuming t_f is comparable to the age of the Universe at $z \sim 0.4$. Using equation (8), we find that $m_{\text{cl}} = (2 - 8) \times 10^2 M_\odot$ for metallicities of between 0.01 solar and solar, respectively. This kinematics-based mass estimate can be compared to the cool clump mass independently constrained from our ionization analysis. Based on a combined bootstrap and Monte Carlo resampling of the full range of inferred cool gas densities and characteristic clump sizes (Section 4.3.2), we estimate that the characteristic clump mass has a median value of $\langle m_{\text{cl}} \rangle = 50 - 1000 M_\odot$, which is consistent with the mass range estimated using equation (8).

Given a mass of cool clump m_{cl} , the characteristic for cloud evaporation due to thermal conduction is given by (Maller & Bullock 2004, equation 35),

$$\tau_{\text{evap}} \approx 1.6 m_{\text{cl}}^{2/3} T_6^{-3/2} (\Lambda_Z t_8)^{-1/3} \text{ Myr}. \quad (9)$$

For typical $m_{\text{cl}} \sim 10^2 - 10^3 M_\odot$ and a metallicity of between 0.01 solar and solar, we find that the evaporation timescale is $\tau_{\text{evap}} \sim 1 - 20$ Myr. The expected evaporation time for typical cool clumps is vastly shorter than the minimum infall time of $\tau_{\text{infall}} \sim 200 - 500$ Myr estimated for cool clumps which condense from the hot gas at

$R_c \sim 100\text{--}200$ kpc. The evaporation time is still significantly less than infall time even for clumps as massive as $m_{cl} \sim 10^4 M_\odot$ (Huang et al. 2016). This exercise suggests that cool clumps travel only a relatively small distance in the halo during their lifetimes, and a majority of clumps originating at large distances will evaporate before reaching the centre of the halo.

The implication that a majority of cool clumps in the gaseous halo of LRGs never reaches the central galaxy could explain a number of observational findings that the cool ISM mass in massive quiescent galaxies remains low, $M_{cool}(ISM) \sim 10^{8\text{--}9} M_\odot$ (e.g. Serra et al. 2012; Zahedy et al. 2017b; Young et al. 2018), despite the existence of a much larger reservoir of cool gas in the halo, $M_{cool}(CGM) \sim 10^{10} M_\odot$. At the same time, the fact that cool gas is routinely observed in the gaseous halo of LRGs suggests that cool clumps are continuously formed and destroyed in the predominantly hot gaseous halo. In this quasi-steady state, $\sim 5\text{--}10$ per cent of the CGM gas by mass resides in cool, $\sim 10^4$ K phase at any given time, a balance which is most likely determined by the amount of additional heating available to offset the increased cooling rate from the cool gas.

Finally, we note although our discussion above is based entirely on considering our observations in the context of thermal instability in a multiphase CGM, it does not exclude the possibility that cool gas in LRG haloes is also generated by other physical processes. These additional mechanisms include cool gas recently accreted from the IGM along filaments (e.g. Churchill et al. 2012; Huang et al. 2016), gas originating in and/or stripped from the CGM or ISM of satellite galaxies (e.g. Gauthier et al. 2010; Huang et al. 2016), and gas ejected by SNe Ia (e.g. Zahedy et al. 2016; 2017a). Indeed, our finding that gas density and metallicity can vary by more than a factor of 10 within individual LRG haloes indicates that the CGM is a multiphase mixture of gas with different chemical enrichment histories, which hints at multiple origins of the cool gas. However, cool clumps in LRG haloes are subject to the same interactions with the hot gas regardless of their physical origin. Therefore, our conclusion above can be applied generally on the nature of cool gas in massive quiescent haloes.

5.3 The nature of O VI absorbers in the CGM: insight from massive haloes

A significant finding in CGM studies over the past decade is the ubiquitous presence of strong O VI absorption with $\log N(\text{O VI})/\text{cm}^{-2} \sim 14.5$ around $\sim L^*$ star-forming galaxies (e.g. Tumlinson et al. 2011). At the same time, O VI-bearing gas is found to be less prevalent in the gaseous haloes of passive galaxies (e.g. Chen & Mulchaey 2009; Tumlinson et al. 2011; Johnson et al. 2015). The apparent dichotomy between O VI absorption properties around late- and early-type galaxies is often ascribed to a direct link between star formation and the observed warm gas properties: recent star formation drive powerful outflows that eject metals to large distances in the CGM. Alternatively, the lower incidence of strong O VI absorption in passive galaxies has been attributed to further ionization of oxygen to higher states (e.g. O VII and O VIII) in the more massive and hotter haloes of passive galaxies (e.g. Oppenheimer et al. 2016).

To gain new insights into the nature of O VI absorbers around galaxies, it is necessary to compare the observed O VI absorption properties around galaxies of different masses. In Fig. 13, we present current observational constraints on circumgalactic O VI absorption spanning over more than three decades in galaxy stellar mass, from $\log M_{star}/M_\odot \sim 8$ to > 11 . O VI measurements for massive




Figure 13. Observational constraints on CGM O VI absorption spanning more than three decades in galaxy stellar mass. Constraints for massive quiescent galaxies are from COS-LRG. Constraints for $\sim L^*$ star-forming galaxies are from Johnson et al. (2015), whereas those for star-forming dwarf galaxies are adopted from Johnson et al. (2017). Top: the mean covering fraction of O VI plotted versus M_{star} at $d < 160$ kpc, for a column density threshold of $\log N(\text{O VI})/\text{cm}^{-2} > 13.5$. For each sample, the median stellar mass is plotted, with the horizontal error bars showing the sample dispersion. The corresponding halo mass for each point is indicated as well, based on the Kravtsov et al. (2018) stellar-to-halo-mass relation. The vertical error bars are calculated assuming binomial statistics. Bottom: the mean and median $N(\text{O VI})$ at $d < 160$ kpc plotted as a function of M_{star} . The vertical error bars represent the 68 per cent confidence intervals for the mean and median $N(\text{O VI})$, calculated using a combined bootstrap and Monte Carlo resampling. For comparison, in grey circles, we plot the mean $N(\text{O VI})$ within $d < 150$ kpc from simulated galaxies in EAGLE zoom simulations (Oppenheimer et al. 2017).

quiescent galaxies are from COS-LRG sample, whereas constraints for $\sim L^*$ star-forming galaxies are from Johnson et al. (2015) and those for star-forming dwarf galaxies are adopted from Johnson et al. (2017). The mean covering fraction of O VI at $d < 160$ kpc is plotted versus M_{star} in the top panel of Fig. 13, for a column density threshold of $\log N(\text{O VI})/\text{cm}^{-2} > 13.5$. In contrast to the near-unity covering fraction of O VI absorbers around L^* star-forming galaxies, passive LRGs, and star-forming dwarf galaxies exhibit lower covering fraction of O VI gas, at $\sim 50\text{--}60$ per cent. While star-formation-driven winds is an attractive scenario to account for the ubiquity of O VI absorption around $\sim L^*$ star-forming galaxies, it does not explain the lower O VI covering fraction around star-forming dwarfs. Furthermore, despite the likely absence of strong outflows in LRGs, they still exhibit a significant incidence of O VI, not to mention comparable covering fractions of lower ionization metals to what have been observed around star-forming galaxies (Paper I).

In the bottom panel of Fig. 13, the mean and median $N(\text{O VI})$ at $d < 160$ kpc are plotted versus M_{star} for the three galaxy samples. It is clear that CGM O VI absorption strength peaks in $\sim L^*$ star-forming galaxies (where $M_{\text{h}} = 10^{11.3-12.1} M_{\odot}$). The mean $N(\text{O VI})$ declines towards both lower and higher mass haloes, where a majority of O VI absorbers around LRGs and star-forming dwarf galaxies have $N(\text{O VI})$ which are 0.5–1 dex lower than typical strong O VI absorbers around $\sim L^*$ galaxies. We note that a similar trend of $N(\text{O VI})$ with galaxy mass (or halo mass) is also present in simulated galaxies from the EAGLE zoom simulations (grey points; Oppenheimer et al. 2017), despite the fact that the predicted $N(\text{O VI})$ are systematically lower than observations over the range of galaxy masses considered (see also Nelson et al. 2018).

The strong dependence of circumgalactic $N(\text{O VI})$ on stellar mass, which in turn correlates with the total halo mass including dark matter, hints at a connection between the dominant ionization state of oxygen and the virial temperature of the halo. In particular, the inferred halo virial temperature for the $\sim L^*$ star-forming galaxy sample is $T_{\text{vir}} = 10^{5.3-5.8}$ K, which is coincident with the narrow range of temperatures where the fractional abundance of O VI is at a maximum in collisionally ionized gas (e.g. Heckman et al. 2002; Gnat & Sternberg 2007; Oppenheimer & Schaye 2013). In contrast, the expected virial temperatures for LRGs ($M_{\text{h}} = 10^{12.6-13.5} M_{\odot}$) and dwarf galaxies ($M_{\text{h}} < 10^{11} M_{\odot}$) are $T_{\text{vir}} = 10^{6.5-7.0}$ and $\lesssim 10^5$ K, respectively. At these temperatures, the expected O VI ionization fractions are very small (< 0.01) under collisional ionization models. The observed peak of $N(\text{O VI})$ versus galaxy mass relation in $\sim L^*$ star-forming haloes supports the interpretation that the high columns of O VI around these galaxies originate in collisionally ionized gas at $T \sim T_{\text{vir}}$ (e.g. Oppenheimer et al. 2016; Werk et al. 2016) or, perhaps more realistically, a gas that follows a temperature distribution centred at T_{vir} (McQuinn & Werk 2018).

For O VI absorbers originating in a radiatively cooling flow of coronal ($T \sim 10^{5.5}$ K) gas, $N(\text{O VI})$ is expected to be related to the flow velocity (e.g. Edgar & Chevalier 1986; Heckman et al. 2002; Bordoloi et al. 2017). A cooling flow develops because as O VI-bearing gas cools in the halo, its density must increase to maintain pressure equilibrium. Consequently, the cooling gas sinks and flow inward. Because bulk motion in the gas broadens its line profile, the observed O VI linewidth is a combination of pure thermal broadening and additional broadening due to cooling-flow velocity. By investigating O VI absorbers in a wide range of environments (Galactic disc and high velocity clouds, the Large and Small Magellanic Clouds, nearby starburst galaxies, and the IGM), Heckman et al. (2002) found a correlation between O VI column density and linewidth that is consistent with the theoretical prediction from the radiative cooling flow model. Later studies have also reported similar trends at both low and high redshifts (e.g. Tripp et al. 2008; Lehner et al. 2014; Werk et al. 2016).

To investigate whether O VI absorbers around LRGs can be explained by a radiatively cooling flow, we plot the observed Doppler linewidth versus column density for COS-LRG O VI absorbers in the left-hand panel of Fig. 14 (red circles). Additional O VI absorbers detected in the vicinity of massive quiescent galaxies ($\log M_{\text{star}}/M_{\odot} > 11$) in Johnson et al. (2015) are shown in black circles. For comparison, O VI measurements around $\sim L^*$ star-forming galaxies from Johnson et al. (2015) are shown in pale grey squares. Finally, the predicted relationship between O VI linewidth and column density for a radiatively cooling flow is shown in solid curve, for temperature $T_{\text{O VI}} = 10^{5.5}$ K. We note that if the gas is radiatively

cooling at a higher/lower temperature, the effect is to shift the prediction curve upward/downward in the parameter space (see e.g. Bordoloi et al. 2017).

It is apparent from the left-hand panel of Fig. 14 that O VI absorbers around $\sim L^*$ star-forming galaxies follow the trend predicted by the cooling flow model. This is consistent with the finding of Werk et al. (2016), who reported a statistically significant correlation between $b_{\text{c}}(\text{O VI})$ and $N_{\text{c}}(\text{O VI})$. For COS-LRG O VI absorbers, a Spearman test on the sample indicates a 2.3σ correlation between $b_{\text{c}}(\text{O VI})$ and $N_{\text{c}}(\text{O VI})$, with a coefficient of $r = 0.73$. While this marginal correlation is suggestive a cooling flow, note that most O VI absorbers around LRGs are situated above the prediction curve for a $T_{\text{O VI}} = 10^{5.5}$ K cooling flow. These vertical displacements imply that if O VI absorbers in LRG haloes trace collisionally ionized gas in a radiatively cooling flow, the gas has to be significantly hotter with $T_{\text{O VI}} \approx 10^6$ K (Bordoloi et al. 2017).

On a superficial level, the existence of a 10^6 K cooling gas may not be that surprising given the expectation that LRGs are surrounded by a hot gaseous halo with $T \sim T_{\text{vir}}$. However, the expected O VI ionization fraction in a 10^6 K gas is very low ($\sim 10^{-3}$) under collisional ionization models (e.g. Gnat & Sternberg 2007; Oppenheimer & Schaye 2013). For a solar metallicity gas at $T = 10^6$ K and density of $\log n_{\text{H}}/\text{cm}^{-3} = -4$, the implied cloud thickness for a $\log N(\text{O VI})/\text{cm}^{-2} = 14$ absorber is in excess of 200 kpc. The absorber size would be even larger for a lower metallicity and/or lower density gas, exceeding the size of typical LRG haloes. For that reason, we consider it unlikely that O VI absorbers in LRG haloes originate in a 10^6 K cooling gas.

Alternatively, we consider the possibility that O VI absorbers around LRGs trace cooler, photoionized gas. The expected O VI thermal linewidth for a $T \approx 10^{4.5}$ K gas is 6 km s^{-1} , which is significantly smaller than the observed O VI linewidths in COS-LRG, $b_{\text{c}}(\text{O VI}) = 20 - 100 \text{ km s}^{-1}$ (Fig. 14). If these O VI absorbers originate in a photoionized gas, then their broad-line profiles are predominantly due to non-thermal motions. At the same time, the implied non-thermal broadening of $b_{\text{nt}} = 20 - 100 \text{ km s}^{-1}$ for O VI gas is significantly higher than the modest non-thermal line broadening seen in lower ionization gas around LRGs, $\langle b_{\text{nt}} \rangle = 7 \pm 5 \text{ km s}^{-1}$ (Section 4.2). Because of this large discrepancy in the implied non-thermal motion and observed kinematic misalignments between O VI and H I as well as lower ionization species (Section 4.2), we conclude that any photoionized O VI gas has a different physical origin from cool gas traced by H I and lower ions (Section 5.2).

In a recent study, Stern et al. (2018) considered the possibility that circumgalactic O VI absorbers trace infalling cool gas which has yet to be virially shocked by the halo. Assuming photoionization and thermal equilibrium with the UVB, the implied absorber size is $\sim a \text{ few} \times 10 \text{ kpc}$ for a gas with $\log N(\text{O VI})/\text{cm}^{-2} = 14$ and a metallicity between 0.1–1 solar (e.g. Oppenheimer & Schaye 2013). Because of the substantial size of the absorber, bulk gravitational infall will produce a velocity shear which broadens the O VI line profile. This gravitational line broadening is expected to grow with increasing absorber size. Because absorber size is proportional to column density for a fixed gas density, a correlation between O VI linewidth and column density is naturally expected. In the right-hand panel of Fig. 14, we show the relationships between the two variables as predicted by Stern et al. (2018) for different densities of O^{+5} ions, $n_{\text{O VI}}$.

Under the gravitational broadening scenario, the observed correlation between $b_{\text{c}}(\text{O VI})$ and $N_{\text{c}}(\text{O VI})$ implies that O VI absorbers around L^* star-forming galaxies have densities of $-9 \lesssim$




Figure 14. O VI component linewidth versus column density in the COS-LRG sample (red circles). The dashed–dotted line in the left-hand panel represents the typical 3σ detection limit in the COS spectra for our sample. O VI absorbers detected in the vicinity of massive quiescent galaxies ($\log M_{\text{star}}/M_{\odot} > 11$) in Johnson et al. (2015) are shown in black circles, whereas absorbers around $\sim L^*$ star-forming galaxies in Johnson et al. (2015) are shown in pale grey squares. The solid curve in the left-hand panel shows the expected behaviour for a radiatively cooling, collisionally ionized gas at $T = 10^{5.5}$ K (Heckman et al. 2002), whereas the dotted horizontal lines indicated the expected thermal broadening at different temperatures. In the right-hand panel, the curves show expected behaviours for a gravitationally broadened O VI-bearing gas (Stern et al. 2018) at different O^{5+} volume densities, calculated for a $T = 10^{4.5}$ K gas in dashed curves and $T = 10^{5.5}$ K in solid curves.

$\log n_{\text{OVI}}/\text{cm}^{-3} \lesssim -8.5$. In contrast, the implied O VI volume density is significantly lower for most O VI absorbers detected around massive quiescent galaxies in both COS-LRG (red circles) and Johnson et al. (2015, black circles) samples, $\log n_{\text{OVI}}/\text{cm}^{-3} \lesssim -9.5$. For a photoionized O VI-bearing gas with a metallicity of $[\text{M}/\text{H}] = -0.7$, which is the median metallicity of lower ionization gas in COS-LRG, the implied upper limit on n_{OVI} corresponds to an upper limit on gas density of $\log n_{\text{H}}/\text{cm}^{-3} \lesssim -4.3$. If one assumes that gas density monotonically declines with increasing galactocentric distance, the lower O VI volume densities around LRGs suggest that these absorbers trace gas which resides at larger distances than typical O VI absorbers around L^* star-forming galaxies. This interpretation is consistent with our understanding that LRG haloes are roughly twice the size of L^* star-forming haloes, and that stable accretion shocks in LRG haloes are expected to be situated further out from the galaxies than accretion shocks in lower mass haloes.

The inferred low density of O VI-bearing gas is also consistent with the lack of detection of NV absorption associated with O VI absorbers in COS-LRG. Coverage of the NV doublet is available for seven out of nine high-ionization absorption components detected in O VI. We do not detect NV absorption associated with any of these O VI absorbers. The typical upper limit on the NV to O VI column density ratio in COS-LRG is $\log N_{\text{c}}(\text{NV})/N_{\text{c}}(\text{O VI}) < -0.4$, estimated from the error array by assuming that NV has the same linewidth as O VI. For a photoionized gas with solar N/O elemental abundance ratio, this upper limit constrains the gas density of O VI-

bearing gas to $\log n_{\text{H}}/\text{cm}^{-3} < -4.1$ under the HM05 UVB.⁶ At these low densities, the gas is highly ionized and little associated absorption is expected from low-ionization states. For instance, in a photoionized gas with $\log n_{\text{H}}/\text{cm}^{-3} \approx -5$ and a typical O VI column of $\log N(\text{O VI})/\text{cm}^{-2} = 14$, the expected column densities in C III and Si III are very low, $\log N/\text{cm}^{-2} < 12$, which is consistent with the lack of lower ionization gas observed to be associated with O VI absorbers in COS-LRG. Therefore, while the current sample of O VI absorbers around LRGs is still small, our observations are suggestive of a physical picture where O VI absorbers around LRGs trace photoionized and low-density gas at large distances from the galaxy (see also Voit et al. in preparation).

6 SUMMARY AND CONCLUSIONS

We carried out a systematic investigation of the physical conditions and elemental abundances in the CGM within $d < 160$ kpc from LRGs. The COS-LRG sample comprises 16 LRGs with $\log M_{\text{star}}/M_{\odot} > 11$ at $z = 0.21\text{--}0.55$, which were selected without prior knowledge of the presence or absence of any CGM absorption features. The primary objectives of the COS-LRG program are: (1) to probe the bulk of cool gas in LRG haloes by obtaining accurate measurements of $N(\text{H I})$; and (2) to constrain the physical properties and chemical enrichment in massive quiescent haloes by observing

⁵It is possible that some of the broad O VI absorbers shown in Fig. 14 are the result of unresolved blending of multiple, narrow O VI components. If unresolved components were present, they would naturally have narrower b_c and lower $N_{\text{c}}(\text{O VI})$ than the measurements shown, and as a consequence the data points in Fig. 14 would move downward and leftward.

⁶Zahedy et al. (2017a) reported that the outer CGM of quiescent galaxies exhibits α -element enhanced abundance patterns that are similar to what have been observed in the high-redshift IGM and damped Ly α absorbers (DLAs). Sub-solar N/O values of $[\text{N}/\text{O}] \lesssim -1$ have been reported in high-redshift DLAs (e.g. Petitjean, Ledoux & Srianand 2008). Therefore, it is possible that the outskirts of LRG haloes have similarly sub-solar N/O ratios. If O VI absorbers in COS-LRG arise in gas with low $[\text{N}/\text{O}] \approx -1$, the resulting constraint on gas density from the lack of NV would be less sensitive, $\log n_{\text{H}}/\text{cm}^{-3} \lesssim -3$.

different ionic metal transitions that probe a wide range of ionization states. In Paper I, we presented the $N(\text{H I})$ measurements for the sample and reported that LRGs contains widespread chemically enriched gas traced by various metal ions. In this paper, we expanded our investigation with a detailed ionization analysis based on resolved component structures of a suite of absorption transitions, including the full H I Lyman series and multiple low-, intermediate-, and high-ionization metal transitions. Resolving the component structures of the various absorption lines was made possible by the high-resolution *HST*/COS FUV spectra and ground-based echelle optical spectra of the background QSOs. Our main findings are summarized below.

(1) LRGs exhibit enhanced absorption in H I, low-ionization (Mg II), and intermediate-ionization (Si III and C III) metals at projected distances $d \lesssim 100$ kpc, compared to absorption at larger d (Fig. 2).

(2) H I-bearing gas detected around LRGs is predominantly cool, with temperatures of $T < 10^5$ K inferred from the H I linewidths. Using the ratios of Doppler linewidths for matched H I and Mg II components, we find that the gas has a mean temperature and dispersion of $\langle T \rangle = 2.0 \times 10^4$ K and $\sigma_T = 1.4 \times 10^4$ K, with a modest inferred non-thermal broadening of $\langle b_{\text{nt}} \rangle = 7 \pm 5$ km s $^{-1}$ (Fig. 4).

(3) The line-of-sight velocity distribution of individual absorption components relative to the systemic redshift of LRGs can be characterized by a mean and dispersion of $\langle \Delta v_{\text{gas-galaxy}} \rangle = 17$ km s $^{-1}$ and $\sigma_{\Delta v_{\text{gas-galaxy}}} = 147$ km s $^{-1}$ (Fig. 3). The observed radial velocity dispersion is consistent with what have been observed in Mg II absorbers around LRGs using low-resolution data (e.g. Huang et al. 2016), but it is only ~ 60 per cent of what is expected from virial motion.

(4) By considering matched absorbing components and comparing the relative abundances of different ions for each component, we find that the underlying gas density and metallicity can vary by more than a factor of 10 within the gaseous halo of an LRG (left-hand panels of Figs 8 and 9). Such large variations in gas density and metallicity within individual sightlines highlight a complex multiphase structure and poor chemical mixing in the gaseous haloes of LRGs. Moreover, they underscore the importance of resolving the component structures of CGM absorbers using high-resolution absorption spectra, because any information on variations in gas metallicity and density within individual haloes is lost in ionization studies utilizing only the integrated H I and metal column densities along individual sightlines.

(5) Over the full sample, the median metallicity of absorbing components is $\langle [M/H] \rangle = -0.7 \pm 0.2$, with an estimated 16–84 percentile range of $[M/H] = (-1.6, -0.1)$. Metal-poor components with $< 1/10$ solar metallicity are seen in 50 per cent of the LRG haloes, while gas with near- and super-solar metallicity is also common (Fig. 8). Furthermore, we find a significant incidence of optically thick components with very low metallicities: 43_{-22}^{+25} per cent of LLSs in the gaseous haloes of LRGs have metallicities lower than a few per cent solar.

(6) The median gas density for individual components in the COS-LRG sample is $\log \langle n_{\text{H}} \rangle / \text{cm}^{-3} = -2.4 \pm 0.1$, with an estimated 16–84 percentile range of $\log n_{\text{H}} / \text{cm}^{-3} = (-3.0, -1.8)$ cm $^{-3}$. The inferred median gas density implies a median ionization parameter of $\log \langle U \rangle = -3.0 \pm 0.1$ under the HM05 UVB. The data points exhibit a trend of rising gas density with increasing H I column density (Fig. 9, right-hand panel).

(7) We infer a density contrast of ~ 100 between optically thick components and the expected gas densities in the hot CGM (Fig. 10). The inferred density contrast indicates that optically thick gas in the CGM of LRGs is roughly in thermal pressure equilibrium with the hot halo at galactocentric radius $r \sim d$. In contrast, only ~ 40 per cent of optically thin components have densities consistent with thermal pressure equilibrium with the hot halo at $r \sim d$, which implies that a majority of optically thin absorbers occur at larger radii, $r > d$.

(8) Cool clumps in LRG haloes are compact. The inferred clump sizes are between 10 pc and ~ 1 kpc thick, with a mode of ~ 100 pc (Fig. 9, right-hand panel). The estimated median clump size for the sample is $\langle l \rangle = 120_{-40}^{+80}$ pc.

(9) We find that high-ionization O VI and low-ionization species (low-ionization metals and H I) exhibit distinct kinematic structures. The median absolute difference in centroid velocity between O VI components and the nearest low-ionization metal and H I components is 24 km s $^{-1}$, with a full range of from $|\Delta v| = 4$ to 71 km s $^{-1}$ (Fig. 5). Furthermore, the implied non-thermal line broadening for O VI gas is high, $b_{\text{nt}} = 20$ – 100 km s $^{-1}$, significantly higher than the modest non-thermal broadening inferred for lower ionization gas. Such kinematic mismatches highlight different physical origins between high-ionization gas traced by O VI and lower ionization gas traced by other metal ions. Based on the observed relation between O VI column density and linewidth, our data suggest that O VI absorbers around LRGs trace photoionized, low-density gas at large distances from the galaxy (Fig. 14).

(10) We calculate the total surface mass density of cool ($T \sim 10^4$ K) gas in the LRG haloes, Σ_{cool} , by applying estimated ionization fraction corrections to the observed H I column densities. The spatial profile of Σ_{cool} is equally well described by an exponential profile in 2D, $\Sigma_{\text{cool}} = (4.1 \pm 1.4) e^{-(d/27 \pm 4 \text{ kpc})} M_{\odot} \text{pc}^{-2}$, and a steep projected Einasto profile with shape parameter $\alpha = 1.0_{-0.2}^{+0.6}$ and scale radius $r_s = 48_{-8}^{+19}$ kpc, consistent with a true exponential profile in 3D (Fig. 11). On the other hand, a projected NFW profile or shallow Einasto profile with $\alpha < 0.3$ is ruled out because they cannot reproduce the steep decline of Σ_{cool} with d . We conclude that the mass distribution of cool gas in the CGM of LRGs is different from the expected mass distribution of the underlying dark-matter halo.

(11) We estimate that typical LRGs at $z \sim 0.4$ harbour at least $M_{\text{cool}} = (1 - 2) \times 10^{10} M_{\odot}$ of photoionized $T \sim 10^4$ K gas at $d < 160$ kpc in their haloes (or as much as $M_{\text{cool}} \approx 4 \times 10^{10} M_{\odot}$ at $d < 500$ kpc), which is comparable to the estimated cool CGM mass of star-forming L^* galaxies. The inferred cool CGM mass is about ~ 6 – 13 per cent of the expected gas mass in the hot phase of the CGM (Fig. 12).

Considering our observations in the context of a multiphase gaseous halo surrounding LRGs, our findings are consistent with a scenario in which cool clumps condense from the hot halo due to local thermal instabilities. The observed distribution of line-of-sight velocities indicates that ram-pressure drag exerted by the hot halo is effective at dissipating the kinetic energy of cool clumps, causing them to fall towards the galaxy. It is likely that a large majority of cool clumps in the CGM of LRGs are destroyed before reaching the central galaxy, thereby explaining the continuing lack of star formation activity in these galaxies despite the existence of a large reservoir of cool gas in the CGM. Interactions with the hot gas (such as thermal conduction) and/or some form of energetic feedback from the galaxy itself (e.g. heating from stellar winds, SNe Ia, or an active nucleus) likely play an active role in preventing an accumulation of cool gas in the ISM the LRG. Moving forward, a

systematic study of the incidence and physical properties of the cool ISM ($d \lesssim 10$ kpc; see e.g. Zahedy et al. 2017b) of LRGs is necessary to connect the observed plethora of cool gas at $d \sim 100$ kpc scales in the CGM with the continuing ‘red and dead’ nature of LRGs over cosmic time.

ACKNOWLEDGEMENTS

The authors thank the anonymous referee for thoughtful comments that helped improve the presentation of this paper. The authors thank Jonathan Stern, Mark Voit, Ann Zabludoff, and Rongmon Bordoloi for illuminating discussions. We are grateful to Ben Oppenheimer for providing OVI data points for galaxies in the EAGLE zoom simulations. This work is based on data gathered under the 6.5 m Magellan Telescopes located at Las Campanas Observatory, the HST-GO-14145.01A observing program using the NASA/ESA *HST* operated by the Space Telescope Science Institute and the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 5-26555. FSZ and HWC acknowledge partial support from NSF grant AST-1715692. SDJ is supported by a NASA Hubble Fellowship (HST-HF2-51375.001-A). This research has made use of the KOA, which is operated by the W. M. Keck Observatory and the NASA Exoplanet Science Institute, under contract with the National Aeronautics and Space Administration.

REFERENCES

- Bernstein R., Shectman S. A., Gunnels S. M., Mochnacki S., Athey A. E., 2003, *Proc. SPIE*, 4841, 1694
- Bordoloi R., Wagner A. Y., Heckman T. M., Norman C. A., 2017, *ApJ*, 848, 122
- Bowen D. V., Chelouche D., 2011, *ApJ*, 727, 47
- Chen H.-W., 2017, in *Astrophysics and Space Science Library*, Vol. 434, *Outskirts of Galaxies*. Springer International Publishing AG, Switzerland, p. 291
- Chen H.-W., Helsby J. E., Gauthier J.-R., Shectman S. A., Thompson I. B., Tinker J. L., 2010, *ApJ*, 714, 1521
- Chen H.-W., Johnson S. D., Zahedy F. S., Rauch M., Mulchaey J. S., 2017, *ApJ*, 842, L19
- Chen H.-W., Lanzetta K. M., Webb J. K., Barcons X., 1998, *ApJ*, 498, 77
- Chen H.-W., Lanzetta K. M., Webb J. K., Barcons X., 2001, *ApJ*, 559, 564
- Chen H.-W., Mulchaey J. S., 2009, *ApJ*, 701, 1219
- Chen H.-W., Zahedy F. S., Johnson S. D., Pierce R. M., Huang Y.-H., Weiner B. J., Gauthier J.-R., 2018, *MNRAS*, 479, 2547
- Chen H.-W. et al., 2018b, *MNRAS*, 484, 431
- Churchill C. W., Kacprzak G. G., Steidel C. C., Spitler L. R., Holtzman J., Nielsen N. M., Trujillo-Gomez S., 2012, *ApJ*, 760, 68
- Conroy C., van Dokkum P. G., Kravtsov A., 2015, *ApJ*, 803, 77
- Crighton N. H. M., Hennawi J. F., Simcoe R. A., Cooksey K. L., Murphy M. T., Fumagalli M., Prochaska J. X., Shanks T., 2015, *MNRAS*, 446, 18
- De Cia A., Ledoux C., Mattsson L., Petitjean P., Srianand R., Gavignaud I., Jenkins E. B., 2016, *A&A*, 596, A97
- Dekel A., Sari R., Ceverino D., 2009, *ApJ*, 703, 785
- Dutton A. A., Macciò A. V., 2014, *MNRAS*, 441, 3359
- Edgar R. J., Chevalier R. A., 1986, *ApJ*, 310, L27
- Einasto J., 1965, *Tr. Astrofiz. Inst. Alma-Ata*, 5, 87
- Eisenstein D. J. et al., 2003, *ApJ*, 585, 694
- Faucher-Giguère C.-A., 2017, in *Astrophysics and Space Science Library*, Vol. 430, *Gas Accretion onto Galaxies*. Springer International Publishing AG, Switzerland, p. 271
- Faucher-Giguère C.-A., Lidz A., Hernquist L., Zaldarriaga M., 2008, *ApJ*, 688, 85
- Ferland G. J. et al., 2013, *Rev. Mex. Astron. Astrofis.*, 49, 137
- Foreman-Mackey D., Hogg D. W., Lang D., Goodman J., 2013, *PASP*, 125, 306
- Gao L., Navarro J. F., Cole S., Frenk C. S., White S. D. M., Springel V., Jenkins A., Neto A. F., 2008, *MNRAS*, 387, 536
- Gauthier J.-R., Chen H.-W., 2011, *MNRAS*, 418, 2730
- Gauthier J.-R., Chen H.-W., Tinker J. L., 2009, *ApJ*, 702, 50
- Gauthier J.-R., Chen H.-W., Tinker J. L., 2010, *ApJ*, 716, 1263
- Gehrels N., 1986, *ApJ*, 303, 336
- Gnat O., Sternberg A., 2007, *ApJS*, 168, 213
- Green J. C. et al., 2012, *ApJ*, 744, 60
- Haardt F., Madau P., 2001, in Neumann D. M., Van J.T.T.eds, *Clusters of Galaxies and the High Redshift Universe Observed in X-rays*, Vol. 64, Savoie, France, p. 64
- Haardt F., Madau P., 2012, *ApJ*, 746, 125
- Hafen Z. et al., 2017, *MNRAS*, 469, 2292
- Hayashi E., White S. D. M., 2008, *MNRAS*, 388, 2
- Heckman T. M., Norman C. A., Strickland D. K., Sembach K. R., 2002, *ApJ*, 577, 691
- Hodge J. A., Zeimann G. R., Becker R. H., White R. L., 2009, *AJ*, 138, 900
- Huang Y.-H., Chen H.-W., Johnson S. D., Weiner B. J., 2016, *MNRAS*, 455, 1713
- Johnson S. D., Chen H.-W., Mulchaey J. S., 2015, *MNRAS*, 449, 3263
- Johnson S. D., Chen H.-W., Mulchaey J. S., Schaye J., Straka L. A., 2017, *ApJ*, 850, L10
- Kereš D., Katz N., Fardal M., Davé R., Weinberg D. H., 2009, *MNRAS*, 395, 160
- Khaire V., Srianand R., 2015, *MNRAS*, 451, L30
- Kollmeier J. A. et al., 2014, *ApJ*, 789, L32
- Kravtsov A. V., Vikhlinin A. A., Meshcheryakov A. V., 2018, *Astron. Lett.*, 44, 8
- Lan T.-W., Mo H., 2018, *ApJ*, 866, 36
- Lehner N., O’Meara J. M., Fox A. J., Howk J. C., Prochaska J. X., Burns V., Armstrong A. A., 2014, *ApJ*, 788, 119
- Lehner N. et al., 2013, *ApJ*, 770, 138
- Liang C. J., Kravtsov A. V., Agertz O., 2016, *MNRAS*, 458, 1164
- Liang C. J., Remming I. S., 2018, preprint ([arXiv:1806.10688](https://arxiv.org/abs/1806.10688))
- Li Y.-P. et al., 2018, *ApJ*, 866, 70
- Lundgren B. F. et al., 2009, *ApJ*, 698, 819
- Maller A. H., Bullock J. S., 2004, *MNRAS*, 355, 694
- Mandelbaum R., Seljak U., Hirata C. M., 2008, *J. Cosmol. Astropart. Phys.*, 8, 006
- McCourt M., Oh S. P., O’Leary R., Madigan A.-M., 2018, *MNRAS*, 473, 5407
- McCourt M., Sharma P., Quataert E., Parrish I. J., 2012, *MNRAS*, 419, 3319
- McQuinn M., Werk J. K., 2018, *ApJ*, 852, 33
- Mo H. J., Miralda-Escudé J., 1996, *ApJ*, 469, 589
- Muzahid S., Fonseca G., Roberts A., Rosenwasser B., Richter P., Narayanan A., Churchill C., Charlton J., 2018, *MNRAS*, 476, 4965
- Navarro J. F., Frenk C. S., White S. D. M., 1997, *ApJ*, 490, 493
- Nelson D., Vogelsberger M., Genel S., Sijacki D., Kereš D., Springel V., Hernquist L., 2013, *MNRAS*, 429, 3353
- Nelson D. et al., 2018, *MNRAS*, 477, 450
- Oppenheimer B. D., Schaye J., 2013, *MNRAS*, 434, 1063
- Oppenheimer B. D., Schaye J., Crain R. A., Werk J. K., Richings A. J., 2017, *MNRAS*, 481, 835
- Oppenheimer B. D. et al., 2016, *MNRAS*, 460, 2157
- Petitjean P., Ledoux C., Srianand R., 2008, *A&A*, 480, 349
- Prochaska J. X., Weiner B., Chen H.-W., Mulchaey J., Cooksey K., 2011, *ApJ*, 740, 91
- Prochaska J. X. et al., 2017, *ApJ*, 837, 169
- Putman M. E., Peek J. E. G., Jong M. R., 2012, *ARA&A*, 50, 491
- Rauch M., Sargent W. L. W., Barlow T. A., 1999, *ApJ*, 515, 500
- Rauch M., Sargent W. L. W., Barlow T. A., Simcoe R. A., 2002, *ApJ*, 576, 45
- Ribaldo J., Lehner N., Howk J. C., Werk J. K., Tripp T. M., Prochaska J. X., Meiring J. D., Tumlinson J., 2011, *ApJ*, 743, 207
- Roseboom I. G. et al., 2006, *MNRAS*, 373, 349
- Rosenwasser B., Muzahid S., Charlton J. C., Kacprzak G. G., Wakker B. P., Churchill C. W., 2018, *MNRAS*, 476, 2258
- Sadler E. M. et al., 2007, *MNRAS*, 381, 211

- Savage B. D., Sembach K. R., 1996, *ARA&A*, 34, 279
 Savage B. D. et al., 2010, *ApJ*, 719, 1526
 Serra P. et al., 2012, *MNRAS*, 422, 1835
 Sharma P., McCourt M., Quataert E., Parrish I. J., 2012, *MNRAS*, 420, 3174
 Shen S., Madau P., Guedes J., Mayer L., Prochaska J. X., Wadsley J., 2013, *ApJ*, 765, 89
 Shull J. M., Moloney J., Danforth C. W., Tilton E. M., 2015, *ApJ*, 811, 3
 Singh P., Majumdar S., Nath B. B., Silk J., 2018, *MNRAS*, 478, 2909
 Stern J., Faucher-Giguère C.-A., Hennawi J. F., Hafen Z., Johnson S. D., Fielding D., 2018, *ApJ*, 865, 91
 Stern J., Hennawi J. F., Prochaska J. X., Werk J. K., 2016, *ApJ*, 830, 87
 Thom C. et al., 2012, *ApJ*, 758, L41
 Tripp T. M., Sembach K. R., Bowen D. V., Savage B. D., Jenkins E. B., Lehner N., Richter P., 2008, *ApJS*, 177, 39
 Tumlinson J., Peebles M. S., Werk J. K., 2017, *ARA&A*, 55, 389
 Tumlinson J. et al., 2011, *Science*, 334, 948
 van de Voort F., Schaye J., Altay G., Theuns T., 2012, *MNRAS*, 421, 2809
 Viel M., Haehnelt M. G., Bolton J. S., Kim T.-S., Puchwein E., Nasir F., Wakker B. P., 2017, *MNRAS*, 467, L86
 Vogt S. S. et al., 1994, *Proc. SPIE*, 2198, 362
 Voit G. M., Donahue M., 2015, *ApJ*, 799, L1
 Voit G. M., Donahue M., Bryan G. L., McDonald M., 2015, *Nature*, 519, 203
 Werk J. K., Prochaska J. X., Thom C., Tumlinson J., Tripp T. M., O'Meara J. M., Peebles M. S., 2013, *ApJS*, 204, 17
 Werk J. K. et al., 2014, *ApJ*, 792, 8
 Werk J. K. et al., 2016, *ApJ*, 833, 54
 Wotta C. B., Lehner N., Howk J. C., O'Meara J. M., Prochaska J. X., 2016, *ApJ*, 831, 95
 York D. G. et al., 2000, *AJ*, 120, 1579
 Young L. M., Serra P., Krajnović D., Duc P.-A., 2018, *MNRAS*, 477, 2741
 Young L. M. et al., 2014, *MNRAS*, 444, 3408
 Zahedy F. S., Chen H.-W., Gauthier J.-R., Rauch M., 2017a, *MNRAS*, 466, 1071
 Zahedy F. S., Chen H.-W., Rauch M., Wilson M. L., Zabludoff A., 2016, *MNRAS*, 458, 2423
 Zahedy F. S., Chen H.-W., Rauch M., Zabludoff A., 2017b, *ApJ*, 846, L29
 Zhu G. et al., 2014, *MNRAS*, 439, 3139

SUPPORTING INFORMATION

Supplementary data are available at [MNRAS](https://www.mnras.org/) online.

Appendix A. Description of Individual LRG Haloes.

Please note: Oxford University Press is not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.

APPENDIX A: DESCRIPTION OF INDIVIDUAL LRG HALOES

Here we describe the observed absorption properties and discuss the physical conditions (density and temperature) and chemical abundances of individual haloes in the COS-LRG sample. The 16 QSO sightlines in COS-LRG are ordered by increasing projected distance from each LRG. Appendix A1 is shown below for the closest QSO sightline in the sample, whereas sections A2 to A16 are published as online material.

A1 SDSS J0946+5123 at $d = 42$ kpc

This LRG is at redshift $z_{\text{LRG}} = 0.4076$. A LLS with a total $N(\text{H I})$ of $\log N(\text{H I})/\text{cm}^{-2} = 17.34 \pm 0.01$ is present near the redshift of the galaxy (Fig. A1a). In addition, the following ionic metal species

are also detected: C II, C III, N II, O VI, Si II, and Si III.

Based on a combined Voigt profile analysis of H I and the corresponding metal absorption profiles, we identify five components in the absorption system (Fig. A1a and Table A1a). The observed velocity spread of the absorber is $\Delta v \approx 200 \text{ km s}^{-1}$ from the bluest to the reddest component. Most (90 per cent) of the H I column density is in component 2 at $z_{\text{abs}} = 0.40701$, or 126 km s^{-1} blueward of the LRG. Two other components have $\log N_c(\text{H I})/\text{cm}^{-2} \sim 16$, components 3 and 4 at $dv_c = +22$ and $+61 \text{ km s}^{-1}$ from the strongest component, respectively. While the bulk of the neutral hydrogen content is in component 2, little metal absorption is associated with it. In contrast, both low-ionization (e.g. C II and Si II) and intermediate-ionization (e.g. C III) absorption are very prominent in components 3 and 4. This particular characteristic of the absorber suggests a large variation in chemical abundances across different components.

The observed Doppler linewidths of individual H I components ($b_c(\text{H I}) \lesssim 25 \text{ km s}^{-1}$ for all but one components) impose a temperature upper limit of $T \lesssim 4 \times 10^4 \text{ K}$ for the gas, under a purely thermal broadening assumption. The other component, component 5, has a very broad H I linewidth of $b_c(\text{H I}) = 71 \text{ km s}^{-1}$. However, similar linewidths are observed for the corresponding Si III and C III absorption in component, which implies that the gas is cool ($T \sim 10^4 \text{ K}$) and the broad line profile is primarily due to non-thermal motion (e.g. turbulence) or the presence of blended narrow components.

As shown in Fig. A1b and Table A1b, our ionization analysis separates the absorbing gas into two different regimes of gas density. For components 2, 3, and 4, which have $\log N_c(\text{H I})/\text{cm}^{-2} \gtrsim 16$, good agreements between observations and models are achieved for a gas density range of from $\log n_{\text{H}}/\text{cm}^{-3} \approx -2.4$ to $\log n_{\text{H}}/\text{cm}^{-3} \approx -2.1$ under HM05, and from $\log n_{\text{H}}/\text{cm}^{-3} \approx -2.7$ to $\log n_{\text{H}}/\text{cm}^{-3} \approx -2.5$ under HM12. On the other hand, models for lower $N_c(\text{H I})$ components 1 and 5 require lower gas densities to match the data: between $\log n_{\text{H}}/\text{cm}^{-3} \approx -3.6$ and $\log n_{\text{H}}/\text{cm}^{-3} \approx -3.0$ under both HM05 and HM12 UVBs.

Similarly, the CLOUDY photoionization models indicate a large variation in metallicities (>1 dex) across different components. For component 2 at $dv_c = 0 \text{ km s}^{-1}$, which has the highest H I column density in the absorber ($\log N_c(\text{H I})/\text{cm}^{-2} = 17.3$) but exhibits little associated metals, the inferred metallicity is very low with an upper limit of $[\text{M}/\text{H}] \lesssim -1.5$ under both HM05 and HM12 UVBs. In contrast, the observed ionic column densities in components 1, 4, and 5 are consistent with the gas having sub-solar metallicities of between $[\text{M}/\text{H}] \approx -0.8$ and $[\text{M}/\text{H}] \approx -0.4$ under HM05, and between $[\text{M}/\text{H}] \approx -0.3$ and $[\text{M}/\text{H}] \approx 0$ under HM05. Finally, for component 3, which has $\log N(\text{H I})/\text{cm}^{-2} \approx 16$ yet shows the strongest metal absorption, solar or super-solar metallicities are required to match the data, $[\text{M}/\text{H}] = 0.4 \pm 0.4$ under HM05 and $[\text{M}/\text{H}] = 0.7 \pm 0.4$ under HM12.

This absorption system is also noteworthy because it is the strongest O VI absorber in the COS-LRG sample. The O VI absorption profile is kinematically complex, comprising three components that extend over $\sim 300 \text{ km s}^{-1}$ in line-of-sight velocity. The measured total O VI column density is $\log N(\text{O VI})/\text{cm}^{-2} = 14.93 \pm 0.02$, which is the highest yet detected in the vicinity of a passive galaxy (cf., Tumlinson et al. 2011; Johnson et al. 2015), and among the highest $N(\text{O VI})$ seen in both star-forming and passive galaxies. The broad and asymmetric O VI absorption profile is in stark contrast to the narrower absorption profiles of the lower-ionization metals, which indicates different physical origins between the low- and high-ionization species.




Figure A1a. Continuum normalized absorption profiles of different transitions along QSO sightline SDSS J0946+5123 at $d = 42$ kpc from the LRG. The absorption transition is identified in the bottom-right corner of each panel. Zero velocity marks the redshift of the strongest H I absorption component identified in the Voigt profile analysis, $z_{\text{abs}} = 0.40701$. The systemic redshift of the LRG is indicated with a blue dotted line. The $1-\sigma$ error spectrum is included in cyan, above the zero-flux level. Contaminating features have been grayed out for clarity. The best-fit Voigt profiles for each individual transition detected are plotted, both for the sum of all components (red curve) and for individual components (different-colored curves). The centroid of each absorption component is marked by a blue tick mark at the top of panels in the first row.

Table A1a. Absorption properties along QSO sightline SDSS J0946+5123 at $d = 42$ kpc from the LRG.

Component	Species	dv_c (km s^{-1})	$\log N_c$	b_c (km s^{-1})	
all	H I	...	17.34 ± 0.01	...	
	C II	...	14.11 ± 0.16	...	
	C III	...	> 14.73	...	
	N II	...	13.90 ± 0.22	...	
	N V	...	< 13.97	...	
	O VI	...	14.93 ± 0.02	...	
	Si II	...	$13.87^{+0.21}_{-0.14}$...	
	Si III	...	$13.48^{+0.19}_{-0.06}$...	
	Fe II	...	< 14.04	...	
	Fe III	...	< 14.14	...	
1	H I	$-62.8^{+2.8}_{-3.0}$	$14.70^{+0.10}_{-0.09}$	18.2 ± 1.7	
	C II	-62.8	< 13.24	10	
	C III	-62.8	$13.67^{+0.13}_{-0.11}$	27.3 ± 8.9	
	N II	-62.8	< 13.29	10	
	Si II	-62.8	< 12.88	10	
	Si III	-62.8	$12.35^{+0.20}_{-0.23}$	$17.3^{+8.6}_{-6.1}$	
	Fe II	-62.8	< 13.48	10	
	Fe III	-62.8	< 13.50	10	
	2	H I	0.0 ± 0.2	17.30 ± 0.02	$10.6^{+0.7}_{-0.6}$
		C II	0.0	< 13.22	10
C III		0.0	$13.20^{+0.28}_{-0.26}$	$13.4^{+5.6}_{-2.0}$	
N II		0.0	< 13.40	10	
Si II		0.0	< 12.92	10	
Si III		0.0	< 12.08	10	
Fe II		0.0	< 13.47	10	
Fe III		0.0	< 13.55	10	
3		H I	$+22.1^{+2.2}_{-2.8}$	$15.97^{+0.30}_{-0.46}$	$10.8^{+1.0}_{-0.9}$
		C II	$+22.1$	14.11 ± 0.16	$10.4^{+5.0}_{-2.5}$
	C III	$+22.1$	> 13.60	< 22.5	
	N II	$+22.1$	13.90 ± 0.22	$9.1^{+5.8}_{-2.4}$	
	Si II	$+22.1$	$13.80^{+0.22}_{-0.20}$	9.9 ± 4.7	
	Si III	$+22.1$	$12.97^{+0.32}_{-0.26}$	$12.0^{+9.5}_{-2.3}$	
	Fe II	$+22.1$	< 13.52	10	
	Fe III	$+22.1$	< 13.49	10	
	4	H I	$+60.8^{+3.9}_{-4.9}$	$15.90^{+0.13}_{-0.12}$	$26.9^{+2.6}_{-1.9}$
		C II	$+60.8$	< 13.31	10
C III		$+60.8$	> 13.68	< 35.7	
N II		$+60.8$	< 13.27	10	
Si II		$+60.8$	$13.07^{+0.20}_{-0.31}$	$9.7^{+6.4}_{-3.1}$	
Si III		$+60.8$	$13.09^{+0.29}_{-0.19}$	$20.0^{+8.3}_{-4.4}$	
Fe II		$+60.8$	< 13.51	10	
Fe III		$+60.8$	< 13.61	10	
5		H I	$+132.3^{+5.2}_{-6.6}$	$15.41^{+0.09}_{-0.07}$	$71.3^{+3.7}_{-3.1}$
		C II	$+132.3$	< 13.16	10
	C III	$+132.3$	$14.34^{+0.22}_{-0.07}$	$69.8^{+5.3}_{-11.8}$	
	N II	$+132.3$	< 13.37	10	
	Si II	$+132.3$	< 12.73	10	
	Si III	$+132.3$	12.81 ± 0.23	$70.0^{+35.1}_{-19.1}$	
	Fe II	$+132.3$	< 13.49	10	
	Fe III	$+132.3$	< 13.59	10	
	high-1	O VI	-69.7 ± 12.7	$14.00^{+0.08}_{-0.17}$	$53.3^{+23.6}_{-14.2}$
	high-2	N V	-69.7	< 13.67	
O VI		$+64.3 \pm 11.0$	$14.38^{+0.08}_{-0.05}$	$58.2^{+22.0}_{-5.0}$	
high-3	N V	$+64.3$	< 13.67		
	O VI	$+202.6 \pm 11.0$	$14.71^{+0.03}_{-0.04}$	$82.9^{+8.6}_{-4.9}$	
	N V	$+202.6$	< 13.82		




Figure A1b. Probability distribution of gas metallicity and density for the individual absorption components detected along QSO sightline SDSS J0946+5123, at $d = 42$ kpc from the LRG. Each component shown here has at least two ionic metal species detected in absorption. The contour levels indicate areas enclosing estimated 68 per cent and 95 per cent probabilities of the model parameters, shown in black for models assuming the HM05 UVB and in blue for the HM12 UVB (see Section 3.2). Not shown here is component 2 at $dv_c = 0$ km s⁻¹, which has the strongest H I absorption in the absorber, with $\log N_c(\text{H I})/\text{cm}^{-2} = 17.3$, yet shows little metal absorption. The weak C III absorption seen in component 2, along with upper limits on the column density of other ions still allows us to constrain the gas metallicity to $[M/H] \lesssim -1.5$.

Table A1b. Ionization modelling results for the absorber along SDSS J0946+5123 at $d = 42$ kpc from the LRG.

Component	N_{metal}	[M/H]		$\log n_{\text{H}}/\text{cm}^{-3}$	
		HM05	HM12	HM05	HM12
SC	5	-0.84 ± 0.16	$-0.73^{+0.16}_{-0.13}$	$-1.90^{+0.08}_{-0.20}$	$-2.28^{+0.10}_{-0.18}$
1	2	$-0.61^{+0.91}_{-0.12}$	$0.04^{+0.47}_{-0.17}$	$-2.84^{+0.04}_{-0.60}$	$-3.20^{+0.08}_{-0.34}$
2	1	< -1.53	< -1.41	$-2.08^{+0.52}_{-0.18}$	$-2.52^{+0.60}_{-0.20}$
3	5	$0.42^{+0.38}_{-0.29}$	$0.70^{+0.26}_{-0.40}$	$-2.44^{+0.16}_{-0.24}$	$-2.84^{+0.14}_{-0.28}$
4	3	$-0.42^{+0.15}_{-0.24}$	$-0.23^{+0.15}_{-0.20}$	$-2.38^{+0.14}_{-0.46}$	$-2.72^{+0.12}_{-0.40}$
5	2	$-0.81^{+0.28}_{-0.19}$	$-0.20^{+0.20}_{-0.22}$	$-3.50^{+0.42}_{-0.16}$	$-3.58^{+0.30}_{-0.16}$

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.