

Proceedings of the Iowa Academy of Science

Volume 8 | Annual Issue

Article 24

1900

Preliminary List of the Flowering Plants of Adair County

James E. Gow

Copyright ©1900 Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

Recommended Citation

Gow, James E. (1900) "Preliminary List of the Flowering Plants of Adair County," *Proceedings of the Iowa Academy of Science*, 8(1), 152-159.

Available at: <https://scholarworks.uni.edu/pias/vol8/iss1/24>

This Research is brought to you for free and open access by the Iowa Academy of Science at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact scholarworks@uni.edu.

PRELIMINARY LIST OF THE FLOWERING PLANTS OF ADAIR COUNTY.

BY JAMES E. GOW.

The collections on which this report is based were made chiefly during the summer of the year 1900, some of the work, however, having been done some years earlier. It is the hope of the author that he may in the course of time be able to supply a complete account of the flora of the county—one which will be exhaustive to the last detail. Heretofore such an undertaking has not been possible for him. The work has been done in the intervals of other work and has taken into account chiefly the more common species. It is here presented as preliminary to the more complete report which, it is hoped, will follow it. The grasses and sedges have been purposely reserved for a separate report.

The nomenclature used is that of the sixth edition of Gray's Botany. While more recent systems may have good claims to superiority, the nomenclature of Gray is more generally known than any other, and is better understood by the majority of amateur botanists.

RANUNCULACEAE.

Clematis virginiana L. Not rare.

Anemone cylindrica Gray. Very common.*

A. virginiana L. Not rare.

Thalictrum purpurascens L.

Ranunculus acris L. Very abundant in low grounds.

R. abortivus L.

Isopyrum biternatum T. and G.

Aquilegia canadensis L.

Delphinium azureum Ait. Low grounds. Common.

D. exaltatum Ait. Very rare. One specimen in the author's collection is certainly of this species.

*In the case of the more common prairie species no attempt is here made to describe the habitat, or abundance of the species, except in cases where Adair county shows features which are novel and unusual. Most of the species are common and generally known. As a rule, woodland species are noted in the text.

IOWA ACADEMY OF SCIENCES.

153

Delphinium tricorne Michx. Very common in low grounds.

BERBERIDACEAE.

Berberis vulgaris L. Escaped from cultivation.

PAPAVERACEAE.

Sanguinaria canadensis L. Common in woodlands.

FUMARIACEAE.

Dicentra cucullaria DC. Very common in woods.

Corydalis aurea Willd. Not uncommon.

CRUCIFERAE.

Capsella bursa-pastoris (L) Moench.

Lepidium virginicum L.

Sisymbrium officinale (L) Scop.

Brassica nigra (L) Koch.

B. sinapistrum Boiss.

Arabis Canadensis L.

Cardamine hirsuta L.

Nasturtium armoracia (L) Fries.

N. officinale R. Br.

Raphanus sativus L. Escaped from cultivation.

CAPPARIDACEAE.

Polanisia trachysperma T. and G.

VIOLACEAE.

Viola pedata L.

V. blanda Willd. Not common.

V. cucullata Ait.

V. pubescens Ait.

CARYOPHYLLACEAE.

Silene stellata Ait.

S. nocturna L.

POTULACACEAE.

Portulaca oleracea L.

Claytonia virginica L. Common in woodlands.

HYPERICACEAE.

Hypericum ascyron L. Common.

MALVACEAE.

Malva rotundifolia L.

Abutilon avicennae Gaertn. Escaped from cultivation, or introduced in grain seed.

TELIACEAE.

Tilia americana L.

LINACEAE.

Linum usitatissimum. Escaped from cultivation.

L. sulcatum Riddell. Not very common.

IOWA ACADEMY OF SCIENCES.

GERANIACEAE.

Geranium maculatum L.

Oxalis violacea L.

O. stricta L.

RUTACEAE.

Xanthoxylum americanum Mill. Not common. Found on steep bluffs along the course of middle river.

CELASTRACEAE.

Celastrus scandens L.

VITACEAE.

Vitis riparia Michx.

Ampelopsis quinquefolia Mx. Common in timber.

SAPINDACEAE.

Aesculus glabra Willd.

Acer dasycarpum Ehrb.

Negundo aceroides Moench.

ANACARDIACEAE.

Rhus glabra L.

R. toxicodendron L. Rare, in dense timber.

LEGUMINOSAE.

Baptisia leucantha T. & G.

B. leucophea Nutt.

Lupinus perennis L. Probably fugitive from gardens.

Trifolium pratense L.

T. repens L.

T. hybridum L.

Melilotus alba Lam. Quite common, only in the western half of the county, where the roadsides are covered with it.

Medicago sativa L.

Amorpha canescens Nutt.

Petalostemon violaceus Michx.

P. candidus Michx.

Tephrosia virginiana Pers.

Astragalus caryocarpus Ker.

A. cooperi Gray. Not common.

Desmodium acuminatum DC. Common on Middle river near northern boundary of county.

D. rigidum D. C.

Lespedeza capitata Michx.

Amphicarpea monoica Nutt. Tolerably common in woods.

Cassia chamaecrista L. Very abundant.

Gleditschia triacanthos L. Rare.

ROSACEAE.

Prunus Americana Marsh.

P. serotina Ehrh.

IOWA ACADEMY OF SCIENCES.

155

- Prunus virginiana* L.
Geum virginianum L.
Rubus villosus Ait. Escaped from cultivation.
Fragaria vesca L. Escaped from cultivation.
F. virginiana Mill.
Potentilla norvegica L.
P. arguta Pursh.
P. paradoxa.
P. canadensis L. Common in low lands.
Agrimonia eupatoria L. Woodlands.
A. parviflora Ait. Woods.
Crataegus coccinea L.
C. tomentosa L. /
Rosa arkansana Porter.
Pyrus coronaria L.

SAXIFRAGACEAE.

- Ribes gracile* Michx. Common in woodlands, and cultivated.

LYTHRACEAE.

- Lythrum alatum* Pursh. Not very common.

ONAGRACEAE.

- Gaura biennis* L.
Oenothera biennis L.
Circaeae lutetiana L. Not common.

CUCURBITACEAE.

- Echinocystis lobata* Torr & Gray.

UMBELLIFERAE.

- Heracleum lanatum* Mx. Low prairie. Common.
Thaspium barbinode Nutt. Banks of streams.
Sium cicutaefolium Gmelin. Common on lowlands.
Zizia aurea Koch. Common on lowlands.
Cicuta maculata L. Common on lowlands.
Osmorrhiza brevistylis DC. Not uncommon on higher land
than preceding.
O. longistylis D. C. Same habitat as preceding.
Eryngium yuccafolium Mx.

CORNACEAE.

- Cornus paniculata*, L'Her. Low thickets. Only tolerably com-
mon.

CAPRIFOLIACEAE.

- Sambucus canadensis* L.
Lonicera glauca Hill. (?)

COMPOSITAE.

- Vernonia fasciculata* Mx.
Eupatorium ageratoides L. Rather common in woods.

IOWA ACADEMY OF SCIENCES.

- Liatris scariosa* Willd.
L. pychnostachya Mx.
Solidago missouriensis Nutt.
S. speciosa var. *angustata*.
S. rigida L.
S. lanceolata L.
Aster multiflorus Ait.
Aster laevis L.
Erigeron strigosus Muhl.
Silphium laciniatum L.
S. integrifolium Mx.
S. perfoliatum L.
Ambrosia trifida L.
A. artemisiæfolia L.
A. psilostachya DC. Less common than the two preceding species.
Xanthium canadense Mill.
Heliopsis scabra Dunal.
Echinacea angustifolia DC.
Rudbeckia subtomentosa Pursh.
Lepachys pinnata T. & G.
Helianthus annuus L.
H. grosse-serratus Marteus.
Bidens frondosa L.
Dysodia chrysanthemoides Lag.
Chrysanthemum leucanthemum L. Abundant in pastures, in scattered localities throughout the county. A very troublesome weed.
Tanacetum vulgare L.
Senecio aureus L.
Cacalia tuberosa Nutt.
Arctium lappa L.
Cnicus arvensis Hoffm. Common only in isolated localities, but spreading.
Taraxacum officinale Weber.
Lactuca scariola L. Very abundant as a weed in gardens, as are also the two following species.
L. canadensis L.
Sonchus asper Vill.

LOBELIACEÆ.

- Lobelia spicata* Lam.
L. syphilitica L.

CAMPANULACEÆ

- Campanula americana* L.

PRIMULACEÆ.

- Steironema ciliatum* Raf. (*Lysimachia ciliata*.)

OLEACEÆ.

- Fraxinus americana* L.
F. rigidis Mx.

ASCLEPIADACEÆ.

- Asclepias tuberosa* L.
A. incarnata L.
A. cornuti Dec.
A. verticillata L.
Acerates longifolia Ell.

GENTIANACEÆ.

- Gentiana alba* Muhl.
G. saponaria L.

POLEMONIACEÆ.

- Phlox pilosa* L.

HYDROPHYLLOACEÆ.

- Hydrophyllum virginicum* L. Woodlands.
Ellisia nyctelea L. Not very common.

BORRAGINACEÆ.

- Lithospermum canescens* Lehm.

CONVOLVULACEÆ.

- Convolvulus sepium* L.
Cuscuta glomerata Choisy. Not common.

SOLANACEÆ.

- Solanum nigrum* L.
S. carolinense L.
S. rostratum Dunal.
Physalis lanceolata Mx.
Datura stramonium L.
D. tatula L.

SCHROPPULARIACEAE.

- Verbascum thapsus* L.
Veronica virginica L.
Catalpa speciosa Warder. (Escaped from cultivation.)

VERBENACEAE.

- Verbena stricta* Vent.
V. urticaefolia L.
V. bracteosa Mx.

LABIATAE.

- Pycnanthemum lanceolatum* Pursh. Not common. Woodlands.
Mentha canadensis L. Low prairies—common.
Monarda fistulosa L.
Nepeta cataria L.
N. glechoma Benth.
Scutellaria lateriflora L. Woods.
Brunella vulgaris L. Woodlands. Common.
Stachys palustris L. Woodlands. Common.

PLANTAGINACEAE.

Plantago major L.

NYCTAGINACEAE.

Oxybaphus hirsutus Sweet.

O. angustifolius Sweet. (?)

ILLECKBRACEAE.

Anychia dichotoma Mx. Woods. Not very common.

AMARANTACEAE.

Amarantus retroflexus L.

CHENOPODIACEAE.

Chenopodium album L.

POLYGONACEAE.

Rumex crispus L. Common everywhere.

R. verticillatus L. Tolerably common.

Polygonum aviculare L.

P. ramoissimum Mx.

P. incarnatum Watson. Sloughs. Only tolerably common.

P. persicaria L.

P. orientale L. Escaped from gardens.

Fagopyrum esculentum Moench. Cultivated species run wild.

EUPHORBIACEAE.

Euphorbia corollata L.

E. maculata L.

E. preslii Guss.

Acalypha virginica var *gracileus* Mueller. Not common.

URTICACEAE.

Ulmus americana L.

U. pubescens Walt. (*U. fulva* Mx.)

Ulmus racemosa Thomas. Reported from the west side of the county, along the course of the Nodaway river, but very doubtful.

Celtis occidentalis L.

Cannabis sativa L. Escaped from cultivation, or adventitious.

Humulus lupulus L. Occasionally fugitive from cultivation in brush and low woody thickets.

Urtica gracilis Ait.

Pilea pumila Gray. Common in all woods.

JUGLANDACEAE.

Juglans nigra L.?

Carya alba Nutt.

C. amara Nutt.

Juglans cinerea occurs in Madison county, but has not been found in Adair county. The sycamore tree has also been found to the east of the line separating the two counties, but never to the west of it.

IOWA ACADEMY OF SCIENCES.

159

CUPULIFERAE.

Corylus americana Walt.

Ostrya virginica Willd.

Quercus macrocarpa Mx. High prairie and bluffs along river.

Q. rubra L.

Q. alba L.

Q. coccinea Wang. All four species common along larger streams.

SALICACEAE.

Salix amygdaloides And.

S. alba L.

S. nigra Marsh.

S. Cordata Muhl. Not common. Discovered on Middle river near Madison county line.

Populus monilifera Ait.

ORCHIDACEAE.

Spiranthes gracilis Bigelow. Very rare. Collected by Mr. J.

G. Culver on the road between Greenfield and Orient.

Cypripedium candidum Willd. Very rare.

Habenaria leucophea Gray. Once very common. Now almost extinct.

IRIDACEAE.

Sisyrinchium augustifolium Mill.

AMARYLLIDACEAE.

Hypoxis erecta L.

LILIACEAE.

Allium canadense Kalm. Abundant in two or three restricted localities.

Polygonatum biflorum Ell. Low woodlands or brush.

Asparagus officinalis L. Escaped from gardens.

Uvularia grandiflora Smith. Woodlands. Not very common.

Erythronium americanum Ker. Woods.

Lillium philadelphicum L.

Trillium nivale Riddell. Woods.

Melanthium virginicum L.

Smilacina racemosa Desf. Woods.

MAYACEAE.

Tradescantia virginica L.

TYPHACEAE.

Typha latifolia L.

ARACEAE.

Arisaema triphyllum Tou.

ALISMACEAE.

Sagittaria variabilis Eng.

¶The birch occurs in Guthrie county but has not been discovered in Adair.