

1 Short Communication

2

3 **Olive pomace oil can be used as an alternative carbon source for clavulanic acid production by**

4 *Streptomyces clavuligerus*

5
6 Taran Young¹, Yingying Li², Georgios Efthimiou^{3*}

7 ¹Life Sciences, University of Nottingham, East Dr, NG7 2TQ, Nottingham, UK.

8 ²Manchester Institute of Biotechnology, University of Manchester, 131 Princess Street, M1 7DN, Manchester,
9 UK

10 ³ School of Life Sciences, Hardy Building, University of Hull, Cottingham Road, HU6 7RX, Hull, UK

11
12 * Corresponding author: Dr Georgios Efthimiou, Lecturer in Microbiology

13 School of Life Sciences, Hardy Building, University of Hull, Cottingham Road, HU6 7RX, Hull, UK

14 E-mail: g.efthimiou@hull.ac.uk; Tel.: (+44) 1482 465970

15 Orcid ID: 0000-0002-5909-6794

16
17 **Abstract**

18 Clavulanic acid is an important drug, both medically and economically. It is used to combat bacterial resistance
19 to β -lactam antibiotics and is on the World Health Organisation's List of Essential Medicines in combination with
20 amoxicillin. An olive oil industry waste product, olive pomace oil (OPO), is a potential alternative carbon source
21 for clavulanic acid production by *Streptomyces clavuligerus*. Olive pomace oil is six times cheaper than glycerol,
22 which is the current industry standard. The aims of this study were to examine if olive pomace oil can be used as
23 a carbon source for clavulanic acid production and to compare the clavulanic acid yield achieved in shake flasks
24 and 1.8 L bioreactors. It was observed that olive pomace oil was efficiently utilised as a sole carbon source by *S.*
25 *clavuligerus* growing in a P-limited medium. The *S. clavuligerus* cells grew faster in olive pomace oil-containing
26 cultures compared to the glycerol-containing cultures (control) and produced comparable levels of clavulanic acid,
27 but much earlier. In cultures with ISP2 medium that contained glycerol or olive pomace oil, higher levels of

28 clavulanic acid were obtained in shake flask cultures with olive pomace oil. Interestingly, the same levels of
29 clavulanic acid were observed in oil-containing cultures in bioreactors, but 48 h earlier. Furthermore, the oil-
30 containing cultures did not need addition of an antifoam agent, while higher levels of cell viability were
31 maintained after 72 h in these fermentations compared to the cultures that contained glycerol. Our results suggest
32 that olive pomace oil can replace glycerol for clavulanic acid production in *S. clavuligerus* fermentations, which
33 will significantly increase the productivity and cut the cost for industrial clavulanic acid biosynthesis. The same
34 carbon source can be tested in other similar fermentation approaches for the production of antibiotics or other
35 valuable bioproducts.

37 **Keywords:** Antibiotics; Bioprocessing, Clavulanic acid; Olive pomace oil; *Streptomyces*; Waste.

39 **Statement of Novelty**

40 Olive pomace oil is an environmentally toxic waste of the olive oil industry. We show that it can replace glycerol
41 in microbial fermentations, leading to higher yield of clavulanic acid production.

43 **Introduction**

44 The most common mechanism by which bacteria acquire resistance to beta-lactam antibiotics is through the
45 production of beta-lactamase enzymes. These enzymes cleave the beta-lactam rings of important antibiotics such
46 as penicillins and cephalosporins, leading to their inactivation. Clavulanic acid, produced by *Streptomyces*
47 *clavuligerus*, is used in conjunction with β -lactam antibiotics, acting as a suicide inhibitor that protects these drugs
48 from the β -lactamases of resistant bacteria [1]. The combination of amoxicillin and clavulanic acid is on the World
49 Health Organisation (WHO) list of essential medicines [2]. Due to the importance of this treatment, much research
50 has been done to improve the growth of *S. clavuligerus*, and the organism's production of clavulanic acid [3].

51 Different from most organisms, *S. clavuligerus* lacks a glucose transport system [4], so it is unable to utilise this
52 sugar. Instead, glycerol is the most popular carbon source used for industrial production of clavulanic acid [5].
53 Glycerol is, however, relatively expensive, so cheaper alternatives such as oils have been looked into. It has been
54 shown previously that olive oil outperforms glycerol (and other oils) in terms of the concentration of clavulanic

1
2 55 acid being produced by *S. clavuligerus*, with olive oil producing a maximum concentration of 47 mg L⁻¹, almost
3 56 double that of the standard glycerol (25 mg L⁻¹) [6].

4 57 Olive mills generate huge waste in short periods of time, which is seen as an environmental problem for
5 58 Mediterranean countries [7]. These wastes have high phytotoxicity, with studies also showing adverse effects on
6 59 soil microbial populations [8] and aquatic ecosystems [9]. While olive oil is cheaper than glycerol, and has been
7 60 shown to work better, it does not seem ethical to add to existing demand, and in turn, increasing waste. One of
8 61 these wastes from the olive oil industry, olive pomace oil (OPO) [10], has its own potential as a carbon source for
9 62 *S. clavuligerus*. Miranda *et al.* [11] show that there is around 52% carbon composition in the pomace, while the
10 63 pulp (where regular olive oil is obtained) has around 55%. This suggests that OPO likely has sufficient energy to
11 64 be a viable carbon source, still outperforming glycerol as olive oil was shown to. The average price of OPO is
12 65 £1.50 per L, which is six times less expensive than the standard carbon source glycerol (£9 per L). In addition to
13 66 being cost effective, OPO utilisation would also create a positive reuse for some of the vast amounts of waste
14 67 found in the olive oil industry. If accepted widely, this could bring huge economic and environmental benefits to
15 68 Mediterranean countries, and lower costs for clavulanic acid producers. Additionally, oil has natural anti-foaming
16 69 properties and can enhance secondary metabolism [12].

17 70 Lots of efforts have been devoted to the development of *S. clavuligerus* strains with the aim to improve the
18 71 production of clavulanic acid. These strain improvements, whether they were done through random mutagenesis
19 72 [13] or targeted modification of metabolic pathways [14], were usually carried out with glycerol in mind as the
20 73 carbon source. With food oils already yielding higher clavulanic acid production, and in some cases culture
21 74 growth, strain improvement of *S. clavuligerus* to increase the utilisation of these oils [15] could greatly improve
22 75 clavulanic acid production.

23 76 Although complex media are commonly used for industrial clavulanic acid production, defined media allow us to
24 77 test varying media ingredients. In this paper, we used defined media to test the growth of *S. clavuligerus* and the
25 78 yield of clavulanic acid using OPO as the sole carbon source. To assess the fermentation efficiency of different
26 79 carbon sources, glycerol was set as control to produce clavulanic acid. Because of the high carbon content, we
27 80 hypothesise that OPO will prove to be a suitable carbon source, likely more effective than the industry standard
28 81 of glycerol for clavulanic acid production. In addition, we believe that OPO will continue to be effective at higher
29 82 volumes as part of complex media.

30 83

84 **Materials and Methods**

85 **Strains and media**

86 The bacterial strain used in this study was *Streptomyces clavuligerus* DM738 (kindly provided by Prof. Paul
87 Hoskisson, University of Strathclyde, UK). The medium used for the pre-culture was SV2, containing (per L):
88 Glycerol, 15 g; Glucose, 15 g; Peptone, 15 g; CaCO₃, 1 g. For the flask cultures, a P-limited medium was used,
89 containing (per L): NH₄Cl, 7 g; KH₂PO₄, 0.25 g; MOPS, 21 g. The pH value was adjusted to 6.8 and the medium
90 was autoclaved at 121°C for 15 min. 10 mL of trace elements stock solution was aseptically added per L of P-
91 limited medium after autoclaving [6]. The trace elements stock solution contained (per L): MgSO₄·7H₂O, 25 g;
92 FeSO₄·7H₂O, 2.5 g; CoCl₂, 0.055 g; CuCl₂ 0.053 g, CaCl₂·2H₂O, 1.4 g; ZnCl₂, 1 g; MnCl₂, 1 g; Na₂MoO₄, 0.03
93 g. For shake flask and bioreactor cultures, ISP2 medium was used, containing (per L): Yeast extract, 4 g; Malt
94 extract, 10 g; Dextrose, 4 g. The carbon source was added at 0.6% v/v by filter sterilisation. All media were
95 autoclaved at 121°C for 15 min unless stated otherwise.

97 **Inoculum preparation and cultures**

98 For the pre-culture, 1 mL of spores was added into 50 mL of SV2 in a 250 mL glass baffled flask and agitated at
99 200 rpm at 30°C for 48 h. The culture was transferred into plastic universals and centrifuged at 6000 xg for 5 min.
100 The supernatant was removed, and the pellets were resuspended in 6 mL of P-limited media. For the shake flask
101 experiments, 2 L glass baffled flasks were used containing 400 mL of medium (P-limited or ISP2). Each flask
102 was inoculated with 1 mL of the re-suspended inoculum. The flasks were incubated at 30°C, with agitation at 150
103 rpm.

104 For the bioreactor cultures, 1.8 L bioreactors (DASGIP parallel system, Eppendorf, UK) were used. Each vessel
105 contained 1 L of ISP2 medium. The inoculum was prepared as described above and introduced aseptically with a
106 syringe (5% v/v). The agitation rate was set at 1000 rpm, with a culturing temperature of 30°C. The pH value was
107 maintained at 7 and the aeration rate was set at 0.25 L min⁻¹. OPO or glycerol were added at the same percentage
108 as above. Filter-sterilised Antifoam 204 (Sigma-Aldrich, UK) was added at 0.001% v/v in the glycerol-containing
109 cultures.

111 **Clavulanic acid assay**

112 Clavulanic acid levels were determined as described by Bird *et al.* [16].

113

114 Biomass determination

115 Biomass levels were determined as described by Efthimiou *et al.* [6].

116

117 **Results and Discussion**

118 Production of many secondary metabolites by *Streptomyces* species, including clavulanic acid, are negatively
119 regulated by phosphate [17]. P-limited media is thus often used to stimulate the production of these compounds.

120 As can be seen in Figure 1A, in P-limited medium, *S. clavuligerus* reached maximal growth five days earlier than
121 in the oil-containing cultures compared to the glycerol containing cultures. The OPO cultures also produced higher
122 levels of clavulanic acid until 264 h with a maximum clavulanic acid concentration of nearly 150 mg L⁻¹. The
123 maximum clavulanic acid production in the glycerol cultures was over 200 mg L⁻¹, however this decreased later.
124 These results are a good first indication that OPO could be a suitable carbon source replacement for clavulanic
125 acid production in *S. clavuligerus*. It should be noted that the method of inoculation ensures a minimal carry over
126 of nutrients.

127 Once it was determined that OPO was able to be utilised in a defined medium, it was then added as a carbon
128 source in a complex media (Figures 2A and 2B), which more accurately replicates the conditions in industrial
129 productions (albeit at a laboratory scale). In the first 24 h of growth in complex media, OD values obtained were
130 very similar between the carbon sources, but OPO cultures achieved higher growth at all time points recorded. In
131 addition, Figure 2B shows that OPO as a carbon source in complex media again led to the production of much
132 larger amounts of clavulanic acid (over double that of glycerol at 48 h). This is similar to previous results obtained
133 comparing clavulanic acid production in olive oil and glycerol [6, 18]. Cell viability was also greater in OPO
134 cultures across all experiments when spread onto plates (data not shown).

135 The next step was to scale up the experiment to 1.8 L bioreactors, further replicating industrial production. Figure
136 2C continues to support OPO as a superior carbon source to glycerol, with the OPO cultures obtaining a higher
137 maximum biomass, earlier than the glycerol culture. In Figure 2D, it can be seen that clavulanic acid production

138 also occurs earlier in OPO cultures with a maximum of 325 mg L⁻¹ at 48 h. This is achieved 48 hours earlier than
139 the maximum 350 mg L⁻¹ found in glycerol cultures.

140 Using data from the bioreactor cultures, Table 1 shows that OPO outcompetes glycerol in all four aspects
141 measured: growth yield, growth rate, clavulanic acid yield, clavulanic acid rate. Growth yield and growth rate
142 values from OPO cultures were almost double that of glycerol cultures. Clavulanic acid yield between carbon
143 sources was closer, but OPO still appears the more favourable than glycerol (59.96 mg g⁻¹ of carbon source and
144 46.16 mg g⁻¹ of carbon source, respectively). Similarly, the rate of clavulanic acid production was 1.47 times
145 higher in OPO cultures than in glycerol cultures.

146 The effectiveness of OPO also has the potential to drastically increase through strain improvement targeted
147 specifically at the use of this carbon source. Industrial strains reportedly have a clavulanic acid production nearing
148 3 g L⁻¹ which is far higher than wild type strains (around 25-120 mg L⁻¹) [3]. Spontaneous mutations with oil as a
149 carbon source have been recorded [19], proving that this is viable. In combination with targeted mutagenesis,
150 there is an even greater potential for OPO. It would be interesting to see the levels of clavulanic acid produced by
151 these industrial strains when OPO is the carbon source.

152 It was also observed in these experiments that OPO acted as a strong antifoam, leading to negligible foaming of
153 the media compared to glycerol. This might partly explain the better results of OPO in the shake flask experiments
154 (antifoam was added to glycerol cultures in the bioreactor experiments). Foam in the glycerol cultures may have
155 impeded on gas exchange, negatively impacting on growth and clavulanic acid production [20]. Foam is generally
156 seen as undesirable in the production of antibiotics [21].

157 All three of these experiments indicate that OPO is a superior carbon source as it led to faster growth, faster
158 clavulanic acid production, and removed the need for an antifoam – an additional cost when glycerol is used. In
159 other studies, fed-batch cultures have been shown to further increase the levels of clavulanic acid production [22].

160 This by-product of the olive oil industry is extracted from undesirable waste such as the skin, pulp, stone, and
161 kernel of olives [23]. By providing these waste products with a use, OPO extraction both alleviates environmental
162 pollution and can provide a further source of revenue for olive oil-producing countries. This has the potential to
163 boost the economies of Spain, Greece, Italy, and other Mediterranean countries. OPO as a carbon source could
164 also be of great economic impact for clavulanic acid producers as a result of its low cost of around £1.50 per L
165 litre, six times cheaper than glycerol.

166 There are disadvantages that should be taken into account. First, there is often oil residue at the end of the culture
167 due to limited interaction of the bacteria and the lipid droplets [12]. However, this problem is often countered by
168 slowly deeding the oil during culture [24]. Secondly, there is often a greater oxygen demand in cultures with oils
169 compared to carbohydrates [12].

170 In conclusion, our study shows that OPO is an effective carbon source for clavulanic acid production by *S.*
171 *clavuligerus* fermentation. This extremely cheap by-product of the olive oil industry outperformed the standard
172 carbon source, glycerol, by promoting faster growth, larger quantities of clavulanic acid production, and natural
173 antifoaming properties. We believe that by switching to OPO, industrial producers could save large amounts of
174 money, while at the same time achieving higher clavulanic acid yield. This would not only hugely benefit the
175 Mediterranean countries that produce OPO, but reduce production costs of clavulanic acid, bringing the price
176 down of an essential drug. Finally, it is possible that the same carbon source can be used in other similar
177 fermentation approaches for the production of antibiotics or other valuable bioproducts.

179 **Conflicts of Interest**

180 The authors declare no conflict of interest.

183 **References**

- 184 1. Drawz, S., Bonomo, R.: Three Decades of β -Lactamase Inhibitors. *Clin. Microbiol. Rev.* **23**, 160-201
185 (2010)
- 186 2. World Health Organisation: WHO Model List of Essential Medicines 20th List. World Health
187 Organisation (2017)
- 188 3. Paradkar, A.: Clavulanic acid production by *Streptomyces clavuligerus*: biogenesis, regulation, and strain
189 improvement. *J. Antibiot.* **66**, 411-420 (2013)
- 190 4. Pérez-Redondo, R., Santamarta, I., Bovenberg, R., Martin, J., Liras, P.: The enigmatic lack of glucose
191 utilization in *Streptomyces clavuligerus* is due to inefficient expression of the glucose permease gene.
192 *Microbiol.* **156**, 1527–1537 (2010)

- 193 5. Costa, C.L.L., Badino, A.C.: Production of clavulanic acid by *Streptomyces clavuligerus* in batch
1
2 194 cultures without and with glycerol pulses under different temperature conditions. *Biochem. Eng. J.* **69**,
3
4 195 1-7 (2012)
- 5
6 196 6. Efthimiou, G., Thumser, A., Avignone-Rossa, C.: A novel finding that *Streptomyces clavuligerus* can
7
8 197 produce the antibiotic clavulanic acid using olive oil as a sole carbon source. *J. Appl. Microbiol.* **105**,
9
10 198 2058-2064 (2008)
- 11
12 199 7. Roig, A., Cayuela, M.L., Sánchez-Monedero, M.A.: An overview on olive mill wastes and their
13
14 200 valorisation methods. *Waste Manag.* **26**, 960-969 (2006)
- 15
16 201 8. Paredes, M.J., Moreno, E., Ramos-Cormenzana, A., Martinez, J.: Characteristics of soil after pollution
17
18 202 with waste waters from olive oil extraction plants. *Chemosphere.* **16**, 1557-1564 (1987)
- 19
20 203 9. DellaGreca, M., Monaco, P., Pinto, G., Pollio, A., Previtiera, L., Temussi, F.: Phytotoxicity of Low-
21
22 204 Molecular-Weight Phenols from Olive Mill Waste Waters. *Bull. Environ. Contam. Toxicol.* **67**, 352-359
23
24 205 (2001)
- 25
26 206 10. Moghaddam, G., Heyden, Y.V., Rabiei, Z., Sadeghi, N., Oveisi, M.R., Jannat, B., Araghi, V., Hassani,
27
28 207 S., Behzad, M., Hajimahmoodi, M.: Characterisation of different olive pulp and kernel oils. *J. Food*
29
30 208 *Compos. Anal.* **28**, 54-60 (2012)
- 31
32 209 11. Miranda, T., Esteban, A., Rojas, S., Montero, I., Ruiz, A.: Combustion Analysis of Different Olive
33
34 210 Residues. *Int. J. M. Sci.* **9**, 512-525 (2008)
- 35
36 211 12. Large, K.P., Ison, A.P., Williams, D.J.: The effect of agitation rate on lipid utilisation and clavulanic
37
38 212 acid production in *Streptomyces clavuligerus*. *J. Biotechnol.* **63**, 111-119 (1998)
- 39
40 213 13. Korbekandi, H., Darkhal, P., Hojati, Z., Abedi, D., Hamed, J., Pourhosein, M.: Overproduction of
41
42 214 Clavulanic Acid by UV Mutagenesis of *Streptomyces clavuligerus*. *Iran. J. Pharm. Res.* **9**, 177-181
43
44 215 (2010)
- 45
46 216 14. Medema, M.H., Alam, M.T., Breitling, R., Takano, E.: The future of industrial antibiotic production:
47
48 217 From random mutagenesis to synthetic biology. *Bioeng. Bugs.* **2**, 230-233 (2011)
- 49
50 218 15. Kim, S., Kim, J., Shin, C., Park, H.W., Kim, C.: An Approach to Strain Improvement and Enhanced
51
52 219 Production of Clavulanic Acid in *Streptomyces clavuligerus*. *Biosci. Biotech. Biochem.* **73**, 160-164
53
54 220 (2009)
- 55
56 221 16. Bird, A.E., Bellis, J.M., Gasson, B.C.: Spectrophotometric assay of clavulanic acid by reaction with
57
58 222 imidazole. *Analyst.* **107**, 1241-1245 (1982)
- 59
60
61
62
63
64
65

223 17. Martin, J.F.: Phosphate Control of the Biosynthesis of Antibiotics and Other Secondary Metabolites Is
1 Mediated by the PhoR-PhoP System: an Unfinished Story. *J. Bacteriol.* **186**, 5197-5201 (2004)
2 224
3
4 225 18. Salem-Berkhit, M.M., Alanazi, F.K., Alsarra, I.A.: Improvement and enhancement of clavulanic acid
5 production in *Streptomyces clavuligerus* using vegetable oils. *Afr. J. Biotech.* **9**, 6806-6812 (2010)
6 226
7
8 227 19. Lee, P.C., Ho, C.C.: Production of clavulanic acid and cephamycin C by *Streptomyces clavuligerus* in
9 palm-oil medium. *World J. Microbiol. Biotechnol.* **12**, 73-75 (1996)
10 228
11
12 229 20. Ser, H., Law, J.W., Chaiyakunapruk, N., Jacob, S.A., Palanisamy, U.D., Chan, K., Goh, B., Lee, L.:
13 Fermentation Conditions that Affect Clavulanic Acid Production in *Streptomyces clavuligerus*: A
14 230 Systematic Review. *Front. Microbiol.* **7**, 522 (2016)
15 231
16
17
18 232 21. Routledge, S.J.: Beyond de-foaming: the effects of antifoams on bioprocess productivity. *Comput. Struct.*
19 *Biotechnol. J.* **3**, e201210014 (2012)
20 233
21
22 234 22. Bellao, C., Antonio, T., Araujo, M.L.G.C., Badino, A.C.: Production of clavulanic acid and cephamycin
23 C by *Streptomyces clavuligerus* under different fed-batch conditions. *Braz. J. Chem. Eng.* **30**, 257-266
24 235 (2013)
25 236
26
27
28 237 23. Moral, P.S., Méndez, M.V.R.: Production of pomace olive oil. *Grasas y Aceites.* **57**, 47-55 (2006)
29
30 238 24. Anderson, R.F., Tornqvist, E.G.M., Peterson, W.H.: Effect of oil in pilot plant fermentations. *J. Agric.*
31 *Food. Chem.* **4**, 556-559 (1956)
32 239
33
34
35 240
36
37
38 241
39
40
41 242
42
43 243
44
45
46 244
47
48
49 245
50
51
52 246
53
54 247
55
56
57 248
58
59
60 249
61
62
63
64
65

250 **Table Legend**

1
2
3 251 **Table 1** Biomass and clavulanic acid production yields and rates (curve slope values), with olive pomace oil and
4
5 252 glycerol as carbon sources (calculated using data from the bioreactor cultures).
6
7 253
8
9
10 254
11
12
13 255
14
15 256
16
17
18 257
19
20
21 258
22
23
24 259
25
26 260
27
28
29 261
30
31
32 262
33
34 263
35
36
37 264
38
39
40 265
41
42
43 266
44
45 267
46
47
48 268
49
50
51 269
52
53
54 270
55
56 271
57
58
59 272
60
61
62
63
64
65

273 **Figure Legends**

1
2
3 274 **Figure 1** Growth (A) and clavulanic acid production (B) in shake flasks with P-limited medium and glycerol (gray
4 line) or olive pomace oil (black line) as sole carbon source ($n = 3$, where n is the number of biological replicates).



5
6 275
7 276 CLA: clavulanic acid. Error bars: \pm Standard Deviation.

8
9 277 **Figure 2** Growth (A) and clavulanic acid production (B) in shake flasks with ISP2 and glycerol (gray line) or
10 olive pomace oil (black line) as additional carbon source. Growth (C) and clavulanic acid production (D) in 1.8 L

11 278
12 279 DASGIP bioreactors with ISP2 and glycerol or olive pomace oil as additional carbon source ($n = 3$, where n is the
13 number of biological replicates). CLA: clavulanic acid. Error bars: \pm Standard Deviation.

14
15 280
16
17
18 281

19
20
21 282
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65









Table 1. Biomass and clavulanic acid production yields of carbon sources

Carbon source	Growth yield (g g^{-1} of carbon source)	Growth rate ($\text{g}\cdot\text{L}^{-1}\cdot\text{h}^{-1}$)	Clavulanic acid yield (mg g^{-1} of carbon source)	Clavulanic acid rate ($\text{mg}\cdot\text{L}^{-1}\cdot\text{h}^{-1}$)
Olive pomace oil	6.52	0.75	59.96	6.87
Glycerol	3.70	0.40	46.16	4.66

