

Supplementary material for the article:

Stojsavljević, A.; Rovčanin, B.; Krstić; Jagodić, J.; Borković-Mitić, S.; Paunović, I.; Živaljević, V.; Mitić, B.; Gavrović-Jankulović, M.; Manojlović, D. Cadmium as Main Endocrine Disruptor in Papillary Thyroid Carcinoma and the Significance of Cd/Se Ratio for Thyroid Tissue Pathophysiology. *Journal of Trace Elements in Medicine and Biology* **2019**, *55*, 190–195. <https://doi.org/10.1016/j.jtemb.2019.06.009>

Supplementary material

Cadmium as main endocrine disruptor in papillary thyroid carcinoma and the significance of Cd/Se ratio for thyroid tissue pathophysiology

Aleksandar Stojšavljević^a, Branislav Rovčanin^b, Đurđa Krstić^a, Slavica Borković-Mitić^c, Ivan Paunović^b, Vladan Živaljević^b, Bojan Mitić^d, Marija Gavrović-Jankulović^a, Dragan Manojlović^{a,e}

^a *University of Belgrade – Faculty of Chemistry, Studentski trg 12-16, 11000 Belgrade, Serbia*

^b *Center for Endocrine Surgery, Clinical Center of Serbia, Koste Todorovica 8, Belgrade, Serbia*

^c *Department of Physiology, Institute for Biological Research “Siniša Stanković”, University of Belgrade, Bulevar despota Stefana 142, Belgrade 11060, Serbia*

^d *University of Belgrade – Faculty of Biology, Studentski trg 16, 11000 Belgrade, Serbia*

^e *South Ural State University, Chelyabinsk, Lenin prospect 76, 454080, Russia*

Table S1. Median and interquartile range (IQR) for metals (in ng/g) and differences between two sets of data by Mann-Whitney U-test for HTT samples.

Table S2. Results of Kruskal-Wallis test according to age, PH type, T and N stage.

Table S1

Results of Kruskal-Wallis test (a difference between two sets of data is significant when p value is less or equal to 0.05).

	Age groups		PH type		T stage		N stage	
	<i>p-value</i>	Fisher (LSD)	<i>p-value</i>	Fisher (LSD)	<i>p-value</i>	Fisher (LSD)	<i>p-value</i>	Fisher (LSD)
Mn	0.3444	/	0.1370	/	0.0468	T1(T2) T2(T3)	0.0226	N0(Nx)
Co	0.5881	/	0.6848	/	0.3979	/	0.0089	N0(Nx)
Ni	0.7609	/	0.6079	/	0.7534	/	0.3556	/
Cu	0.4187	/	0.2923	/	0.2556	/	0.0940	/
Zn	0.2634	/	0.9557	/	0.4340	/	0.1432	/
As	0.4230	/	0.7043	/	0.0121	/	0.3308	/
Se	0.2727	/	0.0481	Follicular (solid)	0.0409	T1(T2)	0.6942	/
Cd	0.0052	Group 1 (Group 3) Group 2 (Group 3)	0.8145	/	0.6058	/	0.2894	/
Pb	0.3118	/	0.6441	/	0.9889	/	0.0226	N0(Nx)
Th	0.5606	/	0.7884	/	0.1965	/	0.0089	N0(Nx)
U	0.0174	Group 1 (Group 2)	0.0434	Follicular (solid)	0.8067	/	0.3556	/
Cu/Zn	0.3301	/	0.2157	/	0.7396	/	0.0940	/
Cd/Se	0.0072	Group 1 (Group 3) Group 2 (Group 3)	0.5404	/	0.1843	/	0.1432	/

Table S2

Median and interquartile range (IQR) for metals (in ng/g) and differences between two sets of data by Mann-Whitney U-test for HTT samples.

	Sex				<i>p</i> *	Smoking habits				<i>p</i> *
	Women		Men			Yes		No		
	median	IQR	median	IQR		median	IQR	median	IQR	
Mn	119	58	130	77	0.346	107	62	124	62	0.583
Co	4.27	2.83	3.42	18.92	0.137	3.94	2.20	3.98	3.22	0.678
Ni	74	118	85	182	0.775	80	137	74	140	0.981
Cu	310	173	372	311	0.185	253	152	352	208	0.020
Zn	5642	3253	7032	3094	0.048	5778	2392	5908	3753	0.666
As	1.23	1.82	1.23	1.31	0.635	0.84	1.23	1.26	1.93	0.243
Se	114	90	136	117	0.521	80	84	136	106	0.059
Cd	30	54	46	137	0.172	72	58	30	47	0.033
Pb	19	13	19	10	0.430	18	14	20	12	0.875
Th	0.12	0.14	0.11	0.09	0.291	0.12	0.21	0.12	0.11	0.383
U	0.050	0.030	0.045	0.050	0.112	0.065	0.050	0.045	0.050	0.014
Cu/Zn	0.05	0.03	0.05	0.05	0.68	0.05	0.02	0.06	0.04	0.093
Cd/Se	0.265	0.330	0.375	0.930	0.336	0.725	1.040	0.245	0.310	0.005

*Differences between two sets of data is significant when p value is less or equal to 0.05