

## Accepted Manuscript

Title: Experimental warming had little effect on carbon-based secondary compounds, carbon and nitrogen in selected alpine plants and lichens

Authors: L. Nybakken, S.M. Sandvik, K. Klanderud

PII: S0098-8472(11)00103-1  
DOI: doi:10.1016/j.envexpbot.2011.04.011  
Reference: EEB 2340

To appear in: *Environmental and Experimental Botany*

Received date: 13-10-2010  
Revised date: 25-3-2011  
Accepted date: 19-4-2011

Please cite this article as: Nybakken, L., Sandvik, S.M., Klanderud, K., Experimental warming had little effect on carbon-based secondary compounds, carbon and nitrogen in selected alpine plants and lichens, *Environmental and Experimental Botany* (2010), doi:10.1016/j.envexpbot.2011.04.011

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



1

2 Experimental warming had little effect on carbon-based secondary  
3 compounds, carbon and nitrogen in selected alpine plants and lichens

4

5 L. Nybakken<sup>1,2\*</sup>, S. M. Sandvik<sup>3</sup>, K. Klanderud<sup>1, 4</sup>

6

7

8 <sup>1</sup>Department of Ecology and Natural Resource Management, Box 5003, Norwegian

9 University of Life Sciences, N-1432 Ås, Norway <sup>2</sup>Natural Product Research Laboratory

10 Department of Biology, University of Eastern Finland, P.O.Box 111, FIN-80101 Joensuu,

11 Finland, <sup>3</sup>University of Agder, Faculty of Engineering and Science, Service Box 422, N-4604

12 Kristiansand, Norway <sup>4</sup>Department of Biology, University of Bergen, P.O. Box 7803, N-5020

13 Bergen, Norway

14

15

16

17

18 \*Corresponding author: L. Nybakken. Fax: +358 132513590. E-mail address:

19 [line.nybakken@uef.fi](mailto:line.nybakken@uef.fi)

20

21

22

23 Running title: Effects of warming on alpine plants and lichens

1 **Abstract**

2 Global warming is expected to change plant defence through its influence on plant primary  
3 resources. Increased temperature (T) will increase photosynthesis, and thus carbon (C)  
4 availability, but may also increase soil mineralization and availability of nitrogen (N). More  
5 access to C and N is expected to mainly increase plant growth, and, according to hypotheses  
6 on resource based defence, this could lower plant concentrations of carbon-based secondary  
7 compounds (CBSCs).

8

9 We used two already established warming experiment with open top chambers (OTCs) and  
10 control plots in alpine south-western Norway, one on a ridge (8 years' treatment) and a one in  
11 a leaside (3 years' treatment), to study the effects of warming on plant and lichen defensive  
12 compound concentrations. The study included five vascular plant and six lichen species.

13

14 One vascular plant species had lower concentration of CBSCs under elevated T, while the  
15 others did not respond to the treatment. In lichens there were no effects of warming on  
16 CBSCs, but a tendency to reduced total C concentrations. However, there were effects of  
17 warming on nitrogen, as the concentration decreased inside OTCs for three species, while it  
18 increased for one lichen species. Lichens generally had higher CBSC and total C  
19 concentrations on the ridge than in the leaside, but no such pattern were seen for vascular  
20 plants.

21

22 No elevated temperature effect on CBSCs is most probably a result of high constitutive  
23 defence under the limiting alpine conditions, suggesting that chemical defence is little subject  
24 to change under climate warming, at least on a short-term basis. We suggest that the driving  
25 forces of plant defence in the arctic-alpine should be tested individually under controlled  
26 conditions, and suggest that competition from other plants may be a greater threat under  
27 climate warming than increased herbivory or disease attacks.

28

29 **Key Words:** Vascular plants, lichens, secondary compounds, CBSC, lichen compound,  
30 temperature, carbon, nitrogen.

## 1 **1. Introduction**

2  
3 Carbon-based secondary compounds (CBSCs), generally phenolics and terpenoids, defend  
4 plants against damaging radiation, herbivores, and competition from other plants. The  
5 variation in CBSC concentration and composition within and between species is only partly  
6 understood, and ecologists have put forward several hypotheses where the CBSC level has  
7 been positively linked to available photosynthates (carbon, C), and negatively linked to  
8 growth and nutrient status (nitrogen, N) in the plants (e.g. the Carbon Nutrient Balance (CNB)  
9 Hypothesis, Bryant et al., 1983, see Stamp, 2003 for an overview). The predictions are, in  
10 short, that plants growing in environments with high resource (nutrient) availability will  
11 prioritize growth (simply because they can), and spend less on defence, while plants in  
12 (nutrient) limiting environments will invest more in C-based defence because growth is  
13 restricted and C may be in excess (Herms and Mattson, 1992 and references therein). In line  
14 with this, it is also expected that slow-growing species and perennials will invest more in  
15 defence than pioneer plants and annuals (Tuomi et al., 1991; Stamp, 2003).

16  
17 Light intensity will also affect the defence level of plants, as it affects photosynthesis and thus  
18 assimilation of C. Shade plants may thus have lowered defence levels (Bryant et al., 1983;  
19 Mole et al., 1988; Nichols-Orians, 1991). Experimental studies have both supported (e.g.  
20 Bryant et al., 1983; Coley et al., 2002; Leser and Treutter, 2005) and opposed (e.g. Baldwin et  
21 al., 1993; Iason and Hester, 1993; Lamontagne et al., 2000) these hypotheses. However,  
22 recent biochemical and molecular studies strongly support the idea that secondary metabolites  
23 are regulated in response to C and nitrogen N status in the plant (Fritz et al., 2006; Matt et al.,  
24 2002), meaning that the availability of resources is central for the level of defence. At the  
25 same time, both hypotheses (Bryant et al., 1983; Tuomi et al., 1988; 1991; Stamp, 2003) and  
26 experimental evidence suggest (Muzika et al., 1989; Holopainen et al., 1995; Koricheva et al.,  
27 1998) that defence levels are not exclusively dependent on resource levels. However, the  
28 CNB hypothesis also predicts that some C-based defence is produced independently of the  
29 resource situation, in conjunction with growth, so that plants, to different extents, have a fixed  
30 level of defence, often called constitutive defence (genetically decided). For example, woody  
31 plants adapted to low resource situations are expected to have a low growth rate and therefore  
32 low capacity for compensatory growth after herbivory, which in turn would favour selection  
33 for maintenance of high defence levels and carbon surplus into storage rather than defence  
34 (little or no plasticity in defence). For genotypes with high plasticity in defence, any effects of

1 resource conditions (shading, nutrient availability, increased photosynthesis) on the  
2 carbon:nutrient ratio can cause changes in the total defence levels (Bryant et al., 1983; Tuomi  
3 et al., 1988; Stamp, 2003).

4  
5 Plant growth in high altitude and latitude environments is limited by low temperatures (T)  
6 (Bliss, 1962; Körner 1999) and lack of nutrients (Chapin et al., 1980; Callaghan and Jonasson,  
7 1995, Klanderud and Totland, 2005). Growing under nutrient limited conditions, arctic-alpine  
8 plants would be expected to invest strongly in defence (according to resource-based  
9 hypotheses, reviewed by Herms and Mattson; 1992 and Stamp, 2003), but T-limited  
10 photosynthesis (C acquisition) and metabolism probably also imply restrictions on the  
11 production of defence compounds. Alpine and arctic habitats are expected to experience  
12 significant future climate warming (ACIA, 2005; IPCC, 2007), and, more specifically, the  
13 mean T increase per decade over Norway is expected to be between 0.2 and 0.5 °C (Hanssen-  
14 Bauer and Førland, 2001). The effect of warming on arctic-alpine plant defence has been little  
15 studied, and with inconsistent results (Dormann, 2003; Hansen et al., 2006; Nybakken et al.,  
16 2008). Previous studies have focused more widely in growth responses to T showing early  
17 stimulation followed by a gradual cessation of effects in the longer term (Arft et al., 1999). In  
18 long term experiments, warming increased height and cover of deciduous shrubs and  
19 graminoids, and decreased cover of mosses and lichens (Walker et al., 2006). In a synthesis of  
20 16 warming studies including lichens, Cornelissen et al. (2001) defended the hypothesis that  
21 lichen-decline in sub- and mid-arctic ecosystems is a function of increases in vascular plant  
22 biomass, but did not find a relationship for the coldest high-arctic and alpine sites. Dormann  
23 and Woodin (2002) reviewed 36 warming experiments of different types in the Arctic, and  
24 also found greatest growth responses for grasses and shrubs, while Richardson et al. (2002)  
25 found no significant effect of warming on plant growth in a synthesis of warming experiments  
26 from sub-Arctic Abisko after 9 years. The varying effects of warming on growth of different  
27 life forms imply that effects on defence compounds should also vary. Furthermore, as the  
28 herbs and cryptogams that grow slowly, faster growing shrubs and graminoids might shade  
29 them (Klanderud and Totland, 2005), resulting in a reduction in C resources for defence.  
30 Lichens also contain CBSCs that function as herbivore and/ or solar defences (Emmerich et  
31 al., 1993; Gauslaa, 2005; Lawrey, 1983; Pöykkö et al., 2005, Solhaug and Gauslaa, 1996;  
32 Solhaug et al., 2010), and the concentrations of some lichen CBSCs have been shown to be a  
33 direct function of available light (Gauslaa and Ustvedt, 2003; Gauslaa and McEvoy, 2005;

1 Nybakken et al., 2007; Solhaug et al., 2003; Solhaug et al., 2009), which suggests that lichen  
2 defence would decrease with warming because of increased shading.

3

4 Increases in air T subsequently increase soil T (e.g. Klanderud and Totland, 2005), and  
5 possibly improve soil mineralization and soil nutrient status (Bonan et al., 1992; Nadelhoffer  
6 et al., 1991; White, 1999), which could also increase growth and decrease defence. According  
7 to Wookey et al. (2009), the ability to take advantage of an increased N availability should  
8 also vary between life forms, as biomass and production per unit of N varies greatly among  
9 tissue types and the relative amount of each tissue type a plant has. Evergreen shrubs have for  
10 example been shown to produce more biomass per unit N than graminoids (Shaver and  
11 Shapin, 1991, Suding et al., 2004). Lichens would probably not get any advantage of  
12 increased soil N at all, as they withdraw most of their nutrients from atmospheric sources  
13 (Nash, 2008).

14

15

16 In the slopes of the mountain Sanddalsnuten (1300 -1550 m a.s.l.) at Finse, south-western  
17 Norway (60°N, 7°E) several warming experiments with Open Top Chambers (OTCs) have  
18 been run since the late 1990s, showing that both T and N limit plant growth in this area, and  
19 that warming and increased nutrient availability increase growth of graminoids and some  
20 forbs at the cost of low stature forbs, club mosses, lichens and mosses (Klanderud and  
21 Totland, 2005, Klanderud, 2008). In the present study, our aim was to measure effects of  
22 warming on total C, N and C-based defence in arctic-alpine lichens and vascular plants of  
23 different functional groups. We sampled plant leaves and lichen thalli from OTCs and control  
24 plots from two different experiments, one on a ridge close to the mountain peak, and one from  
25 the leeward side. The treatments had been running for 8 (ridge) and 3 (leeward side) years when the  
26 current analysis was conducted. In line with hypotheses on resource-based defence, we  
27 expected reduced defence in the OTCs, as warming could reduce growth limitations, by both  
28 increased T and N. We expected differences according to functional groups, as they have been  
29 shown to respond differently to both T and N. Some functional groups, like shrubs, may have  
30 a fixed level of defence, and thus be little subject to change on individual basis. Also, as some  
31 species may show less or no growth response, we expected that increased shading from the  
32 responsive plants would cause defence decreases also for the less responsive ones.

## 2. Material and methods

### 2.1. Study area

The study area is southwest-exposed and located at Sanddalsnuten (60° N, 7°E) at Finse, southern Norway. The climate at Finse is alpine-oceanic. The mean summer temperature from June to August is 6.3°C (Aune, 1993) and the mean monthly precipitation is 89 mm (Førland, 1993). The vegetation consists of a *Dryas octopetala* heath alternating with alpine meadows. We collected lichens and leaves from common vascular plants inside and outside open top chambers (OTCs) in two locations differing in altitude, exposure, moisture, and productivity; ridge (1550 m) and leeward (1450 m). The ridge, close to the summit of Sanddalsnuten, is windy, with a ca 3 weeks longer growing season compared to the leeward, where snow accumulates and melts later. In the leeward, snow accumulation, in addition to water drainage from above results in ca 50 % higher soil moisture, ca 20 % higher content of soil organic matter as well as 6 times higher total C and more than 4 times higher total N (Olsen, 2010). Mean air temperature (July - August) was 8.7 °C at the leeward and 7.5 °C at the ridge and mean soil temperature (-5 cm) was 7.5 °C at the leeward and 7.2 °C at the ridge (Tinytag 12 Plus G data loggers, Intab Interface-Teknik AB, Stenkullen Sweden). The OTCs had been permanently established for 3 (leeward) and 8 (ridge) years prior to the sampling, and increased mean air temperature by ca 1.5 °C and soil temperature by ca 1.0 °C in both the leeward (Sandvik and Eide, 2009) and the ridge (Klanderud and Totland, 2005). These moderate increases in temperature correspond well with the predicted increase in summer temperature for this area the next 50-100 years (Hanssen-Bauer and Førland, 2001; Christensen et al., 2007). Open top chambers are commonly used to increase growth season temperature with minimal unwanted side effects on other environmental factors, such as light, precipitation and gas exchange (Arft et al., 1999; Hollister et al., 2000). Moreover, soil analyses inside and outside OTCs after four treatment years at the ridge site at Finse showed no differences in soil moisture (unpublished data K. Klanderud). Open top chambers may act as a physical barrier for some groups of herbivores, and thus be a potential confounding effect with increased T. We did not register herbivory inside and outside the OTCs systematically, but observed that insect larvae, lemmings and bigger herbivores (hares) occasionally were feeding also on plants inside OTCs (K. Klanderud and S. M. Sandvik, personal observation). For more details on the experimental setups see Klanderud and Totland (2005) and Sandvik and Eide (2009).

### 2.2 Measurements of vegetation height

1 Vegetation height was measured from the ground to the tallest point of the tallest plant at  
2 eight points inside each of 10 OTCs and 10 control plots at each location (ridge and leeside)  
3 in the beginning of August.

### 4 5 2.3 Sampling of leaves and lichens

6 We collected leaves from the five vascular plant species of four functional groups that were  
7 growing in all plots in either leeside and/or ridge: *Saussurea alpina* L. (perennial forb, ridge),  
8 *Tofieldia pusilla* (Michx.) Pers. (perennial forb, both sites), *Carex vaginata* (Tausch.) (sedge,  
9 ridge), *Vaccinium uliginosum* L. (dwarf-shrub, ridge), and *Selaginella selaginoides* L. (club  
10 moss, both sites). Furthermore, we collected thalli of six lichen species; *Flavocetraria nivalis*  
11 (L.) Kärnefelt & Thell, *Cetraria islandica* (L.) Ach, *Cladonia arbuscula* (Wallr.) Flot.,  
12 *Peltigera aphthosa* (L.) Willd., and *Stereocaulon* spp. (all in both sites), and *Thamnolia*  
13 *vermicularis* (Sw.) Schaer. (ridge). *Peltigera aphthosa* is a tripartite lichen with cyanobacteria  
14 in the cephalodia, while the other species have green algal photobionts only. We collected  
15 samples as a mix of three individuals in 10 OTCs and 10 control plots (some exceptions when  
16 species were absent, see Table 1) in each location on August 5th 2007. Plants and lichens  
17 were always sampled from the central part of the OTCs, as plants near the walls may have a  
18 different chemistry due to the UV-resistant Plexiglas (3 mm Lexan®Exell). Leaf and lichen  
19 samples were put in small paper bags, and left to dry in room temperature for two weeks or  
20 two days, respectively. This is the preferred method for drying plant material for later analysis  
21 of phenolic compounds (Julkunen-Tiitto and Sorsa, 2001). The samples were then stored in a  
22 freezer (-18°C) until extraction. Before extraction, the samples were kept at room temperature  
23 over night. We measured the dry weight (DW) and then removed the main veins and stems  
24 from leaves with a scalpel. From *S. selaginoides* we used all material from the upper 1 cm of  
25 one stem. The sample was then transferred to pre-weighed Eppendorf vials containing one  
26 conic stainless steel bead of 5 mm diameter. We crushed the sample to powder for 2 min in a  
27 Retsch mixer mill (Model MM301) at frequency 30.0 before it was weighed into to batches,  
28 one for analyses for C and N and one for extraction of CBSCs.

### 29 30 2.4 Chemical Analyses

31 Carbon and nitrogen concentrations were quantified at the Department of Animal and  
32 Aquacultural Sciences (Norwegian University of Life Sciences, Ås, Norway) using the CHN-  
33 N method with an EA 1108 Elementar Analyser (Fison) (Säntis Analytical Scandinavia AB,  
34 Läby Österby, 75592 Uppsala). Before the analysis of CBSCs (according to Julkunen-Tiitto et



1 al., 1996), leaf samples were extracted by adding 600  $\mu\text{l}$  methanol (MeOH) and mixed with an  
2 Ultra-Turrax homogenizer for 30 sec. The sample was then placed in an ice bath for 15 min,  
3 homogenized for 15 sec, centrifuged 15 000 rpm for 3 min and then the supernatant was  
4 poured into a clean glass tube. The residue was added 600  $\mu\text{l}$  MeOH, homogenized for 15sec  
5 and again centrifuged. The last procedure was repeated twice, and the residue was then totally  
6 colourless. Lichen samples were extracted according to Nybakken and Julkunen-Tiitto (2007)  
7 by adding 500  $\mu\text{l}$  acetone and vortexing the sample for 30 s before it was left to stand for 10  
8 minutes before the supernatant was poured off. This procedure was repeated three times. For  
9 both sample types the supernatants were combined and the MeOH or acetone evaporated with  
10 gaseous nitrogen. The dried extracts were stored at  $-18^{\circ}\text{C}$  until analysis.

11

12 The leaf extracts were dissolved in 300 $\mu\text{l}$  MeOH, added 300 $\mu\text{l}$  Milli-Q water and analysed on  
13 HPLC as described in Julkunen-Tiitto et al. (1996). We identified the compounds according to  
14 retention times and UV-spectra, quantified them at 220, 320 or 360 nm, and calculated the  
15 concentrations using the following commercial standards (supplier in parenthesis): caffeic  
16 acid (Aldrich, Steinheim, Germany), chlorogenic acid (Aldrich), 4-hydroxycinnamic acid  
17 (Aldrich), salidroside (Thieme, Germany), (+) catechin (Aldrich), myricetin-3-rhamnoside  
18 (Apin Chemicals, Abingdon, UK), quercetin-3- glucoside (Extrasynthese), apigenin-7-  
19 glucoside (Roth), luteolin-7-glucoside (Extrasynthese). As compounds within the same  
20 chemical group generally responded similarly to the treatments in the studied species (Table  
21 1), we chose to present concentrations ( $\text{mg g}^{-1}$  DW) and statistics for compound groups, and  
22 not for individual compounds when appropriate (Table 1).

23

24 The lichen extracts were dissolved in 500  $\mu\text{l}$  acetone and analysed on HPLC according to  
25 (Nybakken and Julkunen-Tiitto, 2007). The detection wavelength was 245 nm, and the  
26 identification of compounds was based on retention times, online UV-spectra, co-  
27 chromatography of commercial standards (atranorin, fumarprotocetraric acid (Apin  
28 Chemicals), usnic acid (Sigma)) and standards of baeomycecetic acid, squamatic acid, tenuiorin,  
29 gyrophoric acid and lobaric acid was provided by Dr. H.J. Sipman (Botanischer Garten und  
30 Botanischer Museum Berlin-Dahlem, Berling, Germany). The compounds were quantified  
31 against response curves of the above-mentioned standards. Concentrations of  
32 methylgyrophoric acid were calculated from the response curve of gyrophoric acid.

33

34 *2.5 Statistical analyses*

1 Two-way ANOVAs were run with the statistical package, SPSS 15.0.1 for Windows, with  
2 Treatment (control/OTC), Location (leeside/ridge) and the interaction Treatment  $\times$  Location  
3 as fixed factors, and with concentration of C, N or CBSCs as response variables. One-way  
4 ANOVAs were used when species occurred only in one of the locations. Number of samples  
5 analyzed of the different species from the different treatments and locations can be found in  
6 Table 1.

7

8

### 9 **3. Results**

10

#### 11 *3.1. Vegetation height*

12 The vegetation canopy was taller inside OTCs than in controls (leeside, ca 2.4 cm outside and  
13 4.1 cm inside the OTCs; ridge ca 2.1 cm outside and 2.8 cm inside the OTCs) ( $p = 0.003$ ).

14

#### 15 *3.2. Carbon and nitrogen*

16 The C concentration in the vascular plants varied between 435 (*S. alpina*) and 512  
17 (*Vaccinium uliginosum*)  $\text{mg g}^{-1}$  DW, while the corresponding values for lichens were  
18 between 386 (*Cetraria islandica*, leeside) and 454  $\text{mg g}^{-1}$  (*Peltigera aphthosa*, ridge) (Table  
19 1). The difference in N concentration was much more pronounced; between 15.7 (*Tofieldia*  
20 *pusilla*, ridge and leeside) and 25.2  $\text{mg g}^{-1}$  DW (*V. uliginosum*, ridge) for vascular plants and  
21 as low as between 5 and 10  $\text{mg g}^{-1}$  DW for green algal lichens. The tripartite lichen *P.*  
22 *aphthosa* with cyanobacteria in cephalodia had an N concentration comparable with vascular  
23 plants, varying between 23 and 25  $\text{mg g}^{-1}$  DW (Table 1).

24

25 The experimental warming decreased the N concentration in *Carex vaginata*, *Saussurea*  
26 *alpina* and *Selaginella selaginoides*, while it increased in the lichen *Thamnolia vermicularis*.  
27 In all plants, the carbon concentration was unaffected. The carbon concentration in *P.*  
28 *aphthosa* was lower inside the OTCs, and the same tendency was seen for most of the other  
29 lichens, although not statistically significant. Two plants (*S. selaginoides* and *T. pusilla*) and  
30 five lichens (*C. islandica*, *Flavocetraria nivalis*, *Cladonia arbuscula*, *P. aphthosa* and  
31 *Stereocaulon* spp.) were analyzed from both ridge and leeside. For the plants, there were no  
32 location effects on their total C and N concentrations. In contrast, the C concentration in  
33 lichen thalli from the ridge was significantly higher than in those from the leeside for all

1 species except *F. nivalis* (Table 1). The N concentration was significantly higher at the leeside  
2 for *C. arbuscula*, but was not influenced by location in any of the other lichen species. The  
3 interaction Treatment  $\times$  Location was not statistically significant for any of the studied taxa  
4 (results not shown).

5

### 6 3.3. Carbon based secondary compounds

7 The identified CBSCs of the vascular plants were grouped according to their aglycon or as  
8 phenolic acids in Table 1. In *C. vaginata* and *T. pusilla* the CBSCs constituted around 5 % of  
9 the DW. *Selaginella selaginoides* contained only between 2 and 4 %, while *S. alpina* and *V.*  
10 *uliginosum* had as much as from 12 up to 40 % CBSCs (Table 1, Figure 1).

11

12 Lichens generally contained fewer CBSCs, with the individual compounds identified listed in  
13 Table 1. The studied *C. arbuscula* and *F. nivalis* specimens contained only usnic acid in  
14 measurable amounts. In *C. islandica* we identified fumarprotocetraric acid and one compound  
15 following shortly after it in the chromatogram and with similar UV-spectrum. This compound  
16 was tentatively named "fumarprotocetraric acid derivative". *Peltigera aphthosa* contained  
17 tenuiorin and methylglyphoric acid, while the *Stereocaulon* species contained lobaric acid  
18 and atranorin, and is thus probably *Stereocaulon alpinum* (Krog et al., 1994). The *T.*  
19 *vermicularis* population growing in our experimental field contained squamatic acid and  
20 baecomycesic acid, and thus belonged to the chemotype II according to Krog et al. (1994). The  
21 total concentration of CBSCs of the lichens varied between 1.2 % (*P. aphthosa*, leeside) and  
22 6.0 % (*T. vermicularis*) of the DW (Table 1, Figure 2).

23

24 The warming significantly affected the CBSCs in only one vascular plant species (*T. pusilla*)  
25 and in one lichen species (*C. arbuscula*) (Table 1, Figures 1, 2). Nearly all compounds in *T.*  
26 *pusilla* (except the apigenin-glycosides) decreased inside the OTCs. In *S. selaginoides*, all  
27 individual CBSCs had the highest concentration at the ridge (not statistically significant for  
28 the phenolic acids). For *T. pusilla* the opposite was found; all compounds were highest at the  
29 leeside (not significant for the apigenin-glycosides). In the lichen species, four species had  
30 higher total concentration of secondary compounds at the ridge, while *C. arbuscula* had a  
31 higher concentration at the leeside (Figure 2). If the species contained more than one  
32 secondary compound, the pattern was the same for all compounds that had different  
33 concentration at the two sites. The interaction Treatment  $\times$  Location was not statistically  
34 significant for any studied species (results not shown).

1

Accepted Manuscript

#### 1 4. Discussion

2 Experimental warming in arctic-alpine environments often leads to increased growth of some  
3 plant species, while others are less responsive and often out-competed over the long run (Arft  
4 et al., 1999; Walker et al., 2006). According to resource-based hypotheses on plant defence  
5 (summarized by Herms and Mattson, 1992), we expected that warming would reduce C-based  
6 defence in arctic-alpine plants because of increased growth, and also that less growth-  
7 responsive plants and lichens would have less C resources for defence because of increased  
8 shading from more growth-responsive plants.

9  
10 The CBSC concentrations decreased with warming in one plant (*Tofieldia pusilla*) and one  
11 lichen (*Cladonia arbuscula*). All other plant and lichen species, however, showed no response  
12 in CBSC concentrations, although the vegetation height increased significantly inside OTCs.  
13 There are few earlier published studies of effects of warming on plant defence in the arctic-  
14 alpine, but our results are in line with those that exist, as there were either no effect (*Salix*  
15 *polaris* (Dormann, 2003), *Bistorta vivipara*, *Dryas octopetala* and *Salix reticulata* (Nybakken  
16 et al., 2008) or small decreases (*Cassiope tetragona* and *Salix herbacea* × *polaris* (Hansen et  
17 al., 2006) in CBSCs. So, in contrast to our expectations, many species did not reduce their  
18 defence levels when T increases. One explanation could be that growth did not increase much  
19 in most plants and lichens inside the OTCs. However, three of the plant species (*Saussurea*  
20 *alpina*, *C. vaginata* and *S. selaginoides*) had lower leaf N concentrations in the OTCs  
21 compared to the controls at the ridge, with the same tendency for *T. pusilla*, *Vaccinium*  
22 *uliginosum* and *S. selaginoides* in the leeward side. Comparable experiments with plants in arctic-  
23 alpine environments have either shown no effect of warming on leaf N content (*S. polaris*,  
24 (Dormann, 2003); *Oxyria digyna* and *Carex stans*, (Tolvanen and Henry, 2001) or a decrease  
25 (*C. tetragona*, *S. herbacea* × *polaris* and *Vaccinium vitis-idaea*, (Hansen et al., 2006); *C.*  
26 *tetragona*, *Dryas integrifolia* and *Salix arctica*, (Tolvanen and Henry, 2001); *Cerastium*  
27 *cerastoides*, *Epilobium anagallidifolium*, and *Carex lachenalii* (Sandvik and Eide, 2010).  
28 This suggests that there was no or only minor increase in soil N mineralization, and that  
29 decreased leaf N concentrations were results of dilution when growth increased. Generally, N  
30 mineralization rates are less responsive to warming in tundra than in forested ecosystems  
31 (Rustad et al., 2001), and the duration of our experiments have possibly been too short to see  
32 tissue-effects. Mineralization rates increased after 9 years of experimental warming in tussock  
33 tundra in arctic Alaska (Chapin et al., 1995). One lichen species, *T. vermicularis*, showed  
34 increased N concentrations in the OTCs, which may be a result of improved N uptake (from

1 rainwater or dew) at higher T. Obviously, lichens are not able to take up N from the soil  
2 (Nash, 2008).

3

4 Although C-based plant defence is expected to be resource based, it is also thought that some  
5 level of defence is constitutive (fixed) and would be synthesized in conjunction with growth  
6 (Tuomi et al., 1988, Holopainen et al., 1995; Stamp, 2003). High proportions of constitutive  
7 defence is expected to be more common in slow growing perennials and under limiting  
8 conditions (typically many arctic-alpine plants) than in annuals, pioneer plants and under less  
9 limiting conditions (Tahvanainen et al., 1985; Coley, 1987; Folgarait et al., 1994; 1995).

10 Only one of our species responded to the warming in defence levels, the perennial forb *T.*  
11 *pusilla*. The sedge, *C. vaginata*, and the other forb, *S. alpina*, could be expected to show the  
12 same response, but under the limiting conditions at this mid-alpine site one may probably  
13 expect high proportions of constitutive defence not only in woody species, but also in forbs  
14 and sedges. If growth increased, the increased C requirements to maintain high defence levels  
15 were probably met by T-increased photosynthesis, as none of the plants showed reductions in  
16 total C (Table 1). The C/N varied little between the vascular plant species, but the total CBSC  
17 concentrations did. This could be seen as a further support for a high level of constitutive  
18 defence in at least two of the species, as they differed so much from the others: *Vaccinium*  
19 *uliginosum* had almost 3 times the concentration of *S. alpina*, and more than six times the  
20 concentration of the rest of the species. The high level of (constitutive) defence in the woody  
21 *V. uliginosum* is according to the predictions of the CNB hypothesis (Bryant et al., 1983), but  
22 we have no explanations why *S. alpina* should be better defended/have another strategy than  
23 the rest of the species studied. Further complicating our interpretation is the fact that many  
24 arctic-alpine plants are clonal (in this study: *C. vaginata* and *V. uliginosum*), which means  
25 that resources may be transferred through rhizomes beyond the borders of OTCs, and thus for  
26 example reducing the effect of increased growth on resources available for defence. In  
27 summary, it would be difficult to prove that a defence level is fixed, as we cannot know what  
28 would happen if we for example increased the T with 1 °C or improved the nutrient  
29 availability by fertilization. However, in an earlier study from Sanddalsnuten, where T  
30 increase was combined with fertilization, the CBSC levels were reduced in the dwarf shrub  
31 *Salix reticulata*, while they stayed unchanged in the forb *Bistorta vivipara* and in the dwarf  
32 shrub *Dryas octopetala* (Nybakken et al., 2008). These results suggest that some species may  
33 have a fixed defence, while others are more subject to change, also under limiting conditions.

34

1 Most lichen species had a tendency to reduced total C inside OTCs, although statistically  
2 significant only for *P. aphthosa*, which is probably a result of the increased height of the plant  
3 canopy, leaving the low stature lichens in shade. This may be the first step towards carbon  
4 “starvation” of the lichens, as an earlier warming study from the same mountain slope showed  
5 that lichens decreased in abundance already after four years’ warming (Klanderud and  
6 Totland, 2005), confirming a general trend shown in the arctic-alpine (Cornelissen et al.,  
7 2001; Walker et al., 2006). The effect of shading for the C economy of lichens is clearly seen  
8 if we compare the two experiments from two different habitats; all six species sampled from  
9 both habitats had higher C concentrations on the ridge than in the leese, and the same was  
10 true for CBSCs for five of them (Table 1, Figure 2). As described in Material and Methods,  
11 the ridge is a more exposed habitat than the leese, and the vegetation height in both control  
12 plots and OTCs is on average highest in the leese and adds to the original light gradient.  
13 Cortical lichen CBSCs have been shown to increase along light gradients, both in transplanted  
14 lichens (usnic acid, Nybakken et al., 2007) and in lichens collected from their original habitat  
15 (atranorin, Solhaug et al., 2009). These compounds are situated above the algal layer in the  
16 lichen thallus, where they function as solar screens (e.g. Gauslaa et al., 2001; McEvoy et al.,  
17 2007). Our study shows that also CBSCs situated in the interior of lichens, in the medulla,  
18 have higher concentrations at the more exposed ridge (fumarprotocetraric acid in *Cetraria*  
19 *islandica*, tenuiorin in *P. aphthosa* and lobaric acid in *Stereocaulon*) compared to the leese.  
20 This may suggest that also medullary compounds have functions in solar protection, e.g. as  
21 antioxidants or even as screening compounds for lower layers of the lichen. No such pattern  
22 was seen for vascular plants, which further suggests that shading of plants has not been a  
23 factor in this study (but mark that only two plant species were studied from both habitats and  
24 that habitat is not repeated!).

25  
26 In conclusion, the lack of warming effects on CBSC levels in the studied plants and lichens,  
27 suggests that the defence levels are rather robust against raised temperatures, at least on a  
28 short-term basis. The robustness of plant defence in the arctic-alpine should be tested further,  
29 and a first step could be to grow a set of species from different functional groups under  
30 controlled light and nutrient conditions, searching for an optimum. At the moment, the threat  
31 for lichens, and possibly also for some of the plants, seem to be competition from other plants,  
32 rather than reduced defence in the first place. However, as warming could also improve  
33 conditions for e.g. herbivores and fungal diseases otherwise (milder winters, increased  
34 humidity (Hanssen-Bauer and Førland, 2001; Christensen et al., 2007), attacks may anyway

1 increase in the future, and might require further development of the defence, both  
2 qualitatively and quantitatively.

3

4

### 5 **Acknowledgements**

6 We thank Annie Aasen (Norwegian University of Life Sciences, Ås, Norway) for assistance  
7 with the analyses of CBSCs, Prof. Riitta Julkunen-Tiitto (University of Eastern Finland) for  
8 help with the identification of compounds, Siri L. Olsen (University of Agder, Kristiansand,  
9 Norway) for logging temperatures, and Prof. Yngvar Gauslaa (Norwegian University of Life  
10 Sciences, Ås, Norway) for valuable discussions and comments on the manuscript. We are also  
11 indebted to Finse Alpine Research Center for hospitality during field work. The study was  
12 financially supported by the University of Agder.

13

Accepted Manuscript



1

2 **References**

3 ACIA. 2005. Arctic Climate Impact Assessment. Cambridge: Cambridge University Press.

4 Arft, A.M., Walker, M.D., Gurevitch, J., Alatalo, J.M., Bret-Harte, M.S., Dale, M., Diemer,  
5 M., Gugerli, F., Henry, G.H.R., Jones, M.H. et al., 1999. Responses of tundra plants to  
6 experimental warming: meta-analysis of the International Tundra Experiment. *Ecol. Monogr.*  
7 69, 491-511.

8

9 Arft, A.M., Walker, M.D., Gurevitch, J., Alatalo, J.M., Bret-Harte, M.S., Dale, M., Diemer,  
10 M., Gugerli, F., Henry, G.H.R., Jones, M.H., Hollister, R.D, Jonsdottir, I.S., Laine, K.,  
11 Lévesque, E., Marion, G.M., Molau, U., Mølgaard, P., Nordenhäll, U., Raszhivin, V.,  
12 Robinson, C.H., Starr, G., Stenström, A., Stenström, M., Totland, O., Turner, P.L., Walker, J.,  
13 Webber, P.J., Welker, J.M., Wookey, P.A. 1999. Responses of tundra plants to experimental  
14 warming: Meta-analysis of the international tundra experiment. *Ecological Monographs*, 69:  
15 491-511.

16

17 Aune, B., 1993. Air Temperature Normals, normal period 1961-1990. Oslo. The Norwegian  
18 Meteorological Institute.

19

20 Baldwin, I.T., Oesch, R., Merhige, P., Hayes, K. 1993. Damage-induced root nitrogen  
21 metabolism in *Nicotiana sylvestris*: testing C/N predictions for alkaloid production. *J. Chem.*  
22 *Ecol.* 19, 3029-3043.

23

24 Bliss, L. C., 1962. Adaptation of arctic and alpine plants to environmental conditions. *Arct.*  
25 *Alp. Res.* 15, 117-144.

26

27 Bonan, G.B., Van Cleve, K., 1992. Soil temperature, nitrogen mineralization, and carbon  
28 source-sink relationships in boreal forests. *Can. J. Forest Res.* 22, 629-639.

29

30 Bryant, J.P., 1987. Feltleaf willow-snowshoe hare interactions: plant carbon/nutrient balance  
31 and floodplain succession. *Ecology* 68, 1319-1327.

32

33 Bryant, J.P., Chapin, F.S., Klein, D.R., 1983. Carbon/nutrient balance of boreal plants in  
34 relation to vertebrate herbivory. *Oikos* 40, 357-368.

- 1  
2 Bryant, J.P., Reichardt, P.B., Clausem, T.P., Werner, R.A., 1993. Effects of mineral nutrition  
3 on delayed inducible resistance in Alaska paper birch. *Ecology* 74, 2072-2084.  
4  
5 Callaghan, T.V., Jonasson, S. 1995. Arctic terrestrial ecosystems and environmental change.  
6 *Phil. Trans. Royal Soc. London.* A352, 259-276.  
7  
8 Chapin, F.S., Johnson, D.A., McKendrick, J.D., 1980. Seasonal movement of nutrients in  
9 plants of differing growth form in an Alaskan tundra ecosystem-implications for herbivory. *J.*  
10 *Ecol.* 68, 189-209.  
11  
12 Chapin, F.S. III, Shaver, G.R., Giblin, A.E., Nadelhoffer, K.J., Laundre, J.A., 1995.  
13 Responses of arctic tundra to experimental and observed changes in climate. *Ecology* 76, 694-  
14 711.  
15  
16 Christensen, J.H., Hewittson, B., Busuioc, A., Chen, A., Gao, X., Held, I., Jones, R., Kwon,  
17 W.-T., Laprise, R., Rueda, V.M., Mearns, L.O., Menéndez, C. G., Räisänen, J., Rinke, A.,  
18 Kolli, R.K., Sarr A. and Whetton, P., 2007. Pp 847-940 in D. Qin, M. Manning, Z. Chen, M.  
19 Marquis, K.B. Averyt, M. Tignor and H.L. Miller (eds.). *Regional climate projections.*  
20 *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the*  
21 *Intergovernmental Panel on Climate Change Fourth Assessment Report*, S. Solomon,  
22 Cambridge University Press, Cambridge.  
23  
24 Coley, P.D., 1987. Interspecific variation in plant anti-herbivore properties - the role of  
25 habitat quality and rate of disturbance. *New Phytol.* 106, 251-263.  
26  
27 Coley, P.D., Massa, M., Lovelock, C.E., Winter, K., 2002. Effects of elevated CO<sub>2</sub> on foliar  
28 chemistry of saplings of nine species of tropical tree. *Oecologia* 133, 62-69.  
29  
30 Cornelissen, J.H.C., Callaghan, T.V., Alatalo, J.M., Michelsen, A., Graglia, E., Hartley, A.E.,  
31 Hik, D.S., Hobbie, S.E., Press, M.C., Robinson, C.H., Henry, G.H.R., Shaver, G.R., Phoenix,  
32 G.K., Gwynn-Jones, D., Jonasson, S., Chapin III, F.S., Molau, U., Neill, C., Lee, J.A.,  
33 Melillo, J.M., Sveinbjörnsson, B., Aerts, R. 2001. Global change and arctic ecosystems: is  
34 lichen decline a function of increases in vascular plant biomass ?. *J Ecol.* 89, 984-994.

- 1  
2 Dormann, C.F. 2003. Consequences of manipulations in carbon and nitrogen supply for  
3 concentration of anti-herbivore defence compounds in *Salix polaris*. *Ecoscience* 10, 312-318.  
4
- 5 Dormann, C.F., Woodin, S.J. 2002. Climate change in the Arctic: using plant functional types  
6 in a meta-analysis of field experiments. *Functional Ecol.* 16, 4-17.  
7
- 8 Emmerich, R., Giez, I., Lange, O.L., Proksch, P., 1993. Toxicity and antifeedant activity of  
9 lichen compounds against the polyphagous herbivorous insect *Spodoptera littoralis*.  
10 *Phytochemistry* 33, 1389-1394.  
11
- 12 Folgarait, P.J., Davidson, D.W., 1994. Antiherbivore defences of *Myrmecophytic cecropia*  
13 under different light regimes. *Oikos* 71, 305-320.  
14
- 15 Folgarait, P.J., Davidson, D.W., 1995. *Myrmecophytic cecropia* - antiherbivore defences under  
16 different nutrient treatments. *Oecologia* 104, 189-206.
- 17 Fritz, C., Palacios-Rojas, N., Feil, R., Stitt, M., 2006. Regulation of secondary metabolism by  
18 the carbon-nitrogen status in tobacco: nitrate inhibits large sectors of phenylpropanoid  
19 metabolism. *Plant J.* 46, 533-548.  
20
- 21 Fritz, C., Palacios-Rojas, N., Feil, R., Stitt, M. 2006. Regulation of secondary metabolism by  
22 the carbon-nitrogen status in tobacco: nitrate inhibits large sectors of phenylpropanoid  
23 metabolism. *Plant J.* 46, 533-548.  
24
- 25 Førland, E.J., 1993. Precipitation Normals, Normal Period 1961-90. The Norwegian  
26 Meteorological Institute, Oslo.  
27
- 28 Gauslaa, Y., 2005. Lichen palatability depends on investments in herbivore defence.  
29 *Oecologia* 143, 94-105.  
30
- 31 Gauslaa, Y., McEvoy, M., 2005. Seasonal changes in solar radiation drive acclimation of the  
32 sun-screening compound parietin in the lichen *Xanthoria parietina*. *Basic Appl. Ecol.* 6, 75-  
33 82.  
34

- 1 Gauslaa, Y., Solhaug, K.A., 2001. Fungal melanins as a sun screen for symbiotic green algae  
2 in the lichen. *Lobaria pulmonaria*. *Oecologia* 126, 462-471.  
3
- 4 Gauslaa, Y., Ustvedt, E.M., 2003. Is parietin a UV-B or a blue-light screening pigment in the  
5 lichen *Xanthoria parietina*? *Photochem. Photobiol. Sci.* 2, 424-432.  
6
- 7 Hansen, A.H., Jonasson, S., Michelsen, A., Julkunen-Tiitto, R., 2006. Long-term  
8 experimental warming, shading and nutrient addition affect the concentration of phenolic  
9 compounds in arctic-alpine deciduous and evergreen dwarf shrubs. *Oecologia* 147, 1-11.  
10
- 11 Hanssen-Bauer, I., Førland, E.J. 2001. Verification and analysis of a climate simulation of  
12 temperature and pressure fields over Norway and Svalbard. *Clim. Res.* 16, 225-235.  
13
- 14 Herms, D.A., Mattson, W.J. 1992. The dilemma of plants: to grow or to defend. *Q. Rev. Biol.*  
15 67, 283-335.  
16
- 17 Hollister, R.D., Webber, P.J., 2000. Biotic validation of small open-top chambers in a tundra  
18 ecosystem. *Global Change Biol.* 6, 835-842.  
19
- 20 Holopainen, J.K., Rikala, R., Kainulainen, P., Oksanen, J. 1995. Resource partitioning to  
21 growth, storage and defence in nitrogen-fertilized Scots pine and susceptibility of the  
22 seedlings to the tarnished plant bug *Lygus rugulipennis*. *New Phytol* 131, 521-532.  
23
- 24 Iason, G.R., Hester, A.J. 1993. The response of heather to shade and nutrients: predictions of  
25 the carbon-nutrient hypothesis. *J. Ecol.* 81, 75-80.  
26
- 27 IPCC, 2007. *Climate Change 2007: The Physical Science Basis. Summary for Policymakers.*  
28 WMO 508 and UNEF, Geneva.  
29
- 30 Julkunen-Tiitto, R., Rousi, M., Bryant J., Sorsa S., Keinänen, M., Sikanen, H., 1996.  
31 Chemical diversity of several Betulaceae species: comparison of phenolics and terpenoids in  
32 northern birch stems. *Trees* 11, 16-22.  
33

- 1 Julkunen-Tiitto, R., Sorsa, S. 2001. Testing the effects of drying methods on willow  
2 flavonoids, tannins, and salicylates. *J. Chem. Ecol.* 27, 779-789.  
3
- 4 Klanderud, K., 2008. Species-specific responses of an alpine plant community under  
5 simulated environmental change. *J. Veg. Sci.* 19, 363-U109.  
6
- 7 Klanderud, K., Totland, Ø., 2005. Simulated climate change altered dominance hierarchies  
8 and diversity of an alpine biodiversity hotspot. *Ecology* 86, 2047-2054.  
9
- 10 Koricheva, J., Larsson, S., Haukioja, E., Keinänen, M., 1998: Regulation of woody plant  
11 secondary metabolism by resource availability: hypothesis testing by means of meta-analysis.  
12 *Oikos*, 83: 212-226.  
13
- 14 Krog, H., Østhaugen, H., Tønsberg, T., 1994. *Lavflora*. Oslo: Universitetsforlaget.  
15
- 16 Lamontagne, M., Margolis, H.A., Bauce, E. 2000. Testing the ecophysiology basis for the  
17 control of monoterpene concentrations along canopy profiles in thinned and unthinned balsam  
18 fir stand. *Oecologia* 124, 318-331.  
19
- 20 Lawrey, J.D., 1983. Vulpinic and pinastric acids as lichen antiherbivore compounds-contrary  
21 evidence. *Bryologist* 86, 365-369.  
22
- 23 Leser, C., Treutter, D. 2005. Effects of nitrogen supply on growth, contents of phenolic  
24 compounds and pathogen (scab) resistance of apple trees. *Physiol. Plant.* 123, 49-56.  
25
- 26 McEvoy, M., Solhaug, K.A., Gauslaa, Y., 2007. Solar radiation screening in usnic acid-  
27 containing cortices of the lichen *Nephroma arcticum*. *Symbiosis* 43, 143-150.  
28
- 29 Matt, P., Krapp, A., Haake, V., Mock, H.-P., Stitt, M. 2002. Decreased Rubisco activity leads  
30 to dramatic changes of bitrate metabolism, amino acid metabolism and the levels of  
31 phenylpropanoids and nicotine in tobacco antisense *RBCS* transformants. *Plant J.*, 663-677.  
32

- 1 Molau, U., 1997. Responses to natural climatic variation and experimental warming in two  
2 tundra plant species with contrasting life forms: *Cassiope tetragona* and *Ranunculus nivalis*.  
3 *Global Change Biol.* 3, 97-107.  
4
- 5 Mole, S., Ross, A.M., Waterman, P.G. 1988. Light-induced variation in phenolic levels in  
6 foliage of rain-forest plants. I. chemical changes. *J. Chem. Ecol.* 14, 1-21.  
7
- 8 Muzika, R.M., Pregitzer, K.S., Hanover, J.W. 1989. Changes in terpene production following  
9 nitrogen fertilization of grand fir (*Abies grandis* (Dougl.) Lindl.) seedlings. *Oecologia* 80,  
10 485-489.  
11
- 12 Nadelhoffer, K.J., Giblin, A.E., Shaver, G.R., Laundre, J.A., 1991. Effects of Temperature  
13 and Substrate Quality on Element Mineralization in 6 Arctic Soils. *Ecology* 72, 242-253.  
14
- 15 Nash, T.H. 2008. Nutrients, elemental accumulation, and mineral cycling. In: Nash, T.H.(ed.),  
16 *Lichen Biology*, Cambridge University Press, pp 234-251.  
17
- 18 Nichols-Orians, C.M. 1991. The effect of light on foliar chemistry, growth and susceptibility  
19 of seedlings of a canopy tree to an attine ant. *Oecologia* 86, 552-560.  
20
- 21 Nybakken, L., Asplund, J., Solhaug, K.A., Gauslaa, Y., 2007. Forest successional stage  
22 affects the cortical secondary chemistry of three old forest lichens. *J. Chem. Ecol.* 33, 1607-  
23 1618.  
24
- 25 Nybakken, L., Julkunen-Tiitto, R., 2006. UV-B induces usnic acid in reindeer lichens.  
26 *Lichenologist* 38, 477-485.  
27
- 28 Nybakken, L., Klanderud, K., Totland, Ø. 2008. Simulated Environmental Change Has  
29 Contrasting Effects on Defensive Compound Concentration in Three Alpine Plant Species.  
30 *Arct. Antarct. Alp. Res.* 40, 709-715.  
31
- 32 Olsen, S.L. 2010. Do nitrogen-fixing legumes affect soil nutrient levels, plant growth or  
33 community properties of an alpine ecosystem? Master thesis, Norwegian University of Life  
34 Sciences.

- 1  
2 Pöykkö, H., Hyvärinen, M., Backor, M., 2005. Removal of lichen secondary metabolites  
3 affects food choice and survival of lichenivorous moth larvae. *Ecology* 86, 2623-2632.  
4
- 5 Richardson, S.J., Press, M.C., Parsons, A.N., Hartley, S.E. 2002. How do nutrients and  
6 warming impact on plant communities and their insect herbivores? A 9-year study from a sub-  
7 Arctic heath. *J. Ecol.* 90, 544-556.  
8
- 9 Sandvik, S.M., Eide, W., 2009. Costs of reproduction in circumpolar *Parnassia palustris* L. in  
10 light of global warming. *Plant Ecol.* 205, 1-11.  
11
- 12 Sandvik, S.M. Eide, W. 2010. Long-term experimental warming affects tissue C/N ratios  
13 differently in three strongly chionophilous alpine species. 187-198. *In: Global Warming in the*  
14 *21st Century. Ed.: Cossia, J. M. NOVA Science Publishers, Inc. ISBN: 978-1-61728-980-4.*  
15
- 16 Shaver, G.R., Chapin, F.S. III. 1991. Production: biomass relationships and element cycling in  
17 contrasting arctic vegetation types. *Ecol. Mon.* 61, 1-31.  
18
- 19 Solhaug, K.A., Gauslaa, Y., 1996. Parietin, a photoprotective secondary product of the lichen  
20 *Xanthoria parietina*. *Oecologia* 108, 412-418.  
21
- 22 Solhaug, K.A., Gauslaa, Y., Nybakken, L., Bilger, W., 2003. UV-induction of sun-screening  
23 pigments in lichens. *New Phytol.* 158, 91-100.  
24
- 25 Solhaug, K.A., Lind, M., Nybakken, L., Gauslaa, Y., 2009. Possible functional roles of  
26 cortical depsides and medullary depsidones in the foliose lichen *Hypogymnia physodes*. *Flora*  
27 204, 40-48.  
28
- 29 Solhaug, K.A., Larsson, P., Gauslaa, Y., 2010. Light screening in lichen cortices can be  
30 quantified by chlorophyll fluorescence techniques for both reflecting and absorbing pigments.  
31 *Planta* 231, 1003-1011.  
32
- 33 Stamp, N., 2003. Out of the quagmire of plant defense hypotheses. *Q. Rev. Biol.* 78, 23-55.  
34

- 1 Suding, K.N., Larson, J.R., Thorsos, E., Steltzer, H., Bowman, W.D. 2004. Species effects on  
2 resource supply rates: do they influence competitive interactions? *Plant Ecol.* 175, 47-58.  
3
- 4 Tahvanainen, J., Julkunen-Tiitto, R., Kettunen, J., 1985. Phenolic glycosides govern the food  
5 selection pattern of willow feeding leaf beetles. *Oecologia* 67, 52-56.  
6
- 7 Tolvanen, A., Henry, G.H.R., 2001. Responses of carbon and nitrogen concentrations in high  
8 arctic plants to experimental warming. *Can. J. Bot.* 79, 711-718.  
9
- 10 Tuomi, J., Fagerström, T., Niemelä, P., 1991. Carbon allocation, phenotypic plasticity, and  
11 induced defences. In: Tallamy, D.W., Raupp, M., J. (Eds), *Phytochemical induction by*  
12 *herbivores*. Wiley, New York, pp. 85-104.  
13
- 14 Tuomi, J., Niemelä, P., Chapin, F.S.I., Bryant, J. P., Sirén, S., 1988. Defensive responses of  
15 trees in relation to their carbon/nutrient balance. In: Mattson, W.J., Levieux, J., Bernard-  
16 Dagan, D. (Eds.), *Mechanisms of woody plant defenses against insects: Search for pattern*.  
17 Springer Verlag, New York, pp 57-72.  
18
- 19 White, A, Cannel MGR, Friend AD (1999) Climate change impacts on ecosystems and the  
20 terrestrial carbon sink: a new assessment. *Global Environmental Change*, 9, 21–30.  
21 Tim  
22  
23
- 24 Wookey, P.A., Aerts, R., Bardgett, R.D., Baptists, F., Bråthen, K.A., Cornelissen, J.H.C.,  
25 Gough, L., Hartley, I.P., Hopkins, D.W., Lavorel, S., Shaver, G.R. 2009. Ecosystem  
26 feedbacks and cascade processes: understanding their role in the responses of Arctic and  
27 alpine ecosystems to environmental change. *Glob. Change Biol.* 15, 1153-1172.  
28  
29



1 TABLE 1. Concentrations (mg g<sup>-1</sup> DW) of C, N and CBSCs ±S.E in vascular plants and lichens under ambient (control) and warmed (OTC)  
 2 conditions (treatment) in two different locations in alpine southern Norway<sup>1</sup>. Asterisks (\*) behind the F-values denotes significance levels  
 3 (\*P<0.05; \*\*P<0.01; \*\*\*P<0.001)  
 4

	<b>Ridge</b>		<b>Leaside</b>		<b>Treatment</b>	<b>Habitat</b>
	Control	OTC	Control	OTC	F	F
<b>Vascular plants</b>						
<i>Carex vaginata</i>	N = 10	N = 10				
C	455.1±0.9	453.8±1.3			0.78	
N	21.9±0.7	18.1±0.6			17.2***	
C:N	20.9±0.6	25.3±0.9			16.5***	
Luteolin-glyc.	35.9±4.1	45.6±5.6			1.74	
Apigenin-glyc.	2.2±0.2	3.1±0.5			2.11	
Sum, CBSCs	50.3±4.8	65.0±7.0			2.44	
<i>Saussurea alpina</i>	N = 10	N = 10				
C	434.9±1.7	432.7±1.9			0.71	
N	20.7±1.0	17.5±0.5			8.18**	
C:N	21.4±0.9	24.9±0.7			9.50**	
Phenolic acids	89.9±4.9	91.7±4.9			0.06	
Quercetin-glyc.	31.6±2.1	27.8±1.6			1.81	

Sum, CBSCs	121.5±4.6	119.5±5.4			0.07	
<i>Selaginella selaginoides</i>	N = 10	N = 10	N = 10	N = 10		
C	471.8±2.8	480.6±3.0	483.8±2.8	473.5±4.1	0.05	0.47
N	19.5±0.7	17.6±0.5	19.2±0.7	18.7±0.6	3.87	0.41
C:N	24.5±0.9	27.4±0.8	25.4±0.9	25.6±0.8	3.61	0.34
Phenolic acids	5.3±2.4	2.6±0.3	2.1±0.3	1.6±0.1	0.43	9.46
Apigenin der	0.3±0.1	0.4±0.1	0.1±0.02	0.2±0.03	1.53	2.71**
Kaempferol der	33.9±4.3	29.2±1.9	23.4±2.0	17.7±0.9	3.65	16.65**
Coumaryl-Kaempferols	2.8±0.3	3.3±0.3	2.1±0.2	1.7±0.1	0.08	20.63***
Sum, CBSCs	42.3±6.4	35.4±2.3	27.6±2.3	21.2±1.0	0.01	14.23***
<i>Tofieldia pusilla</i>	N = 4	N = 4	N = 10	N = 10		
C	448.3±2.9	445.6±3.6	450±1.2	446.9±1.9	1.83	0.48
N	14.9±0.5	15.7±1.0	14.9±0.5	15.4±0.9	0.19	0.02
C:N	30.2±1.2	28.9±2.1	30.5±1.0	29.6±1.5	0.22	0.01
Apigenin-glyc.	2.7±1.3	1.4±0.5	2.7±0.7	2.4±0.7	0.94	0.48
Quercetin-glyc.	14.0±2.5	7.6±1.5	16.3±0.9	12.8±2.0	8.84**	5.28*
Quercetin-diglyc.	14.9±2.7	9.6±1.3	17.4±0.6	14.9±2.0	5.31*	5.86*
Luteolin-glyc.	18.8±3.4	11.6±2.0	21.6±1.7	19.3±2.4	4.54	5.24*
Sum, CBSCs	50.3±8.8	30.1±4.6	58.1±2.9	49.3±5.8	7.01*	6.04*

<i>Vaccinium uliginosum</i>	N = 10	N = 10	
C	506.9±1.1	511.9±14.0	0.13
N	25.2±0.4	23.7±0.7	3.35
C:N	20.2±0.3	21.7±0.4	8.93**
Catechin der.	88.9±34.7	198.1±152	0.53
Phenolic acids	56.7±27.9	53.6±21.2	0.13
Myricetrin	19.9±7.8	11.5±7.7	0.02
Isoquercetin	143.1±68.3	110.5±52.3	2.57
Kaempferol der	26.6±11.1	24.4±11.9	1.44
Isorhamnetin	3.9±2.1	3.6±1.6	0.01
Sum, CBSCs	339.1±150.8	401.7±239.4	0.04

### Lichens

<i>Cetraria islandica</i>	N = 10	N = 10	N = 10	N = 10		
C	410.2±2.6	406.0±3.3	389.0±2.9	386.8±4.6	0.89	34.94***
N	5.7±0.2	5.5±0.1	5.6±0.2	6.0±0.4	0.24	0.58
C:N	72.8±2.3	73.9±2.0	70.1±1.6	67.2±4.7	0.09	2.57
Fumarprotocetraric acid	16.7±1.6	17.2±3.9	9.2±1.6	3.9±1.0	2.00	16.70***
Fumarprotocetraric acid der	8.7±0.9	8.3±1.9	6.2±1.3	10.4±1.6	1.39	0.01
Sum, CBSCs	25.4±2.5	25.5±5.7	15.2±1.8	14.3±1.0	0.65	16.47***

<i>Cladonia arbuscula</i>	N = 10	N = 10	N = 10	N = 10		
C	429±1.6	426.4±2.0	422.4±2.3	423.0±1.5	0.26	7.32**
N	5.6±0.3	5.5±0.3	5.8±0.4	6.5±0.2	1.09	4.39*
C:N	78.9±4.5	80.0±4.7	76.0±6.3	65.7±1.9	1.07	3.76
Usnic acid	36.4±1.3	32.0±1.4	46.6±4.9	39.8±12.6	4.13*	10.91*
<i>Flavocetraria nivalis</i>	N = 10	N = 10	N = 6	N = 6		
C	409.0±1.9	407.7±2.3	402.9±3.1	396.4±2.9	2.26	11.46**
N	5.1±0.3	4.7±0.3	5.4±0.4	5.3±0.2	0.54	1.63
C:N	82.5±5.3	91.6±7.2	76.0±5.3	75.3±3.7	0.41	3.04
Usnic acid	53.7±2.2	51.2±2.4	49.6±2.0	43.1±5.4	2.19	4.03
<i>Peltigera aphthosa</i>	N = 5	N = 5	N = 9	N = 9		
C	454.4±1.1	438.6±1.5	430.0±2.1	427.4±2.5	16.80***	61.98***
N	24.2±0.8	23.7±1.5	24.9±1.0	22.8±1.1	1.17	0.001
C:N	18.9±0.6	18.9±1.1	17.5±0.7	19.1±0.9	0.75	0.48
Methylglyphoric acid	1.7±0.1	1.7±0.5	1.3±0.3	1.4±0.2	0.02	1.97
Tenuiorin	18.5±1.7	17.0±2.1	11.7±0.9	10.5±0.9	1.00	25.35***
Sum, CBSCs	20.2±3.4	18.7±1.6	12.4±0.9	12.1±1.2	1.14	34.05***

<i>Stereocaulon</i> spp.	N = 10	N = 10	N = 10	N = 10		
C	423.4±3.3	422.7±2.2	412.9±1.5	414.7±1.6	0.05	17.02***
N	9.4±0.9	9.3±0.6	10.3±0.4	9.1±0.3	1.22	0.36
C:N	47.7±3.2	47.3±3.6	40.6±1.7	45.9±1.4	0.84	2.54
Lobaric acid	4.6±0.4	5.7±1.5	4.4±0.4	2.8±0.3	0.10	3.77
Atranorin	21.0±1.4	20.5±2.2	14.4±0.8	13.4±1.2	0.25	18.06***
Sum, CBSCs	25.6±2.1	26.2±3.4	18.8±1.1	16.1±1.3	0.004	14.80***
<i>Thamnia vermicularis</i>	N = 10	N = 10				
C	409.9±5.1	421.3±7.5			1.59	
N	5.7±0.4	7.0±0.2			8.92**	
C:N	76.3±6.3	60.7±2.9			5.08*	
Squamatic acid	23.0±0.6	22.1±1.3			0.16	
Baeomycesic acid	37.2±1.9	32.1±2.2			2.50	
Sum, CBSC	60.2±2.2	54.2±3.3			1.59	

1

2

1 **Figure legends**

2

3

4 Figure 1. Total concentration ( $\text{mg g}^{-1} \pm \text{S.E.}$ ) of phenolic compounds in plant leaves ( $\text{mg g}^{-1} \pm$   
5  $\text{S.E.}$ ) from OTCs (black bars) and controls (grey bars) at the ridge and the leese. Significant  
6 difference between controls and OTCs according to a one-way ANOVA is marked by \* =  
7  $p < 0.100$ , \*\* =  $p < 0.050$  and \*\*\* =  $p < 0.001$ .

8

9 Figure 2. Total concentration ( $\text{mg g}^{-1} \pm \text{S.E.}$ ) of phenolic compounds in lichen thalli from  
10 OTCs (black bars) and controls (grey bars) at the ridge and the leese. Significant difference  
11 between controls and OTCs according to a one-way ANOVA is marked by \* =  $p < 0.100$ , \*\* =  
12  $p < 0.050$  and \*\*\* =  $p < 0.001$ .

13

14 **Research Highlights**

15

16

17

18

19

20

21

22

23

24

25

- Defensive compounds in arctic-alpine vascular plants and lichens are little subject to change under increased temperature
- Plant competition and shading effects caused by elevated temperatures are likely to be more ecologically important than impacts on plant defence
- Defensive compounds in arctic-alpine lichens are strongly responsive to solar exposure, and this holds also for compounds situated in the medulla (probably no function as solar screen).




Fig. 2




Fig. 3