



UNIVERSITETET I AGDER

The Extent of Volatility Predictability

Evaluation of forecasting accuracy dependent on time, distribution and model order

Authors

Robert Dadson and Marta Slonka

Supervisor

Valeriy Zakamulin

This master's thesis is carried out as a part of the education at the University of Agder and is therefore approved as a part of this education. However, this does not imply that the University answers for the methods that are used or the conclusions that are drawn.

University of Agder, May 2015

School of Business and Law

Department of Economics and Finance

Abstract

This thesis focuses on the accuracy and ability of out-of-sample volatility forecasting over different time horizons. Using data at daily frequency we forecast the future volatility over multiple time horizons (1, 3, 6, 9 and 12 months) and evaluate the goodness of forecasting by comparing the Naïve, ARCH, GARCH, EGARCH and GJR-GARCH models using the MSE and the Predictive Power (P). We include different probability distributions for the error terms in an attempt to improve the models accuracy. The research is conducted using three indices: FTSE 100, S&P 500 and the Hang Seng. We find that the goodness of forecasting accuracy decreases dramatically after the 3 month horizon and the selection of a more representative error distribution improves the accuracy of the short term forecasts. The results also show that the higher order GARCH models, beyond (1,1), do not improve the forecasting accuracy.

Contents

CONTENTS	3
1. INTRODUCTION.....	4
2. LITERATURE REVIEW.....	7
3. DATA ANALYSIS	11
3.1 INDICES.....	11
3.2 DESCRIPTIVE STATISTICS	13
3.3 DISTRIBUTION	15
3.4 LJUNG-BOX Q-TEST	19
3.5 VOLATILITY.....	20
4. METHODOLOGY.....	23
4.1 NAIVE FORECASTING MODEL	23
4.2 ROLLING WINDOW	23
4.3 ARCH.....	24
4.4 GARCH.....	25
4.5 EGARCH	26
4.6 GJR-GARCH	27
4.7 DISTRIBUTION STATISTICS	27
Normal distribution (Gaussian)	27
Student t-distribution	28
Generalized error distribution.....	28
4.8 FORECAST EVALUATION	29
4.9 PREDICTIVE POWER.....	30
5. RESULTS	31
5.1 FTSE100	32
5.2 S&P 500	39
5.3 HSI	45
6. DISCUSSION	52
7. CONCLUSION.....	54
ACKNOWLEDGEMENTS	56
REFERENCES	57

1. Introduction

Volatility forecasting is of known importance for risk management, portfolio optimization and option pricing through its contribution to the measurement of potential future losses. Volatility is the uncertainty of asset prices over a given investment time horizon. When forecasted well the investor or financial institution is able to assess the investment risk accurately and achieve a level of risk they are willing to accept. The information provided by volatility is indispensable and was recognised in 2003 with the Nobel Memorial Prize in Economic Sciences being shared by Robert F. Engle and Clive Granger. In his prize lecture at the University of Stockholm, Engle went on to say:

“There are some risks we choose to take because the benefits from taking them exceed the possible costs. Optimal behaviour takes risks that are worthwhile. This is the central paradigm of finance; we must take risks to achieve rewards but not all risks are equally rewarded. Both the risks and the rewards are in the future, so it is the expectation of loss that is balanced against the expectation of reward. Thus we optimize our behaviour, and in particular our portfolio, to maximize rewards and minimize risks” (Engle (2003))

Volatility is estimated from historical data on asset returns while implied volatility uses observed option prices which, from a theoretical perspective, are a good estimate of future realized volatility as they include all relevant and available information. Investors and financial institutions use volatility as a key input in risk management models when forecasting. Much research has been devoted to modelling and forecasting the volatility of financial returns where the modelling of volatility for asset pricing purposes relies on the accurate assessment of future returns and risk. Model risk, as discussed in Figlewski (1998), suggests that there exists a risk that any given model may be misspecified. It is of necessity, when modelling, to specify the stochastic process of the underlying asset. Most existing papers use the log-normal process assumed by the original option-pricing model of Black and Scholes (1973), whereas the empirically correct distribution of the actual financial markets under analysis are likely to be non-normal. This thesis directly compares three different error distributions in order to overcome the difficulty in the judgment of the accuracy

of such models. Empirical finance literature has shown that volatility series develop over time in non-linear fashion due to the mean reverting properties of the models. We investigate the performance of volatility forecasting within the class of Autoregressive Conditional Heteroskedastic (ARCH) models due to their capability of capturing the non-linear features of financial time series and the possibility of extending these models to consider additional features.

The Autoregressive Conditional Heteroskedasticity (ARCH) model introduced by Engle (1982) was one of the first models that provided a way to model conditional heteroskedasticity in volatility. Bollerslev (1986) extended the ARCH model to the Generalized Autoregressive Conditional Heteroskedasticity (GARCH), which has the same basic properties as the ARCH model but requires far less parameters to satisfactorily model the volatility process. Both the ARCH and GARCH models are able to model the persistence and clustering of volatility but are unable to distinguish between the different influences that positive and negative shocks have on the volatility. To be able to model this behaviour and overcome the weaknesses of the ARCH and GARCH models, Nelson (1991) proposed an extension to the GARCH model called the Exponential GARCH (EGARCH), which is capable of taking into account the asymmetric effects of positive and negative asset returns. An additional extension of the GARCH model is the GJR-GARCH which was proposed by Glosten, Jagannathan and Runkle (1993).

The purpose of this thesis is to analyze the methods of forecasting volatility over long term horizons, the longest being 12 months, and compare the results with shorter forecasting time horizons. This thesis adds to the existing literature in several ways. First of all, we examine the importance of assumptions regarding the distributions of the error term. Previous studies have shown that the normal error distribution, which is widely applied in the majority of research, does not necessarily fit the models, nor give the most accurate forecasts (Speight, McMillan and Gwilym (2000)). Thus, in this thesis, three different error distributions are applied (Normal, Student t and Generalized Error Distribution). Secondly, we use various (from 1 month up to 12 months) forecasting horizons. There exist only a few papers that cover longer than 3 months forecasting periods because in practice, variability in volatility is often used in portfolio management and these are rebalanced at least once a month. The longer

horizons are thus of interest to those who, for example, try to find the correct price for financial instruments such as long-term options (called LEAPS - Long-term Equity Anticipation Securities), or warrants which mostly have an expiration of up to 1 year but in some countries up to 3 years. Much of the existing research only covers one time horizon and therefore the comparison of results directly between different indices and time horizons is difficult to establish.

Forecasting the future level of volatility over any horizon is far from straightforward and evaluating the forecasting performance presents an additional challenge. The majority of the 93 papers reviewed by Poon and Granger (2003) use the Mean Squared Error (MSE), or the Root Mean Squared Error (RMSE), which uses the actual and forecasted volatilities to compare the differences. Our third difference is that we also assess the Predictive Power (P) in order to find out how much of the future variability in volatility our models are able to predict. We do this by calculating the proportion of explained variability as proposed by Blair, Poon and Taylor (2010). The comparison of forecasting models for indices from three different continents is also rare since the majority of existing papers focus on multiple markets of individual countries. We have chosen to study the following three indices, FTSE 100, S&P 500 and Hang Seng.

We find that the level of forecasting accuracy decreases as the time horizon increases. The forecasting horizon which produces the most accurate statistics is that of 1 month for the FTSE 100 and S&P 500, and 3 months for the Hang Seng index. The GARCH models with a larger number of properties were found to be better predictors than the more restrictive models and higher order models beyond (1,1) do not significantly improve the accuracy. With the use of three different error distributions we generally see an improvement in accuracy when using the two non-normal distributions compared to that of the Gaussian distribution when considering shorter forecasting horizons.

The rest of the thesis is organized as follows. In Section 2, we review relevant literature and familiarize the reader with the existing research papers regarding volatility forecasting over long term horizons, as well as those that cover similar models and error distributions. In Section 3 we examine the characteristics of the data and analyze the descriptive statistics of the returns. Furthermore, we investigate the

distributions and continue with graphs and statistics of the realized monthly volatility. In Section 4 we describe the methodology for each model including how the forecasts are performed and the three error distributions used. We also explain how the models are going to be compared. In Section 5 we present and interpret the results of each index. The models are presented individually based on the shortest to longest forecast horizon. We then, in Section 6, discuss the important findings and contrast our results with those of literature reviewed. Section 7 provides a concise summary of our approach and main findings.

2. Literature review

The majority of the papers regarding volatility forecasting study the predictability over horizons shorter than 3 months (usually 1 month). Moreover, the model selection varies where not one individual model is preferred over others. Even when empirical work uses the same models and markets, the observations and evaluation methods differ, thus, the conclusions rarely agree. In this thesis we focus on the length of horizon predictability using the ARCH/GARCH family models whilst considering the model's sensitivity to distribution assumptions and time horizons. Below we review some relevant papers regarding these matters.

Poon and Granger (2003) make a review of papers regarding volatility forecasting. They compare 93 papers under various models, distributions and forecasting horizons. The review is clear evidence that researchers mostly prefer to forecast volatility for short time horizons yet there are a few relevant papers which focus on longer horizons and these are revealed below.

Scott and Tucker (1989) study the predictive accuracy of standard deviations and the constant elasticity of variance (CEV) implied in American currency call options traded on the Philadelphia Stock Exchange (PHLX). Five currencies are used, namely the British pound, Canadian dollar, German deutschmark, Japanese yen, and the Swiss franc. The authors accumulate daily closing prices and forecast subsequent currency return volatility for 3, 6 and 9 months. The obtained forecasting results are evaluated using the MSE. The authors conclude that the MSEs are statistically

indistinguishable and suggest that the simple Black-Scholes predictions are as good as those achieved from the CEV model.

Kroner, Kneafsey and Claessens (1995), using daily data, derive long-term horizon (225 calendar days ahead) forecasts of various commodity price volatilities. The authors use seven models to forecast volatility (ISD (Implied Standard Deviation), ISDAT (ISD from the at-the-money option), ISDAVG (weighted average of ISDs), HV (historical volatility), GARCH, COMB (which combines the ISD and GARCH forecasts) and GR (Granger and Ramanathan regression weighted combined forecast)). The results are compared by means of the MSE and ME. The two combined methods, namely GR and COMB, are found to give the most accurate predictions and outperform other forecasts.

Figlewski (1997) collects results from his earlier research papers regarding volatility forecasting for option pricing applications and investigates long-term horizon volatility forecasting using daily and monthly historical data. The most relevant for our thesis is where he investigates the long-term (1, 3, 6, 12, 24 months) forecasting accuracy of Historical Volatility (HV) and GARCH models by means of the RMSE. He finds the GARCH (1,1) model to be the best at forecasting volatility for the S&P 500 index over all time horizons, but in all other markets (3M US T-Bill, 20Y T-Bond, Deutschemark Exchange Rate (DM per \$)) its performance is very poor.

Green and Figlewski (1999), forecast volatility of the S&P 500 stock index, 3-month LIBOR (short-term interest rate), 10-year T-Bond yield (long-term interest rate) and DM exchange rate, by means of HV and Exponential Smoothing (ES). While using daily data, the forecast horizon varies from 1 to 12 month. Comparing RMSE, the authors find that ES is a better tool than HV for the short term S&P 500 (1–3 months) and the LIBOR, while bond yield, exchange rate and long-term S&P 500 (12 months) volatility are forecasted more accurately by means of the HV. They do importantly note that when the forecasted horizon is expanded the estimation period should also be expanded, in order to give reasonable long-term forecasts. In the second part of their studies they lengthen the forecast horizon to 2 and 5 years, and use monthly instead of daily data. In this case the HV outperforms ES for all four underlying assets.

Ederington and Guan (2002) wrote a paper in which they investigate whether the implied volatility is an informationally efficient and effective predictor of future volatility. The interesting aspect for us is in regards to the accuracy of the applied models (Implied Volatility, GARCH (1,1) and HV). In this study they use daily data for options on S&P 500 futures and forecast volatility of overlapping option maturity for 7-90, 91-180, 181-365 and 7-365 days ahead. The authors show that GARCH (1,1) outperforms the HV but at the same time it performs worse than the Implied Volatility. Additionally their findings confirm, as per the properties of the model, that the volatility forecasted by GARCH tends to converge to a constant at longer horizons.

Li (2002), uses Implied Volatility and the Autoregressive Fractionally Integrated Moving Average (ARFIMA) model to forecast volatility of forward options on 3 different currencies, for 1, 2, 3 and 6 months. The five-minute and daily returns on the German deutschemark, the Japanese yen, and the British pound vs. the US dollar are used. The Mean Absolute Error (MAE) reveals that the Implied Volatility provides more an accurate prediction at shorter time horizons while ARFIMA is more suitable for longer forecasts.

Since we compare four ARCH/GARCH models there is an additional aspect in Poon and Granger (2003), which we are interested in, namely, the analysis of 17 studies, which compare alternative versions of ARCH and GARCH models and their ability to forecast. The authors remark that, most of the time, papers use different data frequencies and do not explicitly suggest a clear rank of the models, thus it is difficult to point to an ultimate “winner”. They are however able to come to a main conclusion that GARCH models tend to outperform the more restrictive ARCH model.

Cao and Tsay (1992) use stock market data at daily and monthly frequencies to test which of the following models: Threshold Autoregressive (TAR), Autoregressive–Moving-Average model ARMA (1,1), ARCH (1,1) and EGARCH (1,0), give the most accurate volatility forecasts. They compare the forecasting ability over multiple horizons (from 1 to 30 months) by means of the MSE and MAE. The most relevant

finding is that the EGARCH (1,0) model gives the best volatility forecasts over long-time horizons, which they suggest is due to the leverage effect¹.

The use of different distribution assumptions in existing empirical works is rather poor, yet we review two papers which investigate this issue. Liu and Morley (2009), research whether forecasts based on the GARCH family models can outperform simple historical averaging models. In their paper, they use nine different models and apply daily and weekly sample frequencies. The authors test the forecasts sensitivity by applying three different error distributions (Normal, Student-t and Generalized Error Distribution). They reveal that the volatility forecasting is indeed sensitive to the assumption made about the use of different error distribution statistics. Moreover, they show that whilst not all of the models give better outcomes than the simple historical averaging, the use of non-normal distributions in the application of GARCH models, fits the in-sample data better and gives a better one-period ahead volatility forecast than the more commonly used normal distribution.

In the master thesis by Wei (2012), the author predicts the conditional variance of the rate of return in five different markets, including three used in this thesis, in terms of daily data. Three different forecasting models are applied, namely: GARCH (1,1), EGARCH (1,1) and GJR-GARCH (1,1) (2,1) (1,2) (2,2). By means of these models, the author studies forecasting performance under two different error distributions (Normal and Student t-distribution) and reports their accuracy with the use of the RMSE over a forecasting horizon of 1 year. The results exhibit that the normal distribution improves the models and the higher order GJR-GARCH outperforms other models in all cases except for the Hang Seng Index, where the GARCH (1,1) with a normal error distribution has the smallest RMSE.

¹ The leverage effect is generally a negative correlation between the return of an asset and its changes of volatility.

3. Data Analysis

3.1 Indices

The Financial Times Stock Exchange (FTSE) 100 index is a market capitalization-weighted index of United Kingdom-listed blue chip² companies and is the most widely used UK stock market indicator. The index was launched on the 3rd of January 1984 with a base date of the 30th December 1983 and a base value of 1000. The FTSE 100 measures the performance of the 100 largest companies, which are traded on the London Stock Exchange, and represents 81% of the entire market capitalization. The index is part of the FTSE UK Series and was designed for use in the creation of index tracking funds, derivatives and also as a performance benchmark. The top 5 constituents are HSBC Holdings, Royal Dutch Shell, BP Oil & Gas, GlaxoSmithKline Pharmaceuticals & Biotechnology and British American Tobacco .



Figure 1: FTSE 100 index 1988-2014.

The Standard and Poor (S&P) 500 (also referred to as GSPC) is a capitalization-weighted index composed of the 500 leading companies listed on the New York Stock

² Blue Chip: large companies which are characterized by high capitalization and liquidity. These companies are known for their high-quality, reliability and ability to perform profitably even in the face of adverse economic conditions, thus can be bought in order to provide steady portfolio growth.

Exchange and NASDAQ. The index was established on the 4th of March 1957 and was the first American based market capitalization-weighted stock index. Nowadays, the S&P 500 captures about 80% coverage of available market capitalization and is the basis of numerous listed and over-the-counter investment instruments. The S&P 500 was intended to measure performance of the domestic economy through changes in the aggregate market value of 500 stocks which represent a diverse selection of major industries. The top 5 constituents are Apple Inc., Exxon Mobil Corp., Microsoft Corp., Johnson & Johnson and Berkshire Hathaway B. The three largest sectors of the index are information technology (IT), financials and health care of which together they represent about 50% of all of the sectors contained in the index .




Figure 2: S&P 500 index 1988-2014.

The Hang Seng Index (HSI) is a free-float capitalization-weighted index and is the most widely quoted gauge for the Stock Exchange of Hong Kong composing of the 50 largest and most liquid stocks. The index was established with a base level of 100 as of the 31st of July 1964, however it was not launched until the 24th of November 1969. The HSI consists of companies from various types of industries, but the financial sector occupies almost half of the industry weights in the index (47,31%). In order to better reflect the price movements of the main sectors of the market, the

index constituents stocks that are grouped into four sub-indices, namely: Finance, Utilities, Properties & Commerce and Industry. The top 5 companies listed on the HSI are HSBC Holdings, Tencent³, China Mobile, China Construction Bank and AIA, of which three of these belong to the financial sector.



Figure 3: HSI index 1988-2014.

3.2 Descriptive statistics

Our historical data has been collected from the Yahoo Finance website and covers the daily period from January 4th 1988 until December 31st 2014 (over 6800 observations). We forecast volatility for different out of sample time horizons using ARCH/GARCH models, but for now our descriptive statistics help us to begin to describe the historical data using the realized daily returns and monthly volatility.

We have firstly computed the daily returns by means of the adjusted close prices of the three indices and we then go on to describe the data using statistics and graphs. We continue by calculating and analyzing the realized monthly volatility.

³ Tencent: a Chinese investment holding company, which provides a wide range of internet services, mass media, entertainment and mobile phone value-added services.

Statistic	FTSE 100	S&P 500	HSI
Observations	7035	6805	6717
Mean	0.0002496	0.0003701	0.0004674
Min	-0.0884834	-0.0903498	-0.2174532
Max	0.0983867	0.1158004	0.1882374
Variance	0.0001174	0.0001272	0.0002667
Standard Dev.	0.0108363	0.0112787	0.0163310
Skewness	0.0099	-0.1131	-0.1240
Kurtosis	9.601	11.97	17.58

Table 1: Summary statistics of returns (FTSE, S&P500, HSI).

Since we are using three different indices, from three different countries, the number of daily observations varies due to a diverse number of trading days. We see that during the period under observation there is a larger amount of trading days in the UK and fewest in the HSI data. We estimated the returns using the following equation:

$$\mathbf{r}_t = \frac{\mathbf{p}_t - \mathbf{p}_{t-1}}{\mathbf{p}_{t-1}} \quad (1)$$

Assuming that (y_1, y_2, \dots, y_T) is a random sample of Y and T denotes the number of observations, we can then calculate the mean as:

$$\hat{\mu}_y = \frac{1}{T} \sum_{t=1}^T y_t \quad (2)$$

The mean daily return for the HSI (0.047%) is twice as large as that of the FTSE 100 (0.025%). We observe the biggest range for the Hang Seng index, which extends from -0.2175 to 0.1882 and the smallest for FTSE 100 index (from -0.0885 to 0.0984). The standard deviations for the “western” indices are quite similar, 0.0108 and 0.0113, respectively. In the case of the HSI, the standard deviation is clearly larger (0.0163). We are specifically interested in the skewness and kurtosis since it is a measure of the shape of data distribution. Skewness measures the symmetry of a distribution and when the coefficient of skewness equals 0 (no skewness), we can say that the distribution is symmetric. If skewness < 0 , the median is often greater than the mean

and the distribution is negatively skewed (distribution skewed left). The opposite holds when the skewness > 0 (distribution skewed to the right). Kurtosis, on the other hand measures the peakedness of a distribution, meaning how extreme the observations in a given data set are. The kurtosis coefficient of a normal distribution is equal to 3 (Black (2009)). We distinguish between three types, namely leptokurtic (high and thin), platykurtic (flat and spread) and mesokurtic (the most "normal" shape). By means of the same assumptions, which were used while calculating the mean return, we compute the skewness as:

$$\hat{S}(\mathbf{y}) = \frac{1}{(T-1)\hat{\sigma}_y^3} \sum_{t=1}^T (y_t - \hat{\mu}_y)^3 \quad (3)$$

and the kurtosis as:

$$\hat{K}(\mathbf{y}) = \frac{1}{(T-1)\hat{\sigma}_y^4} \sum_{t=1}^T (y_t - \hat{\mu}_y)^4 \quad (4)$$

From the marginally negative skewness of the S&P 500 and HSI and close to zero for FTSE 100, along with large kurtosis (>3) for all, we can deduce that the data is all similarly distributed, which is to be expected since we are only examining indices. The obtained values for skewness and kurtosis indicate that the data of returns may display the leptokurtic property (high and thin) and we approach this issue by analyzing the distribution further in the next section.

3.3 Distribution

Since the distribution is of special interest in this thesis we investigate this issue through various means. Firstly, we study the histograms.




Figure 4: Histogram, FTSE 100.




Figure 5: Histogram, S&P 500.




Figure 6: Histogram, HSI.

We see in Figures 4, 5 and 6 that the leptokurtic property exists since compared to a normal distribution (Blue line), its central peak is higher and sharper, and its tails are longer and fatter. This observation is in line with the high values of kurtosis, which we obtained for each index in the previous subsection. This is consistent also with findings which states that “researchers have long noticed that stock returns have heavy tailed probability distributions” and this is “common with financial markets data” (Ruppert (2011)). We use also the Jarque-Bera (JB) test of normality to investigate the distribution of the data. This test incorporates the squared value of the Skewness (S) as well as the Kurtosis (K) when it differs from its normal distribution value of 3, and is appropriate to use when there is a large number of data observations (n):

$$JB = \frac{n}{6} \left(S^2 + \frac{1}{4} (K - 3)^2 \right) \quad (5)$$


The JB is tests the null hypothesis, which states that the data is normally distributed. The closer the Jarque-Bera statistic is to 0, the better the assumption regarding normality of distribution (Gujarati (2014)).

Jarque-Bera Normality Test			
	FTSE100	S&P500	HSI
Statistic	12771.3206	22834.2915	59522.4236
p-value	2.2e-16	2.2e-16	2.2e-16


Table 2: Jarque-Bera test statistics. If the p-values are smaller than the critical value (5%), we can reject the null hypothesis, which states that the examined indices are normally distributed. Else, fail to reject.

Such small p-values (<0.05) related to the JB statistic suggests that we can reject the null hypothesis, meaning, the data is not normally distributed. We have also included the quantile plots (Figure 7). The points should fall approximately on a straight line if the data is normally distributed. We do not observe such a pattern in any of the plots presented below. This kind of point’s location, obtained for all three indices, suggests that the data is heavily tailed compared to that of the normal distribution.

Normal Q-Q Plot



Normal Q-Q Plot



Normal Q-Q Plot




Figure 7: Normal Q-Q Plots for FTSE 100, S&P 500 and HSI, respectively. In the case where the data follows the assumed (normal) distribution, the points should fall approximately on a straight line.

3.4 Ljung-Box Q-Test

The autocorrelation function is applied to collect information about the properties of a time series (Koop (2006)). The best way to test for quantification of the autocorrelation is using the Ljung-Box Q-Test, which is able to check whether the data is random and independent. In the case where the observations are not independent, we can see a correlation between adjacent observations (autocorrelation). The Ljung-Box Q-test examines the null hypothesis, which states that the data is independently distributed up to lag ℓ (no autocorrelation), versus the alternative hypothesis where the data is not random and independent (autocorrelation is present). The Ljung-Box test statistic is as follow:

$$Q(m) = T(T + 2) \sum_{\ell=1}^m \frac{\hat{\rho}_{\ell}^2}{T - \ell} \quad (6)$$

Where,

$\hat{\rho}_{\ell}$ - estimated autocorrelation of the series at lag ℓ ,

m - denotes degrees of freedom (the largest lag),

T - observed data point.

Statistic	Index		
	FTSE100	S&P500	HSI
(Q) Statistic	0.713	22.54	0.1075
Prob > Chi2 (1)	2.2e-16	2.2e-16	2.2e-16

Table 3: Ljung-Box test. If the Q -value is larger than the critical value (5%), we can reject the null hypothesis that states that there is no autocorrelation in the returns. Else, fail to reject.

We can reject the null hypothesis when Q is larger than $\chi_{1-\alpha-h}^2$ (the chi-square distribution value with h degrees of freedom and significance level α). Therefore, all three indices exhibit autocorrelation (Box (1978)).

3.5 Volatility

We forecast volatility for 1, 3, 6, 9 and 12 months thus, for the sake of simplicity, we adjust the computed daily returns by assuming that there are 21 trading days in a month.

In order to calculate the realized (actual) volatility for month i , we select the returns for a given month r_t , where $t = \{1, 2, \dots, N\}$, and N represents the number of days in a month. To calculate the daily-realized volatility, we use the following formula, which squares the difference between the return and the mean return before dividing it by the number of observation minus one and then applying the square-root function:

$$\sigma_i^{daily} = \sqrt{\frac{\sum_{t=1}^T (r_t - E[r_i])^2}{N - 1}} \quad (7)$$

Where:

$E[r_i]$ - mean daily return for i^{th} month,

σ_i - realized volatility for i^{th} month.

In practice it is more common to use a simpler formula to calculate the realized volatility:

$$\sigma_i^{daily} = \sqrt{\frac{\sum_{t=1}^T r_t^2}{N}} \quad (8)$$

The main reason for this is that $E[r_i]$ is virtually zero. Moreover, the difference between dividing by N and $N - 1$ is insignificant when working with a large number of observations.

Both formulas presented above give us the daily volatility. In order to adjust it to our needs for forecasting horizons we re-compute equation (8) to the monthly volatility by means of the square root rule and based on our assumption of trading days outlined

above. Moreover, we computed the summary statistics of realized monthly volatility of all three indices.

Index	Min	1st Quantile	Median	Mean	3rd Quantile	Max
FTSE 100	5.63	10.20	12.70	15.20	17.90	75.90
S&P 500	4.91	10.20	12.90	15.50	18.20	82.10
Hang Seng	6.99	14.30	18.60	22.30	26.40	110.00

Table 4: Realized monthly volatility statistics.

We observe that the volatility statistics for the FTSE 100 and S&P 500 are very similar, whereas the HSI is similar only in the minimums. A larger difference can be seen in the mean. It takes a value of around 15% for the FTSE 100 and S&P 500, while the mean of the HSI is around 42% larger (22%). Another noticeable difference can be seen in the maximums where the HSI is extreme compared to that of the FTSE 100 and S&P 500.

Regressing the volatility on the lagged volatility for one period results in a $R^2 = 0.4993, 0.5220$ and 0.2530 for the three indices respectively. This indicates that the variation in volatility is not greatly explained by the previous period's volatility for the HSI index whereas for the FTSE and S&P the variation is around 50%. For all of the indices we increase the lag by one period at a time until the last periods coefficient are no longer statistically significant ($p > 0.05$). It become not significant for the variable lagged twice for the FTSE, three times for the S&P 500 and four time for the HSI index.

Figures 8, 9 and 10, present the realized monthly historical volatility for the three indices. There are no obvious common patterns that can be observed. Yet, there are some similarities in the volatility of the FTSE and the S&P indices, which are almost identical over the whole period (this is in line with the summary statistics from Table 4. For these two indices, the most sudden changes are observed at the turn of 2000-2002 (dot-com bubble crash) and during the most recent financial crises (2007/2008). The HSI peaks in 1987, 1997 and 2007/2008 and are all caused by financial crises. An interesting observation is that the most recent financial crisis seems to have resulted in

two after effects for the FTSE and S&P where as the volatility of the HSI has only dramatically increased once since.




Figure 8: Realized monthly historical volatility of the FTSE 100 index.




Figure 9: Realized monthly historical volatility of the S&P 500 index.




Figure 10: Realized monthly historical volatility of the Hang Seng Index.

4. Methodology

The methodology section presents the naïve model and the forecasting model that is used by the ARCH/GARCH family models along with the individually described error distributions before presenting the forecasting evaluation and predictive power statistics. The ARCH/GARCH models are used in the forecasting of volatility and we have introduced the model with the largest number of restrictions first. We have then expanded our model selection with the motivation of including models that remove some restrictions and limitations with the hope of achieving an insight into whether the more complicated models offer better forecasting predictions over our selected horizons.

4.1 Naive Forecasting Model

This straightforward and simple model forecasts based on the previous known observations. It treats the last period's volatility as this period's volatility (Giusti, Ritter and Vichi (2012)).

$$\hat{\sigma}_{t+1} = \sigma_t \quad (9)$$

The biggest weakness of this model is that the volatility is too persistent and does not perform well with long-run predictions (Zakamulin (2014)). We include the naive model as a benchmark for which to measure the more technical models against.

4.2 Rolling window

A rolling window forecast generates a series of forecasted observations for a given number of steps ahead. By using this method we define a fixed length of the in-sample period and a specific step size, meaning that the first and last date increase by the assigned value of observations/steps at each time. We use a 5-year rolling window and for simplicity assume that there is 21 trading days in a month (regardless of the country), which gives us 1260 observations. There are over 6800 daily observations for each index, so for example the first rolling window of the FTSE 100 contains data from day 1 to 1260. Then, by means of the model we obtain the forecast for the next day (observation number 1261) and thus, the second sample covers the observations from day 2 until day 1261. Based on the second sample we forecast observation number 1262 and so on. The last sample, for instance for the FTSE 100 index, will

contain observations from 5775 to 7035. We repeat this operation for all three indices whilst varying the step size based on the desired forecasting horizon.

4.3 ARCH

The Autoregressive Conditional Heteroskedasticity (ARCH) model was developed by Engle (1982). The main idea behind this model is that the mean corrected asset return is serially uncorrelated and dependent, meaning that ARCH is based on a linear regression where the variable, in time series form, varies around the mean (μ) randomly. Moreover, the model can be defined by the quadratic function of its lagged values. We assume that the return on assets can be computed as:

$$r_t = \mu + \sigma_t \epsilon_t \quad (10)$$

Where ϵ_t is a sequence of $N(0,1)$ independent and identically distributed (i.i.d.) random variables. We can then describe the residual return ($r_t - \mu$) at time t , as:

$$Z_t = \sigma_t \epsilon_t \quad (11)$$

Therefore, the ARCH model, which presents conditional variance as σ_t^2 , can be expressed as follows:

$$\sigma_t^2 = \alpha_0 + \sum_{k=1}^p \alpha_k Z_{t-k}^2 \quad (12)$$

Where $\alpha_0 > 0$ and $\alpha_k \geq 0$.

The lag length ($t - k$) of the ARCH model to the in-sample data is chosen in this thesis by the AIC, SBIC, HQIC in section 5. Lower order ARCH models may not be the most accurate in forecasting volatility due to the number of restrictions. We still include the ARCH model as it provides the basis of the parameters for the GARCH models that we wish to study further due to their handling of the previously stated issues. The main weakness is that the squared shocks model the conditional variance, and this leads to the ARCH model not being able to distinguish between positive and

negative effects, otherwise known as asymmetric effects. Another drawback is that recent evidence suggests that there exists substantial persistence in the variance of such models (Lamoureux and Lastrapes (1990)). The ARCH model also carries mean reverting properties which suggests that the process reverts to its mean slowly therefore remaining highly persistent for long periods before returning to the equilibrium, but due to its restrictive intervals it causes the volatility to often be over predicted owing to a slow response to shocks.

The ARCH model's strengths lie in the fact that it can model volatility clustering through the lagged squared returns of Z_{t-k}^2 which is described as "large changes tend to be followed by large changes, of either sign, and small changes tend to be followed by small changes" (Mandelbrot (1997)).

4.4 GARCH

Bollerslev (1987) developed the Generalized Autoregressive Conditional Heteroskedasticity (GARCH) model, which is an extension of the ARCH model where the order of ARCH (t) ($\alpha_t Z_{t-k}^2$), as established previously, is built upon to include the GARCH term $\beta_j \sigma_{t-j}^2$. The model is as follows:

$$\sigma_t^2 = \alpha_0 + \sum_{k=1}^p \alpha_k Z_{t-k}^2 + \sum_{j=1}^q \beta_j \sigma_{t-j}^2 \quad (13)$$

The GARCH model is said to be "much more flexible and capable of matching a wide variety of patterns of financial volatility" (Koop (2006)). The GARCH again deals with volatility clustering and is characterized by a symmetric response of current volatility to positive and negative returns. It also has the addition of σ_{t-j}^2 , the lagged conditional variance, which helps to increase the accuracy of the ARCH model by avoiding excessive lags of the squared returns. Both ARCH's and GARCH's biggest weakness lies in the fact that the models assume that the shocks (both positive and negative) have the same effects on volatility which is generally accepted not to be true.

Unlike the conditional variance, which is changing, the unconditional variance of Z_t is constant if the assumption of mean reversion is maintained ($\alpha + \beta < 1$) and can be expressed as:

$$\bar{\sigma}^2 = \frac{\alpha_0}{1 - \alpha_1 - \beta_1} \quad (14)$$

Otherwise, when $\alpha + \beta \geq 1$, we say that the unconditional variance of Z_t is not defined, meaning we face a “non-stationarity” in variance. Moreover, in the case where $\alpha + \beta = 1$, we have a “unit root” in variance (Brooks (2014)).

4.5 EGARCH

In order to deal with the weaknesses of the GARCH model, which assumes that positive and negative shocks affect volatility equally, we use the Exponential GARCH (EGARCH), which was introduced and modeled by Nelson (1991). This is an asymmetric form of the GARCH model. The EGARCH is able to respond asymmetrically to positive and negative effects as well as the volatility persistence and mean reversion. The model is formally written as, an EGARCH (p, q):

$$\log(\sigma_t^2) = \alpha_0 + \sum_{k=1}^p [\alpha_k Z_{t-k} + \gamma_k (|Z_{t-k}| - E(|Z_{t-k}|))] + \sum_{j=1}^q \beta_j \log(\sigma_{t-j}^2) \quad (15)$$

The section in the squared brackets illustrates the models ability to deal with the asymmetric effects of positive and negative returns. The function has a mean of zero and is uncorrelated. It can be re-written as:

$$(\alpha_1 + \gamma_1)Z_t I(Z_t > 0) + (\alpha_1 - \gamma_1)Z_t I(Z_t < 0) - \gamma_1 E(|Z_{t-1}|) \quad (16)$$

Where $(\alpha_1 + \gamma_1)$ takes the positive shocks into consideration and $(\alpha_1 - \gamma_1)$ deals with the negative shocks. This gives rise to the leverage effect which states that negative shocks have a larger impact than positive shocks since $\alpha_1 < 0, 0 \leq \gamma_1 \leq 1$ and $\beta_1 + \gamma_1 < 1$. The shocks impact the logarithm of the conditional variance

$\log(\sigma_t^2)$. Since $\log(\sigma_t^2)$ is the log of the variance, the conditional variance is guaranteed to be positive, therefore not artificially imposing non-negativity on the model parameters and the model allows for asymmetries (Brooks (2014)). A possible drawback of the EGARCH model is that it does not fit the normal distribution and under normality it tends to overestimate the impact of outliers. This is an interesting aspect and the findings between the normal, student-t and generalized error distributions will be of special interest for this model.

4.6 GJR-GARCH

We use the GJR-GARCH as another asymmetric model since we would still like to take into consideration the different impacts that positive and negative innovations have and since the EGARCH model is highly dependent on the data not being normally distributed we will closely compare these two models.

The model was presented by Glosten, Jagannathan and Runkle (1993) and the difference between the EGARCH and the GJR-GARCH model is a slight change in the conditional variance function which is no longer a logarithm and the positive and negative shocks are seen through γ_k and α_k respectively.

$$\sigma_t^2 = \alpha_0 + \sum_{k=1}^p [\alpha_k Z_{t-k}^2 (1 - I[Z_{t-k} > 0]) + \gamma_k Z_{t-k}^2 I[Z_{t-k} > 0]] + \sum_{j=1}^q \beta_j \log(\sigma_{t-j}^2) \quad (17)$$

4.7 Distribution statistics

Normal distribution (Gaussian)

The Normal or Gaussian distribution is an absolute continuous distribution where the mean is equal to zero and the variance is equal to one ($N(0,1)$). For the normal distribution we observe zero skewness and zero excess kurtosis. It is the most

commonly used distribution within financial analysis since it is symmetric and requires only the mean and variance.

$$f(x, \mu, \sigma) = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{(x-\mu)^2}{2\sigma^2}} \quad (18)$$

Student t-distribution

As we have seen in (Section 3.3) our in-sample data does not fit a standard normal distribution. We therefore consider the student t-distribution, first derived by Helmer (1876) but formally written by Gosset (1908) under the pseudonym ‘‘Student’’. This distribution can work with smaller sample sizes, which are considered through $\nu = n - 1$ degrees of freedom.

$$f(t) = \frac{\Gamma\left(\frac{\nu+1}{2}\right)}{\sqrt{\nu\pi}\Gamma\left(\frac{\nu}{2}\right)} \left(1 + \frac{t^2}{\nu}\right)^{-\frac{\nu+1}{2}} \quad (19)$$

The student t-distribution has, similarly to the normal distribution, zero skewness (it is symmetric) but excess kurtosis equal to $\frac{6}{\nu-4}$ for $\nu > 4$ (leptokurtic property).

Generalized error distribution

The generalized error distribution (GED) was introduced by Subbotin (1923) (Agro (1995)). This distribution is used when the errors of the data are not necessarily normally distributed. Vasudeva and Kumari (2013), discuss the first class of GED (GED-I) from the parametric family of symmetric distributions, which we have used, since it can cope with the heavier tailed symmetric distributions of our data and can become a useful way to parameterize (Nadarajah (2005)).

$$dF(x|\mu, \sigma, \kappa) = \frac{e^{-\frac{1}{2}\left|\frac{x-\mu}{\sigma}\right|^{\kappa}}}{2^{k+1}\sigma\Gamma(k+1)} dx \quad (20)$$

Where x is the domain of the p.d.f and the distribution is defined by μ, σ, k , which relates to the mode, dispersion and skewness of the data respectively. The generalized error distribution has symmetric properties but allows for variation in the kurtosis which are considered to be close to the normal distribution (Giller (2005)).

4.8 Forecast Evaluation

There exist various methods that allow us to compare how well the models can predict the dependent variable. We use the mean squared error (MSE) to analyze the quality of the ARCH and GARCH family models so that our findings can be directly compare to existing research.

The MSE is often used in comparing the difference between things that vary from an accepted standard. It is highly dependent on the scale of the dependent variable with the smaller the MSE of the model suggesting the better the models ability. We first predict the volatility, and then compute the realized volatility, at the end we compare our results by means of the following formula:

$$MSE = \frac{1}{n} \sum_{i=1}^n (\sigma_i - \hat{\sigma}_i)^2 \tag{21}$$

$$MSE = \frac{1}{n} \sum_{i=1}^n e_i^2 \tag{22}$$

Where:

σ_i - actual value of the i^{th} observation,

$\hat{\sigma}_i$ - forecasted value of the i^{th} observation,

e_i - forecasting error of the i^{th} observation,

n - number of observations.

The main weakness of this method is that the prediction error variance varies across time. This problem is also common among other similar measurement techniques (Cipan (2004)). The forecast accuracy is considered to be increasing as the MSE

approaches zero (0 means that there is no difference between what was predicted and what actually happened). We compare the findings for each index over the three chosen distribution statistics.

4.9 Predictive Power

Together with a method that helps us to evaluate the goodness of forecast, we will also compute the predictive power of each model over the different forecasted time horizons. To do this, we use a method proposed by Blair, Poon and Taylor (2010). The predictive power is a comparison between the prediction errors and the variation in σ_i , which are both squared. The closer the predictive power ratio is to one hundred the better the accuracy of the forecasted volatility. Any deviation suggests that there exists greater variation of the errors of the forecasted volatility than those of the realized volatility. The predictive power (P) can be both positive and negative (Poon (2005)):

$$P = \left(1 - \frac{\sum_{i=1}^n (\sigma_i - \hat{\sigma}_i)^2}{\sum_{i=1}^n (\sigma_i - \bar{\sigma})^2} \right) \times 100 \quad (23)$$

Where:

n - number of out-of-sample observations,

σ_i - realized volatility,

$\hat{\sigma}_i$ - forecasted volatility,

$\bar{\sigma}$ - mean value of volatility.

The mean value of volatility in the out-of-sample period can be computed as:

$$\bar{\sigma} = \frac{1}{n} \sum_{i=1}^n \sigma_i \quad (24)$$

5. Results

We first provide an initial look at the of lag length selection through observation of the auto and partial-auto correlation functions based on the daily returns where we distinguish a rough estimation of the ARMA (p,q). We then acquire the (p,q) more accurately using the Akaike's information criterion (AIC), Schwarz's Bayesian information criterion (BIC), and the Hannan and Quinn information criterion (HQIC). We then look at the significance of the parameters of GARCH, E-GARCH and GJR-GARCH (1,1) models and look each of the parameters individually. For the higher order models ((1,2) (2,1) (2,2)) we compare AIC statistics. We present the MSE and predictive power values for 1, 3 6, 9 and 12 months forecasting periods. We also compare the models on the time horizon with the best forecast evaluation with the higher order models selected from the lowest AIC statistic, in order to gain an insight into the advantages or disadvantages of using higher order models.

5.1 FTSE100




Figure 11: Auto and Partial-Auto correlation functions for the FTSE 100 index.

The ACF and PACF exhibit significant autocorrelations in the residuals and both have their first significant autocorrelation visible already at lag 2. The ACF is significant at lags 2-6 and 8, after which it gradually fades away. The PACF is significant in lags 2-6 and also has the tendency of fading away. This would suggest that an ARMA (p,q) is present and a higher order model could be necessary. These results are also confirmed in the test statistics of lag length selection models presented below.

	ARMA(1,0)			ARMA(0,1)			ARMA(1,1)		
	Normal	Student-t	GED	Normal	Student-t	GED	Normal	Student-t	GED
AIC	-6,4765	-6,562	-3,1759	-6.4827	-6.5620	-3.1761	-6,4803	-6,5672	-3,1685
BIC	-6,4663	-6,5416	-3,1555	-6.4663	-6.5416	-3.1557	-6,46	-6,5428	-3,144
HQIC	-6,4765	-6,5544	-3,1682	-6.4765	-6.5544	-3.1684	-6,4727	-6,558	-3,1593
	ARMA(2,1)			ARMA(1,2)			ARMA(2,2)		
	Normal	Student-t	GED	Normal	Student-t	GED	Normal	Student-t	GED
AIC	-3,7095	-6,7988	-3,7308	-7,1066	-6,3598	0,33505	-6,2762	-6,7869	-3,2224
BIC	-3,6851	-6,7702	-3,7022	-7,0821	-6,3313	0,3636	-6,2476	-6,7543	-3,1897
HQIC	-3,7003	-6,7881	-3,7200	-7,0974	-6,3491	0,34578	-6,2654	-6,7747	-3,2101

Table 5: FTSE 100 Information Criteria for model selection of the Auto Regressive Moving Average (ARMA).

The specifications for the conditional variance, given in Table 5, contain parameters for the auto regressive and moving average, denoted by p and q . We have included six combinations of ARMA lengths. For the normally distributed model, p and q are the lowest at ARMA (1,2) and for the t-distribution ARMA (2,1). The GED also shows an improvement when, p and q ARMA order is (2,1). The AIC, BIC and HQIC for the normal distribution become positive at (3,2), (2,3) and (3,3) and increase for the other two distributions. In order for our ARCH/GARCH models to remain reasonable and comparable and to forecast the same models on each index we keep the ARMA length at (1,1). Since models with large amounts of lags rarely give more accurate results than the same models with fewer lags, it is rational to assume, that this small modification regarding ARMA (p,q) will not result in any disturbance in our analysis. Using the lag length of (1,1) also keeps our research in-line and comparable with the work which were summarized in the literature review section of this thesis (Section 2).

Using ARMA (1,1) we compare the GARCH, EGARCH and GJR-GARCH models under the three error distributions and select the optimal order of models beginning at (1,1), whilst also extending them to higher order models if necessary.

FTSE 100	GARCH(1,1)			EGARCH(1,1)			GJR-GARCH(1,1)		
	Normal	Student-t	GED	Normal	Student-t	GED	Normal	Student-t	GED
mu	0,00048	0,000251	0,000456	0,000068	-0,000053	0,000187	0,00013	-0,000073	0,000219
s.e	0,000226	0,000152	0,00021	0,000084	0,000171	0,000194	0,000216	0,000197	0,000169
p-value	0,033449	0,097789	0,029637	0,41487	0,757603	0,33579	0,54601	0,711519	0,19472
omega	0,000002	0,000001	0,000002	-0,36703	-0,119368	-0,37034	0,000003	0,000001	0,000003
s.e	0,000002	0,000004	0,000003	0,007283	0,003255	0,00569	0	0,000002	0
p-value	0,354979	0,778516	0,529729	0	0	0	0	0,361531	0
alpha1	0,096327	0,10161	0,096325	-0,14235	-0,120732	-0,15779	0	0	0
s.e	0,030046	0,063243	0,043353	0,009031	0,003255	0,016323	0,004826	0,021445	0,006231
p-value	0,001346	0,108127	0,026291	0	0	0	1	0,999995	0,99998
beta1	0,879383	0,892257	0,883401	0,960921	0,987434	0,961371	0,876258	0,913898	0,871061
s.e	0,036675	0,061892	0,051184	0,000763	0,000028	0,000123	0,014238	0,024565	0,017622
p-value	0	0	0	0	0	0	0	0	0
shape		12,93365	1,322303		21,069995	1,397429		18,675424	1,409927
s.e		1,199575	0,054445		8,820335	0,070228		8,858992	0,082652
p-value		0	0		0,016904	0		0,035024	0
ar1	-0,94889	0,868502	-0,96733	-0,94844	0,530585	-0,96139	-0,95245	0,612553	-0,96724
s.e	0,010101	0,138921	0,006196	0,003954	0,044616	0,002677	0,008485	0,224405	0,005997
p-value	0	0	0	0	0	0	0	0,00634	0
gamma1				0,145409	0,075922	0,149203	0,186358	0,142047	0,201142
s.e				0,043415	0,012064	0,015416	0,027435	0,030648	0,032638
p-value				0,00081	0	0	0	0,000004	0
ma1	0,9684	-0,91144	0,984307	0,967237	-0,5849	0,978779	0,971924	-0,680342	0,984318
s.e	0,003588	0,118393	0,000643	0,00465	0,042505	0,000715	0,002585	0,208171	0,000639
p-value	0	0	0	0	0	0	0	0,001082	0

FTSE 100	GARCH(1,1)			EGARCH(1,1)			GJR-GARCH(1,1)		
	Normal	Student-t	GED	Normal	Student-t	GED	Normal	Student-t	GED
AIC	-6,6383	-6,4165	-6,6736	-6,6848	-6,4539	-6,7115	-6,6896	-6,4347	-6,7139
BIC	-6,6138	-6,3879	-6,6450	-6,6562	-6,4212	-6,6788	-6,6611	-6,4020	-6,6813
HQIC	-6,6291	-6,4057	-6,6629	-6,6741	-6,4416	6,6992	-6,6789	-6,4224	-6,7016

Table 6: GARCH (1,1) family models statistics and information criteria for the FTSE100 index. The values which are not statistically different from zero are highlighted in blue.

Our results show that based on the assumption of 5% significance the parameters of the GARCH models are all significantly different from zero apart from omega for all

distributions and mu and omega for the student t-distribution. The EGARCH is not significant in its mean for all three distributions. The GJR parameters are not significance in the mean and alpha. These findings suggest that higher order model would maybe be more appropriate. We have therefore looked at the AIC of the higher order models and our findings suggest that the GARCH (2,2), EGARCH (2,1) and GJR-GARCH (1,1) should be used, however to keep our results comparable to previous research and due to the minor differences in the AIC statistic we have chosen to forecast all of our models as (1,1). We do however present the results for the higher models over a 1 month forecasting horizon in order to prove that the higher order models selected based on the lower AIC statistic do not significantly enhance the results.

FTSE 100	GARCH			EGARCH			GJR-GARCH		
	Normal	Student-t	GED	Normal	Student-t	GED	Normal	Student-t	GED
AIC(1,1)	-6,6383	-6,6603	-6,6736	-6,6848	-6,7059	-6,7115	-6,6896	-6,7061	-6,7139
AIC(2,1)	-6,6383	-6,6613	-6,6738	-6,6876	-6,7081	-6,7126	-6,6866	-6,703	-6,7107
AIC(1,2)	-6,6367	-6,6587	-6,6720	-6,6832	-6,7042	-6,7098	-6,6881	-6,7045	-6,7123
AIC(2,2)	-6,6387	-6,6623	-6,6741	-6,6860	-6,7066	-6,7111	-6,6850	-6,7014	-6,7091

Table 7: Akaike's Information Criteria of higher order GARCH family models for the FTSE 100 index.

In Table 8 and Figure 12 below we present our findings and observe that the MSE gives the best results for a 1 month forecast. The 1 month forecast is better than the longer forecasted horizons and all of the ARCH/GARCH models give more accurate results when using the student t-distribution whereas the naïve model performs best with the normal and generalized distribution. For the 1 month forecast the benchmark naïve model is outperformed by all other models.

nDays	Distribution	MSE				
		Naive	ARCH	GARCH(1,1)	EGARCH(1,1)	GJR-GARCH(1,1)
21	Normal	0,000355	0,012210	0,0003214	0,0002826	0,0002741
	Student t	0,001148	0,000289	0,0001309	0,0001091	0,0001144
	GED	0,000355	0,002392	0,0003198	0,0002844	0,0002734
63	Normal	0,001148	0,028010	0,0009224	0,0008503	0,0007975
	Student t	0,000473	0,000851	0,0004323	0,0003219	0,0003373
	GED	0,001148	0,007381	0,0009239	0,0008789	0,0008092
126	Normal	0,002759	0,03401	0,002148	0,001808	0,001677
	Student t	0,001260	0,00195	0,000935	0,000801	0,000690
	GED	0,002759	0,01497	0,002097	0,001887	0,001694
189	Normal	0,003526	0,13810	0,002975	0,002754	0,002453
	Student t	0,003526	0,02750	0,002836	0,002862	0,002501
	GED	0,003526	0,02257	0,002837	0,002871	0,002511
252	Normal	0,006815	0,08425	0,008209	0,005658	0,005823
	Student t	0,006815	0,02884	0,007566	0,005617	0,005725
	GED	0,006815	0,03060	0,007848	0,005746	0,005820

Table 8: MSE forecasting results for the FTSE 100 index over multiple time horizons.




Figure 12: Clustered Column Chart of FTSE 100 MSE results.

The most accurate forecasting horizon for the predictive power is 1 month where all student t-distribution models outperform that of the normal and generalized distribution. The naïve model outdoes ARCH (1,1) and is only slightly worse than GARCH (1,1). The most accurate results are given by the EGARCH (1,1) model. Although the Predictive power does decrease as the time horizon increases, this is seen to only be marginal in the case of the models with the student t-distribution , up to and including the 6 month (126 day) forecasting horizon. These results seem to be disproportionately high and have been retested multiple times to check there accuracy.

nDays	Distribution	Predictive Power				
		Naive	ARCH	GARCH(1,1)	EGARCH(1,1)	GJR-GARCH(1,1)
21	Normal	44,4	-1811	49,68	55,75	57,09
	Student t	80,09	59,47	81,64	84,69	83,95
	GED	44,4	-274,5	49,93	55,48	57,19
63	Normal	28,55	-1643	42,61	47,09	50,38
	Student t	77,12	58,81	79,07	84,41	83,67
	GED	28,55	-359,2	42,52	45,31	49,65
126	Normal	3,021	-1095	24,49	36,46	41,06
	Student t	69,08	52,12	77,06	80,35	83,07
	GED	3,021	-426,2	26,29	33,67	40,47
189	Normal	6,044	-3580	20,73	26,62	34,64
	Student t	6,044	-632,7	24,44	23,73	33,35
	GED	6,044	-501,3	24,42	23,51	33,1
252	Normal	-33,1	-1545	-60,33	-10,50	-13,74
	Student t	-33,1	-463,3	-47,77	-9,701	-11,82
	GED	-33,1	-497,6	-53,27	-12,22	-13,66

Table 9: Predictive Power forecasting results for the FTSE 100 index over multiple time horizons.

When comparing the results against the higher order models selected based on the lowest AIC, we observe no improvement in the models MSE or predictive power where there was only a slight loss for GARCH (2,2).

nDays	Distribution	MSE				
		Naive	ARCH	GARCH(2,2)	EGARCH(2,1)	GJR-GARCH(1,1)
21	Normal	0,0003552	0,009101	0,0003214	0,0002817*	0,0002741
	Student t	0,0001418	0,0002888	0,0001315	0,0001091	0,0001144
	GED	0,0003552	0,002392	0,0003174*	0,0002833*	0,0002734
nDays	Distribution	Predictive Power				
		Naive	ARCH	GARCH(2,2)	EGARCH(2,1)	GJR-GARCH(1,1)
21	Normal	44,4	-1325	49,75*	55,9*	57,1
	Student t	80,09	59,47	81,55	84,69	83,95
	GED	44,4	-274,5	50,31*	55,65*	57,19

Table 10: MSE and Predictive Power forecasting results for higher order GARCH AND EGARCH models over a 21 day forecasting horizon. Values marked with an asterisk (*) show an improvement.

5.2 S&P 500




Figure 13: Auto and Partial-Auto correlation functions for the S&P 500 index.

The ACF cuts off after the second lag. This behavior indicates an MA (2) process. The PACF also cuts off after the second lag, indicating an AR (2) process but the correlogram shows the larger correlation in higher lags such as 5, 7, 10 and 12. All three information criteria statistics suggest that in the case of the normal and student t-distributions that ARMA (1,1) produces the lowest statistics. For the generalized distributions the best is ARMA (2,1). The values obtained in ARMA (2,2) for the normal and student t-distributions vary significantly and become positive or are not obtainable in R. We therefore select to continue our analysis with the ARMA (1,1) model.

	ARMA(1,0)			ARMA(0,1)			ARMA(1,1)		
	Normal	Student-t	GED	Normal	Student-t	GED	Normal	Student-t	GED
AIC	-6.3978	-6.5661	-2.6646	-6.3977	-6.5660	-2.6612	-6.4039	-6.5715	-2.7159
BIC	-6.3815	-6.5457	-2.6442	-6.3814	-6.5456	-2.6408	-6.3835	-6.5470	-2.6914
HQIC	-6.3917	-6.5584	-2.6569	-6.3916	-6.5584	-2.6535	-6.3962	-6.5623	-2.7067
	ARMA(2,1)			ARMA(1,2)			ARMA(2,2)		
	Normal	Student-t	GED	Normal	Student-t	GED	Normal	Student-t	GED
AIC		-6.5710	-2.7135	-2.9071	-6.5639	-2.7260		385.84	-2.7234
BIC		-6.5424	-2.6850	-2.8827	-6.5353	-2.6975		385.88	-2.6908
HQIC		-6.5603	-2.7028	-2.8979	-6.5532	-2.7153		385.86	-2.7111

Table 11: S&P500 Information Criteria for model selection of the Auto Regressive Moving Average (ARMA).

In Table 12 our results show that EGARCH (1,1) is significantly different from zero for all distributions and GARCH is again not significant in omega. GJR-GARCH is not statistically significant in alpha for all distributions and AR (1) and MA (1) with the normal distribution.

S&P 500	GARCH(1,1)			EGARCH(1,1)			GJR-GARCH(1,1)		
	Normal	Student-t	GED	Normal	Student-t	GED	Normal	Student-t	GED
mu	0,000681	0,000765	0,000981	0,000312	0,000727	0,000662	0,000471	0,000682	0,000662
s.e	0,000031	0,000043	0,000122	0,000106	0,000176	0,000033	0,000201	0,000032	0,000033
p-value	0	0	0	0,003201	3,50E-05	2,00E-06	0,018936	0	0,0000E+00
omega	0,000003	0,000003	0,000003	-0,49701	-0,46257	-0,51395	0,000003	0,000004	0,000004
s.e	0,000002	0,000002	0,000002	0,005284	0,007278	0,008199	0	0	0
p-value	0,15606	0,089678	0,15793	0	0	0	0	0	0
alpha1	0,14213	0,152062	0,143579	-0,22036	-0,24961	-0,23122	0	0	0
s.e	0,020458	0,028887	0,026431	0,01651	0,024397	0,026379	0,010072	0,010431	0,009414
p-value	0	0	0,031085	0	0	0	0,999999	0,99999	1,00E+00
beta1	0,82469	0,821367	0,822839	0,947161	0,95257	0,947264	0,838845	0,800636	0,808844
s.e	0,025721	0,028626	0,031085	0,000229	0,000476	0,001038	0,01455	0,019685	0,017864
p-value	0	0	0	0	0	0	0	0	0
shape		5,548586	1,272639		5,821013	1,331867		6,111765	1,361226
s.e		0,935857	0,070512		1,004069	0,069237		1,032129	0,074012
p-value		0	0		0	0		0	0,00E+00
ar1	0,961876	0,961661	0,918547	-0,87803	0,526459	0,750844	-0,84899	0,969699	0,971911
s.e	0,00425	0,005266	0,015172	0,031641	0,057167	0,06747	1,709544	0,003455	0,003817
p-value	0	0	0	0	0	0	0,619456		0
gamma1				0,152124	0,143476	0,157328	0,244052	0,292037	0,270568
s.e				0,01125	0,015698	0,010672	0,036007	0,048059	0,043926
p-value				0	0	0	0	0	0
ma1	-0,99349	-0,99135	-0,95319	0,867427	-0,55909	-0,77742	0,838975	-0,99295	-0,992798
s.e	0,000125	0,00012	0,007955	0,03299	0,054959	0,062389	1,764539	0,000103	0,000087
p-value	0	0	0	0	0	0	0,634456	0	0

S&P 500	GARCH(1,1)			EGARCH(1,1)			GJR-GARCH(1,1)		
	Normal	Student-t	GED	Normal	Student-t	GED	Normal	Student-t	GED
AIC	-6,6656	-6,7025	-6,708	-6,7112	-6,7496	-6,7515	-6,7027	-6,7469	-6,7500
BIC	-6,6411	-6,6740	-6,6794	-6,6827	-6,7170	-6,7188	-6,6741	-6,7143	-6,7174
HQIC	-6,6564	-6,6918	-6,6972	-6,7005	-6,7374	-6,7392	-6,6919	-6,7347	-6,7378

Table 12: GARCH (1,1) family models statistics and information criteria for the S&P 500 index. The values which are not statistically different from zero are highlighted in blue.

The AIC for the higher order models suggest an improvement could be found through higher order models. The AIC suggests (2,1) for the GARCH and EGARCH models and due to no improvement in the AIC for the GJR GARCH, (1,1) is selected despite the non-significant alpha, AR (1) and MA (1) in the normal distribution. The results are again calculated using the (1,1) on all models and directly compared with the results of the best AIC models.

S&P 500	GARCH			EGARCH			GJR-GARCH		
	Normal	Student-t	GED	Normal	Student-t	GED	Normal	Student-t	GED
AIC(1,1)	-6,6656	-6,7025	-6,708	-6,7112	-6,7496	-6,7515	-6,7027	-6,7469	-6,75
AIC(2,1)	-6,6756	-6,7096	-6,7153	-6,731	-6,7647	-6,7669	-6,7022	-6,7458	-6,7486
AIC(1,2)	-6,6647	-6,7013	-6,7068	-6,7114	-6,7495	-6,7511	-6,7026	-6,7462	-6,7493
AIC(2,2)	-6,6752	-6,5376	-6,7141	-6,7296	-6,5838	-6,7656	-6,7109	-6,5471	-6,7535

Table 13: Akaike's Information Criteria of higher order GARCH family models for the S&P 500 index.

nDays	Distribution	MSE				
		Naive	ARCH	GARCH(1,1)	EGARCH(1,1)	GJR-GARCH(1,1)
21	Normal	0,0004123	0,023860	0,0003609	0,0003643	0,0003248
	Student t	0,0004123	0,006222	0,0003610	0,0003462	0,0003251
	GED	0,0004123	0,002632	0,0003606	0,0775100	0,0003219
63	Normal	0,001524	0,04183	0,001291	0,001503	0,001250
	Student t	0,001524	0,05242	0,001295	0,001437	0,001219
	GED	0,001524	0,008115	0,001295	0,001502	0,001233
126	Normal	0,003708	0,03713	0,003348	0,003486	0,003251
	Student t	0,003708	0,1531	0,003597	0,003453	0,003223
	GED	0,003708	0,01637	0,003495	0,003586	0,003254
189	Normal	0,005624	0,07937	0,003961	0,00497	0,004226
	Student t	0,005624	0,4221	0,004125	0,005161	0,0045
	GED	0,005624	0,02466	0,004121	0,005483	0,004352
252	Normal	0,005843	0,09134	0,006248	0,006571	0,006153
	Student t	0,005843	0,1242	0,008113	0,006902	0,006073
	GED	0,005843	0,03276	0,007228	0,007228	0,006292

Table 14: MSE forecasting results for the S&P 500 index over multiple time horizons.

The MSE results suggest that the most accurate forecasting horizon is again 1 month using the student t-distribution. In terms of longer time horizons we see that the 1 month horizon is more accurate than the longer horizons.




Figure 14: Clustered Column Chart of S&P 500 MSE results.

For the predictive power the 21-day forecast is the most accurate. With the GARCH (1,1) we obtain very similar results for all three distributions. The student t-distribution improves the EGARCH model compared to the other distributions and the GED offers a small improvement in the GJR-GARCH model. We also noticed that the Naïve model is improved over a 1 year (252 day) forecasting period compared to the predictive power of both 6 and 9 months (126 and 189 days). The P value for the 9 month forecast using a normal distribution is higher than the shorter 6 month forecast when using the GARCH (1,1) and the GJR-GARCH(1,1).

nDays	Distribution	Predictive Power				
		Naive	ARCH	GARCH(1,1)	EGARCH(1,1)	GJR-GARCH(1,1)
21	Normal	47,67	-2929	54,2	53,77	58,77
	Student t	47,67	-689,7	54,19	56,06	58,75
	GED	47,67	-234	54,24	55,01	59,14
63	Normal	25,71	-1939	37,08	26,74	39,07
	Student t	25,71	-2455	36,89	29,94	40,6
	GED	25,71	-295,6	36,87	26,78	39,89
126	Normal	1,025	-891,1	10,63	6,95	13,23
	Student t	1,025	-3986	3,973	7,835	13,96
	GED	1,025	-336,9	6,713	4,282	13,13
189	Normal	-12,18	-1483	20,98	0,8708	15,71
	Student t	-12,18	-8318	17,72	-2,944	10,25
	GED	-12,18	-392	17,8	-9,36	13,19
252	Normal	1,532	-1439	-5,3	-10,74	-3,699
	Student t	1,532	-1993	-36,73	-16,31	-2,341
	GED	1,532	-452,1	-21,82	-24,69	-6,027

Table 15: Predictive Power forecasting results for the S&P 500 index over multiple time horizons.

Using the AIC selected higher order GARCH models and the predictive power statistic we observe that with GARCH (2,1) the results are improved but not dramatically. The student t-distribution is the most accurate for the EGARCH (1,1) and is found to be slightly improved when using the higher order model EGARCH (2,1). GJR-GARCH is the most accurate of all models and gives the best results regardless of distribution with the generalized distribution performing marginally better whilst still remaining at the lower order GJR-GARCH (1,1) model.

nDays	Distribution	MSE				
		Naive	ARCH	GARCH(2,1)	EGARCH(2,1)	GJR-GARCH(1,1)
21	Normal	0,0004123	0,009268	3,57E-04*	3,58E-04*	0,0003248
	Student t	0,0004123	6,22E-03	3,60E-04*	3,36E-04*	0,0003251
	GED	0,0004123	2,63E-03	3,57E-04*	3,47E-04*	0,0003219
nDays	Distribution	Predictive Power				
		Naive	ARCH	GARCH(2,1)	EGARCH(2,1)	GJR-GARCH(1,1)
21	Normal	47,67	-1164	54,67*	54,58*	58,77
	Student t	47,67	-689,7	54,35*	57,32*	58,75
	GED	47,67	-234	54,72*	55,97*	59,14

Table 16: MSE and Predictive Power forecasting results for higher order GARCH AND EGARCH models over a 21 day forecasting horizon. Values marked with an asterisk (*) show an improvement.

5.3 HSI




Figure 16: Auto and Partial-Auto correlation functions for the Hang Seng index.

The ACF is statistically significant between lags 3 and 5 and gradually fades away indicating an MA (q) process. The partial auto correlation function is also firstly significantly different at lag three and again fades away slowly. When examining the ARMA selection only the student t-distribution with p and q equal to (1,1) is statistically different from zero in all of its parameters, however the AIC suggests a (1,0) or (0,1) ARMA. The AIC for the normal distribution is lowest at ARMA (2,1) and ARMA (0,1) is suggested for the generalized distribution. Again we will continue our analysis with ARMA (1,1).

HSI	ARMA(1,0)			ARMA(0,1)			ARMA(1,1)		
	Normal	Student-t	GED	Normal	Student-t	GED	Normal	Student-t	GED
AIC	-0.6437	-6.1343	-0.9814	-0.95971	-6.1343	-0.98211	-2.749	-6.1182	-0.92683
BIC	-0.6274	-6.1139	-0.9609	-0.94340	-6.1139	-0.96172	-2.729	-6.0938	-0.90236
HQIC	-0.6376	-6.1266	-0.9737	-0.95358	-6.1266	-0.97445	-2.743	-6.1090	-0.91763
	ARMA(2,1)			ARMA(1,2)			ARMA(2,2)		
	Normal	Student-t	GED	Normal	Student-t	GED	Normal	Student-t	GED
AIC	-6.0585	-6.1169	-0.9245	-0,41512	-6,1218	-0.92441	1,6957	-6,1098	-0.92520
BIC	-6.0340	-6.0884	-0.8959	-0,39065	-6,0933	-0.89586	1,7243	-6,0771	-0.89258
HQIC	-6.0493	-6.1062	-0.9137	-0,40592	-6,1111	-0.91368	1,7065	-6,0975	-0.91294

Table 17: HSI Information Criteria for model selection of the Auto Regressive Moving Average (ARMA).

GARCH (1,1) is not significantly different from zero in its omega for the normal and generalized distribution but is significant for all parameters for the student t-distribution. The GJR-GARCH is non-significant in omega for the normal and generalized distribution and the mean for the student t-distribution. EGARCH is not significantly different from zero in four of its 6 parameters under the normal distribution. Whereas the student t and generalized error distribution have only one parameter which is non-significantly different from zero, AR2 and Omega, respectively.

HSI	GARCH(1,1)			EGARCH(1,1)			GJR-GARCH(1,1)		
	Normal	Student-t	GED	Normal	Student-t	GED	Normal	Student-t	GED
mu	0,013435	0,000147	0.000162	-0.00009	0.000148	0.000162	0,007967	0,000431	0.000169
s.e	0,00008	0	0	0.000329	0.000027	0	0,000003	0,000301	0
p-value	0	0	0	0.776326	0	0	0	0,152412	0
omega	0	0,000128	0	0.000120	0.000128	0	0	0,00013	0
s.e	0,000001	0,000009	0	0.000006	0.000009	0	0,00001	0,00001	0
p-value	0,88592	0	0.75384	0	0	0.75385	0,89829	0	0.74845
alpha1	0,998999	0,068696	0.050000	0.133760	0.074889	0.050000	1	0,071931	0.050000
s.e	0,019545	0,005217	0.000019	0.038939	0.035902	0.000019	0,000361	0,027139	0.000019
p-value	0	0	0	0.000592	0.036983	0	0	0,008039	0
shape		5,580338	2.000000		5.567047	2.000000		5,335257	2.000000
s.e		0,927316	0.000816		0.935582	0.000816		0,8272	0.000798
p-value		0	0		0	0		0	0
ar1	-0,03299	0,983322	0.006803	0.421737	0.996365	0.006860	-0,19247	-0,27528	-0.18955
s.e	0,00388	0,000503	0.000002	0.723044	0.008090	0.000002	0,000064	0,005287	0.000069
p-value	0	0	0	0.559705	0	0	0	0	0
ar2				-0.02149	-0.01289	-0.00453	-0,03843	-0,99144	-0.95843
s.e				0.019657	0.006691	0.000002	0,000012	0,00242	0.000347
p-value				0.274312	0.053951	0	0	0	0
ma1	1,668469	-0,98765	0.006691	-0.39497	-1.00000	0.006743	-1,92329	0,278844	0.203802
s.e	0,000198	0,00002	0.000002	0.723353	0.000005	0.000002	0,000486	0,002515	0.000074
p-value	0	0	0	0.585052	0	0	0	0	0
ma2	0,671011	-0,01781	-0.00429				0,980848	0,997148	0.980824
s.e	0,000078	0,000029	0.000001				0,000299	0,000051	0.000356
p-value	0	0	0				0	0	0

HSI	GARCH(1,1)			EGARCH(1,1)			GJR-GARCH(1,1)		
	Normal	Student-t	GED	Normal	Student-t	GED	Normal	Student-t	GED
AIC	-6,1797	-6,1943	-6,2017	-6,2057	-6,2161	-6,2223	-6,2014	-6,2115	-6,2187
BIC	-6,1552	-6,1657	-6,1732	-6,1771	-6,1835	-6,1897	-6,1729	-6,1789	-6,186
HQIC	-6,1705	-6,1835	-6,191	-6,1949	-6,2039	-6,21	-6,1907	-6,1992	-6,2064

Table 18: GARCH (1,1) family models statistics and information criteria for the Hang Seng index. The values which are not statistically different from zero are highlighted in blue.

The AIC for the higher order models suggest an improvement may be possible. The AIC suggests (2,1) for the GARCH and EGARCH models and due to no improvement in the AIC for the GJR GARCH, (1,1) is selected despite the non-significant omega and mu (student t-distribution). The results are again calculated

using the (1,1) on all models and directly compared with the results of the best AIC models.

HSI	GARCH			EGARCH			GJR-GARCH		
	Normal	Student-t	GED	Normal	Student-t	GED	Normal	Student-t	GED
AIC(1,1)	-6,1797	-6,1943	-6,2017	-6,2057	-6,2161	-6,2223	-6,2014	-6,2115	-6,2187
AIC(2,1)	-6,1856	-5,3337	-6,2069	-6,2289	-5,3691	-6,2412	-6,1989	-5,3764	-6,2165
AIC(1,2)	-6,1781	-6,1927	-6,2001	-6,2043	-6,2148	-6,2209	-6,200	-6,2100	-6,2172
AIC(2,2)	-6,1841	-5,5639	-6,2053	-6,2278	-5,6079	-6,2400	-6,1975	-5,5797	-6,215

Table 19: Akaike's Information Criteria of higher order GARCH family models for the Hang Seng index.

The MSE gradually increases as the forecasting horizon increases. The most accurate forecasting horizon is 1 month where the naïve model outperforms ARCH and is very similar to that of GARCH. ARCH improves with the use of the student t distribution whereas the GARCH models all perform best under the generalized error distribution.

nDays	Distribution	MSE				
		Naive	ARCH	GARCH(1,1)	EGARCH(1,1)	GJR-GARCH(1,1)
21	Normal	0,001199	0,0184	0,001198	0,0009143	0,00105
	Student t	0,001199	0,003331	0,001181	0,0009138	0,001048
	GED	0,001199	0,005217	0,001166	0,0009027	0,001047
63	Normal	0,002988	0,04255	0,002345	0,002259	0,002303
	Student t	0,002988	0,02052	0,002365	0,002353	0,002443
	GED	0,002988	0,01618	0,002372	0,002332	0,002404
126	Normal	0,005718	0,06676	0,005920	0,005127	0,005374
	Student t	0,005718	0,05959	0,005821	0,005200	0,005638
	GED	0,005718	0,03287	0,005860	0,005135	0,005595
189	Normal	0,007221	0,17990	0,009283	0,007169	0,007646
	Student t	0,007221	0,11120	0,009964	0,006719	0,008178
	GED	0,007221	0,04973	0,009545	0,006358	0,008009
252	Normal	0,01394	0,02080	0,01405	0,01230	0,01373
	Student t	0,01394	0,41630	0,01353	0,01210	0,01381
	GED	0,01394	0,06597	0,01349	0,01183	0,01372

Table 20: MSE forecasting results for the Hang Seng index over multiple time horizons.




Figure 17: Clustered Column Chart of HSI MSE results.

The results of the predictive power show that when using the naïve model a forecasting horizon of 3 months (63 days) offers a slightly larger predictive power than that of 1 month. The ARCH performs poorly over all time horizons and all distributions and actually improves when forecasting over 1 year and using the normal error distribution. The GARCH model performs very similarly to that of the naïve model under the normal distribution when using a 1 month forecasting horizon, but it improves slightly when using one of the non-normal distributions. The predictive power of GARCH (1,1) is twice as large as the best naïve model when considering the 3 month forecasting horizon where the EGARCH (1,1) and GJR-GARCH (1,1) are also only slightly improved compared to the simpler GARCH (1,1) model. The 3 month forecast gives the best predictive power under the normal distribution for all three GARCH (1,1) models. As the forecasting horizon increases the predictive power decreases (becomes worse) and even becomes negative for EGARCH and GJR-GARCH over the 9 month (depending on distribution) and 1 year horizons.

nDays	Distribution	Predictive Power				
		Naive	ARCH	GARCH(1,1)	EGARCH(1,1)	GJR-GARCH(1,1)
21	Normal	17,11	-1172	17,18	36,81	27,44
	Student t	17,11	-130,2	18,41	36,84	27,57
	GED	17,11	-260,6	19,43	37,61	27,66
63	Normal	17,38	-1076	35,16	37,54	36,34
	Student t	17,38	-467,3	34,61	34,95	32,45
	GED	17,38	-347,3	34,43	35,52	33,53
126	Normal	11,11	-937,8	7,966	20,3	16,45
	Student t	11,11	-826,3	9,514	19,16	12,35
	GED	11,11	-410,9	8,906	20,17	13,03
189	Normal	10,85	-2121	-14,61	11,49	5,597
	Student t	10,85	-127	-23,02	17,05	-0,9612
	GED	10,85	-514	-17,85	21,5	1,117
252	Normal	-27,36	-90,01	-28,34	-12,38	-25,39
	Student t	-27,36	-3703	-23,6	-10,5	-26,14
	GED	-27,36	-502,6	-23,27	-8,014	-25,35

Table 21: Predictive Power forecasting results for the Hang Seng index over multiple time horizons.

When using the higher order models, as suggested by the lowest AIC statistic, over 1 month the predictive power of GARCH (2,1) and EGARCH (2,1) are not improved under any of the distributions. However the higher order models do result in a slight improvement of the GARCH and EGARCH models under the Gaussian distribution (and student t for EGARCH). Since the 3 month forecasting period produced the best predictive power statistics for all statistics under GARCH (1,1) and GJR-GARCH (1,1) we decided to run the higher order models on the 3 month forecast as well. The MSE did not improve for the higher order EGARCH model and only improved for the higher order GARCH model using the normal distribution. The higher order models improved under all distributions for the GARCH and EGARCH models except for the normally distributed higher order GARCH model when using the predictive power statistic.

nDays	Distribution	MSE				
		Naive	ARCH	GARCH(2,1)	EGARCH(2,1)	GJR-GARCH(1,1)
21	Normal	0,001199	0,01498	1,21E-03	9,16E-04	0,00105
	Student t	0,001199	3,33E-03	1,18E-03	9,15E-04	0,001048
	GED	0,001199	5,22E-03	1,17E-03	9,03E-04	0,001047
Predictive Power						
nDays	Distribution	Naive	ARCH	GARCH(2,1)	EGARCH(2,1)	GJR-GARCH(1,1)
21	Normal	17,11	-935	16,44	36,72	27,44
	Student t	17,11	-130,2	18,35	36,76	27,57
	GED	17,11	-260,6	19,18	37,58	27,66
MSE						
nDays	Distribution	Naive	ARCH	GARCH(2,1)	EGARCH(2,1)	GJR-GARCH(1,1)
63	Normal	0,002988	0.04294	0.002396	0.002244*	0.002303
	Student t	0,002988	0.02052	0.002358*	0.002334*	0.002443
	GED	0,002988	0.01618	0.002372	0.002310*	0.002404
Predictive Power						
nDays	Distribution	Naive	ARCH	GARCH(2,1)	EGARCH(2,1)	GJR-GARCH(1,1)
63	Normal	17,38	-1087	33.75	37.96*	36.34
	Student t	17,38	-467.3	34.82*	35.47*	32.45
	GED	17,38	-347.3	34.49*	36.15*	33.53

Table 22: MSE and Predictive Power forecasting results for higher order GARCH AND EGARCH models over a 21 and 63 day forecasting horizon. Values marked with an asterisk (*) show an improvement.

6. Discussion

Our results show that the longest reasonable forecasting horizon of interest was that of 3 months for the HSI. The 1 month forecasting period was of most relevance and gave the best predictive power and lowest MSE results for the FTSE 100 and S&P 500. Previous papers, as suggested in the literature review, have focused on short-term (< 3 months) forecasting horizons. Our findings have also shown that relevance of forecasting over longer time horizons diminishes after 3 months.

To complement Cao and Tsay (1992) we have compared our findings using the EGARCH (1,1) model and found that over longer time horizons (12 months) our EGARCH model outperforms all other models tested when considering both the MSE and predictive power for the FTSE and HSI indices over all three distributions. The same cannot be said for the S&P 500 where EGARCH (1,1) only outperforms ARCH and GARCH (1,1) whilst using a student t-distribution. EGARCH (1,1) is outperformed by the naïve model, which actually performs the best for the 12 month forecasting horizon out of all the models analyzed with the S&P 500.

Kroner, Kneafsey and Claessens (1995) found that the COMB which combines the ISD and GARCH forecasts, outperformed the other five forecasts when using the MSE as an evaluation technique. This supports our findings that GARCH, and the extensions of GARCH, outperform the other models studied in this thesis over all three indices.

The findings of Figlewski (1997) are confirmed for the S&P 500 where the GARCH (1,1) outperforms both the naïve and ARCH models with forecasting horizons shorter than 12 months using both the MSE and predictive power measurements. Since we also include EGARCH (1,1) and GJR-GARCH (1,1) we find that the GARCH (1,1) is often outperformed by either of these. For the 12 month forecasting horizon GARCH does however outperform EGARCH (1,1) for the normal and generalized distributions. Although Figlewski looked at other markets (3M US T-Bill, 20Y T-Bond, Deutschemark Exchange Rate (DM per \$)) and we compared pure indices, we can conclude that the GARCH (1,1) outperforms the most basic models (Naïve and ARCH) on all forecasted horizons shorter than 12 months with the FTSE 100 index.

In terms of the HSI index, GARCH (1,1) only ever slightly outperforms the naïve model and with longer than 6 months forecasting the GARCH (1,1) model is outperformed. The only instance where GARCH (1,1) outperforms using a 12 month time horizon is when the student t-distribution is used. These results hold for both the MSE and the predictive power.

Our results agree with those of Poon and Granger (2003), when they reviewed 17 different papers and stated that the GARCH (1,1) outperforms the more restrictive ARCH model. This is clear for all three indices regardless of the distribution and is a consequence of the restrictions of ARCH.

Liu and Morley (2009) suggest that the non-normal GARCH model fits better than the normally distributed GARCH models over 1 period forecasting. From our results we see that there exists sensitivity to distribution functions for the 1 month forecasts where the normal error distribution is outperformed. Since our findings are extended over multiple forecasting periods we have looked to see if Lui and Morley's findings also hold over longer forecasting periods. We find that for the FTSE the student t continues to outperform the normal and GED, which tend to compete and converge towards zero as the forecasting horizon increases. The same cannot be said for the S&P 500 index where the GED is closer to the student t-distribution and both are outperformed by the normal distribution on forecasting time horizons longer than 1 month. For longer time horizons with the HSI the distributions are ranked closely when using both the MSE and predictive power. The non-normal distributions outperform the normal distribution over the 1, 6 and 12 month forecasting horizons.

Wei (2012) finds that the normal distribution is the best error distribution over a 12 month forecasting horizon regardless of the model. Our results do not confirm this and we find that the student t-distribution is the best performer for both the FTSE and S&P 500 over the same period. With these two indices the best performing models are found to be EGARCH (1,1) and GJR-GARCH (1,1) respectively. The HSI index performs best over the 252 day forecasting horizon using the generalized error distribution and our findings point to the EGARCH (1,1) as the best performer using both the MSE and predictive power.

It would appear from our results that since the in-sample data is not normally distributed, the student t-distribution is able to cope better with models of a lower order. We see this through selection of the models using AIC, that the greater the model parameters (i.e. GJR- GARCH) the lower the model selected and generally the better results are achieved with the time horizon of 3 months or shorter.

We used ARMA (1,1) for all three indices. The calculations and results with the ARMA and Information criteria statistics are all based on a 1 month forecasting horizon. This could be studied individually for each time horizon and each index. The comparisons of our results with those in the literature review seem, for the majority of papers, to agree with only a few small digressions. Most existing papers only study one time horizon or if multiple time horizons are studied they usually lack different error distributions or a suitable number of models in order to make the necessary comparisons. We also noticed that higher order models had often not been used in previous research. Our research has proven that they do not offer a considerable amount of additional information and in fact most researchers document that higher order GARCH models are often found to be unstable and unreliable.

7. Conclusion

This thesis covers four main areas. First the basic structure of the ARMA modelling framework was investigated with correlation functions and information criteria's. Secondly, we looked at the properties of the GARCH model and its extensions with respect to error distributions. Thirdly, we looked at higher order models beyond (1,1), and finally the results of the MSE and P-value of the out-of-sample forecasting for all models, time horizons and distributions were analysed.

We found that the error distribution does play a part in the accuracy of the model over a 1 month forecasting horizon where the non-normal distributions gave the best results. Over the longer term forecasting horizons the error distribution plays a lesser part in the accuracy, as can be seen for the 3 month forecasting for HSI where the normally distributed models give the best evaluation results. For longer forecasting

time horizons it is not necessarily the model with the best in-sample fit that provides the best out-of-sample forecast.

The out-of-sample forecasting performance results of the FTSE 100 and S&P 500 over 1, 3 and 6 month forecasting periods select the same preferred model (GJR-GARCH). There is not a single volatility model that is preferred based on the 9 month forecasting horizon. With a 12 month forecasting horizon the FTSE 100 and HSI select the EGARCH model as the most accurate whereas the naïve model was selected for the S&P 500. The HSI is the only index for which the same model is selected (EGARCH) as the best performer, regardless of the forecasting horizon. The MSE and P value offered the same results and the inclusion of both was only necessary for direct comparisons with previous literature.

When we tested the higher order models over the most accurately forecasted time horizons, we found the results of most models only improved marginally. The additional time and work involved with the analysis of higher order models for such little improvement was far from optimal. We also find that models with a larger number of parameters required a lower order but still resulted in relevant and competitive forecasting accuracy statistics.

Acknowledgements

We are especially grateful to our supervisor Professor Valeriy Zakamulin for his helpful comments, suggestions and corrections towards the realization of this work.

We are also indebted to all the lecturers who have taught us throughout the entire program and we wish to express a heartfelt thank you to our families for their help and encouragement throughout our education.

References

- Agro, G, 1995. Maximum likelihood estimation for the exponential power function parameters, *Communications in Statistics-Simulation and Computation* 24, 523-536.
- Black, K, 2009. *Business Statistics: Contemporary Decision Making*. 2009, 6th edition edition.(John Wiley & Sons).
- Blair, Bevan J, Ser-Huang Poon, and Stephen J Taylor, 2010. *Handbook of Quantitative Finance and Risk Management Forecasting S&P 100 volatility: the incremental information content of implied volatilities and high-frequency index returns*.(Springer).
- Bollerslev, Tim, 1986. Generalized autoregressive conditional heteroskedasticity, *Journal of Econometrics* 31, 307-327.
- Bollerslev, Tim, 1987. A conditionally heteroskedastic time series model for speculative prices and rates of return, *The review of economics and statistics*, 542-547.
- Box, G. M. Ljung and G. E. P., 1978. On a measure of lack of fit in time series models, *Biometrika* 65, 297-303.
- Brooks, Chris, 2014. *Introductory econometrics for finance*, Third edition. edition.(Cambridge University Press, New York).
- Cao, Charles Q, and Ruey S Tsay, 1992. Nonlinear time - series analysis of stock volatilities, *Journal of applied econometrics* 7, S165-S185.
- Cipan, Woschnagg, 2004. *Evaluating Forecast Accuracy*(University of Vienna).
- Ederington, Louis H, and Wei Guan, 2002. Measuring implied volatility: is an average better? Which average?, *Journal of Futures Markets* 22, 811-837.
- Engle, Robert, 2003. Risk and volatility: Econometric models and financial practice, *American Economic Review*, 405-420.
- Engle, Robert F. , 1982. Autoregressive Conditional Heteroskedasticity with Estimates of the Variance of United Kingdom Inflation, *Econometrica: Journal of the Econometric Society* 50, pp. 987-1007.
- Figlewski, Stephen, 1997. Forecasting volatility, *Financial markets, institutions & instruments* 6, 1-88.
- Figlewski, Stephen, 1998. Derivatives risks, old and new.
- Giller, Graham L, 2005. A generalized error distribution.
- Giusti, Antonio, Gunter Ritter, and Maurizio Vichi, 2012. *Classification and Data Mining*(Springer Science & Business Media).
- Glosten, Lawrence R, Ravi Jagannathan, and David E Runkle, 1993. On the relation between the expected value and the volatility of the nominal excess return on stocks, *The journal of finance* 48, 1779-1801.
- Gosset, "Student" William Sealy, 1908. The probable error of a mean, *Biometrika*, 1-25.
- Green, T Clifton, and Stephen Figlewski, 1999. Market risk and model risk for a financial institution writing options, *The journal of finance* 54, 1465-1499.
- Gujarati, Damodar N, 2014. *Econometrics by example*(Palgrave Macmillan Hampshire, UK).
- Helmert, FR, 1876. Die Genauigkeit der Formel von Peters zur Berechnung des wahrscheinlichen Beobachtungsfehlers director Beobachtungen gleicher Genauigkeit, *Astronomische Nachrichten* 88, 300-303.

- Koop, Gary, 2006. *Analysis of financial data* (John Wiley & Sons Inc., Chichester ; Hoboken, NJ).
- Kroner, Kenneth F, Kevin P Kneafsey, and Stijn Claessens, 1995. Forecasting volatility in commodity markets, *Journal of Forecasting* 14, 77-95.
- Lamoureux, Christopher G, and William D Lastrapes, 1990. Persistence in variance, structural change, and the GARCH model, *Journal of Business & Economic Statistics* 8, 225-234.
- Li, Kai, 2002. Long-memory versus option-implied volatility predictions, *The Journal of Derivatives* 9, 9-25.
- Liu, Wei, and Bruce Morley, 2009. Volatility forecasting in the hang seng index using the GARCH approach, *Asia-Pacific Financial Markets* 16, 51-63.
- Mandelbrot, Benoit B, 1997. *The variation of certain speculative prices* (Springer).
- Nadarajah, Saralees, 2005. A generalized normal distribution, *Journal of Applied Statistics* 32, 685-694.
- Nelson, Daniel B, 1991. Conditional heteroskedasticity in asset returns: A new approach, *Econometrica: Journal of the Econometric Society*, 347-370.
- Poon, Ser-Huang, 2005. *A practical guide to forecasting financial market volatility* (John Wiley & Sons).
- Poon, Ser-Huang, and Clive WJ Granger, 2003. Forecasting volatility in financial markets: A review, *Journal of economic literature* 41, 478-539.
- Ruppert, David, 2011. *Statistics and data analysis for financial engineering* (Springer, New York).
- Scott, Elton, and Alan L Tucker, 1989. Predicting currency return volatility, *Journal of Banking & Finance* 13, 839-851.
- Speight, Alan EH, David G McMillan, and Owain Gwilym, 2000. Intra - day volatility components in FTSE - 100 stock index futures, *Journal of Futures Markets* 20, 425-444.
- Vasudeva, R, and J Vasantha Kumari, 2013. On general error distributions.
- Wei, Jiang, 2012. Modeling and predicting of different stock markets with GARCH model.
- Zakamulin, Valeriy, 2014. *Forecasting Volatility - Lecture Notes* (UiA).