

RESEARCH ARTICLE

Dynamics of triacylglycerol and EPA production in *Phaeodactylum tricornutum* under nitrogen starvation at different light intensities

Ilse M. Remmers^{1*}, Dirk E. Martens¹, René H. Wijffels^{1,2}, Packo P. Lamers¹

1 Bioprocess Engineering & AlgaePARC, Wageningen University and Research, Wageningen, The Netherlands, **2** Biosciences and Aquaculture, Nord University, Bodø, Norway

* ilse.remmers@wur.nl


 OPEN ACCESS

Citation: Remmers IM, Martens DE, Wijffels RH, Lamers PP (2017) Dynamics of triacylglycerol and EPA production in *Phaeodactylum tricornutum* under nitrogen starvation at different light intensities. PLoS ONE 12(4): e0175630. <https://doi.org/10.1371/journal.pone.0175630>

Editor: Adrianna Ianora, Stazione Zoologica Anton Dohrn, ITALY

Received: February 22, 2017

Accepted: March 29, 2017

Published: April 12, 2017

Copyright: © 2017 Remmers et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: This work was carried out within the EU project Fuel4Me. Fuel4Me was funded by the European Union's Seventh Programme for Research and Technology Development (EU-FP7) under grant agreement No. 308983. The funder had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Abstract

Lipid production in microalgae is highly dependent on the applied light intensity. However, for the EPA producing model-diatom *Phaeodactylum tricornutum*, clear consensus on the impact of incident light intensity on lipid productivity is still lacking. This study quantifies the impact of different incident light intensities on the biomass, TAG and EPA yield on light in nitrogen starved batch cultures of *P. tricornutum*. The maximum biomass concentration and maximum TAG and EPA contents were found to be independent of the applied light intensity. The lipid yield on light was reduced at elevated light intensities ($>100 \mu\text{mol m}^{-2} \text{s}^{-1}$). The highest TAG yield on light ($112 \text{ mg TAG mol}_{\text{ph}}^{-1}$) was found at the lowest light intensity tested ($60 \mu\text{mol m}^{-2} \text{s}^{-1}$), which is still relatively low to values reported in literature for other algae. Furthermore, mass balance analysis showed that the EPA fraction in TAG may originate from photosynthetic membrane lipids.

Introduction

Lipid production by phototrophic microalgae is only economically feasible today for high-value products, such as pigments and ω -3-fatty acids [1]. One of the major hurdles for commercial bulk microalgal lipid production is the low lipid yield on light [1–3].

Typically, microalgal lipid production is induced by nitrogen deficiency. When exposed to nitrogen deficient conditions, some microalgal species can accumulate triacylglycerol (TAG) up to 30–60% of dry weight [4–6]. However, nitrogen deficiency is often accompanied with impaired or even fully inactivated photosynthesis [4,7] and therefore results in a reduced TAG yield on light. For commercial production of TAGs, the TAG yield on light should be increased.

One of the most advertised microalgae for commercial lipid production is *Phaeodactylum tricornutum*. *P. tricornutum* is a salt-water diatom that is already cultivated at large scale for aquaculture. This microalgae produces large amounts of eicosapentaenoic acid (EPA; a long chain polyunsaturated fatty acid) and can produce TAG up to 30% of dry weight [4]. It is

Competing interests: The authors have declared that no competing interests exist.

generally acknowledged, however, that high light exposure can severely affect the biomass and lipid productivity of this diatom [8–12]. However, clear consensus regarding the impact of light intensity on TAG and EPA productivity in *P. tricornutum* is still lacking. The aim of this work is to obtain insight in the relation between incident light intensity and the accumulation of TAG and EPA under nitrogen starvation.

Material and methods

Strain, medium and precultivation

Phaeodactylum tricornutum SAG1090-1b was obtained from the Culture Collection of Algae Goettingen University, Germany. Cultures were maintained in 250 mL shake flasks containing 100 mL filter sterilized (pore size 0.2 μm) medium. The medium was designed to reach a biomass concentration of 2.5 g L^{-1} and consisted out of 252 mM NaCl, 16.8 mM KNO_3 , 3.5 mM Na_2SO_4 , 5 mM $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 2.4 mM $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$, 2.5 mM K_2HPO_4 , 10 mM NaHCO_3 , 28 μM NaFeEDTA, 80 μM $\text{Na}_2\text{EDTA} \cdot 2\text{H}_2\text{O}$, 19 μM $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$, 4 μM $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$, 1.2 μM $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$, 1.3 μM $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, 0.1 μM $\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}$, 0.1 μM Biotin, 3.7 μM vitamin B1 and 0.1 μM vitamin B12. The medium contained 100 mM, 2-[4-(2-hydroxyethyl)piperazin-1-yl]ethanesulfonic acid (HEPES) as a pH buffer. Prior to cultivation, the pH was adjusted with KOH to pH 7.2.

Cultures were maintained in a cultivation chamber at 25°C, orbitally shaken at 120 rpm, illuminated at an incident light intensity of 40–60 $\mu\text{mol m}^{-2} \text{s}^{-1}$ and a 16:8h light:dark (LD) cycle under enriched air (2% CO_2). Three days prior to the start of the experiment, cultures were transferred to an orbital shaker incubator with an incident light intensity of 160 $\mu\text{mol m}^{-2} \text{s}^{-1}$, 16:8h LD cycles and 2.5% CO_2 enriched air.

Photo bioreactor setup and operation

Experiments were performed in aseptic, flat-panel, airlift-loop reactors (Algaemist, Technical Development Studio, Wageningen University, the Netherlands). The reactor has a working volume of 400 mL, a light path of 14 mm and an illuminated area of 0.028 m^2 (Schematic illustration of the reactor design is given in Breuer et al. [9]). Illumination was provided using 6 LED lamps with a warm white light spectrum, (BXRA W1200, Bridgelux, USA, emission spectrum given in the S1 Fig) under a 16:8h LD cycle. The incident photon flux density (PFD_{in}) was measured with a LI-COR 190-SA 2π PAR (400–700 nm) quantum sensor (PAR, USA) and determined as the average over 28 points evenly distributed over the illuminated surface of the inside of the front glass panel of the reactor. The culture was continuously supplied with sterile moisturized air at 0.4 L min^{-1} . pH was maintained at 7.2 by on-demand sparging of CO_2 to the airflow and temperature was kept constant at 20°C. Reactor medium was similar to the preculture medium, with modifications to the amount of KNO_3 (5 mM instead of 16.8 mM) and exclusion of HEPES. 5 mM of KNO_3 is sufficient to reach a biomass concentration of 1.2 g L^{-1} (assuming a cellular nitrogen fraction of 10% per dry weight, measurements not shown). Per day, a maximum of 2 drops of 1% (v/v) antifoam (Antifoam B, J.T. Baker) were added manually to the reactor when foaming was visible.

Reactors were heat sterilized (60 min at 121°C) and inoculated at a biomass concentration of approximately 0.5 g L^{-1} of dry weight and grown at an incident light intensity of 100 $\mu\text{mol m}^{-2} \text{s}^{-1}$. Once the biomass concentration reached 1 g L^{-1} , which was typically three days after inoculation, the set points for incident light intensity were changed to the desired values (60, 100, 250, 500, or 750 $\mu\text{mol m}^{-2} \text{s}^{-1}$ incident light intensity). The day at which the light intensity was changed to the experimental set point was regarded as start of the experiment, and therefore referred to as $t = 0$. Periodically, samples (approximately 20 mL culture volume) were

taken aseptically and analysed for dry weight concentration, triacylglycerol (TAG) and total fatty acid (TFA) concentration and composition, dissolved (residual) nitrate Photosystem II (PSII) maximum quantum yield (F_v/F_m) ratio, and absorption cross section. After taking a sample, the culture volume was restored by adding 0.2 μm -filtered reactor medium to ensure that the pH probe remained submerged. Depending on whether the culture was in N-replete phase or N-depleted phase, this medium contained either 5 mM KNO_3 or no nitrogen, respectively.

Five different incident light intensities were selected that cover both low light and light saturation (light saturation point for *P. tricornutum* was found at 150–200 $\mu\text{mol m}^{-2} \text{s}^{-1}$ [13–15]). Three photo bioreactor experiments were conducted in biological duplicate to test reproducibility [16].

Measurements for dry weight, F_v/F_m and dissolved nitrogen were performed in technical triplicates. Measurements for the light absorption spectrum, lipid content and composition were performed in technical duplicate.

Biomass analysis

Samples were taken at regular time points to monitor biomass dry weight concentration gravimetrically as described by [17]. Light absorption spectrum was measured in a double-beam spectrophotometer (UV-2600, Shimadzu, Japan) equipped with an integrated sphere (ISR-2600). The average dry weight-specific optical cross section (α_C in $\text{m}^2 \text{g}^{-1}$) was calculated from the obtained absorption spectrum as described by [18]. The F_v/F_m ratio was measured using an AquaPen-C AP-C 100 FluorPen fluorometer (Photon Systems Instruments, Czech Republic) equipped with an orange LED emitter (excitation light 620 nm) as described by [19].

The triacylglycerol (TAG), membrane lipid (ML) and total fatty acid (TFA) content and fatty acid composition were determined by lipid extraction (chloroform:methanol mixture), lipid class separation using a silica column, transesterification and subsequently quantified using gas chromatography (GC-FID) as described by Breuer et al. [9]. The dissolved nitrate concentration was measured with an AQ2 nutrient analyser (Seal Analytical, USA) as described by Benvenuti et al. [20].

Calculating yield and productivity

The time averaged volumetric productivity ($r_i(t_0 \rightarrow t)$) was calculated by dividing the amount of formed product (i: biomass, triacylglycerol, EPA) over cultivation time according to Eq 1;

$$r_i(t_{N=0} \rightarrow t) = \frac{C_i(t) - C_i(t_{N=0})}{t - t_{N=0}} \quad (1)$$

The time averaged yield of TAG on light ($Y_{TAG,ph}(t_0 \rightarrow t)$) on each time point was calculated by dividing the amount of TAG produced in that period by the amount of light supplied in that period [21]. In addition, the photon costs for initial biomass production until nitrogen depletion were incorporated by assuming that this biomass was made with a theoretical net photosynthetic yield, $Y_{x,ph}$, of 1g mol_{ph} ;

$$Y_{TAG,ph}(t) = \frac{TAG(t) - TAG(t_{N=0})}{\frac{I_0 \cdot (t - t_{N=0})}{z} + \frac{C_{x,N=0}}{Y_{x,ph}}} \quad (2)$$

In Eq 2, TAG(t) and TAG($t_{N=0}$) represent the TAG concentration at time point t and the moment of nitrogen starvation, respectively. The exact time of nitrogen starvation ($t_{N=0}$) was estimated by interpolation of the biomass concentration to 1.2g L^{-1} . I_0 is the incident light

intensity, z is the reactor depth and $C_{x,N=0}$ is the biomass concentration at the onset of nitrogen starvation.

Results and discussion

Effect of light on biomass growth

The impact of five different light intensities on biomass, TAG and EPA productivity during nitrogen starvation was assessed in batch-wise operated flat-panel photo bioreactors that were subjected to 16:8h light:dark (LD) cycles. Results for dry weight, quantum yield and TAG content are shown in Fig 1.

Sufficient nitrogen was added to reach a biomass concentration of 1.2 g DW L^{-1} . After that biomass concentration was reached, all nitrogen was used and the culture was exposed to nitrogen starvation. The biomass concentration further increased up to 2.2 g L^{-1} of dry weight, which was due to the accumulation of TAG (Fig 1) and carbohydrates (S1 Table). The maximum achieved biomass concentration was reached 5 days after nitrogen starvation started, except for the culture exposed to the lowest light conditions, which reached the maximum biomass 14 days after nitrogen starvation started. The maximum reached biomass concentration was $2.15 \pm 0.1 \text{ g L}^{-1}$, independent of the supplied light intensity (Fig 1A). Other studies showed similar findings for the freshwater microalgae *Acutodesmus obliquus* under nitrogen starvation [4].

Cultures that are light limited are expected to show an increase in biomass productivity upon increasing light intensities [9]. However, our results show no differences in productivities for the cultures exposed to $100 \mu\text{mol m}^{-2} \text{ s}^{-1}$ and higher (Fig 1). This similarity in productivity might be explained by photo inhibition that occurred from $100 \mu\text{mol m}^{-2} \text{ s}^{-1}$ onwards. As soon as the photosynthetic machinery becomes inhibited, further increasing light intensity would not enhance the productivity anymore [22]. According to Fig 1B, a strong decline in F_v/F_m was observed (from 0.6 to 0) for cultures exposed to light intensities above $100 \mu\text{mol m}^{-2} \text{ s}^{-1}$ immediately after nitrogen starvation. A decline in F_v/F_m is generally acknowledged with reduced photosynthetic efficiency or photo damage [19,23,24]. Geider et al. [13] presented a light curve for *P. tricornutum*. Similar to our findings, they reported that from $100 \mu\text{mol m}^{-2} \text{ s}^{-1}$ onwards light saturation occurs in *P. tricornutum*. In addition, Ación Fernández et al. [25] performed outdoor growth experiments with *P. tricornutum*, where the maximum biomass productivity was reached at average irradiances of $50\text{--}150 \mu\text{mol m}^{-2} \text{ s}^{-1}$ and decreased at light intensities higher than $150 \mu\text{mol m}^{-2} \text{ s}^{-1}$. Altogether, it is likely that in this work *P. tricornutum* becomes photo inhibited at light intensities above $100 \mu\text{mol m}^{-2} \text{ s}^{-1}$.

Changes in TAG accumulation, productivity and yields

Independent of incident light intensity, the TAG concentration increased from 0.05 g L^{-1} (3% of dry weight) under nitrogen replete conditions to $\sim 0.41 \text{ g L}^{-1}$ ($\sim 25\%$ of dry weight) after prolonged nitrogen starvation (Fig 1 and S2 Table). These maximum TAG contents are in agreement with observations from literature (Table 1).

Except for the culture exposed to the lowest light intensity ($60 \mu\text{mol m}^{-2} \text{ s}^{-1}$), TAG accumulation did not differ significantly for the tested conditions (Fig 1). The culture exposed to high incident light intensity ($750 \mu\text{mol m}^{-2} \text{ s}^{-1}$) showed a decline in TAG content from day 11 onwards, which might indicate photo damage or even cell death.

The TAG yield on light is directly proportional to areal productivity and is therefore also strongly related to investment and variable production costs. The maximum time-averaged TAG yield on light is often used as indicator to determine optimal harvest moments [7]. Our study shows that large differences were observed in the maximum time-averaged TAG yield




Fig 1. Impact of five different incident light intensities on (A) the biomass concentration, (B) F_v/F_m ratio and (C) the TAG content over time. Light was supplied at intensities of 60 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (blue circles), 100 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (red squares), 250 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (green triangles), 500 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (black circles) and 750 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (open diamonds). Sufficient nitrogen was added to reach biomass concentrations of 1.2 g L^{-1} , thereafter cells were exposed to nitrogen starvation. Average values of duplicate reactor runs are shown, except for the cultures exposed to 250 and 750 $\mu\text{mol m}^{-2} \text{s}^{-1}$ ($n = 1$). Error bars indicate the standard deviation of both cultures from the average value.

<https://doi.org/10.1371/journal.pone.0175630.g001>

on light among the different tested incident light intensities (Fig 2). Of all tested conditions, the cultures exposed to low light (60 $\mu\text{mol m}^{-2} \text{s}^{-1}$) reached the highest maximum TAG yield on light (112 $\text{mg TAG mol}_{\text{ph}}^{-1}$). Increasing the incident light intensity only resulted in lower TAG yields on light: 96 $\text{mg TAG mol}_{\text{ph}}^{-1}$ for the culture exposed to 100 $\mu\text{mol m}^{-2} \text{s}^{-1}$ and

Table 1. Comparison of the maximum TAG and total fatty acid content obtained in this study to literature values. *TAG contents were estimated from the total fatty acid content, assuming 8% of membrane lipids per dry weight.

TAG content (% dry weight)	Total fatty acid content (% dry weight)	Strain	Reference
25	31	<i>P. tricornutum</i>	This study
22*	30	<i>P. tricornutum</i>	[4]
20*	28	<i>P. tricornutum</i>	[26]
26*	34	<i>P. tricornutum</i>	[27]

<https://doi.org/10.1371/journal.pone.0175630.t001>

36–48 mg TAG mol_{ph}⁻¹ for cultures exposed to even higher light intensities. In comparison with other studies (Table 2), *P. tricornutum* shows relative low TAG yields on light compared to other strains at similar or higher incident light intensities.

Sustained biomass productivity upon nitrogen starvation is a key selection parameter for efficient TAG production. As shown in Fig 1A, *P. tricornutum* only showed a 2–2.2-fold biomass increase upon nitrogen starvation. Other oleaginous green microalgae (e.g. *Acutodesmus obliquus*, *Chlorella zofingiensis*) showed a 6–8 fold increase in biomass upon nitrogen starvation [4,5,19]. High photon to TAG conversion yields are only possible when light energy is efficiently used to fix carbon (e.g. measured as biomass productivity) and when the majority of fixed carbon is channelled towards storage lipids (e.g. TAG content). Literature reports that diatoms are specifically known to mobilize alternative electron transport (AET) pathways, that can eradicate up to 50% of the electrons released by PS II [28–30]. Also, increased Non-Photochemical Quenching (NPQ) rates were found for *P. tricornutum* when exposed to




Fig 2. Time averaged TAG yield on light for five different incident light intensities over time. Incident light intensities of 60 μmol m⁻² s⁻¹ (blue circles), 100 μmol m⁻² s⁻¹ (red squares), 250 μmol m⁻² s⁻¹ (green triangles), 500 μmol m⁻² s⁻¹ (black circles) and 750 μmol m⁻² s⁻¹ (open diamonds). Sufficient nitrogen was added to reach biomass concentrations of 1.2 g L⁻¹, thereafter cells were exposed to nitrogen starvation. Error bars indicate the standard deviation of both cultures from the average value.

<https://doi.org/10.1371/journal.pone.0175630.g002>

Table 2. Comparison of the maximum time-averaged TAG yield on light obtained in this study to literature values. CL: continuous light, LD16:8: light dark cycles with 16h of light followed by 8h of darkness. ¹ was calculated by assuming that the total lipid fraction existed for 90% out of TAG. ² Assumed that 42% (J/J) of the total irradiance has a wavelength in the photosynthetic active radiation spectrum with an energy content of 0.217 MJ mol_{ph}⁻¹. ³ Light was supplied with a sinus function with a solar noon at 1500 μmol m⁻² s⁻¹ and a daily averaged light intensity of 636 μmol m⁻² s⁻¹. ⁴ the calculated range is depends on initial biomass concentration at the moment of nitrogen deprivation and includes measurement errors. ⁵ calculation includes the costs for inoculum production (assuming 1 g biomass mol_{ph}⁻¹).

Species	TAG yield on light	Incident light intensity	Operational strategy	Reference
	(g mol ⁻¹)	(μmol m ⁻² s ⁻¹)		
<i>Phaeodactylum tricornutum</i>	0.12	60	Batch, LD16:8	This study
<i>Acutodesmus obliquus</i>	0.24, 0.13, 0.09, 0.04	200, 500, 800, 1500	Batch, CL	[9]
<i>Acutodesmus obliquus</i>	0.15	500	Batch, CL	[39]
<i>Acutodesmus obliquus slm1</i>	0.22	500	Batch, CL	[39]
<i>Acutodesmus obliquus</i>	0.16	500	Batch, LD16:8	Remmers et al. (submitted)
<i>Acutodesmus obliquus slm1</i>	0.20	500	Batch, LD16:8	Remmers et al. (submitted)
<i>Clorella zofingiensis</i>	0.32	245	Batch, CL	[5]
<i>Chlorella vulgaris</i>	0.16 ¹	615 ²	Batch, outdoor	[40]
<i>Chlorococcum litrorale</i>	0.16	636 ³	Batch, sinusoidal LD 16:8	[41]
<i>Chlorococcum litrorale</i> , S5 sorted population	0.32	636 ³	Batch, sinusoidal LD 16:8	[41]
<i>Neochloris oleoabundans</i>	0.16	218	Batch, CL	[6]
<i>Nannochloropsis oculata</i>	0.17–0.25 ⁴	250	Batch, CL	[42]
<i>Nannochloropsis sp.</i>	0.10	636 ³	Batch, sinusoidal LD 16:8	[20]
<i>Nannochloropsis sp.</i>	0.14 ⁵	636	Batch, CL	[7]

<https://doi.org/10.1371/journal.pone.0175630.t002>

increased light intensities [31–33]. Both processes are known to protect the cells from photo inhibition but to also result in inherent productivity losses. For *P. tricornutum*, these mechanisms might explain the reduced photosynthetic efficiency compared to other microalgae under low light nitrogen starvation.

Considering the abovementioned light sensitivity of *P. tricornutum* during nitrogen starvation, precautions should be taken to prevent the culture from extensive photo inhibition when working in high irradiance conditions [34]. Examples are to increase the culture cell density to create a dark zone in which the photo inhibitory damage can be repaired [35,36] or to execute strain improvement for reduced antenna size [37,38].

FA composition

The fatty acid composition of TAG and membrane lipids (ML) did not substantially change over time nor with incident light intensity (the complete fatty acid composition of both fractions during the time-course of the experiments can be found in Table 3). The most abundant fatty acids in TAG were found to be C16:1 (44% of total TAG fatty acids) and C16:0 (30% of total TAG fatty acids). Only a small fraction of C20:5 (EPA) was found in the TAG pool (6–10% of total TAG fatty acids). The membrane lipid fraction was primarily comprised of C16:0 (20% of total ML fatty acids), C16:1 (25–30% of total ML fatty acids) and C20:5 (25–30% of total ML fatty acids) (Table 3). Literature reports similar fatty acid profiles in *P. tricornutum* [4,43–45]. As a consequence of TAG accumulation (Fig 1) and the lower EPA content of TAG compared to membrane lipids, the EPA fraction of the total fatty acid pool in the biomass dropped from 25% to 9% of the total fatty acids (S3 Table). Because the biomass productivity declined at the onset of nitrogen depletion, EPA productivity was highest (6–12 mg L⁻¹ day⁻¹) under nitrogen replete growth conditions.

Table 3. Average fatty acid composition of *P. tricornutum* in the TAG and ML lipid fraction for five different light intensities: 60, 100, 250, 500 and 750 $\mu\text{mol m}^{-2} \text{s}^{-1}$. Values are averaged over the entire time course of the experiment (n = 7). Standard deviations are shown between brackets.

	% of fatty acids in TAG fraction					% of fatty acids in ML fraction				
	60	100	250	500	750	60	100	250	500	750
C14:0	5.1	5	5.5	5.2	5.8	6.5	5.7	5.5	6.4	6
	(0.7)	(0.5)	(0.9)	(0.5)	(0.4)	(0.3)	(0.7)	(0.8)	(0.4)	(1.2)
C16:0	24.8	26.3	31.8	29.8	25	19.1	21.2	25.7	26.3	31.1
	(3.6)	(4.5)	(1.4)	(3.7)	(0.4)	(3.3)	(3.7)	(5.6)	(7.9)	(8.5)
C16:1	40.4	44.8	43.1	42	49.1	26.4	26.5	26.6	25.8	21.8
	(5.1)	(3.2)	(3.1)	(5.5)	(3.4)	(2.3)	(2.1)	(2.2)	(1.5)	(4.3)
C16:2	0.7	1.3	1.9	0.7	2.1	3	2.5	1.2	1.7	1.2
	(0.5)	(1.3)	(0.4)	(0.3)	(3.1)	(1.1)	(1.9)	(0.6)	(1)	(0.5)
C16:3	1.1	1.3	1.2	0.9	1.4	5.2	4.3	3.7	4.1	3.4
	(0.5)	(0.8)	(0.6)	(0.3)	(0.7)	(2.3)	(1.7)	(1.9)	(2.5)	(2.3)
C18:0	1.8	1.7	1.6	1.4	1.5	2.4	3.6	5.2	3.4	7.6
	(0.2)	(0.3)	(0.5)	(0.2)	(0.5)	(1)	(1.5)	(2.8)	(1.3)	(5.3)
C18:1	5.8	5.7	5.1	4.2	4.6	2.7	3.8	4.1	4	4.5
	(1.2)	(1.4)	(1.1)	(0.6)	(1.1)	(1.1)	(2.4)	(1.7)	(1.7)	(1.6)
C18:2w6	1.4	1.3	1.3	1.1	1.3	2	1.4	2.2	2	2
	(0.3)	(0.3)	(0.2)	(0.3)	(0.2)	(0.6)	(0.4)	(0.8)	(0.8)	(1.1)
C18:3w3	6.7	3.3	2.8	5.6	2.7	1.5	1.5	1.7	1.3	1.5
	(4.3)	(2.4)	(0.6)	(5.4)	(0.6)	(0.4)	(0.6)	(0.5)	(0.3)	(0.8)
C20:3'(n3)	0.9	0.4	0.1	1.7	0.4	5.8	5.7	3.8	3.7	3
	(0.8)	(0.3)	(0.1)	(4)	(0.8)	(1.7)	(2.7)	(1.3)	(0.7)	(2.1)
C20:5'(n3)	7.8	7.1	6.7	7.4	7.2	25.5	23.8	20.3	21.3	18
	(1.4)	(1.4)	(0.8)	(1)	(1.1)	(5.7)	(7)	(5.5)	(7.1)	(8.9)

<https://doi.org/10.1371/journal.pone.0175630.t003>

EPA incorporation in membrane lipids and triacylglycerol

Fig 3 shows the EPA concentration per litre culture with respect to the TAG en membrane lipid fraction. Apart from a small increase until day 5, neither a substantial net synthesis nor breakdown of EPA upon N starvation is observed. However, the distribution of EPA over TAG and membrane lipids changes over time. Upon nitrogen depletion, a substantial increase in EPA distribution towards TAG can be observed. Simultaneously, we observed a decrease in EPA that was bound to membrane lipids. Therefore, we hypothesize that the EPA found in TAG under nitrogen starvation could partially be derived from EPA-containing membrane lipids that were restructured upon nitrogen depletion. Similar observations have been made for *Nannochloropsis oceanica* [46], *Nannochloropsis oculata* and *Pavlova lutheri* [47].

For many species, nitrogen depletion results in an inherent degradation of photosynthetic membrane lipids with simultaneous accumulation of TAG. Part of the fatty acids originating from membrane lipids could thus be used as fatty acyl donors for TAG synthesis under nutrient deprivation (Fig 3). However, detailed knowledge on the mechanisms behind carbon partitioning in microalgae under nitrogen deprivation is still lacking [47–51] and further research using ¹³C labelling should confirm these observations. To further improve EPA productivity by means of metabolic engineering or process development [52], it is crucial to understand the regulation of EPA towards different lipids classes (e.g. TAG or structural membrane lipids) under different environmental conditions.




Fig 3. EPA concentration in both TAG and ML lipids over time at an incident light intensity of $100 \mu\text{mol m}^{-2} \text{s}^{-1}$. The moment of nitrogen starvation is indicated with a dashed line.

<https://doi.org/10.1371/journal.pone.0175630.g003>

Conclusion

Elevated light intensities ($>100 \mu\text{mol m}^{-2} \text{s}^{-1}$) reduce the photosynthetic efficiency and the TAG yield on light during nitrogen starvation, likely caused by photo inhibition. Compared to other microalgae strains, relative low TAG yields on light were observed for *P. tricornutum* as the maximum yield was only $112 \text{ mg TAG mol}_{\text{ph}}^{-1}$ (at $60 \mu\text{mol m}^{-2} \text{s}^{-1}$). TAG contents were similar for all light intensities tested (i.e. 25% of dry weight). Further analysis on the fatty acid composition revealed that the TAG fraction of *P. tricornutum* contains 6–10% of EPA. Mass balance analysis revealed this EPA fraction may originate partly from photosynthetic membrane lipids that were restructured after nitrogen depletion.

Supporting information

S1 Fig. Emission spectrum of the Algaemist Bridgelux LED panel. Data obtained from de Mooij et al (2014).

(TIF)

S1 Table. Carbohydrate content over time for five different incident light intensities. For all experiments $n = 2$, except for incident light intensity of 250 and $750 \mu\text{mol m}^{-2} \text{s}^{-1}$ with $n = 1$. Values in brackets represent the standard deviation from the two biological duplicates.

(DOCX)

S2 Table. TAG concentration over time for five different incident light intensities. For all experiments $n = 2$, except for incident light intensity of 250 and $750 \mu\text{mol m}^{-2} \text{s}^{-1}$ with $n = 1$. Values in brackets represent the standard deviation from the two biological duplicates.

(DOCX)

S3 Table. EPA content in the total lipid fraction over time for five different incident light intensities. For all experiments $n = 2$, except for incident light intensity of 250 and 750 $\mu\text{mol m}^{-2} \text{s}^{-1}$ with $n = 1$. Values in brackets represent the standard deviation from the two biological duplicates.
(DOCX)

Author Contributions

Conceptualization: IR DM RW PL.

Formal analysis: IR.

Funding acquisition: RW.

Investigation: IR.

Methodology: IR DM RW PL.

Project administration: IR RW PL.

Resources: RW.

Supervision: DM RW PL.

Validation: IR PL.

Visualization: IR.

Writing – original draft: IR.

Writing – review & editing: IR DM RW PL.

References

1. Ruiz J, Olivieri G, de Vree J, Bosma R, Willems P, Reith JH, et al. Towards industrial products from microalgae. *Energy Environ Sci*. 2016; 9: 3036–3043.
2. Chauton MS, Reitan KI, Norsker NH, Tveterås R, Kleivdal HT. A techno-economic analysis of industrial production of marine microalgae as a source of EPA and DHA-rich raw material for aquafeed: Research challenges and possibilities. *Aquaculture*. 2015; 436: 95–103.
3. Norsker N-H, Barbosa MJ, Vermuë MH, Wijffels RH. Microalgal production—A close look at the economics. *Biotechnol Adv*. 2011; 29: 24–27. <https://doi.org/10.1016/j.biotechadv.2010.08.005> PMID: 20728528
4. Breuer G, Lamers PP, Martens DE, Draaisma RB, Wijffels RH. The impact of nitrogen starvation on the dynamics of triacylglycerol accumulation in nine microalgae strains. *Bioresour Technol*. 2012; 124: 217–26. <https://doi.org/10.1016/j.biortech.2012.08.003> PMID: 22995162
5. Mulders KJM, Janssen JH, Martens DE, Wijffels RH, Lamers PP. Effect of biomass concentration on secondary carotenoids and triacylglycerol (TAG) accumulation in nitrogen-depleted *Chlorella zofingiensis*. *Algal Res*. 2014; 6: 8–16.
6. Santos AM, Wijffels RH, Lamers PP. pH-upshock yields more lipids in nitrogen-starved *Neochloris oleoabundans*. *Bioresour Technol*. 2014; 152: 299–306. <https://doi.org/10.1016/j.biortech.2013.10.079> PMID: 24296123
7. Benvenuti G, Lamers PP, Breuer G, Bosma R, Cerar A, Wijffels RH, et al. Microalgal TAG production strategies: why batch beats repeated-batch. *Biotechnol Biofuels*. 2016; 9: 64. <https://doi.org/10.1186/s13068-016-0475-4> PMID: 26985237
8. Bitaubé Pérez E, Caro Pina I, Pérez Rodríguez L. Kinetic model for growth of *Phaeodactylum tricornutum* in intensive culture photobioreactor. *Biochem Eng J*. 2008; 40: 520–525.
9. Breuer G, Lamers PP, Martens DE, Draaisma RB, Wijffels RH. Effect of light intensity, pH, and temperature on triacylglycerol (TAG) accumulation induced by nitrogen starvation in *Scenedesmus obliquus*. *Bioresour Technol*. 2013; 143: 1–9. <https://doi.org/10.1016/j.biortech.2013.05.105> PMID: 23774290

10. Molina-Grima E, Pérez JAS, Camacho FG, Fernández FGA, Sevilla JMF, Sanz FV. Effect of dilution rate on eicosapentaenoic acid productivity of *Phaeodactylum tricornutum* utex 640 in outdoor chemostat culture. *Biotechnol Appl Biochem*. Kluwer Academic Publishers; 1994; 16: 1035–1040.
11. Reis A, Gouveia L, Veloso V, Fernandes HL, Empis JA, Novais JM. Eicosapentaenoic acid-rich biomass production by the microalga *Phaeodactylum tricornutum* in a continuous-flow reactor. *Bioresour Technol*. 1996; 55: 83–88.
12. Yongmanitchal W, Ward OP. Growth and eicosapentaenoic acid production by *Phaeodactylum tricornutum* in batch and continuous culture systems. *J Am Oil Chem Soc*. Springer-Verlag; 1992; 69: 584–590.
13. Geider J. R, Osborne BA, Raven JA. Light Dependence of Growth and Photosynthesis in *Phaeodactylum Tricornutum* (bacillariophyceae). *J Phycol*. 1985; 21: 609–619.
14. Ritchie RJ. Fitting light saturation curves measured using modulated fluorometry. *Photosynth Res*. 2008; 96: 201–215. <https://doi.org/10.1007/s11120-008-9300-7> PMID: 18415696
15. Wiegman S, Barranguet C, Spijkerman E, Kraak MHS, Admiraal W. The role of ultraviolet-adaptation of a marine diatom in photoenhanced toxicity of acridine. *Environ Toxicol Chem*. 2003; 22: 591–598. PMID: 12627647
16. Postma PR, Miron TL, Olivieri G, Barbosa MJ, Wijffels RH, Eppink MHM. Mild disintegration of the green microalgae *Chlorella vulgaris* using bead milling. *Bioresour Technol*. 2015; 184: 297–304. <https://doi.org/10.1016/j.biortech.2014.09.033> PMID: 25280602
17. Lamers PP, van de Laak CCW, Kaasenbrood PS, Lorier J, Janssen M, De Vos RCH, et al. Carotenoid and fatty acid metabolism in light-stressed *Dunaliella salina*. *Biotechnol Bioeng*. 2010; 106: 638–48. <https://doi.org/10.1002/bit.22725> PMID: 20229508
18. de Mooij T, Janssen M, Cerezo-Chinarro O, Mussgnug JH, Kruse O, Ballottari M, et al. Antenna size reduction as a strategy to increase biomass productivity: a great potential not yet realized. *J Appl Phycol*. 2014; 27: 1063–1077.
19. Benvenuti G, Bosma R, Cuaresma M, Janssen M, Barbosa MJ, Wijffels RH. Selecting microalgae with high lipid productivity and photosynthetic activity under nitrogen starvation. *J Appl Phycol*. 2014; 27: 1425–1431.
20. Benvenuti G, Bosma R, Ji F, Lamers P, Barbosa MJ, Wijffels RH. Batch and semi-continuous microalgal TAG production in lab-scale and outdoor photobioreactors. *J Appl Phycol*. 2016; 28: 3167–3177. <https://doi.org/10.1007/s10811-016-0897-1> PMID: 28035172
21. Breuer G, Lamers PP, Janssen M, Wijffels RH, Martens DE. Opportunities to improve the areal oil productivity of microalgae. *Bioresour Technol*. 2015; 186: 294–302. <https://doi.org/10.1016/j.biortech.2015.03.085> PMID: 25836038
22. Dillschneider R, Steinweg C, Rosello-Sastre R, Posten C. Biofuels from microalgae: Photoconversion efficiency during lipid accumulation. *Bioresour Technol*. 2013; 142: 647–654. <https://doi.org/10.1016/j.biortech.2013.05.088> PMID: 23777817
23. Parkhill J-P, Maillet G, Cullen JJ. Fluorescence-based maximal quantum yield for PSII as a diagnostic of nutrient stress. *J Phycol*. 2001; 37: 517–529.
24. Zhao Y, Wang Y, Quigg A. Comparison of population growth and photosynthetic apparatus changes in response to different nutrient status in a diatom and a coccolithophore. Wetherbee R, editor. *J Phycol*. 2015; 51: 872–884. <https://doi.org/10.1111/jpy.12327> PMID: 26986884
25. Acién Fernández FG, Hall DO, Cañizares Guerrero E, Krishna Rao K, Molina Grima E. Outdoor production of *Phaeodactylum tricornutum* biomass in a helical reactor. *J Biotechnol*. 2003; 103: 137–152. [http://dx.doi.org/10.1016/S0168-1656\(03\)00101-9](http://dx.doi.org/10.1016/S0168-1656(03)00101-9) PMID: 12814873
26. Griffiths MJ, van Hille RP, Harrison STL. Lipid productivity, settling potential and fatty acid profile of 11 microalgal species grown under nitrogen replete and limited conditions. *J Appl Phycol*. 2011; 24: 989–1001.
27. Santos-Ballardo DU, Rendón-Unceta MDC, Rossi S, Vázquez-Gómez R, Hernández-Verdugo S, Valdez-Ortiz A. Effects of outdoor cultures on the growth and lipid production of *Phaeodactylum tricornutum* using closed photobioreactors. *World J Microbiol Biotechnol*. 2016; 32: 128. <https://doi.org/10.1007/s11274-016-2089-1> PMID: 27339309
28. Jallet D, Caballero MA, Gallina AA, Youngblood M, Peers G. Photosynthetic physiology and biomass partitioning in the model diatom *Phaeodactylum tricornutum* grown in a sinusoidal light regime. *Algal Res*. 2016; 18: 51–60.
29. Wagner H, Jakob T, Lavaud J, Wilhelm C. Photosystem II cycle activity and alternative electron transport in the diatom *Phaeodactylum tricornutum* under dynamic light conditions and nitrogen limitation. *Photosynth Res*. 2015; 128: 151–161. <https://doi.org/10.1007/s11120-015-0209-7> PMID: 26650230

30. Waring J, Klenell M, Bechtold U, Underwood GJC, Baker NR. Light-Induced Responses of Oxygen Photoreduction, Reactive Oxygen Species Production and Scavenging in Two Diatom Species. *J Phycol.* 2010; 46: 1206–1217.
31. Bailleul B, Rogato A, de Martino A, Coesel S, Cardol P, Bowler C, et al. An atypical member of the light-harvesting complex stress-related protein family modulates diatom responses to light. *Proc Natl Acad Sci.* 2010; 107: 18214–18219. <https://doi.org/10.1073/pnas.1007703107> PMID: 20921421
32. Chauton MS, Winge P, Brembu T, Vadstein O, Bones AM. Gene regulation of carbon fixation, storage, and utilization in the diatom *Phaeodactylum tricornutum* acclimated to light/dark cycles. *Plant Physiol.* 2013; 161: 1034–48. <https://doi.org/10.1104/pp.112.206177> PMID: 23209127
33. Lavaud J, Rousseau B, van Gorkom HJ, Etienne A-L. Influence of the Diadinoxanthin Pool Size on Photoprotection in the Marine Planktonic Diatom *Phaeodactylum tricornutum*. *Plant Physiol.* 2002; 129: 1398–1406. <https://doi.org/10.1104/pp.002014> PMID: 12114593
34. Richmond A. *Handbook of Microalgal Culture: Biotechnology and Applied Phycology.* John Wiley & Sons; 2008.
35. Olivieri G, Salatino P, Marzocchella A. Advances in photobioreactors for intensive microalgal production: configurations, operating strategies and applications. *J Chem Technol Biotechnol.* 2014; 89: 178–195.
36. Tredici MR, Zittelli GC. Efficiency of sunlight utilization: Tubular versus flat photobioreactors. *Biotechnol Bioeng.* 1998; 57: 187–197. PMID: 10099193
37. de Mooij T, Schediwy K, Wijffels RH, Janssen M. Modeling the competition between antenna size mutant and wild type microalgae in outdoor mass culture. *J Biotechnol.* 2016; 240: 1–13. <https://doi.org/10.1016/j.jbiotec.2016.10.009> PMID: 27746308
38. Perin G, Bellan A, Segalla A, Meneghesso A, Alboresi A, Morosinotto T. Generation of random mutants to improve light-use efficiency of *Nannochloropsis gaditana* cultures for biofuel production. *Biotechnol Biofuels.* 2015; 8.
39. Breuer G, de Jaeger L, Artus VPG, Martens DE, Springer J, Draaisma RB, et al. Superior triacylglycerol (TAG) accumulation in starchless mutants of *Scenedesmus obliquus*: (II) evaluation of TAG yield and productivity in controlled photobioreactors. *Biotechnol Biofuels.* 2014; 7: 70. <https://doi.org/10.1186/1754-6834-7-70> PMID: 24883102
40. Münkler R, Schmid-Staiger U, Werner A, Hirth T. Optimization of outdoor cultivation in flat panel airlift reactors for lipid production by *Chlorella vulgaris*. *Biotechnol Bioeng.* 2013; 110: 2882–2893. <https://doi.org/10.1002/bit.24948> PMID: 23616347
41. Cabanelas ITD, van der Zwart M, Kleinegris DMM, Wijffels RH, Barbosa MJ. Sorting cells of the microalga *Chlorococcum littorale* with increased triacylglycerol productivity. *Biotechnol Biofuels.* 2016; 9: 183. <https://doi.org/10.1186/s13068-016-0595-x> PMID: 27582875
42. Van Vooren G, Le Grand F, Legrand J, Cuiñé S, Peltier G, Pruvost J. Investigation of fatty acids accumulation in *Nannochloropsis oculata* for biodiesel application. *Bioresour Technol.* 2012; 124: 421–432. <https://doi.org/10.1016/j.biortech.2012.08.009> PMID: 23018107
43. Abida H, Dolch L-J, Meř C, Villanova V, Conte M, Block MA, et al. Membrane Glycerolipid Remodeling Triggered by Nitrogen and Phosphorus Starvation in *Phaeodactylum tricornutum*. *Plant Physiol.* 2015; 167: 118–136. <https://doi.org/10.1104/pp.114.252395> PMID: 25489020
44. Alonso DL, Belarbi E-H, Fernández-Sevilla JM, Rodríguez-Ruiz J, Grima EM. Acyl lipid composition variation related to culture age and nitrogen concentration in continuous culture of the microalga *Phaeodactylum tricornutum*. *Phytochemistry.* 2000; 54: 461–471. [http://dx.doi.org/10.1016/S0031-9422\(00\)00084-4](http://dx.doi.org/10.1016/S0031-9422(00)00084-4) PMID: 10939349
45. Gatenby CM, Orcutt DM, Kreeger DA, Parker BC, Jones VA, Neves RJ. Biochemical composition of three algal species proposed as food for captive freshwater mussels. *J Appl Phycol.* 2003; 15: 1–11.
46. Jia J, Han D, Gerken HG, Li Y, Sommerfeld M, Hu Q, et al. Molecular mechanisms for photosynthetic carbon partitioning into storage neutral lipids in *Nannochloropsis oceanica* under nitrogen-depletion conditions. *Algal Res.* 2015; 7: 66–77.
47. Guihéneuf F, Stengel DB. LC-PUFA-enriched oil production by microalgae: accumulation of lipid and triacylglycerols containing n-3 LC-PUFA is triggered by nitrogen limitation and inorganic carbon availability in the marine haptophyte *Pavlova lutheri*. *Mar Drugs.* 2013; 11: 4246–66. <https://doi.org/10.3390/md11114246> PMID: 24177672
48. Juergens MT, Disbrow B, Shachar-Hill Y. The Relationship of Triacylglycerol and Starch Accumulation to Carbon and Energy Flows during Nutrient Deprivation in *Chlamydomonas reinhardtii*. *Plant Physiol.* 2016; 171: 2445–2457. <https://doi.org/10.1104/pp.16.00761> PMID: 27325664
49. Klein U. Intracellular Carbon Partitioning in *Chlamydomonas reinhardtii*. *Plant Physiol.* 1987; 85: 892–897. PMID: 16665826

50. Mühlroth A, Li K, Røkke G, Winge P, Olsen Y, Hohmann-Marriott MF, et al. Pathways of lipid metabolism in marine algae, co-expression network, bottlenecks and candidate genes for enhanced production of EPA and DHA in species of Chromista. *Mar Drugs*. 2013; 11: 4662–97. <https://doi.org/10.3390/md11114662> PMID: 24284429
51. Tonon T, Harvey D, Larson TR, Graham IA. Long chain polyunsaturated fatty acid production and partitioning to triacylglycerols in four microalgae. *Phytochemistry*. 2002; 61: 15–24. PMID: 12165297
52. Adarme-Vega TC, Lim DKY, Timmins M, Vernen F, Li Y, Schenk PM. Microalgal biofactories: a promising approach towards sustainable omega-3 fatty acid production. *Microb Cell Factories*. 2012; 11: 96.