

## Mitochondrial DNA Part B Resources

ISSN: (Print) 2380-2359 (Online) Journal homepage: <http://www.tandfonline.com/loi/tmdn20>

# The complete mitochondrial genome of the copepod *Calanus glacialis*

Marvin Choquet, Homère J. Alves Monteiro, Johan Bengtsson-Palme & Galice Hoarau

To cite this article: Marvin Choquet, Homère J. Alves Monteiro, Johan Bengtsson-Palme & Galice Hoarau (2017) The complete mitochondrial genome of the copepod *Calanus glacialis*, *Mitochondrial DNA Part B*, 2:2, 506-507, DOI: [10.1080/23802359.2017.1361357](https://doi.org/10.1080/23802359.2017.1361357)

To link to this article: <https://doi.org/10.1080/23802359.2017.1361357>



© 2017 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 06 Aug 2017.



Submit your article to this journal [↗](#)



Article views: 335



View related articles [↗](#)




View Crossmark data [↗](#)



Citing articles: 1 View citing articles [↗](#)

## The complete mitochondrial genome of the copepod *Calanus glacialis*

Marvin Choquet<sup>a</sup> , Homère J. Alves Monteiro<sup>a</sup>, Johan Bengtsson-Palme<sup>b</sup> and Galice Hoarau<sup>a</sup>

<sup>a</sup>Faculty of Biosciences and Aquaculture, Nord University, Bodø, Norway; <sup>b</sup>Department of Infectious Diseases, Institute of Biomedicine, The Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden

### ABSTRACT

*Calanus glacialis*, a marine planktonic copepod, is a keystone species in the Arctic Ocean. In this study, we shotgun sequenced the total DNA of one *C. glacialis* individual, using the NextSeq<sup>®</sup> Illumina platform, in order to determine its mitochondrial genome sequence. We successfully assembled and annotated this 20,674 bp long sequence, which included 13 protein-coding genes, 2 rRNA genes and 22 tRNA genes. Common gene-coding regions of 19 other species were used to reconstruct a phylogenetic tree, using mitogenomes of the phylogenetically closest copepods available. The new resource described here constitutes a tool of interest for better understanding the structure and dynamics of *C. glacialis* populations.

### ARTICLE HISTORY

Received 4 July 2017  
Accepted 26 July 2017

### KEYWORDS

*Calanus glacialis*; copepod; mitochondrion; mitogenome; zooplankton

The genus *Calanus* consists of 26 distinct marine copepod species (WoRMS Editorial Board, 2017), present in every ocean in the world as part of the zooplankton. Despite their ecological importance, only two mitochondrial genomes have been reported within the *Calanus* genus: *C. sinicus* (Minxiao et al. 2011) and *C. hyperboreus* (Kim et al. 2013). *C. glacialis* is one of the key species of the Arctic Ocean, as the crucial link between primary production and higher trophic levels such as fishes, invertebrates, marine mammals and birds (Falk-Petersen et al. 1990).


In this study, we report the complete sequence of the mitochondrial genome of *C. glacialis*. We selected one *C. glacialis* individual from Sørfolda (Norwegian coast: 67°30 N, 15°28 E), which we identified as such using a set of nuclear molecular markers (Smolina et al. 2014). Total DNA was extracted using the E.Z.N.A.<sup>®</sup> Insect DNA Kit and was shotgun sequenced on the NextSeq<sup>®</sup> Illumina platform. Given the amount of DNA recovered from a single individual, everything was used for the library construction. *De novo* assembly of the filtered reads was performed using Ray version 2.3.1 (Boisvert et al. 2010) with a k-mer length of 31. Contigs that matched the mitochondrial genomes of *C. hyperboreus* or *C. sinicus* in a BLAST (Altschul et al. 1997) search (e-value cutoff  $10^{-10}$ ) were extracted. To potentially further merge these contigs, they were used as seeds in a Peacat search (<http://microbiology.se/sw/petkit>) against all assembled contigs (Bengtsson-Palme et al. 2014). The resulting consensus sequences were tested for circularity using Pemap

(<http://microbiology.se/software/petkit/>), but no evidence of circularity was found.

We mapped the annotated mitochondrial genomes of *C. hyperboreus* and *C. sinicus* to the longest contig obtained from the assembly and were able to identify all expected mitochondrial genes.

The mitochondrial sequence of *C. glacialis* is 20,674 bp long and contains 13 protein-coding genes (total of 3458 amino acids), 2 rRNA genes, 22 tRNA genes and 1 putative control region. The sequence is composed of 31.7% base A, 28.8% base T, 19.6% base C and 19.9% base G. Ribosomal 12S and 16S RNA are 656 bp and 1138 bp long, respectively. The sequence has been deposited in GenBank under the accession number MF422146.

A phylogenetic analysis was performed using all coding genes for 18 species of Crustaceans (including 10 copepods species) and two hexapods as out-groups (*Japyx solifugus* and *Campodea fragilis*). The phylogenetic tree was reconstructed with a maximum likelihood method using PHYML (Guindon and Gascuel 2003) (GRT+I+G model, 1000 bootstraps) (Figure 1). All copepods formed a monophyletic group and *C. glacialis* clustered with the other two *Calanus* mitogenome (100% support). Given the ecological importance of *C. glacialis* within the Arctic ecosystem, the newly determined mitogenome will be useful for investigating the history of *C. glacialis* populations and their spatiotemporal variability.



**Figure 1.** Phylogenetic tree of *C. glacialis* and 19 other species, with *Japyx solifugus* and *Campodea fragilis* as out-groups. ML bootstrap values (1000 replications) are indicated in front of each node.

## Acknowledgements

We would like to acknowledge Morten Krogstad for sampling, Anusha Dhanasiri and Martina Kopp for technical support in the lab.

## Disclosure statement

The authors have no conflict of interest to declare.

## Funding

This project was supported by the Norwegian Research Council (216578) and Nord University.

## ORCID

Marvin Choquet  <http://orcid.org/0000-0001-6719-2332>

## References

- Altschul S, Madden T, Schäffer A, Zhang J, Zhang Z, Miller W, Lipman D. 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res.* 25:3389–3402.
- Bengtsson-Palme J, Boulund F, Fick J, Kristiansson E, Larsson DJ. 2014. Shotgun metagenomics reveals a wide array of antibiotic resistance genes and mobile elements in a polluted lake in India. *Front Microbiol.* 5:648.
- Boisvert S, Laviolette F, Corbeil J. 2010. Ray: simultaneous assembly of reads from a mix of high-throughput sequencing technologies. *J Comput Biol.* 17:1519–1533.
- Falk-Petersen S, Hopkins C, Sargent J. 1990. Trophic relationships in the pelagic, Arctic food web. Trophic relationships in the marine environment. Aberdeen: Aberdeen University Press; p. 315–333.
- Guindon S, Gascuel O. 2003. A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. *Syst Biol.* 52:696–704.
- Kim S, Lim BJ, Min GS, Choi HG. 2013. The complete mitochondrial genome of Arctic *Calanus hyperboreus* (Copepoda, Calanoida) reveals characteristic patterns in calanoid mitochondrial genome. *Gene.* 520:64–72.
- Minxiao W, Song S, Chaolun L, Xin S. 2011. Distinctive mitochondrial genome of Calanoid copepod *Calanus sinicus* with multiple large non-coding regions and reshuffled gene order: useful molecular markers for phylogenetic and population studies. *BMC Genomics.* 12:73.
- Smolina I, Kollias S, Poortvliet M, Nielsen TG, Lindeque P, Castellani C, Moller EF, Blanco-Bercial L, Hoarau G. 2014. Genome- and transcriptome-assisted development of nuclear insertion/deletion markers for *Calanus* species (Copepoda: Calanoida) identification. *Mol Ecol Resour.* 14:1072–1079.
- WoRMS Editorial Board. 2017. World Register of Marine Species. [accessed 2017 Jul 31]. Available from <http://www.marinespecies.org> at VLIZ.