

# 1 Management routines influencing piglet survival in loose-housed sow 2 herds

3 Ellen Marie Rosvold\*‡, Camilla Kielland\*, Marko Ocepek\*, Tore Framstad\*, Bente  
4 Fredriksen\*\*, Ina Andersen-Ranberg†, Geir Næss‡, Inger Lise Andersen\*

5 \* Norwegian University of Life Sciences, Department of Animal and Aquacultural Sciences, PO Box 5003, 1432  
6 Ås, Norway

7 \*\* Norwegian Meat and Poultry Research Centre (Animalia), PO Box 396, 0513 Oslo, Norway

8 † Topigs Norsvin, PO Box 504, 2304 Hamar, Norway

9 ‡ Nord University, PO Box 2501, 7729 Steinkjer, Norway

10

11 Corresponding author: Ellen Marie Rosvold. E-mail: [ellen.m.rosvold@nord.no](mailto:ellen.m.rosvold@nord.no)

12

## 13 Abstract

14 Piglet mortality is still a significant welfare and ethical matter in pig production, as well as an  
15 economical challenge for the farmer. Most of the mortality occurs early after farrowing, and previous  
16 studies have shown that the farm`s management routines, especially around farrowing, are important  
17 factors to reduce it. When sows are loose-housed at farrowing and in the following lactation period, it  
18 puts higher demands on management input from the farmer to keep piglet mortality low. The objective  
19 of this study was to assess the importance of different management routines around the time of  
20 farrowing, and other farm qualities for piglet survival in loose-housed herds. To study risk factors for  
21 herd piglet mortality, a cross-sectional field survey was carried out in Norway in the year 2013, and  
22 included 52 commercial herds with hybrid LY sows (Norwegian Landrace x Swedish Yorkshire). The  
23 farms were visited once, and the farmers answered a questionnaire about their management practices.  
24 The outcome was the average herd pre-weaning mortality in the years of 2012-2013. To include as many  
25 management factors as possible into the multivariable linear regression model, we generated a new  
26 variable based on 4 management routines: 3 routines at farrowing (presence at 80-100% of the  
27 farrowings, drying newborn piglets, and practice split suckling), and one concerning farmer`s contact  
28 with the sows. This variable was called “Management type” (M), and were divided into 4 categories  
29 with increasing effort; M1 herds without any of the 4 mentioned routines, M2 had contact with sows >2  
30 times per day, M3 performed the 3 routines at farrowing, and M4 combined the high sow contact and  
31 the 3 routines. The predicted values of mean herd piglet mortality for M1, M2, M3 and M4 were 20.1%,  
32 17.0%, 16.2% and 13.3% respectively. The farmer`s increased management effort was associated with  
33 lower piglet mortality (P<0.05). The farmer`s effort at critical times together with systematic and  
34 important routines, and having frequent contact with the sows, makes a huge difference for piglet  
35 survival. The farmers are credited for this work by having lower piglet mortality as a result.

36 Keywords: Piglet mortality, Management, Farrowing, Loose housed sows, Field survey

37 Highlights:

- 38 • Piglet mortality is multifactorial as many factor together leads to reduction.
- 39 • Presence at farrowing, drying newborn piglets and split suckling are routines associated with  
40 lower piglet mortality.
- 41 • Frequent contact with sows is associated with lower piglet mortality.

## 42 1. Introduction

43 High piglet mortality is still an ethical and economical challenge in pig production. As much as 50-80%  
44 of the piglet mortality is caused by crushing and starvation (English and Morrison, 1984; Dyck and  
45 Swiestra, 1987; Marchant et al., 2000), and this mainly occurs within the first two or three days after  
46 farrowing (Dyck and Swiestra, 1987; Cronin et al., 2000; Marchant et al., 2000; Andersen et al., 2005;  
47 Westin et al., 2015). A field survey from Norwegian farms reported that the mortality of live born piglets  
48 ranged from 5 to 24%, and management was suggested to be an important factor (Andersen et al., 2007).  
49 In a review by Kirkden et al. (2013), it was concluded that piglet mortality can be reduced by a range of  
50 management routines, especially around farrowing. One important procedure is the supervision of  
51 farrowing by trained staff, and also attending sows a couple of days postpartum, which can reduce piglet  
52 mortality (Holyoake et al., 1995; White et al., 1996). While being present, the farmer could more easily  
53 detect animals that are in need of assistance, and for instance save piglets from near-crushing incidents.  
54 Some management routines, such as drying and placing piglets under a heat source immediately after  
55 birth can all reduce mortality (White et al., 1996; Christison et al., 1997; Andersen et al., 2009).

56 Rearing piglets in loose housing systems demands sows with good maternal abilities (Wechsler and  
57 Hegglin, 1997; Andersen et al., 2005; Johnson et al., 2007). But as litter size has increased over the  
58 years, and sows have a limited biological capacity related to number of functional teats and maternal  
59 investment, these larger litters demands more management input from the farmer to keep piglet mortality  
60 low (English, 1993). For instance, one experiment demonstrated that litters with more than 12 piglets,  
61 on average one piglet had no teat during a nursing bout in the first couple of days after farrowing, a  
62 factor that could lead to starvation (Rosvold, 2006). Management options when litters are large are for  
63 instance cross-fostering, split-suckling and nurse sow systems (Baxter et al., 2013). A good relationship  
64 between humans and animals is another factor important for welfare, health and production. For  
65 instance, in a study by Andersen et al. (2006), sows with low confidence that were positively handled  
66 the last two weeks prior farrowing, had increased confidence score, shorter farrowing duration, and also  
67 tended to give birth to fewer mummified or immature stillborn piglets compared to control sows. Ravel  
68 et al. (1996) found in their farm survey that the stockperson factors constitutes 26-27% of the variance  
69 in pre-weaning mortality.

70 The pre-weaning mortality of live born piglets on herd level is frequently used when evaluating a farms`  
71 production result from one year to another. This is a number that most pig farmers are familiar with, and  
72 refers to high survival rate among the piglets. A field survey was carried out to obtain information about  
73 pig farms, their management, especially around farrowing, and their production results. In this study we  
74 will investigate pre-weaning mortality of live born piglets (%) on herd level (HPM). All farm  
75 information are factors on herd level. The objective of this survey was to identify and assess the  
76 importance of systematic management routines around the time of farrowing for piglet survival in loose-  
77 housed sow herds.

## 78 2. Materials and methods

### 79 2.1 Farm selection and study population

80 This field survey was planned to include 60 commercial sow herds, with 20 farms representing each out  
81 of three major pig production regions in Norway (East, West and Middle). Inclusion criteria were breed  
82 (LY; sows of Norwegian Landrace x Swedish Yorkshire) and a consistent practice of keeping the sows  
83 loose during farrowing. The farms also had to keep regular recordings of production results to Ingris  
84 (The National Efficiency Control Database, administrated by Animalia (Norwegian Meat and Poultry  
85 Research Centre) and Norsvin (Norwegian Pig Breeding Association)). Information from Ingris  
86 concerning the numbers of litters born per year in each herd, gave us a possibility to select herds with a  
87 variety in size. Farmers were initially invited to participate in the study by letter in February 2013,  
88 followed by phone call for a second invitation. Fifty-two herds that complied with the inclusion criteria  
89 accepted to participate in the field survey. Before the onset of the study, the selected farmers were well  
90 prepared and we explained the importance of assessing the causes of death while they were present  
91 during farrowing.

### 92 2.2 Collecting of farm data

93 During spring and summer 2013, one of two trained researchers visited the farms once. The visit was  
94 carried out during the lactation period, with a compulsory tour in the pig house. Farmers answered  
95 questions about management practice and routines before, during and immediately after farrowing.  
96 Questions, categories and responses are presented in the results, including Table 2-4. The farms`  
97 production results for 2012 and 2013 were extracted from Ingris, and are presented in Table 5 and Figure  
98 1. In 2013, there were 281 commercial herds in Ingris with registrations on LY sows and piglets, and  
99 the herds in the field survey (52) constitute 18.5% of these herds.

### 100 2.3 Data analysis

101 Data handling and statistical analyses were performed in Stata (Stata SE/11, Stata Corp., College Station,  
102 TX, USA) and SPSS (IMB SPSS Statistics Version 22, SPSS Inc. Chicago, USA).

103 For multiple choice questions distribution of the answers were calculated. Questions with answers given  
104 as continuous variables were reported by mean, standard error (S.E.) and range. The outcome were the  
105 average HPM in the years 2012 and 2013, and the average of two years was chosen to even out potential  
106 bad or good years. A multivariable linear regression model was used to evaluate which and how  
107 explanatory herd level factors were associated with HPM.

108 Descriptive statistics to assess the assumptions were made using a multivariable regression model, where  
109 evaluated using various techniques. Linearity between the continuous outcome and dichotomous  
110 variables was investigated with graphs using a “logit” function in Stata, creating a lowess line between  
111 the two variables. In addition, probability plots, best linear fit, and R<sup>2</sup> were used to explore how  
112 continuous explanatory variables explained the variation in HPM.

113 Several management factors were recorded during the farm visit, i.e. split suckling, drying piglets (for  
114 more details see Table 2-4). The challenge regarding the various managements registered, was that some  
115 farms had similar management routines, but several farms had their own unique routines. The regression  
116 analysis made many 2x2 tables, and we needed enough numbers in each box to give sensible estimates.  
117 Therefore, we had to cluster the farms into groups with similar management systems. After identifying  
118 management variables from the univariate analyses during the model building process, a new variable  
119 were generated using the Stata command “egen concat”, concatenate routines, categorizing farms based  
120 on four routines (concatenate commands are normally used to join two or more text strings into one  
121 string). This variable was called “Management type” (M), and was based on four management routines.  
122 Three of the management routines were conducted at farrowing (being present at 80-100% of the  
123 farrowings, drying and massaging newborn piglets, and performing split suckling), and the fourth  
124 routine was contact with the sows >2 times per day (Table 1). Contact was defined as touching, talking  
125 to and/or being present near the sow in the farrowing pen. This new variable had four categories; M1  
126 herds did not perform any of the four management routines displayed in Table 1. These herds had all  
127 unique combinations of the management routines from Table 2-4, and could not be grouped. M2 herds  
128 had contact with sows >2 times per day, M3 herds performed the three mentioned routines at farrowing,  
129 and M4 herds combined contact and the routines. Management types were ordinal categories, and M1  
130 meant low management effort, with increasing effort by M2, M3 and M4. The latter therefore meant  
131 high management effort.

132 When building the final model, a forward stepwise technique was used, exploring variables with a P-  
133 value <0.20 from the univariable analysis, according to the method described by Dohoo et al. (2009).  
134 Distortion and confounding could be observed as each variable was included. Biologically plausible  
135 first-order interactions of the predictor variables were evaluated and included if the interaction was  
136 significant. Normal probability plots was evaluated, and Shapiro-Wilks statistic used to test for normal  
137 distribution of these residuals. When exploring influencing values and leverage points, no values were

138 deleted from the analysis due to high influence if the value was within reasonable boundaries. If  
139 variables were highly correlated with each other ( $|\rho| > 0.8$ ) (Dohoo et al., 2009), only one of these  
140 variables was included. The model was tested for heteroscedasticity using Breusch-Pagan / Cook-  
141 Weisberg test. The varying inflating factors also explored to evaluate the final model. For each variable  
142 included, the model was evaluated and the best model chosen. The best model was the model with the  
143 lowest mean square error. In all analyses, statistical significance was considered with a P-value  $< 0.05$   
144 and borderline significance with a P-value  $< 0.10$ .

### 145 3. Results

#### 146 3.1. Descriptive statistics of farms

147 Of the 52 farmers, 30 (57.7%) were men and 9 (17.3%) women. Twenty-three (44.2%) of the farms had  
148 more than one person working, and 13 (25.3%) farms had both men and women involved in the daily  
149 routines. Twenty (38.5%) farms were situated in the East, 13 (25.0%) in the West and 19 (36.5%) in the  
150 Middle of Norway. When dividing farmers in age groups, 5 (9.6%) were between 20-30 years old, 34  
151 (65.4%) were between 30-50 years old and 13 (25.0%) were more than 50 years old. Three (5.8%)  
152 farmers had no education above primary school, 40 (76.9%) had finished high school, and 9 (17.3%)  
153 had been to university. When asked about pig farming experience, 4 (7.7%) farmers had less than 5  
154 years of experience, 6 (11.5%) had 5-10 years, 16 (30.8%) had 10-20 years, and 26 (50.0%) had more  
155 than 20 years of experience.

156 The farms had different systems of batch farrowing. Four (7.7%) farmers had the system of farrowing  
157 every 2.5-3<sup>rd</sup> week, 8 (15.4%) every 5.5 weeks, 34 (65.4%) every 7<sup>th</sup> week, 3 (5.8%) every 11<sup>th</sup> week  
158 and 3 (5.7%) had farrowing every 22-26<sup>th</sup> week. Mean number of sows in one farrowing batch was  $26.3$   
159  $\pm 1.9$  (10-65), and the sows spent on average  $9.5 \pm 0.7$  (2-21) days in the farrowing pen before farrowing.  
160 The number of litters born at the farms (mean  $\pm$  S.E.) during 2012 and 2013 was  $178.6 \pm 13.0$  (ranging  
161 from 57.5-498.0).

162 At the farm visit, type and amount of nest-building material, as well as timing of distribution to sows  
163 prior to farrowing were reported. Long stemmed straw was given by 24 (46.2%) farmers, 4 (7.7%)  
164 farmers gave chopped straw, 19 (36.5%) gave wood-shavings, 4 (7.7%) gave long-stemmed straw and  
165 wood-shavings in combination, and only one (1.9%) farmer gave hay as nest-building material. The  
166 mean amount of nest-building material given was  $2.6 \pm 0.5$  kg (ranging from 0.1-20.0), distributed on  
167 average  $28.7 \pm 2.8$  hours (ranging from 3.0-96.0) before farrowing. Also, feeding of roughage during  
168 gestation and lactation were reported. Five (9.6%) farmers did not provide roughage (hay, silage and  
169 straw) at all to their pregnant sows, 17 (32.7%) farmers fed  $< 200$  g roughage daily, 22 (42.3%) fed 200-  
170 500 g, and 8 (15.4%) farmers fed their pregnant sows roughage ad libitum. When the sows were in  
171 lactation, 15 (28.8%) farmers did not provide roughage at all, 18 (34.6%) fed  $< 200$  g roughage daily,  
172 16 (30.8%) fed 200-500 g, and 3 (5.8%) farmers fed their lactating sows roughage ad libitum.

173 Characterizations of management and routines around farrowing can be found in Table 2. At farrowing,  
174 24 (46.2%) of the farmers moved newborn piglets both to the udder and to the creep area. There were  
175 also combinations of the routines dried/massaged followed by moving piglets to udder and/or creep area.  
176 Number of farmers that dried/massaged and moved piglets to the udder was 6 (11.5%), number of  
177 farmers who dried/massaged and moved piglets to creep area was also 6 (11.5%), and dried/massaged  
178 piglets and moved to both places was 16 (30.8%). Management routines during the first 48 hours had  
179 also some combinations identified. Twelve (23.1%) farmers moved piglets both to the udder and to the  
180 creep area. Only one farmer (1.9%) massaged and laid the piglets to the creep area, and another two  
181 farmers (3.8%) massaged and moved to both places. All farmers conducted cross-fostering, but to what  
182 extent it was done and which criteria that were used varied considerably (Table 3).

183 The farmers were asked if good relationship with the sows was important on a scale from 1 (not  
184 important) to 10 (very important), and 32 farmers (61.5 %) scored it to 10. How often farmers had  
185 contact with their sows in general, and the farmer`s opinion about ease of handling were reported and  
186 shown in Table 4.

### 187 3.2 Factors associated with HPM

188 The results from the investigated 52 farms are presented in Table 5 and Figure 1, and demonstrates some  
189 similarities to the national averages in Ingris (Table 5). However, in the 52 survey farms, mean values  
190 of number of live born, stillborn and HPM were higher compared to Ingris.

191 All the factors concerning farm demographics, management and routines described in section 3.1 were  
192 explored in relation to HPM. Significant factors associated with HPM were batch system, number of  
193 sows per batch, management type as described in Table 1 (i.e. the routines of being present at 80-100%  
194 of the farrowings, drying and massaging newborn piglets, conduct split suckling, and having contact  
195 with the sows >2 times per day), and time of cross-fostering. Table 6 shows the details of these factors.

196 From the predicted model in Table 6, one can compare predicted HPM between farms with different  
197 size (number of sows in each batch) and management type. As the intercept were an average of baseline,  
198 a farm with system and management like the categories in baselines would have 20.1% as predicted  
199 HPM. Farms with higher management effort than M1 (baseline) would have a lower value of predicted  
200 HPM. The respective predicted HPM values of M2 (having contact with the sows >2 times/day), M3  
201 (having three management routines at farrowing; being present at 80-100% of the farrowings, drying  
202 and massaging newborn piglets, and performing split suckling) and M4 (combination of contact and the  
203 three farrowing routines), were 17.0%, 16.2% and 13.3% (Figure 2). Cross-fostering conducted at 13-  
204 24 hours after farrowing had predicted value of HPM of 20.1% (baseline). Having no systematic routine  
205 would make a higher predicted HPM, 24.2% (Figure 3).

### 206 3.3.1 *Diagnostics*

207 No heteroscedasticity was detected. Variation inflating factors was low both in total and at each variable  
208 included in the regression analysis. Normality plots of standardized residuals did not display potential  
209 outliers. No influencing points were identified.

## 210 4. Discussion

211 The main purpose of this field survey was to identify management factors that could be associated with  
212 low HPM, and were therefore important to give a higher piglet survival. We found that several  
213 management factors together lead to a reduction in HPM in commercial farms. Farmers with high  
214 management effort (M4: i.e. presence at 80-100% of the farrowings, drying and massaging newborn  
215 piglets, split suckling and contact with the sows > 2 per day) were credited for this work by having 6.8  
216 percentage-points lower HPM than the baseline herds (M1 versus M4). One of the reasons for several  
217 factors acting together were also that farmers that achieve good results appear to have a more systematic  
218 way of managing their farm and their routines, and a good system is important when many sows farrow  
219 in batches at the same time even though farmers may focus on slightly different factors. Systematic  
220 routines also become predictable routines for the animals themselves, and will most likely give positive  
221 effects on the human-animal relationship as well. We also found a high variation from the farm with the  
222 lowest losses to the farm with the highest, and this range was in accordance with a previous survey in  
223 Norway (Andersen et al., 2007).

224 As predicted, a high degree of presence during farrowing was one of the factors identified as important  
225 to reduce HPM. Other studies have shown that piglet mortality due to stillbirths, crushing by the sow,  
226 low viability and starvation were reduced when farrowing was attended (Holyoake et al., 1995; White  
227 et al., 1996). However, it is not only about being present, but also having systematic routines that are  
228 done while attending the farrowing. For instance, while present, the farmer could more easily detect  
229 sows that are in need of birth assistance, remove mucus from the nose and mouth, remove the placental  
230 envelopes around newborn piglets to prevent suffocation, dry the piglet and tie the umbilical cord  
231 (Holyoake et al., 1995; White et al., 1996). Also putting the piglets under a heat source or at the udder  
232 to suckle colostrum could be routines done while present at farrowing, as well as having the possibility  
233 to save piglets that are near crushed or savaged by their mother sow.

234 In our study, it was the combination of being present at 80-100% of the farrowings, drying newborn  
235 piglets, and practice split suckling in addition to being in contact with the sows more than two times per  
236 day resulted in the lowest HPM. However, in order to decide whether all this extra effort pays off for  
237 the farmer, we would have to calculate the benefit in terms of how many extra piglets are saved per hour  
238 extra effort made in the farm compared to the baseline herds. Although experiments on drying and  
239 placing the piglets under the heat lamp have resulted in a much higher piglet survival in controlled  
240 experiments conducted on one particular farm (e.g. Andersen et al., 2009), these data were difficult to



241 reproduce when studying a large number of farms differing in so many ways, i.e. stockmanship, feeding,  
242 management and physical environment. Furthermore, we were not able to control how many litters that  
243 actually were subjected to the specific routines that they claim to have. In our study, we had no  
244 knowledge of how many piglets that were dried or placed in the creep within a litter, as our data are on  
245 herd level, not on piglet or sow level. This is also why we decided to focus on a combination of factors  
246 that separately had been documented as successful in earlier studies, and the present data shows quite  
247 clearly that an increased number of routines in combination produces a steady decline in HPM. This is  
248 also an important message to give to the farmers that want to improve their production results.

249 Split suckling was also one of the routines in combination with others that resulted in lower HPM in this  
250 survey. This routine of having the larger piglets in a litter enclosed for approximately an hour so that the  
251 smaller piglets could have full access to the udder, should allow all the piglets access to colostrum, and  
252 therefore acquire passive immunity (Baxter et al., 2013). However, Donovan and Dritz (2000) found no  
253 effect of split suckling on mortality or serum immunoglobulin concentrations, but found a reduced  
254 heterogeneity of weight gain in larger litters ( $\geq 9$  piglets). Considering the large work load put on the  
255 farmer, this routine could better be viewed as a last strategy to save piglets in extremely large litters  
256 rather than a common everyday routine.

257 The frequency of the farmer`s contact with the sows had an effect on HPM. As suggested in the review  
258 by Kirkden et al. (2013), improved human-animal relationship, by reducing negative behaviours and  
259 increasing positive behaviours, could reduce the sow`s fear level. Positive contact or handling means  
260 that the animals` behavioural response is positive when being approached, touched and/or talked to by  
261 humans (Andersen et al., 2006). In our field study, contact could be neutral or positive as it was defined  
262 as touching, talking to or being in close proximity of the sow in the pen. By being more present in a  
263 predictive way, the sows habituates to the stockperson, may perhaps also develop some positive  
264 expectations to this presence, thereby reducing the level of fear. An increased confidence and calmness  
265 in the presence of humans may benefit the overall maternal behaviour of the sows (e.g. Lensink et al.,  
266 2009a; Lensink et al., 2009b; Marchant Forde, 2002) and most likely increase the ease of handling  
267 whenever this is necessary, for instance during birth assistance. By being more present, the farmer is  
268 also likely to discover problems with individual sows earlier and for instance act earlier in near crushing  
269 events or when sows are having birth problems.

270 Number of sows per batch had influence on HPM, as 20 or more sows in a batch were associated with  
271 lower HPM. This effect could be caused by higher professionalism, more systematically routines, and  
272 higher level of focus on what was happening in the pig house. Also, in the model, a batch system with  
273 frequent farrowing (2.5-3 weeks) tended to be associated with lower HPM.

274 All farmers conducted cross-fostering to a certain degree, and with variations in routines. As the number  
275 of newborn piglets in a litter often exceed the number of functional and accessible teats, cross-fostering



276 has been a method with aim to secure milk to the piglets. A recommended fostering strategy is to leave  
277 the weaker and smaller piglets with the mother and foster off the strong ones, but also to foster off  
278 weaker piglets to a newly farrowed sow who has a smaller litter. It is also recommended that fostering  
279 of piglets should occur as early as possible after farrowing, provided that they have an adequate intake  
280 of colostrum before taken from the mother (English, 1993). In an experiment by Heim et al. (2012),  
281 cross-fostering was performed within 24 hours after farrowing, and the results indicated that the adopted  
282 piglets had neither reduced survival rate nor growth. Another experiment, with piglets cross-fostered  
283 within 48 hours, concluded that cross-fostered piglets had lower survival rates than those not cross-  
284 fostered (Neal and Irvin, 1991). In our study, most of the farmers cross-fostered within the first 24 hours  
285 after farrowing, and within a 12 hours “time-window”, but when farmers had no systematic routine of  
286 this (i.e. conducted cross-fostering for a longer time period than 12 hours and with variation of timing  
287 after farrowing), it was associated with higher HPM.

## 288 **Conclusions**

289 Piglet mortality in commercial pig herds are affected by several management factors, and some of these  
290 may, if combined in a systematic way, increase piglet survival. Based on our results, we can recommend  
291 that farmers are more present during farrowings, have a systematic and frequent contact with the sows,  
292 dry newborn piglets whenever some need special attention and conduct split-suckling in large litters.

## 293 **Conflicts of interest**

294 The authors have no conflicts of interest to declare.

## 295 **Acknowledgements**

296 The Norwegian Research Council, Norsvin, Animalia and Nortura financed this project.

297

298   References

- 299   Andersen, I.L., Berg, S., Bøe, K.E., 2005. Crushing of piglets by the mother sow (*Sus scrofa*)—purely  
300   accidental or a poor mother? *Applied Animal Behaviour Science* 93, 229-243.
- 301   Andersen, I.L., Berg, S., Bøe, K.E., Edwards, S., 2006. Positive handling in late pregnancy and the  
302   consequences for maternal behaviour and production in sows. *Applied Animal Behaviour Science* 99,  
303   64-76.
- 304   Andersen, I.L., Haukvik, I.A., Bøe, K.E., 2009. Drying and warming immediately after birth may  
305   reduce piglet mortality in loose-housed sows. *Animal* 4, 592-597.
- 306   Andersen, I.L., Tajet, G.M., Haukvik, I.A., Kongsrud, S., Bøe, K.E., 2007. Relationship between  
307   postnatal piglet mortality, environmental factors and management around farrowing in herds with  
308   loose-housed, lactating sows. *Acta Agriculturae Scandinavica, Section A* 57, 38-45.
- 309   Baxter, E.M., Rutherford, K.M.D., D'Eath, R.B., Arnott, G., Turner, S.P., Sandoe, P., Moustsen, V.A.,  
310   Thorup, F., Edwards, S.A., Lawrence, A.B., 2013. The welfare implications of large litter size in the  
311   domestic pig II: management factors. *Animal Welfare* 22, 219-238.
- 312   Christison, G.I., Wenger, I.I., Follensbee, M.E., 1997. Teat seeking success of newborn piglets after  
313   drying or warming. *Canadian Journal of Animal Science* 77, 317-319.
- 314   Cronin, G.M., Lefébure, B., McClintock, S., 2000. A comparison of piglet production and survival in  
315   the Werribee Farrowing Pen and conventional farrowing crates at a commercial farm. *Australian*  
316   *Journal of Experimental Agriculture* 40, 17-23.
- 317   Dohoo, I., Martin, W., Stryhn, H., 2009. *Veterinary Epidemiologic Research*. AVC Inc.,  
318   Charlottetown, Prince Edward Island, Canada.
- 319   Donovan, T.S., Dritz, S.S., 2000. Effect of split nursing on variation in pig growth from birth to  
320   weaning. *Journal of the American Veterinary Medical Association* 217, 79-86.
- 321   Dyck, G.W., Swiestra, E.E., 1987. Causes of piglet death from birth to weaning. *Canadian Journal of*  
322   *Animal Science* 67, 543-547.
- 323   English, P.R., 1993. Factors affecting neonatal piglet losses and management practices to minimize  
324   such losses. *Veterinary Annual* 33, 107-119.
- 325   English, P.R., Morrison, V., 1984. Causes and prevention of piglet mortality. *Pig News and*  
326   *Information* 5, 369-376.
- 327   Heim, G., Mellagi, A.P.G., Bierhals, T., de Souza, L.P., de Fries, H.C.C., Piuco, P., Seidel, E.,  
328   Bernardi, M.L., Wentz, I., Bortolozzo, F.P., 2012. Effects of cross-fostering within 24 h after  
329   birth on pre-weaning behaviour, growth performance and survival rate of biological and adopted  
330   piglets. *Livestock Science* 150, 121-127.
- 331   Holyoake, P.K., Dial, G.D., Trigg, T., King, V.L., 1995. Reducing pig mortality through supervision  
332   during the perinatal period. *Journal of animal science* 73, 3543-3551.

333 Johnson, A.K., Morrow, J.L., Dailey, J.W., McGlone, J.J., 2007. Prewaning mortality in loose-  
334 housed lactating sows: Behavioral and performance differences between sows who crush or do not  
335 crush piglets. *Applied Animal Behaviour Science* 105, 59-74.

336 Kirkden, R.D., Broom, D.M., Andersen, I.L., 2013. INVITED REVIEW: Piglet mortality:  
337 Management solutions. *Journal of Animal Science* 91, 3361-3389.

338 Lensink, B.J., Leruste, H., De Bretagne, T., Bizeray-Filoché, D., 2009a. Sow behaviour towards  
339 humans during standard management procedures and their relationship to piglet survival. *Applied*  
340 *Animal Behaviour Science* 119, 151-157.

341 Lensink, B.J., Leruste, H., Le Roux, T., Bizeray-Filoché, D., 2009b. Relationship between the  
342 behaviour of sows at 6 months old and the behaviour and performance at farrowing. *animal* 3, 128-  
343 134.

344 Marchant Forde, J.N., 2002. Piglet- and stockperson-directed sow aggression after farrowing and the  
345 relationship with a pre-farrowing, human approach test. *Applied Animal Behaviour Science* 75, 115-  
346 132.

347 Marchant, J.N., Rudd, A.R., Mendl, M.T., Broom, D.M., Meredith, M.J., Corning, S., Simmins, P.H.,  
348 2000. Timing and causes of piglet mortality in alternative and conventional farrowing systems.  
349 *Veterinary Record* 147, 209-214.

350 Neal, S.M., Irvin, K.M., 1991. The effects of crossfostering pigs on survival and growth. *Journal of*  
351 *animal science* 69, 41-46.

352 Ravel, A., D'Allaire, S., Bigras-Poulin, M., 1996. Influence of management, housing and personality  
353 of the stockperson on preweaning performances on independent and integrated swine farms in Québec.  
354 *Preventive Veterinary Medicine* 29, 37-57.

355 Rosvold, E.M., 2006. Kullstørrelse og morsinvestering hos gris - innvirkning på diætferd, tap og vekt  
356 (Translated title: Litter size and maternal investment in pigs - the effect on nursing behaviour, piglet  
357 mortality and weight). Master Thesis. University of Life Sciences, Ås, Norway, p. 51.

358 Wechsler, B., Hegglin, D., 1997. Individual differences in the behaviour of sows at the nest-site and  
359 the crushing of piglets. *Applied Animal Behaviour Science* 51, 39-49.

360 Westin, R., Holmgren, N., Hultgren, J., Ortman, K., Linder, A., Algers, B., 2015. Post-mortem  
361 findings and piglet mortality in relation to strategic use of straw at farrowing. *Preventive Veterinary*  
362 *Medicine* 119, 141-152.

363 White, K.R., Anderson, D.M., Bate, L.A., 1996. Increasing piglet survival through an improved  
364 farrowing management protocol. *Canadian Journal of Animal Science* 76, 491-495.

365

366 **Figure captions**

367 **Table 1:** Definition of four different management routines, number (n) and percentage (%) of farmers  
368 grouped within different types of management.

369 **Table 2:** Percentage of 52 farms that conducted the different management routines at farrowing and  
370 within the first 48 hours after farrowing.

371 **Table 3:** Distribution of different routines and criteria of cross fostering at the 52 farms.

372 **Table 4:** Distribution of farmer/ sow relationship and farmer`s opinion of the sows at 52 farms.

373 **Table 5:** Production results of study herds (n=52) extracted from Ingris (The National Efficiency  
374 Control Database) (n=290 in 2012 and 281 in 2013).

375 **Table 6:** Factors significantly associated with pre-weaning herd piglet mortality (HPM %).  
376 Multivariable adjusted estimated coefficients from a linear regression model. Number of observations  
377 (n), estimates ( $\beta$ ), standard error (SE), *P*- value and the 95% confidence intervals (CI).

378 **Figure 1:** Frequency of Herd live born piglet mortality (HPM) in the study herds (n=52).

379 **Figure 2:** Distribution of predicted values of herd piglet mortality (HPM) from different management  
380 types (M) in the regression model, % (mean  $\pm$  S.E). Management effort increases from M1 to M2  
381 (contact with sows >2 times per day), to M3 (presence at 80-100% of the farrowings, drying newborn  
382 piglets, and practice split suckling) and to M4 (combined M2 and M3). M1 herds were baseline without  
383 any of the four previously mentioned routines.

384 **Figure 3:** Distribution of predicted values of herd piglet mortality (HPM), % (mean  $\pm$  S.E), with  
385 different routines of cross-fostering timing in the regression model (Differences between predicted  
386 HPM are indicated by letters: a and b: *P* = 0,01, ab: NS).

387

388 **Table 1.** Definition of four different management routines, number (n) and percentage (%) of farmers  
 389 grouped within different types of management.

Management type (M)	n	%	Present at 80-100 % of the farrowings	Drying and massaging	Split suckling	Contact sows >2 times per day	with
M1	28	53.8	-	-	-	-	-
M2	11	21.2	-	-	-	+	+
M3	9	17.3	+	+	+	-	-
M4	4	7.7	+	+	+	+	+

390

391

392 **Table 2.** Percentage of 52 farms that conducted the different management routines at farrowing and  
 393 within the first 48 hours after farrowing.

	Farrowing n (%)	First 48 h n (%)
Farmers presence;	<sup>1</sup>	<sup>2</sup>
80-100%	22 (42.3)	12 (23.1)
60-80%	12 (23.1)	14 (26.9)
40-60%	14 (26.9)	16 (30.8)
20-40%	3 (5.8)	5 (9.6)
0-20%	1 (1.9)	5 (9.6)
Piglets are		
dried/massaged	28 (53.8)	4 (7.7)
moved to the udder	36 (69.2)	15 (28.8)
moved to the creep	35 (67.3)	25 (48.1)
Split suckling	29 (55.8)	-
Additional milk <sup>3</sup>	28 (53.8)	-
Piglets closed inside creep at feeding	-	28 (53.8)
Sow exercise <sup>4</sup>	-	16 (30.8)
Tooth grinding	-	42 (80.8)
Other routines <sup>5</sup>	23 (44.2)	15 (28.8)

394 <sup>1</sup>Farmers` presence in % of all farrowings in a typical batch.

395 <sup>2</sup>Farmers` presence in % of the time the first 48 hours after farrowing.

396 <sup>3</sup>Milk replacer

397 <sup>4</sup>Sows are taken out from the farrowing pen for a short walk in the farrowing unit.

398 <sup>5</sup>Routines done regularly by farmer but not asked for specifically in the survey.

399

400 **Table 3:** Distribution of different routines and criteria of cross fostering at the 52 farms.

Cross fostering	n (%)
Proportions of litters where cross-fostering is performed	
80-100%	4 (7.7)
60-80%	7 (13.5)
40-60%	18 (34.6)
20-40%	17 (32.7)
00-20%	6 (11.5)
Criteria for cross fostering	
Even out number of piglets between litters	23 (44.2)
Homogeneity in piglet size within litter	4 (7.7)
According to number of functional teats	8 (15.4)
Two of the criteria <sup>1</sup>	11 (21.0)
Three of the criteria <sup>1</sup>	5 (9.6)
Other criteria <sup>1</sup>	1 (1.9)
Which piglets are cross-fostered	
The biggest	25 (48.1)
The smallest	2 (3.8)
The medium	2 (3.8)
No preference	2 (3.8)
The biggest + smallest <sup>1</sup>	15 (28.8)
Biggest + other <sup>1</sup>	2 (3.8)
Biggest + medium <sup>1</sup>	2 (3.8)
Biggest + smallest + medium <sup>1</sup>	2 (2.8)
Timing of cross fostering after farrowing	
First 12 hours	7 (13.5)
13-24 hours	21 (40.4)
25-36 hours	15 (28.8)
<12-24 hours <sup>1</sup>	2 (3.8)
13-36 hours <sup>1</sup>	4 (7.7)
<12-36 hours <sup>1</sup>	2 (3.8)
<12- > 48 hours <sup>1</sup>	1 (1.9)

401 <sup>1</sup>Farmers with more than one routine or other routine(s) than the existent answer categories.

402



403 **Table 4:** Distribution of farmer/ sow relationship and farmer`s opinion of the sows at 52 farms.

Farmer/ sow relationships	n (%)
Contact with the sow	
> 2 times/day	15 (28.8)
2 times/day (at feeding)	29 (55.8)
≤ 7 times/week	8 (15.4)
Farmer`s opinion of handling sows during pregnancy	
80-100% of sows easy to handle	44 (84.6)
60-80% of sows easy to handle	4 (7.7)
40-60% of sows easy to handle	3 (5.8)
20-40% of sows easy to handle	0 (0)
0-20% of sows easy to handle	1 (1.9)
Farmer`s opinion of handling sows at farrowing/lactation	
80-100% of sows easy to handle	42 (80.8)
60-80% of sows easy to handle	7 (13.5)
40-60% of sows easy to handle	1 (1.9)
20-40% of sows easy to handle	2 (3.8)
0-20% of sows easy to handle	0 (0)

404

405 **Table 5:** Production results of study herds (n=52), extracted from Ingris (The National Efficiency  
 406 Control Database), and national results from Ingris (n=290 in 2012 and 281 in 2013).

	Year	Study herds		National results, Ingris
		Mean $\pm$ S.E.	(Min-Max)	Mean
Live born	2012+2013	13.6 $\pm$ 0.1	(11.6-15.1)	-
	2012	13.6 $\pm$ 0.1	(11.7-15.3)	13.3
	2013	13.6 $\pm$ 0.1	(11.2-15.0)	13.2
Stillborn	2012+2013	1.7 $\pm$ 0.1	(0.6-2.7)	-
	2012	1.6 $\pm$ 0.1	(0.6-2.9)	1.2
	2013	1.7 $\pm$ 0.1	(0.5-2.8)	1.2
Weaned	2012+2013	11.3 $\pm$ 0.1	(9.3-13.3)	-
	2012	11.3 $\pm$ 0.1	(9.2-13.4)	11.3
	2013	11.3 $\pm$ 0.1	(9.4-13.2)	11.2
Herd piglet mortality (HPM), % <sup>1</sup>	2012+2013	16.9 $\pm$ 0.6	(5.5-28.3)	-
	2012	16.9 $\pm$ 0.7	(6.4-29.3)	15.0
	2013	16.9 $\pm$ 0.7	(4.3-27.6)	15.3

407 <sup>1</sup> Herd piglet mortality (HPM): ((Live born - weaned)/Live born)\*100%.

408 **Table 6:** Factors significantly associated with pre-weaning herd piglet mortality (HPM %).  
 409 Multivariable adjusted estimated coefficients from a linear regression model. Number of observations  
 410 (n), estimates ( $\beta$ ), standard error (SE), *P*- value and the 95% confidence intervals (CI).

Variables	n	$\beta$	SE	<i>P</i>	[95% Conf. Interval]	
Intercept	52	20.07	1.41	<0.01	17.22 22.91	
Batch system <sup>1</sup>						
7 wk	34	0.00	(base)			
2.5-3 wk	4	-4.31	2.22	0.06	-8.81 0.18	
5.5 wk	8	1.30	1.70	0.45	-2.15 4.75	
11/22-26 wk	6	3.13	1.85	0.10	-0.62 6.88	
Number of sows/batch						
<20	17	0.00	(base)			
20	5	-5.33	2.03	0.01	-9.33 -1.12	
21-39	21	-3.58	1.27	0.01	-6.16 -1.01	
40-65	9	-5.16	1.77	0.01	-8.75 -1.57	
Management type <sup>2</sup>						
1 (M1)	28	0.00	(base)			
2 (M2)	11	-3.05	1.40	0.04	-5.88 -0.21	
3 (M3)	9	-3.85	1.54	0.02	-6.98 -0.73	
4 (M4)	4	-6.77	2.12	<0.01	-11.07 -2.48	
Time of cross-fostering						
13-24 h after farrowing	21	0.00	(base)			
< 12 h after farrowing	7	0.8	1.67	0.63	-2.57 4.17	
25-36 h after farrowing	15	1.61	1.39	0.25	-1.20 4.42	
No systematic routine <sup>3</sup>	9	4.11	1.51	0.01	1.07 7.16	
Herd litter size centered around the mean <sup>4</sup>	52	0.73	0.95	0.45	-1.19 2.65	


411 <sup>1</sup>Time interval between farrowings, for instance 7 wk means farrowing every 7<sup>th</sup> week.

412 <sup>2</sup>Management type 1-4 are ordinal categories, where management type 1 (M1) means low management  
 413 effort and management type 4 (M4) means high management effort.

414 <sup>3</sup>Farmers had a longer “time-window” than 12 hours for cross-fostering, and it was done with variation  
 415 of timing after farrowing.


416 <sup>4</sup>2012 and 2013 results on herd level and centered around mean to get a more biological constant, even  
 417 out results that could be too good or bad that it cannot represent the herd in overall.

418




419

420 **Figure 1.** Frequency of Herd live born piglet mortality (HPM) in the study herds (n=52).



421

422 **Figure 2.** Distribution of predicted values of herd piglet mortality (HPM) from different management  
 423 types (M) in the regression model, % (mean ± S.E). Management effort increases from M1 to M2  
 424 (contact with sows >2 times per day), to M3 (presence at 80-100% of the farrowings, drying newborn  
 425 piglets, and practice split suckling) and to M4 (combined M2 and M3). M1 herds were baseline without  
 426 any of the four previously mentioned routines.



427

428 **Figure 3.** Distribution of predicted values of herd piglet mortality (HPM), % (mean  $\pm$  S.E), with  
 429 different routines of cross-fostering timing in the regression model (Differences between predicted  
 430 HPM are indicated by letters: a and b: P = 0,01, ab: NS).