

Structural and transport properties of hydrogen in ZIF-22

Uthumporn Arsawang^{1*}, Siegfried Fritzsche¹, Wolfhard Janke¹, Jürgen Caro²,
Tawun Remsungnen³, Supot Hannongbua^{4,5}

¹Universität Leipzig, Institute for Theoretical Physics, Leipzig, Germany

²Leibniz University Hannover, Institute of Physical Chemistry and Electrochemistry,
Hannover, Germany

³Khon Kaen University, Department of Mathematics, Khon Kaen, Thailand

⁴Chulalongkorn University, Computational Chemistry Unit Cell (CCUC),
Department of Chemistry, Bangkok, Thailand

⁵Chulalongkorn University, Metallurgy and Materials Science Research Institute (MMRI),
Bangkok, Thailand

*uarsawang@gmail.com

ZIF-22 which has a structure similar to A zeolites [1, 2] is investigated in this work because of its high selectivity for hydrogen with respect to other gases [1]. By using Molecular Dynamics simulations with the DL_POLY package, structural and dynamical properties of hydrogen adsorbed in ZIF-22 are examined.

Moreover, the adsorption of hydrogen in ZIF-22 is simulated by Gibbs-ensemble Monte-Carlo simulation studies.




Figure 1: The density plot of hydrogen in one of the cavities of ZIF-22

The small window size turns out to be the most important reason for the high selectivity. The results are compared with experiments [1] and they are discussed by means of spatial density distributions, radial correlation functions and mean square displacements.

References

- [1] A. Huang, H. Bux, F. Steinbach, J. Caro: *Molecular-sieve membrane with hydrogen permselectivity: ZIF-22 in LTA topology prepared with 3-aminopropyltriethoxysilane as covalent linker*. *Angewandte Chemie International Edition* **49**, 4958–4961 (2010)

- [2] H. Hayashi, A.P. Côté, H. Furukawa, M. O’Keeffe: *Zeolite A imidazolate frameworks*. *Nature Materials* **6**, 501–506 (2007)