

Short-term Associations between Fine and Coarse Particulate Matter and Hospitalizations in Southern Europe: Results from the MED-PARTICLES Project

Massimo Stafoggia,¹ Evangelia Samoli,² Ester Alessandrini,¹ Ennio Cadum,³ Bart Ostro,^{4,5} Giovanna Berti,³ Annunziata Faustini,¹ Benedicte Jacquemin,^{4,6} Cristina Linares,^{7,8} Mathilde Pascal,⁹ Giorgia Randi,¹⁰ Andrea Ranzi,¹¹ Elisa Stivanello,¹² Francesco Forastiere,¹ and the MED-PARTICLES Study Group

¹Department of Epidemiology, Lazio Region Health Service, Rome, Italy; ²Department of Hygiene, Epidemiology and Medical Statistics, Medical School, University of Athens, Athens, Greece; ³Department of Epidemiology and Environmental Health, Regional Environmental Protection Agency, Piedmont, Italy; ⁴Centre for Research in Environmental Epidemiology, Barcelona, Spain; ⁵Air Pollution Epidemiology Section, Office of Environmental Health Hazard Assessment, California Environmental Protection Agency, Oakland, California, USA; ⁶INSERM U1018, CESP—Centre for research in Epidemiology and Population Health, UMRS U1018, Respiratory and Environmental Epidemiology Team, University Paris Sud, Villejuif, France; ⁷CIBER Epidemiología y Salud Pública (CIBERESP), Madrid, Spain; ⁸Cancer and Environmental Epidemiology Unit, National Centre for Epidemiology, Carlos III Institute of Health, Madrid, Spain; ⁹Unité Air Eau et Climat, Département Santé Environnement Institut de Veille Sanitaire, Saint-Maurice Cedex, France; ¹⁰Epidemiology Unit, Local Health Authority, Milan, Italy; ¹¹Regional Centre for Environment and Health, Regional Environmental Protection Agency—Emilia-Romagna, Modena, Italy; ¹²Epidemiology Observatory, Department of Public Health, Local Health Authority, Bologna, Italy

BACKGROUND: Evidence on the short-term effects of fine and coarse particles on morbidity in Europe is scarce and inconsistent.

OBJECTIVES: We aimed to estimate the association between daily concentrations of fine and coarse particles with hospitalizations for cardiovascular and respiratory conditions in eight Southern European cities, within the MED-PARTICLES project.

METHODS: City-specific Poisson models were fitted to estimate associations of daily concentrations of particulate matter with aerodynamic diameter $\leq 2.5 \mu\text{m}$ ($\text{PM}_{2.5}$), $\leq 10 \mu\text{m}$ (PM_{10}), and their difference ($\text{PM}_{2.5-10}$) with daily counts of emergency hospitalizations for cardiovascular and respiratory diseases. We derived pooled estimates from random-effects meta-analysis and evaluated the robustness of results to co-pollutant exposure adjustment and model specification. Pooled concentration–response curves were estimated using a meta-smoothing approach.

RESULTS: We found significant associations between all PM fractions and cardiovascular admissions. Increases of $10 \mu\text{g}/\text{m}^3$ in $\text{PM}_{2.5}$, $6.3 \mu\text{g}/\text{m}^3$ in $\text{PM}_{2.5-10}$, and $14.4 \mu\text{g}/\text{m}^3$ in PM_{10} (lag 0–1 days) were associated with increases in cardiovascular admissions of 0.51% (95% CI: 0.12, 0.90%), 0.46% (95% CI: 0.10, 0.82%), and 0.53% (95% CI: 0.06, 1.00%), respectively. Stronger associations were estimated for respiratory hospitalizations, ranging from 1.15% (95% CI: 0.21, 2.11%) for PM_{10} to 1.36% (95% CI: 0.23, 2.49) for $\text{PM}_{2.5}$ (lag 0–5 days).

CONCLUSIONS: $\text{PM}_{2.5}$ and $\text{PM}_{2.5-10}$ were positively associated with cardiovascular and respiratory admissions in eight Mediterranean cities. Information on the short-term effects of different PM fractions on morbidity in Southern Europe will be useful to inform European policies on air quality standards.

CITATION: Stafoggia M, Samoli E, Alessandrini E, Cadum E, Ostro B, Berti G, Faustini A, Jacquemin B, Linares C, Pascal M, Randi G, Ranzi A, Stivanello E, Forastiere F, the MED-PARTICLES Study Group. 2013. Short-term associations between fine and coarse particulate matter and hospitalizations in Southern Europe: results from the MED-PARTICLES project. *Environ Health Perspect* 121:1026–1033; <http://dx.doi.org/10.1289/ehp.1206151>

Introduction

The European air quality standards are under revision, and a new directive will be delivered by the European Union (EU) in the next few years. As part of this process, the EU has indicated several specific issues of concern. Among the open issues is the extent of short-term health effects of fine and coarse particle concentrations and components in Europe, and the shape of the concentration–response relationships between fine particles and mortality and morbidity. Much of the evidence about short-term associations between fine particles [particulate matter with aerodynamic diameter $\leq 2.5 \mu\text{m}$ ($\text{PM}_{2.5}$)] and health end points comes from studies conducted in the United States, where a 24-hr National Ambient Air Quality Standard for $\text{PM}_{2.5}$ was first introduced in 1997 (U.S.

Environmental Protection Agency 2012). From that time, data on fine particles have been collected in many parts of the United States, and evidence has accumulated suggesting significant effects of fine particles on both mortality (Ostro et al. 2006; Zanobetti and Schwartz 2009) and hospital admissions (Bell et al. 2008; Dominici et al. 2006). In contrast, EU legislation currently has a single limit value for exposure to $\text{PM}_{2.5}$ based on an annual averaging period, without regulatory standards for daily concentrations. Only a few studies on the short-term effects of $\text{PM}_{2.5}$ on mortality or morbidity have been conducted in Europe, with most conducted in a single city (Anderson et al. 2001; Atkinson et al. 2010; Belleudi et al. 2010; Halonen et al. 2009; Linares and Diaz 2010), and only one based on multiple cities in one country

(France) (Host et al. 2008). Therefore, it is unclear whether previous findings can be generalized to all of Europe.

Another topic under debate in the EU is the role of other PM fractions on human health and, more specifically, whether coarse particles (particles with diameter 2.5–10 μm ; $\text{PM}_{2.5-10}$) are associated with health, and whether they should be monitored by European policies. This discussion was stimulated by a systematic review published by Brunekreef and Forsberg (2005), in which the authors concluded that “coarse PM has a stronger or as strong short-term effect” on respiratory health (based on emergency hospitalizations for respiratory outcomes) as fine PM. In addition, they concluded that there

Address correspondence to M. Stafoggia, Department of Epidemiology of the Lazio Region Health Service, Via Santa Costanza 53, 00198, Rome, Italy. Telephone: 39-0683060474. E-mail: m.stafoggia@deplazio.it

Supplemental Material is available online (<http://dx.doi.org/10.1289/ehp.1206151>).

MED-PARTICLES Study Group: Italy: E. Alessandrini, P. Angelini, G. Berti, L. Bisanti, E. Cadum, M. Catrambone, M. Chiusolo, M. Davoli, F. de’ Donato, M. Demaria, M. Gandini, M. Groza, A. Faustini, S. Ferrari, F. Forastiere, P. Pandolfi, R. Pelosini, C. Perrino, A. Pietrodangelo, L. Pizzi, V. Poluzzi, G. Priod, G. Randi, A. Ranzi, M. Rowinski, C. Scarinzi, M. Stafoggia, E. Stivanello, S. Zauli-Sajani; Greece: K. Dimakopoulou, K. Eleftheriadis, K. Katsouyanni, A. Kelesiss, T. Maggos, N. Michalopoulos, S. Pateraki, M. Petrakakis, S. Rodopoulou, E. Samoli, V. Sytsa; Spain: J. Alguacil, B. Artiñano, J. Barrera-Gómez, X. Basagaña, J. de la Rosa, J. Diaz, B. Jacquemin, A. Karanasiou, C. Linares, B. Ostro, J. Pey, X. Querol, J. Sunyer, A. Tobias; France: M. Bidondo, C. Declercq, A. Le Tertre, P. Lozano, S. Medina, L. Pascal, M. Pascal.

We thank the Consorci Sanitari de Barcelona (Cat-Salut) for providing hospitalization data for Barcelona.

This research was supported by the European Union under the grant agreement LIFE+ ENV/IT/327.

The authors declare they have no actual or potential competing financial interests.

Received: 18 October 2012; Accepted: 17 June 2013; Advance Publication: 18 June 2013; Final Publication: 1 September 2013.

was some evidence supporting effects of $PM_{2.5-10}$ on cardiovascular hospitalizations, whereas, for overall mortality, the evidence was stronger for an effect of $PM_{2.5}$, and limited for coarse particles.

The MED-PARTICLES (“Particles size and composition in Mediterranean countries: geographical variability and short-term health effects”) project was specifically designed to address these and other related questions. It is financed by the EU under the LIFE+ framework, and aims to describe and compare the composition of airborne particles across Mediterranean cities and to estimate health effects associated with exposure to PM concentrations, PM components and sources, and Saharan dust and forest fires (MED-PARTICLES 2010).

Here we present the results of an investigation of short-term associations of daily concentrations of $PM_{2.5}$, $PM_{2.5-10}$, and PM_{10} (PM with aerodynamic diameter $\leq 10 \mu m$) with emergency hospitalizations for cardiovascular and respiratory diseases in eight Southern European cities.

Methods

Study population. Daily counts of emergency hospital admissions were collected from national or regional health information systems for 10 European cities, between 2001 and 2010: Milan, Turin, Bologna, Parma, Reggio Emilia, Modena, and Rome, Italy; Marseille, France; and Madrid and Barcelona, Spain. Because Parma, Reggio Emilia, and Modena are very close and share common environmental and sociodemographic characteristics, they have been analyzed altogether as a single conurbation called “Emilia Romagna.” Hospitalizations only of residents ≥ 15 years of age were considered. Two study outcomes were defined on the basis of the primary discharge diagnosis: cardiovascular hospitalizations [*International Classification of Diseases, 9th Revision* (ICD-9) codes 390–459; and *10th Revision* (ICD-10) codes I00–I99] (WHO 1999), and respiratory hospitalizations (ICD-9 codes 460–519; ICD-10 codes J00–J99). All data were extracted and collected according to a common protocol. Because data were anonymous and collected as daily counts, no informed consent was needed. And because the analysis on daily counts was conducted by a public health institute, there was no need for approval by an institutional review board.

Environmental variables. Daily concentrations of $PM_{2.5}$, PM_{10} , nitrogen dioxide (NO_2), and ozone (O_3) were collected in each city from multiple monitors belonging to air quality monitoring networks. Only monitors with complete data for $\geq 75\%$ of the study period were accepted. When a monitor had a missing value for a specific day, it was

replaced by the average of the values of the remaining stations for that day multiplied by a factor equal to the ratio of the annual mean for the missing station over the corresponding annual mean for the other stations (Katsouyanni et al. 2001). When data from all monitors were missing on a day, the daily value was left as missing. Daily mean concentrations were computed for PM and NO_2 , and daily maximum 8-hr running means were calculated for O_3 . Daily $PM_{2.5-10}$ was calculated for each station as the difference between PM_{10} and $PM_{2.5}$, provided that both PM fractions were measured at the same station using the same sampling methodology. Nine of 10 cities sampled PM concentrations using the gravimetric method or an equivalent one, whereas only one city, Marseille, provided uncorrected concentrations from tapered element oscillating microbalance (TEOM) samplers. Daily mean air temperature was collected from each center, using airport meteorological stations where available.

Other confounders. Time-varying confounders were constructed according to a common protocol, under the assumption of similar behavior of the study populations during holidays and vacation periods. They include holidays (a four-level variable assuming a value of 3 on Christmas and Easter; 2 in the periods surrounding Christmas and Easter; 1 on isolated holidays; 0 on other days); summer population decrease (a three-level variable assuming value 2 in the 2-week period around 15 August; 1 from 16 July to 31 August, except for the aforementioned period; 0 for all other days); and influenza epidemics (a two-level variable assuming 0 on normal days and 1 on days with particularly high influenza episodes). Influenza epidemics were identified using national influenza surveillance systems, where available, or were identified based on daily counts of hospitalizations for influenza (ICD-9 code 487; ICD-10 codes J09–J11).

Statistical analysis. The analyses were carried out using a two-stage approach. In the first stage, city-specific overdispersed Poisson regression models were fitted, in which the dependent variable was the daily count of hospitalizations, the exposure was the (lagged) daily concentration of PM_x , and variables added for confounding adjustment. The adjustment model was defined *a priori* and, to reduce the potential for heterogeneity in the city-specific results, was the same for all cities. The confounders were long-term and seasonal time trends, air temperature, holidays, summer population decrease, and influenza epidemics. A directed acyclic graph (DAG) of the causal relationships under study is depicted in Supplemental Material, Figure S1.

Holidays, summer population decreases, and influenza epidemics were modeled using

indicator variables. Time trend was adjusted for by introducing a three-way interaction term among year, month, and day of the week. This method has been demonstrated to be equivalent to a case-crossover design using a time-stratified approach to select control days (Levy et al. 2001; Lu and Zeger 2007; Maclure 1991). Two sensitivity analyses were designed to check the robustness of the results to time trend specification: *a*) a penalized spline of time trend with 50 knots per year and a smoothing parameter tuned to approximate 8 effective degrees of freedom (df) per year (Samoli et al. 2003); *b*) a penalized spline for time trend with the smoothing parameter set to minimize the absolute value of the sum of the partial autocorrelation functions (PACF) of residuals from lag 0 to 30, but requiring a minimum of 3 df per year (Katsouyanni et al. 2009). In both cases, indicator variables for days of the week were included in the models. We performed sensitivity analyses of seasonality control because a consensus on the method best suited for time trend adjustment has not been reached in the literature (Janes et al. 2005; Whitaker et al. 2007).

We controlled for the effect of temperature by modeling high and low temperatures separately. For high temperatures we calculated the average temperature on the current and previous day (lag 0–1) and fit a natural spline with 3 df on the lagged variable only for days on which the lag 0–1 temperature was higher than the median annual temperature for the city as a whole. Similarly, we adjusted for low temperatures by fitting a natural spline with 2 df for the average temperature on previous 6 days (lag 1–6) only for days on which the lag 1–6 temperature was below the median annual value for the city (Chiusolo et al. 2011). This method accounts for differences in the lag structures and effects of cold and warm temperatures on hospitalizations while reducing the correlation between the two spline terms. We also performed an analysis that adjusted for potentially prolonged effects of warm temperatures on hospitalizations by replacing the lag 0–1 temperature term from the base model with the lag 0–6 average.

Individual pollutants were added to city-specific regression models adjusted for the *a priori* covariates described above. We evaluated the lag structure of the association between PM_x concentrations and daily counts of cause-specific hospitalizations by applying cubic polynomial distributed lag models, with individual daily lags from lag 0 to 5 entered simultaneously in the model and constrained to follow a polynomial shape. In addition, we defined three cumulative lag structures *a priori* to represent immediate (lag 0–1), delayed (lag 2–5), and prolonged (lag 0–5) effects, and chose one of these structures as

the default lag for each PM–outcome combination based on the meta-analytical results of the distributed and cumulative lag models. This choice has been applied previously (Stafoggia et al. 2010) as a compromise between *a priori* and data-driven approaches for selecting the lag structure for different exposure/outcome combinations.

The robustness of associations to adjustment for co-exposure to other pollutants was evaluated by fitting two-pollutant models with the co-pollutant modeled using the cumulative lag selected as the default for the primary exposure–outcome combination. In addition, models that included O₃ as the co-pollutant were fit using data restricted to the warm season only (April–September), in addition to year-round data. Finally, we investigated concentration–response functions between PM_x and hospitalizations by fitting, for each city, a natural spline model for the exposure with two equally spaced inner knots, and pooling the city-specific estimates using a meta-smoothing approach (Schwartz and Zanobetti 2000).

In the second stage of the analysis, we pooled the city-specific results using random-effects meta-analytical procedures according to the method proposed by Jackson et al. (2010). We tested heterogeneity among the city-specific results by applying the chi-square test from Cochran's Q statistic, and estimated the amount of heterogeneity by computing the I² statistic (Higgins and Thompson 2002), which represents the proportion of total variation in effect estimates due to between-cities heterogeneity. We considered city-specific effect estimates to be significantly heterogeneous when I² was > 50% and the χ² p-value was < 0.10.

All results are expressed as percent increases in hospitalizations, with 95% CI, relative to fixed increments in each PM fraction: 10 μg/m³ for PM_{2.5}, 6.3 μg/m³ for PM_{2.5–10}, and 14.4 μg/m³ for PM₁₀. These increments were chosen to represent a comparable amount of daily variability across pollutants and facilitate comparisons of effect estimates among the

pollutants. More specifically, we computed the interquartile range (IQR) of each PM fraction from the joint distribution for the eight cities (six cities for PM_{2.5–10}), and scaled the three resulting IQRs to express PM_{2.5} effects per 10-μg/m³ increase. In addition, we report selected associations in the text using a common increment of 10 μg/m³ for all three PM fractions.

All first-stage analyses were fit using R, version 2.15.0 (R Development Core Team; <http://R-project.org>). Meta-analyses were conducted using Stata, version 11 (StataCorp, College Station, TX, USA).

Results

The population base comprised residents ≥ 15 years of age in eight Southern European cities, totaling > 11 million inhabitants (Table 1). The mean daily counts of hospitalizations ranged from 18 in the Emilia Romagna conurbation to 116 in Madrid for cardiovascular diseases, and from 8 in Bologna to 97 in Madrid for respiratory conditions. The study periods were recent and comparable across cities, except for Marseille with 2001–2003 data.

Pollutant concentrations and air temperature data for each city are summarized in Supplemental Material, Tables S1 (year-round) and S2 (separate PM concentration distributions for the cold and warm seasons). Daily concentrations of PM₁₀ and PM_{2.5} were highest in Milan and Turin. Madrid had the smallest daily PM_{2.5} concentrations and the largest PM_{2.5–10} concentrations, resulting in a fine:coarse particles ratio that was much smaller for this city (< 1) than the others (ranging from 1.5 to 2.2). NO₂ and O₃ concentrations were similar among the cities, whereas temperature displayed a mild increasing north–south gradient. Pearson correlation coefficients between PM_{2.5} and PM_{2.5–10} were close to 0 in Barcelona, Marseille, and Rome (the cities closest to the sea and therefore likely to be more affected by sea winds), but were ≥ 0.5 in the other cities (data not shown). NO₂ concentrations were highly

correlated with both PM_{2.5} and PM₁₀ (> 0.6 for all cities except Barcelona), whereas correlations between NO₂ and PM_{2.5–10} ranged from 0.17 in Marseille to 0.57 in Madrid.

Pooled estimates from polynomial distributed lag models (Supplemental Material, Figure S2) clearly show evidence of an immediate effect of the three PM fractions on cardiovascular hospitalizations, up to lag 1, whereas associations with respiratory admissions were evident until day 5.

Table 2 reports associations between PM and hospitalizations for the three cumulative lags (0–1, 2–5, and 0–5 days). Significant positive associations of a comparable magnitude were estimated for all PM fractions with cardiovascular hospitalizations: increments of 10 μg/m³ in PM_{2.5}, 6.3 μg/m³ in PM_{2.5–10}, and 14.4 μg/m³ in PM₁₀ (lag 0–1) were associated with 0.51% (95% CI: 0.12, 0.90%), 0.46% (95% CI: 0.10, 0.82%), and 0.53% (95% CI: 0.06, 1.00%) increases in cardiovascular admissions, respectively. When expressed per 10 μg/m³, corresponding estimates for PM_{2.5–10} and PM₁₀ were 0.73% (95% CI: 0.16, 1.30%) and 0.36% (95% CI: 0.04, 0.69%), suggesting an effect of coarse particles around 40% higher than that of PM_{2.5} for the same increment. Associations between cardiovascular hospitalizations and PM were null for lag 2–5, and smaller or null for lag 0–5 (Table 2). Given these results and the results of the distributed lag models, we used lag 0–1 for subsequent analyses of cardiovascular hospitalizations.

Associations with respiratory hospitalizations were strongest for all three PM fractions at the cumulative lag 0–5, though the association was not statistically significant for PM_{2.5–10}, and effect estimates were highly heterogeneous across cities. Specifically, lag 0–5 effect estimates were 1.36% (95% CI: 0.23, 2.49%) for a 10-μg/m³ increase in PM_{2.5}, 1.24% (95% CI: –0.32, 2.82%) for a 6.3-μg/m³ increase in PM_{2.5–10}, and 1.15% (95% CI: 0.21, 2.11%) for a 14.4-μg/m³ increase in PM₁₀. Associations were more homogeneous among cities for

Table 1. Study populations and emergency hospital admissions for cardiovascular and respiratory causes among residents ≥ 15 years of age in the eight cities of the MED-PARTICLES project.

City ^a	Study period	Date	Population		Cardiovascular admissions		Respiratory admissions		
			<i>n</i>	<i>n</i>	<i>n</i> per 1,000 person-years	Daily mean	<i>n</i>	<i>n</i> per 1,000 person-years	Daily mean
Milan	2006–2010	1 January 2008	1,299,633	71,779	11.0	39.3	34,427	5.3	18.9
Turin	2006–2010	1 January 2008	908,263	48,967	10.8	26.8	21,761	4.8	11.9
Emilia Romagna ^b	2008–2010	1 January 2009	529,699	19,717	12.4	18.0	10,164	6.4	9.3
Bologna	2006–2010	1 January 2008	372,256	34,568	18.6	18.9	14,103	7.6	7.7
Marseille	2001–2003	Census 1999	796,525	46,905	19.6	42.8	18,069	7.6	16.5
Rome	2006–2010	1 January 2008	2,718,768	153,176	11.3	83.9	53,825	4.0	29.5
Barcelona	2003–2010	1 January 2007	1,595,110	151,426	11.9	51.8	139,062	10.9	47.6
Madrid	2004–2009	1 January 2007	3,132,503	201,041	13.5	115.9	167,850	11.3	96.7
Total	—	—	11,352,757	727,579	12.6	51.4	459,261	7.9	32.5

^aCities are ordered by latitude, north to south. ^bEmilia Romagna includes Parma, Reggio Emilia, and Modena.

lag 0–1, and statistically significant for the coarse fraction (0.60%; 95% CI: 0.08, 1.13%) and PM₁₀ (0.65%; 95% CI: 0.20, 1.10%). When expressed for increments of 10 µg/m³, the association with respiratory admissions was strongest for coarse particles (1.95%; 95% CI: –0.51, 4.48% for lag 0–5). We used lag 0–5 for subsequent analyses of association between the PM fractions and respiratory hospitalizations.

Pooled estimates from one- and two-pollutant models are reported in Table 3. When evaluated simultaneously, the association between fine particles and cardiovascular hospitalizations (lag 0–1) was unaffected by adjustment for PM_{2.5–10} (0.49% increase per 10 µg/m³ PM_{2.5}; 95% CI: 0.06, 0.91%) but adjusting for PM_{2.5} decreased the association between cardiovascular hospitalizations and coarse PM (0.28% increase per 6.3 µg/m³ PM_{2.5–10}; 95% CI: –0.09, 0.66%). Associations of fine and coarse particles with respiratory admissions (lag 0–5) both decreased considerably with mutual adjustment (to 0.55%; 95% CI: –0.29, 1.40 for PM_{2.5}, and 0.66%; 95% CI: –0.78, 2.13 for PM_{2.5–10}). Adjustment for NO₂ decreased association between PM_{2.5–10} and both outcomes, whereas it decreased the association between PM_{2.5} and cardiovascular hospitalizations, but strengthened the association with respiratory admissions. However, given the high correlation between PM_{2.5} and NO₂ in all cities, results from two-pollutant models must be interpreted with caution. No confounding from O₃ was apparent in the all-year analysis, nor in the analysis restricted to the warm period. However, associations of PM_{2.5} and PM_{2.5–10} with both outcomes were much stronger and always statistically significant when the analyses were restricted to the warm period (April–September) both with and without adjustment for O₃.

The robustness of the main findings was checked against model specification in three sensitivity analyses, whose results are summarized in Table 4. In general, the case-crossover method adopted in the base model for the adjustment of time trend provided effect estimates quite consistent with the spline method with minimization of the PACF of residuals, whereas associations based on models that used the spline method with 8 df per year were weaker for all particle metrics and both study outcomes. In addition, associations between all three PM fractions and respiratory admissions were strongest when estimated using the spline method with the PACF criterion. Effect estimates from the sensitivity model adjusted for prolonged effects of warm temperatures were consistent with base model estimates.

Finally, Figure 1 reports the concentration–response relationships between PM_{2.5}, PM_{2.5–10}, and PM₁₀ with cardiovascular

Table 2. Associations between PM and hospitalizations per IQR increases of 10, 6.3, and 14.4 µg/m³ for PM_{2.5}, PM_{2.5–10} and PM₁₀, respectively.

Pollutant	Lag	Cardiovascular admissions			Respiratory admissions		
		Percent increase (95% CI)	χ ² p-value	I ²	Percent increase (95% CI)	χ ² p-value	I ²
PM _{2.5}	0–1	0.51 (0.12, 0.90)	0.20	29	0.49 (–0.12, 1.09)	0.16	34
	2–5	0.15 (–0.22, 0.53)	0.82	0	1.07 (0.04, 2.11)	0.00	71
	0–5	0.49 (0.03, 0.95)	0.70	0	1.36 (0.23, 2.49)	0.01	65
PM _{2.5–10}	0–1	0.46 (0.10, 0.82)	0.33	13	0.60 (0.08, 1.13)	0.26	23
	2–5	–0.38 (–0.89, 0.12)	0.23	28	0.75 (–0.71, 2.23)	0.00	77
	0–5	0.05 (–0.68, 0.78)	0.09	47	1.24 (–0.32, 2.82)	0.00	74
PM ₁₀	0–1	0.53 (0.06, 1.00)	0.06	48	0.65 (0.20, 1.10)	0.38	6
	2–5	–0.06 (–0.43, 0.31)	0.47	0	0.95 (–0.03, 1.94)	0.00	68
	0–5	0.30 (–0.14, 0.74)	0.49	0	1.15 (0.21, 2.11)	0.04	52

Table 3. Associations between PM and hospitalizations from one- and two-pollutant models, per increases of 10, 6.3, and 14.4 µg/m³ for PM_{2.5}, PM_{2.5–10}, and PM₁₀, respectively.

Pollutant	Period	Cardiovascular admissions (lag 0–1)			Respiratory admissions (lag 0–5)		
		Percent increase (95% CI)	χ ² p-value	I ²	Percent increase (95% CI)	χ ² p-value	I ²
PM _{2.5}	All year	0.51 (0.12, 0.90)	0.20	29	1.36 (0.23, 2.49)	0.01	65
+ PM _{2.5–10}	All year	0.49 (0.06, 0.91)	0.48	0	0.55 (–0.29, 1.40)	0.47	0
+ NO ₂	All year	0.21 (–0.28, 0.69)	0.21	28	1.63 (0.30, 2.97)	0.05	51
+ O ₃	All year	0.49 (0.06, 0.91)	0.18	31	1.13 (0.05, 2.23)	0.04	53
PM _{2.5}	April–September	1.76 (0.68, 2.84)	0.18	32	4.49 (1.72, 7.35)	0.04	51
+ O ₃	April–September	2.11 (1.13, 3.10)	0.34	11	4.51 (1.35, 7.77)	0.03	56
PM _{2.5–10}	All year	0.46 (0.10, 0.82)	0.33	13	1.24 (–0.32, 2.82)	0.00	74
+ PM _{2.5}	All year	0.28 (–0.09, 0.66)	0.85	0	0.66 (–0.78, 2.13)	0.07	51
+ NO ₂	All-year	0.25 (–0.13, 0.62)	0.42	0	0.99 (–0.71, 2.73)	0.02	61
+ O ₃	All year	0.48 (0.14, 0.83)	0.41	2	1.17 (–0.45, 2.81)	0.01	70
PM _{2.5–10}	April–September	0.90 (0.28, 1.52)	0.33	13	3.01 (0.68, 5.39)	0.07	51
+ O ₃	April–September	0.93 (0.37, 1.49)	0.50	0	3.03 (0.44, 5.68)	0.05	54
PM ₁₀	All year	0.53 (0.06, 1.00)	0.06	48	1.15 (0.21, 2.11)	0.04	52
+ NO ₂	All year	0.19 (–0.42, 0.79)	0.05	50	1.33 (–0.02, 2.69)	0.07	47
+ O ₃	All year	0.50 (–0.03, 1.03)	0.04	52	0.99 (0.13, 1.86)	0.17	32
PM ₁₀	April–September	2.07 (1.30, 2.85)	0.61	0	4.33 (2.59, 6.11)	0.32	14
+ O ₃	April–September	2.25 (1.45, 3.06)	0.83	0	4.98 (1.98, 8.06)	0.07	47

Table 4. Association between PM and hospitalizations for different model specifications, per increases of 10, 6.3, and 14.4 µg/m³ for PM_{2.5}, PM_{2.5–10}, and PM₁₀, respectively.

Pollutant/model	Cardiovascular admissions (lag 0–1)			Respiratory admissions (lag 0–5)			
	Percent increase (95% CI)	χ ² p-value	I ²	Percent increase (95% CI)	χ ² p-value	I ²	
PM _{2.5}	Base model	0.51 (0.12, 0.90)	0.20	29	1.36 (0.23, 2.49)	0.01	65
	Trend, 8 df/year	0.28 (–0.12, 0.68)	0.11	41	0.87 (0.03, 1.72)	0.08	46
	Trend, minimum PACF ^a	0.47 (0.14, 0.80)	0.23	25	1.64 (0.63, 2.65)	0.00	66
	Temperature, lag 0–6 for warm period	0.52 (0.13, 0.91)	0.17	32	1.48 (0.29, 2.69)	0.00	69
PM _{2.5–10}	Base model	0.46 (0.10, 0.82)	0.33	13	1.24 (–0.32, 2.82)	0.00	74
	Trend, 8 df/year	0.34 (0.05, 0.63)	0.91	0	1.10 (0.01, 2.20)	0.05	55
	Trend, minimum PACF ^a	0.53 (0.25, 0.81)	1.00	0	1.69 (0.26, 3.13)	0.00	72
	Temperature, lag 0–6 for warm period	0.57 (0.26, 0.89)	0.50	0	1.40 (–0.27, 3.10)	0.00	76
PM ₁₀	Base model	0.53 (0.06, 1.00)	0.06	48	1.15 (0.21, 2.11)	0.04	52
	Trend, 8 df/year	0.31 (–0.05, 0.67)	0.18	31	0.96 (0.31, 1.61)	0.26	22
	Trend, minimum PACF ^a	0.56 (0.28, 0.84)	0.40	4	1.72 (0.85, 2.59)	0.03	55
	Temperature, lag 0–6 for warm period	0.57 (0.12, 1.02)	0.07	47	1.29 (0.28, 2.32)	0.02	58

^aEstimated degrees of freedom per year were 3 (Milan, Turin, Emilia-Romagna, Bologna, Barcelona), 4 (Rome), and 5 (Marseille, Madrid) for cardiovascular hospitalization; and 3 (Turin, Bologna), 4 (Marseille, Madrid), 5 (Rome), 6 (Milan, Emilia-Romagna), and 9 (Barcelona) for respiratory hospitalizations.

hospitalizations (lag 0–1) and respiratory hospitalizations (lag 0–5). All estimates are reported as percentage increases in hospital admissions (95% CI) associated with increasing concentrations of PM_{2.5} and PM_{2.5–10} relative to 5 µg/m³, and of PM₁₀ relative to 10 µg/m³. Associations of PM_{2.5} and PM₁₀ with cardiovascular hospitalizations seemed to be consistent with two linear functions, with a steeper slope for lower concentrations. There was no visual evidence of a departure from linearity for PM_{2.5–10} and cardiovascular

hospitalizations, or for any of the PM metrics and respiratory admissions.

Discussion

We investigated the association between daily concentrations of fine and coarse particles and hospitalizations for cardiorespiratory conditions in eight Mediterranean cities, finding evidence of harmful effects of the PM metrics on both study outcomes.

Evidence of short-term effects of particulate matter on mortality and morbidity

has been described widely in the literature. Although the first multicenter studies focused on PM₁₀ as the main exposure (Katsouyanni et al. 2001; Samet et al. 2000), more recently interest has shifted to specific PM fractions, constituents, and sources, with the aim of better elucidating the underlying biological mechanisms. It has been argued that the fine fraction of PM is the one more responsible for the health effects, because it includes toxic components such as nitrates, sulfates, acids, and metals originating from combustion




Figure 1. Concentration–response relationship between PM_{2.5} (A), PM_{2.5–10} (B), and PM₁₀ (C) with cardiovascular hospitalizations, lag 0–1 (left), and respiratory hospitalizations, lag 0–5 (right). Values are percent increase (95% CI) of hospital admissions associated with increases of PM_{2.5} and PM_{2.5–10} levels relative to 5 µg/m³, and of PM₁₀ levels relative to 10 µg/m³.

processes, and it can deposit more deeply into the lungs (Pope and Dockery 2006). In contrast, coarse particles are dominated by crustal materials, resuspended dust, sea salts, desert dust, and biogenic components including pollen, spores, and other plant parts (Pope and Dockery 2006). A systematic review conducted by Brunekreef and Forsberg (2005) pointed toward a need to further investigate the health effects of $PM_{2.5-10}$ based on evidence that coarse and fine PM exert similar effects on respiratory outcomes, and on evidence supporting an association between $PM_{2.5-10}$ and cardiovascular outcomes.

Recent multicenter studies focusing on the association between $PM_{2.5}$ and hospitalizations in adult populations have been mainly conducted in North America (Bell et al. 2008; Dominici et al. 2006; Stieb et al. 2009; Zanobetti et al. 2009). Dominici et al. (2006) investigated the effects of $PM_{2.5}$ on cardiovascular and respiratory hospital admissions in 204 U.S. urban counties within the National Morbidity, Mortality, and Air Pollution Study (NMMAPS). They estimated statistically significant increases in hospitalizations associated with a $10\text{-}\mu\text{g}/\text{m}^3$ increase in $PM_{2.5}$ for a number of specific cardiovascular conditions, ranging from 0.44% (95% CI: 0.02, 0.86%) for ischemic heart disease hospitalizations to 1.28% (95% CI: 0.78, 1.78%) for heart failure admissions. Positive associations were also reported for hospital admissions for specific respiratory outcomes (0.91%; 95% CI: 0.18, 1.64% for $PM_{2.5}$ on the same day and chronic obstructive pulmonary disease, and 0.92%; 95% CI: 0.41, 1.43% for respiratory tract infection, with a 2-day lag). Similar results for cardiovascular hospitalizations were reported for another NMMAPS study (Bell et al. 2008) of 202 U.S. counties, which estimated a 0.8% increase in cardiovascular admissions (95% posterior interval: 0.59, 1.01) per $10\text{-}\mu\text{g}/\text{m}^3$ increase in lag 0 $PM_{2.5}$. The same study reported a 0.41% increase (95% posterior interval: 0.09, 0.74) in all respiratory admissions associated with $10\text{-}\mu\text{g}/\text{m}^3$ increase in lag 2 $PM_{2.5}$, a weaker association than reported by Dominici et al. (2006) for specific respiratory conditions. Zanobetti et al. (2009) investigated associations of $PM_{2.5}$ concentrations and components with cause-specific admissions in 26 U.S. communities, and estimated stronger associations with cardiac and respiratory diseases than previous studies: specifically, a 1.89% increase (95% CI: 1.34, 2.45) in cardiac admissions and a 2.07% increase (95% CI: 1.20, 2.95) in respiratory admissions in association with $10\text{-}\mu\text{g}/\text{m}^3$ increase in lag 0–1 $PM_{2.5}$. The authors concluded that “particles originating from industrial combustion processes or traffic may, on average, have greater toxicity.” However, evidence from European studies has been weaker.

One six-city study conducted in France (Host et al. 2008) reported large and statistically significant associations between $PM_{2.5}$ and cardiovascular admissions (0.9% increase; 95% CI: 0.1, 1.8 with a $10\text{-}\mu\text{g}/\text{m}^3$ increase in lag 0–1 $PM_{2.5}$), but no associations between $PM_{2.5}$ and respiratory diseases. Results from single-city European investigations (Anderson et al. 2001; Atkinson et al. 2010; Belleudi et al. 2010; Halonen et al. 2009; Linares and Diaz 2010) have been inconsistent.

Epidemiological studies on the short-term effects of $PM_{2.5-10}$ on hospitalizations are few and inconsistent. In a large U.S. study of 108 counties (Peng et al. 2008), a $10\text{-}\mu\text{g}/\text{m}^3$ increase in $PM_{2.5-10}$ was associated with a 0.36% (95% posterior interval: 0.05, 0.68) increase in cardiovascular disease admissions on the same day. However, when adjusted for $PM_{2.5}$, the association was still positive but no longer statistically significant. Two studies conducted in the United Kingdom (Anderson et al. 2001; Atkinson et al. 2010) did not identify any evidence of effects of coarse particles on hospital admissions for either cardiovascular or respiratory causes. A study conducted in six French cities (Host et al. 2008) found no association between coarse PM and cardiovascular or respiratory admissions, and reported only one statistically significant association with ischemic heart disease in the elderly. We identified positive associations of coarse particles with both cardiovascular (0.46% increase; 95% CI: 0.10, 0.82 with a $6.3\text{-}\mu\text{g}/\text{m}^3$ increase in lag 0–1 $PM_{2.5-10}$) and respiratory admissions (1.24% increase; 95% CI: –0.32, 2.82) with a $6.3\text{-}\mu\text{g}/\text{m}^3$ increase in lag 0–5 $PM_{2.5-10}$. Corresponding associations with a $10\text{-}\mu\text{g}/\text{m}^3$ increase in $PM_{2.5-10}$ were higher than previously reported (0.73%; 95% CI: 0.16, 1.30%, and 1.95%; 95% CI: –0.51, 4.48%, respectively). However, because $PM_{2.5-10}$ is obtained as the difference between PM_{10} and $PM_{2.5}$, it is affected by measurement error from two sources, and the direction and magnitude of the resulting bias in the associations with hospitalizations cannot be predicted.

Our study is, to our knowledge, the first study involving cities from multiple countries in Southern Europe and investigating the health effects of fine and coarse particles on cause-specific hospitalizations. Our estimates of associations between $PM_{2.5}$ and cardiovascular admissions are slightly smaller than those reported by U.S. studies, whereas associations between $PM_{2.5}$ and respiratory admissions, and associations between both outcomes and $PM_{2.5-10}$, are more similar. Several differences must be acknowledged between our study and the North American ones. First, the chemical mixture of $PM_{2.5}$ in Europe is likely to be different from that in the United States. For

example, diesel powered cars make up 50% of the fleet in Europe and only about 2% in the United States. Consistent with this, elemental carbon concentrations estimated for 187 counties in the United States were about $0.6\text{ }\mu\text{g}/\text{m}^3$ (Bell 2012) compared with about $1.5\text{ }\mu\text{g}/\text{m}^3$ for Madrid and Barcelona (unpublished data). Second, PM concentrations for most U.S. studies were measured every 3rd or 6th day, whereas daily PM measurements were available for the eight cities in the present study. In addition, our study, along with the entire MED-PARTICLES project, comprises Southern European cities characterized by highly urbanized areas with intense traffic congestion, elevated sea traffic due to tourist and shipping activities over the Mediterranean area, mild meteorological conditions favoring outdoor activities during most of the year, enhanced formation of secondary pollutants owing to intense solar radiation, high frequency of wildfires, and Saharan dust advection episodes, especially in summer and spring.

Associations between fine particles and cardiovascular hospitalizations in our study were not affected by coarse PM co-exposure, but the associations of both fine and coarse PM with respiratory admissions decreased to nonsignificance when evaluated together in two-pollutant models. Because of moderate to high correlations between the two exposures, it is difficult to disentangle their potential effects. This problem was even more apparent in two-pollutant models involving $PM_{2.5}$ and NO_2 . The respiratory effects were highly heterogeneous, and there are no simple explanations for that. These heterogeneous effects were already seen in a large Italian study (Faustini et al. 2013). City-specific prevalence of patients with chronic respiratory conditions may be different, as may their individual level of response to air pollutants on different seasons.

An additional contribution of the present study is the strategy for time-trend adjustment. We applied three different approaches to adjust for the confounding effect of long-term and seasonal time trends: a case-crossover approach that used three-way interaction terms among year, month, and day of the week (84 df per year) to adjust for time trends; a penalized spline and a smoothing parameter with 8 df per year; and a penalized spline with df per year selected to minimize PACF residuals (3–5 df per year for cardiovascular admissions, and 3–9 df per year for respiratory admissions). Despite the variation in the degrees of control, the three methods provided consistent results, suggesting little residual confounding due to long-term and seasonal time trends.

We estimated much stronger associations of fine and coarse particles with hospitalizations for both cardiovascular and respiratory conditions during the warm period compared

with the cold period. This is consistent with the worldwide literature (Faustini et al. 2013; Strieb et al. 2009; Strickland et al. 2010). Two complementary hypotheses have been suggested: First, exposures to air pollutants may increase during warm months because of increased outdoor activities and open windows (Stafoggia et al. 2008); second, the air pollution mixture may include a greater proportion of toxic components during warmer months (Peng et al. 2005). Both explanations are plausible during the warm season in the Mediterranean area because of seasonal differences in PM composition (Perrino et al. 2009; Querol et al. 2009; van Drooge et al. 2012) and an increased frequency of wildfires and Saharan dust episodes, whose particles have been related to health effects (Hänninen et al. 2009; Karanasiou et al. 2012). These aspects have been already documented in the Southern Mediterranean countries (Querol et al. 2009), and further results will be available from the MED-PARTICLES project.

To our knowledge, this is the first study to investigate the concentration–response functions of fine and coarse particles with cardiovascular and respiratory morbidity. Epidemiological studies of the short-term effects of PM₁₀ on cause-specific mortality (Daniels et al. 2004; Faustini et al. 2011; Katsouyanni et al. 2009) have reported no apparent departures from linearity. An extension of the Harvard Six Cities Study also reported no evidence of a departure from linearity for the relationship between long-term exposure to PM_{2.5} and survival (Schwartz et al. 2008). In the present study we found a suggestion of linear effects of all PM metrics on respiratory hospitalizations, and of PM_{2.5–10} on cardiovascular admissions. However, the slopes of estimated exposure–response curves for cardiovascular admissions in association with PM₁₀ and PM_{2.5} were steeper for low to moderate concentrations (approximately 30 µg/m³ for PM_{2.5} and 60 µg/m³ for PM₁₀) than higher concentrations, consistent with the dose–response curve previously estimated for the long-term effects of PM_{2.5} on mortality from ischemic heart disease (Pope et al. 2009). These results need further investigation in other locations and with other analytical approaches, but they support effects of all PM metrics on cardiovascular and respiratory conditions at low concentrations, including concentrations below the current EU air quality standards.

We believe that our results have clear implications for the EU policy on air quality. Specifically, they suggest that the current 24-hr daily limit value for PM₁₀ in Europe is not sufficient to protect the population from short-term effects, and that a daily limit value for PM_{2.5} is clearly warranted. Both PM_{2.5} and PM₁₀ daily limits are needed to control

air pollution generated from different sources: vehicle exhausts and combustion sources for fine particles, natural sources together with resuspension of road dust (containing a mixture of soil, tire wear, and brake wear), non-exhaust emissions, and commercial and industrial residues for coarse particles. Because coarse particles of natural origin cannot be controlled, policy measures aimed at controlling anthropogenic sources of coarse particles would be advisable. Alternatively, a specific short-term limit value for coarse particles (PM_{2.5–PM10}) may be considered, but only in addition to an effective PM_{2.5} daily limit.

Conclusions

In summary, we estimated significant short-term effects of fine and coarse particles on cardiovascular and respiratory hospitalizations in eight Southern European cities. Associations were similar across PM metrics, stronger but more heterogeneous for respiratory admissions, much more pronounced during the warm period, and robust to time-trend specification. In addition, we did not find strong evidence of departure from linearity in the range of pollutant concentrations measured for the associations under investigation. These findings will help inform planned revisions of EU air quality standards and support legislation on daily PM_{2.5} concentrations to better target policies on air pollution in Europe.

REFERENCES

- Anderson HR, Bremner SA, Atkinson RW, Harrison RM, Walters S. 2001. Particulate matter and daily mortality and hospital admissions in the west Midlands conurbation of the United Kingdom: associations with fine and coarse particles, black smoke and sulphate. *Occup Environ Med* 58:504–510.
- Atkinson RW, Fuller GW, Anderson HR, Harrison RM, Armstrong B. 2010. Urban ambient particle metrics and health: a time-series analysis. *Epidemiology* 21:501–511.
- Bell ML. 2012. Assessment of the health impacts of particulate matter characteristics. HEI Health Review Committee. *Res Rep Health Eff Inst* 161:5–38.
- Bell ML, Ebisu K, Peng RD, Walker J, Samet JM, Zeger SL, et al. 2008. Seasonal and regional short-term effects of fine particles on hospital admissions in 202 US counties, 1999–2005. *Am J Epidemiol* 168:1301–1310.
- Belleudi V, Faustini A, Stafoggia M, Cattani G, Marconi A, Perucci CA, et al. 2010. Impact of fine and ultrafine particles on emergency hospital admissions for cardiac and respiratory diseases. *Epidemiology* 21:414–423.
- Brunekreef B, Forsberg B. 2005. Epidemiological evidence of effects of coarse airborne particles on health. *Eur Respir J* 26:309–318.
- Chiusolo M, Cadum E, Stafoggia M, Galassi C, Berti G, Faustini A, et al. 2011. Short-term effects of nitrogen dioxide on mortality and susceptibility factors in 10 Italian cities: the EpiAir Study. *Environ Health Perspect* 119:1233–1238; doi:10.1289/ehp.1002904.
- Daniels MJ, Dominici F, Zeger SL, Samet JM. 2004. The National Morbidity, Mortality, and Air Pollution Study. Part III: PM₁₀ concentration-response curves and thresholds for the 20 largest US cities. *Res Rep Health Eff Inst* 94 (pt 3):1–21.
- Dominici F, Peng RD, Bell ML, Pham L, McDermott A, Zeger SL, et al. 2006. Fine particulate air pollution and hospital admission for cardiovascular and respiratory diseases. *JAMA* 295:1127–1134.
- Faustini A, Stafoggia M, Berti G, Bisanti L, Chiusolo M, Cernigliaro A, et al. 2011. The relationship between ambient particulate matter and respiratory mortality: a multicenter study in Italy. *Eur Respir J* 38:538–547.
- Faustini A, Stafoggia M, Colais P, Berti G, Bisanti L, Cadum E, et al. 2013. Air pollution and multiple acute respiratory outcomes. *Eur Respir J*; doi:10.1183/09031936.00128712 [Online 11 January 2013].
- Halonen JI, Lanki T, Yli-Tuomi T, Tiittanen P, Kulmala M, Pekkanen J. 2009. Particulate air pollution and acute cardiorespiratory hospital admissions and mortality among the elderly. *Epidemiology* 20:143–153.
- Hänninen O, Salonen RO, Koistinen K, Lanki T, Barregard L, Jantunen M. 2009. Population exposure to fine particles and estimated excess mortality in Finland from an East European wildfire episode. *J Expo Sci Environ Epidemiol* 19:414–422; doi:10.1038/jes.2008.31.
- Higgins JP, Thompson SG. 2002. Quantifying heterogeneity in a meta-analysis. *Stat Med* 21:1539–1558.
- Host S, Larrieu S, Pascal L, Blanchard M, Declercq C, Fabre P, et al. 2008. Short-term associations between fine and coarse particles and hospital admissions for cardiorespiratory diseases in six French cities. *Occup Environ Med* 65:544–551.
- Jackson D, White IR, Thompson SG. 2010. Extending DerSimonian and Laird's methodology to perform multivariate random effects meta-analyses. *Stat Med* 29:1282–1297.
- Janes H, Sheppard L, Lumley T. 2005. Case-crossover analyses of air pollution exposure data: referent selection strategies and their implications for bias. *Epidemiology* 16:717–726.
- Karanasiou A, Moreno N, Moreno T, Viana M, de Leeuw F, Querol X. 2012. Health effects from Sahara dust episodes in Europe: literature review and research gaps. *Environ Int* 47:107–114.
- Katsouyanni K, Samet JM, Anderson HR, Atkinson R, Le Tertre A, Medina S, et al. 2009. Air Pollution and Health: a European and North American Approach (APHENA). HEI Health Review Committee. *Res Rep Health Eff Inst* 142:5–90.
- Katsouyanni K, Touloumi G, Samoli E, Gryparis A, Le Tertre A, Monopoli Y, et al. 2001. Confounding and effect modification in the short-term effects of ambient particles on total mortality: results from 29 European cities within the APHEA2 project. *Epidemiology* 12:521–531.
- Levy D, Lumley T, Sheppard L, Kaufman J, Checkoway H. 2001. Referent selection in case-crossover analyses of acute health effects of air pollution. *Epidemiology* 12:186–192.
- Linares C, Diaz J. 2010. Short-term effect of PM_{2.5} on daily hospital admissions in Madrid (2003–2005). *Int J Environ Health Res* 20:129–140.
- Lu Y, Zeger SL. 2007. On the equivalence of case-crossover and time series methods in environmental epidemiology. *Biostatistics* 8:337–344.
- Maclure M. 1991. The case-crossover design: a method for studying transient effects on the risk of acute events. *Am J Epidemiol* 133:144–153.
- MED-PARTICLES (Particles size and composition in Mediterranean countries: geographical variability and short-term health effects). 2010. MED-PARTICLES. Available: <http://www.epidemiologia.lazio.it/medparticles/index.php/en/> [accessed 24 September 2012].
- Ostro B, Broadwin R, Green S, Feng WY, Lipssett M. 2006. Fine particulate air pollution and mortality in nine California counties: results from CALFINE. *Environ Health Perspect* 114:29–33; doi:10.1289/ehp.8335.
- Peng RD, Chang HH, Bell ML, McDermott A, Zeger SL, Samet JM, et al. 2008. Coarse particulate matter air pollution and hospital admissions for cardiovascular and respiratory diseases among Medicare patients. *JAMA* 299:2172–2179.
- Peng RD, Dominici F, Pastor-Barriuso R, Zeger SL, Samet JM. 2005. Seasonal analyses of air pollution and mortality in 100 US cities. *Am J Epidemiol* 161:585–594.
- Perrino C, Catrambone M, Esposito G, Lahav D, Mamane Y. 2009. Characterisation of gaseous and particulate atmospheric pollutants in the east Mediterranean by diffusion denuder sampling lines. *Environ Monit Assess* 152:231–244.
- Pope CA III, Burnett RT, Krewski D, Jerrett M, Shi Y, Calle EE, et al. 2009. Cardiovascular mortality and exposure to airborne fine particulate matter and cigarette smoke: shape of the exposure-response relationship. *Circulation* 120:941–948.

- Pope CA III, Dockery DW. 2006. Health effects of fine particulate air pollution: lines that connect. *J Air Waste Manag Assoc* 56:709–742.
- Querol X, Alastuey A, Pey J, Cusack M, Pérez N, Mihalopoulos N, et al. 2009. Variability in regional background aerosols within the Mediterranean. *Atmos Chem Phys* 9:4575–4591.
- Samet JM, Dominici F, Currier FC, Coursac I, Zeger SL. 2000. Fine particulate air pollution and mortality in 20 U.S. cities, 1987–1994. *N Engl J Med* 343:1742–1749.
- Samoli E, Touloumi G, Zanobetti A, Le Tertre A, Schindler C, Atkinson R, et al. 2003. Investigating the dose-response relation between air pollution and total mortality in the APHEA-2 multicity project. *Occup Environ Med* 60:977–982.
- Schwartz J, Coull B, Laden F, Ryan L. 2008. The effect of dose and timing of dose on the association between airborne particles and survival. *Environ Health Perspect* 116:64–69; doi:10.1289/ehp.9955.
- Schwartz J, Zanobetti A. 2000. Using meta-smoothing to estimate dose-response trends across multiple studies, with application to air pollution and daily death. *Epidemiology* 11:666–672.
- Stafoggia M, Forastiere F, Faustini A, Biggeri A, Bisanti L, Cadum E, et al. 2010. Susceptibility factors to ozone-related mortality: a population-based case-crossover analysis. *Am J Respir Crit Care Med* 182:376–384.
- Stafoggia M, Schwartz J, Forastiere F, Perucci CA, SISTI Group. 2008. Does temperature modify the association between air pollution and mortality? A multicity case-crossover analysis in Italy. *Am J Epidemiol* 167:1476–1485.
- Stieb DM, Szyszkowicz M, Rowe BH, Leech JA. 2009. Air pollution and emergency department visits for cardiac and respiratory conditions: a multi-city time-series analysis. *Environ Health* 8:25; doi:10.1186/1476-069X-8-25.
- Strickland MJ, Darrow LA, Klein M, Flanders W, Sarnat JA, Waller LA, et al. 2010. Short-term associations between ambient air pollutants and pediatric asthma emergency department visits. *Am J Respir Crit Care Med* 182:307–316.
- U.S. Environmental Protection Agency. 2012. National Ambient Air Quality Standards (NAAQS). Available: <http://www.epa.gov/air/criteria.html> [accessed 12 July 2013].
- van Drooge BL, Lopez JF, Grimalt JO. 2012. Influences of natural emission sources (wildfires and Saharan dust) on the urban organic aerosol in Barcelona (Western Mediterranean Basin) during a PM event. *Environ Sci Pollut Res Int* 19:4159–4167.
- Whitaker HJ, Hocine MN, Farrington P. 2007. On case-crossover methods for environmental time series data. *Environmetrics* 18:157–171.
- WHO (World Health Organization). 1999. International Classification of Diseases (ICD). Available: <http://www.who.int/classifications/icd/en/> [accessed 12 June 2013].
- Zanobetti A, Franklin M, Koutrakis P, Schwartz J. 2009. Fine particulate air pollution and its components in association with cause-specific emergency admissions. *Environ Health* 8:58; doi:10.1186/1476-069X-8-58.
- Zanobetti A, Schwartz J. 2009. The effect of fine and coarse particulate air pollution on mortality: a national analysis. *Environ Health Perspect* 117:898–903; doi:10.1289/ehp.0800108.