

Meta-Analysis of TPB/TRA studies predicting screening attendance

Running Head. Meta-Analysis of TPB/TRA studies predicting screening attendance

How well do the theory of reasoned action and theory of planned behaviour predict intentions and attendance at screening programmes? A meta-analysis

Dr Richard Cooke, Lecturer in Health Psychology, School of Life & Health Sciences, Psychology, Aston University, Aston Triangle, Birmingham, B4 7ET.

Dr David P French, Reader in Health Behaviour Interventions, Applied Research Centre in Health & Lifestyle Interventions, Faculty of Health & Life Sciences, Coventry University, Priory Street, Coventry CV1 5FB

Please reference as **Cooke, R., & French, D. P.** (2008). How well do the theory of reasoned action and the theory of planned behaviour predict intentions and attendance at screening programmes? A meta-analysis. *Psychology & Health, 23*, 745-765.

Please direct correspondence to:

Richard Cooke

School of Life & Health Sciences, Psychology,

Aston University

Birmingham

B4 7ET

r.cooke@aston.ac.uk

Meta-Analysis of TPB/TRA studies predicting screening attendance

ABSTRACT

Meta-analysis was used to quantify how well the Theories of Reasoned Action and Planned Behaviour have predicted intentions to attend screening programmes and actual attendance behaviour. Systematic literature searches identified 33 studies that were included in the review. Across the studies as a whole, attitudes had a large-sized relationship with intention, while subjective norms and perceived behavioural control (PBC) possessed medium-sized relationships with intention. Intention had a medium-sized relationship with attendance whereas the PBC-attendance relationship was small-sized. Due to heterogeneity in results between studies, moderator analyses were conducted. The moderator variables were (a) type of screening test, (b) location of recruitment, (c) screening cost and (d) invitation to screen. All moderators affected TPB relationships. Suggestions for future research emerging from these results include targeting attitudes to promote intention to screen, a greater use of implementation intentions in screening information, and examining the credibility of different screening providers.

Keywords: Screening attendance, Theory of Reasoned Action, Theory of Planned Behaviour, intention, mammography, cervical

Meta-Analysis of TPB/TRA studies predicting screening attendance

How well do the theory of reasoned action and theory of planned behaviour predict intentions and attendance at screening programmes? A meta-analysis

The main aim of screening programmes is to identify people at high risk of a particular condition, to allow interventions to prevent that condition developing or progressing. A major determinant of the effectiveness of all programmes is the level of uptake of those programmes. A high level of attendance at screening programmes is a pre-requisite if screening programmes are to have a significant impact on population morbidity and mortality.

Despite the importance of attendance, rates of attendance for screening programmes vary widely, and are often low, even when screening is offered free of charge. For example, breast screening is offered free of charge to all women between the ages of 50 and 64 years in England, and achieved nearly 75% coverage in 2003-04 (Department of Health, 2005). However, coverage was less than 60% in 24 out of 303 primary health care organisations, and less than 50% in 8 of these. A good deal of research has been conducted to explore the demographic, economic, motivational and organisational factors which predict variation in attendance at screening programmes (Jepson, Clegg, Forbes, Lewis, Sowden & Kleijnen, 2000). The focus of this paper is on motivational factors which predict attendance behaviour.

Two models of human behaviour which have been extensively utilised to predict health behaviour (Johnston, French, Bonetti & Johnston, 2004), such as screening attendance, are the theory of planned behaviour (TPB: Ajzen, 1991) and its predecessor the theory of reasoned action (TRA: Ajzen & Fishbein, 1980; Fishbein & Ajzen, 1975). Both models propose that the most important determinant of any behaviour is a person's

Meta-Analysis of TPB/TRA studies predicting screening attendance

intention to perform the behaviour. In the TPB, perceived behavioural control (PBC), i.e., an individual's perceptions of control over behavioural performance in the face of internal and external barriers, is also specified as an additional predictor of behaviour. Intentions are predicted by attitudes, subjective norms and, in the TPB, PBC. Attitudes are an individual's positive or negative evaluation of performing the behaviour. Subjective norms reflect an individual's perceptions of social approval for performing the behaviour.

Meta-analytic reviews have supported the capacity of both the TPB (Armitage & Conner, 2001; Godin & Kok, 1996) and the TRA (Shepherd, Hartwick & Warshaw, 1988) to predict intentions and behaviour across a wide range of contexts. Godin and Kok (1996) reviewed the literature applying the TPB to health behaviours, including a supplementary analysis of the TPB applied to attendance at screening programmes. They found that intentions to attend screening correlated strongly with attitudes ($r_+ = .51$), but more moderately with subjective norms ($r_+ = .33$) and PBC ($r_+ = .46$) in eight studies. Across six studies, they reported sample-weighted average correlations of $r_+ = .35$ between intention and behaviour and $r_+ = .29$ between PBC and behaviour. These results are similar to those reported by meta-analyses of the TPB with all behaviours, based on many more studies (Armitage & Conner, 2001). In contrast, other models of health-related behaviour, such as Protection Motivation Theory and the Health Belief Model, tend to account for less variance in intentions and behaviour (see Conner & Norman, 2005 for a review of each theory).

However, there are three critical limitations with Godin and Kok's review. First, the results are based on papers available up to 1994. Therefore, the review does not cover

Meta-Analysis of TPB/TRA studies predicting screening attendance

the last decade of research, which has seen a dramatic increase in the quantity and quality of studies employing the TPB to study attendance at screening programmes. Second, due to the small number of papers included, the review of Godin and Kok did not have sufficient power to investigate potentially important moderator variables, e.g., whether relationships differ between systems of organising screening programmes. Third, not all papers included in the review examined attendance at a screening programme: One examined doctors' decisions to perform a clinical exam on HIV positive patients (Godin, Boyer, Duval, Fortin & Nadeau, 1992), one tested the use of oral rehydration therapy (Hounsa, Godin, Alihonou, Valois & Girard, 1993) and one investigated dental hygiene behaviours (McCaul, O'Neill, & Glasgow, 1988).

The main aim of the present study is to provide a comprehensive and up-to-date quantitative review of applications of the TRA/TPB in the domain of screening attendance. Specifically, the present study examines the strength of five relationships within the TRA/TPB—attitude-intention, subjective norm-intention, PBC-intention, intention-behaviour, PBC-behaviour—in the context of individuals attending a health screening programme. The second aim is to investigate the extent to which several moderator variables affect the size of the relationships just listed: (a) the type of screening test carried out (e.g., mammogram, cervical smear), (b) the location of recruitment for the study (e.g., general practice (GP), hospital), (c) the cost of screening, either screening was paid for as part of a privately organised system of healthcare (as in the United States) or free as part of a national healthcare system (e.g., in the UK) and (d) whether or not participants received an invitation to screen.

METHOD

Meta-Analysis of TPB/TRA studies predicting screening attendance

Literature review and inclusion/ exclusion criteria

Several methods were used to identify relevant studies: (a) computerised searches of social scientific and medical databases (BIDS ISI, PubMed, Psychinfo and Web of Science) from January 1981 to the time of writing (July 2006), (b) studies cited in each article retrieved, and (c) the authors of published articles were contacted and requests were made for studies in press. Searches were conducted by combining the following keywords ‘screening’, ‘mammograph*’, ‘cervical’, ‘health check/screening’, and ‘attend*’ with ‘theory of reasoned action’ and ‘theory of planned behavior*’ (to allow for US/UK differences in spelling of behaviour). These searches yielded 156 independent papers, which were considered against the following inclusion criteria:

1. Studies had to report data on predicting attendance at screening programmes or predicting intention to attend screening. Hence, we included studies that looked at screening for cancer (breast, cervical, colon) as well as at health checks, genetic screening, prenatal screening, diabetes screening and screening for tuberculosis. However, we did not include studies of behaviours such as breast self examination, which did not involve attending for a screening test. Attendance was measured objectively from medical records in every study except Poss (2000) where participants were followed up two days after completing measures of TPB variables.
2. As a minimum, studies had to include both TRA constructs as predictors of intention. This led to the exclusion of (a) review papers, (b) papers that measured TRA/TPB variables but did not report the relationships between variables (e.g., Fernbach, 2002) and (c) papers that tested only the attitude-intention relationship

Meta-Analysis of TPB/TRA studies predicting screening attendance

- (e.g., McCaffery, Wardle & Waller, 2003). Papers that only reported the intention-behaviour relation were also excluded.
3. A bivariate statistical relationship between cognitions and intention (or behaviour) had to be retrievable. Where reports did not include relevant statistics (e.g., Barling & Moore, 1996), the authors of the study were contacted and requests were made for bivariate associations.

Using these inclusion criteria, the following were included: 33 tests of the association between attitudes and intentions, 31 tests of the association between subjective norm and intention and 25 tests of the relationship between PBC and intention. There were 19 tests of the relationship between intentions and behaviour and 18 tests of the PBC-behaviour relationship. Table 1 presents the bivariate correlations of intention with attitude, subjective norm and PBC, and Table 2 presents the bivariate correlations between attendance and intention, and attendance and PBC.

Meta-analytic strategy

The effect size estimate employed here was an average of the sample correlation coefficients, r_{+} , weighted by sample size. In this way correlations based on larger samples receive greater weight than those from smaller samples (see Hedges & Olkin, 1985). We calculated 'Fail-Safe N ' (FSN; Rosenthal, 1984) values for each relationship, which provide an estimate of the number of unpublished studies comparable in size but containing null results that would be required to invalidate the conclusion that a relationship is statistically significant.

Homogeneity analyses were conducted using the chi-square statistic (Hunter et al., 1985) to determine whether variation in the correlations between studies was greater than

Meta-Analysis of TPB/TRA studies predicting screening attendance

chance. If the chi-square statistic is non-significant, then the correlations are homogeneous and the average weighted effect size, \bar{r}_+ , can be said to represent the best estimate of population effect size. If the overall χ^2 statistic was significant, pairwise Z tests were conducted between all pairs of tests to determine which screening tests in particular yielded significant different effect size estimates.

Computation of the weighted average correlations and homogeneity analyses were all conducted using Schwarzer's (1988) Meta computer program. All analyses are described in line with Cohen's (1992) recommendations where a correlation of $r = .50$ is a large-sized effect, $r = .30$ is a medium-sized effect and $r = .10$ a small-sized effect.

Multiple Samples and Multiple Measures

Where studies reported separate statistical tests for more than one sample, then the correlation from each sample was used as the unit of analysis (e.g., Braithwaite, Sutton & Steggle, 2002). Where studies had measured attendance on multiple occasions (e.g., Drossaert, Boer, & Seydel, 2003) we employed the conservative strategy of using the weighted average of the sample correlations and the smallest N in the analysis to determine the overall effect size for the study.

RESULTS

For the prediction of intention, all relationships were large or medium-sized, with the strongest relationship between attitudes and intentions and the weakest between subjective norms and intentions (Table 3). For predicting screening attendance, intention had a medium-sized relationship whereas PBC had a small-sized relationship (Table 3). All of the FSN values were large, and since it is unlikely so many unpublished studies exist, we can conclude that each population relationship is genuinely different from zero.

Meta-Analysis of TPB/TRA studies predicting screening attendance

Moderator analyses

Tests for heterogeneity for each of the correlations reported were significant (see Table 3), suggesting greater variability in effect size estimates between studies than would be expected on the basis of random sampling error alone. This encourages a search for moderator variables, which can explain this variation. We conducted four moderator analyses, specifically were TPB relationships affected by (a) the type of screening the participant was invited to, (b) the location of recruitment, (c) cost of screening and (d) whether participants received an invitation to screen or not

(a) Type of Screening Test

The TRA/TPB has been applied to mammography ($k = 10$), cervical smear ($k = 8$), health check ($k = 4$), genetic ($k = 3$), colorectal ($k = 3$) and prenatal ($k = 2$) screening attendance. Only Poss' (2000) study of tuberculosis inoculation and Orbell and Hagger's (2006a) study of diabetes testing did not fit into one of these categories. Data were analysed separately according to test context for the five TRA/TPB relationships tested above. We first tested whether the effect sizes were heterogeneous for each relation (e.g., attitude-intention) between the different types of test (Table 4).

There was a significant chi-square value associated with the attitude-intention relationships reported for the six different tests ($\chi^2(5) = 345.32, p < .001$). The effect reported for prenatal screening ($r_+ = .79$) was significantly larger than all the other attitude-intention relationships reported. The effect sizes associated with genetic tests and health checks were significantly larger than the three remaining tests. Finally, there were no differences between the effect sizes reported in mammography, cervical smear and colorectal screening studies.

Meta-Analysis of TPB/TRA studies predicting screening attendance

The size of the subjective norm-intention relationship was different according to the test context considered ($\chi^2 (5) = 350.77, p < .001$). Prenatal subjective norms were a significantly better predictor of screening intentions than for any other test ($r_+ = .69$). There were no differences between the size of the relationships reported for genetic tests, health checks and colorectal screening although these relationships were all significantly larger than those reported for mammography and cervical smear. The effect size reported for cervical smear was significantly higher than that reported for mammography.

The size of the relationship between PBC and intention differed depending on test type ($\chi^2 (4) = 168.51, p < .001$). Effect sizes for cervical smear and colorectal screening were significantly greater than the effects reported for other tests, which did not differ from each other. Effect sizes for health check and mammography were identical and both were significantly greater than the effect reported in prenatal contexts.

Intention differed in its effectiveness as a predictor of behaviour depending on the test studied ($\chi^2 (4) = 494.56, p < .001$). Prenatal intentions were a significantly better predictor of attendance ($r_+ = .78$) compared to all other intentions. Colorectal and mammography intentions were both better predictors of attendance than health check and cervical smear intentions. There were no other differences. PBC did not differ in its prediction of attendance depending on the type of screening ($\chi^2 = 6.86, ns$).

(b) Location of Recruitment

Further analyses examined whether the size of TPB relationships varied as a result of where participants were recruited. In some studies, participants were contacted by their general practice (GP) surgery. Alternatively, participants were recruited as part of a national screening programme, or when they attended hospital for another procedure.

Meta-Analysis of TPB/TRA studies predicting screening attendance

Finally, in some studies, participants were recruited from other settings such as universities, churches or community centres. All studies could be categorised as either GP ($k = 8$), hospital ($k = 9$), health authority ($k = 6$) or other ($k = 10$) based on information provided in the method section of the primary studies (Table 5).

Attitudes had the strongest relationship with intentions in GP studies ($r_+ = .60$), followed by hospitals ($r_+ = .53$), and health authorities ($r_+ = .50$), with other studies having the weakest relationship ($r_+ = .40$). The largest subjective norm-intention relationship was also found in GP settings ($r_+ = .54$). The next largest relationship was hospitals ($r_+ = .49$), which was significantly larger than other studies ($r_+ = .41$) and those recruited through health authorities ($r_+ = .27$). Other studies had the greatest PBC-intention consistency ($r_+ = .60$). There was no difference between health authorities ($r_+ = .46$) or GP practices ($r_+ = .44$) in the size of their PBC-intention relation, though both of these relationships were stronger than those reported in hospital studies ($r_+ = .39$). Intention predicted attendance best when participants were recruited via hospitals ($r_+ = .57$). This relationship was significantly larger than the effects reported for health authorities ($r_+ = .34$) or GP practices ($r_+ = .23$), which also significantly differed from each other. There were no differences in the results for PBC as a predictor of attendance.

(c) Screening Cost

In the majority of studies included in the review, particularly those conducted in Europe, participants did not have to pay for screening. By contrast, in the United States, screening is not provided free of charge by a national healthcare system and individuals must have health insurance to qualify for tests.

Meta-Analysis of TPB/TRA studies predicting screening attendance

All of the studies included in the review could be classed as measuring the impact of the TPB in free ($k = 25$) or paid contexts ($k=8$) based on information provided in the method section of the primary studies (Table 6). In free contexts, attitude had a large-sized relationship with intention ($r_+ = .53$) which was significantly greater than the medium-to-large-sized relationship reported in paid contexts ($r_+ = .42$). There was no difference in the subjective norm-intention relationship found in paid ($r_+ = .43$) and free ($r_+ = .40$) contexts. The PBC-intention relationship was medium-to-large sized in free contexts ($r_+ = .47$) which was significantly greater than the medium-sized relationship found in paid contexts ($r_+ = .38$).

Both intention and PBC were better predictors of attendance in paid rather than free contexts. For intention, the large-sized relationship with attendance in paid contexts ($r_+ = .58$) was significantly greater than the medium-to-large-sized relationship found in free contexts ($r_+ = .40$). The PBC-attendance relationship was significantly greater in paid contexts ($r_+ = .34$) relative to free contexts ($r_+ = .17$).

(d) Invitation to screen

Studies differed in whether participants were sent an invitation to attend screening ($k = 20$) or were not sent an invitation ($k = 13$). Receiving an invitation may act as a helpful cue to action, so we were interested to see if this factor moderated TPB relationships (see Table 7). We report the results for the three predictors of intention because, in all but one study (Poss, 2000) that measured behaviour, all participants were sent an invitation. Attitudes were significantly stronger predictors of intentions in the invitation studies ($r_+ = .55$) compared to the no invitation studies ($r_+ = .39$). For subjective norms, there was a stronger relationship with intentions in the no invitation studies ($r_+ =$

Meta-Analysis of TPB/TRA studies predicting screening attendance

.44) compared to the invitation studies ($r_+ = .39$), and this pattern was repeated for the PBC-intention relationship, with greater PBC-intention consistency in the no invitation studies ($r_+ = .57$) compared to the no invitation studies ($r_+ = .44$).

There was considerable heterogeneity in the results for the invitation studies. This variation could be due to differences in when participants received the questionnaire. Examination of the method sections of the papers indicated that questionnaires were either sent before the invitation ($k = 10$), at the same time as the invitation ($k = 3$) or after the invitation ($k = 6$). We performed additional analyses comparing the TPB relationships by timing of questionnaire in relation to screening invitation (see Table 8).

When participants completed questionnaires after, or at the same time as, receiving an invitation there was a stronger attitude-intention relationship (After $r_+ = .60$, Same Time $r_+ = .61$) compared to participants who answered questionnaires before their invitation ($r_+ = .51$). Receiving the invitation at the same time as completing the questionnaire produced a large subjective-norm intention relationship ($r_+ = .51$), with medium sized effects for questionnaires before ($r_+ = .42$) and after ($r_+ = .37$). PBC was a significantly better predictor of intentions when measured after receiving an invitation ($r_+ = .45$) opposed to completing the questionnaire before receiving the invitation ($r_+ = .40$). Completing questionnaires after receiving an invitation lead to greater intention-behaviour consistency (After $r_+ = .47$ vs. Before $r_+ = .31$) and PBC-behaviour consistency (After $r_+ = .22$ vs. Before $r_+ = .15$)

DISCUSSION

The present paper describes a meta-analysis of research that has applied the TRA/TPB to predicting screening intentions and screening attendance. Overall, attitudes

Meta-Analysis of TPB/TRA studies predicting screening attendance

had a large-sized relationship with intention, and subjective norms and PBC both had medium sized relationships with intention. In addition, intention had a medium-sized relationship with attendance whereas PBC had a small-sized relationship with attendance.

These relationships are consistent with those reported in previous meta-analyses of the TPB/TRA (e.g., Armitage & Conner, 2001; Godin & Kok 1996). For example, all three meta-analyses reported a large sized attitude-intention relationship and medium-sized subjective norm-intention, PBC-intention, and intention-behaviour relationships. The only difference between the papers was the size of the PBC-behaviour relationship, which was small-sized in the present paper and medium-sized in previous papers.

Type of screening test

The attitude-intention relationship appears strongest for tests which could be seen as having the largest effect on people other than those being screened (prenatal screening and genetic testing). In both these cases, values other than personal health are at stake, i.e., those to do with unborn children and other family members. Given that health can be seen as an unambiguous good, the wider variation in these other values that are considered in these contexts appear to leads to a greater impact of attitudes.

Subjective norms also appear particularly strong predictors of intentions in prenatal and genetic test settings. This may be because there is an ethical dimension to these decisions; prenatal decisions affect the woman, her partner and her baby, while genetic testing may occur because of previous family experience of a condition that is more likely due to genetic similarity. Hence, the impact of normative pressure is strongest for these types of screening.

Meta-Analysis of TPB/TRA studies predicting screening attendance

The medium-sized subjective norm-intention relation for mammography was the smallest for any of the types of screening test. One explanation for this may be that women do not vary much in their perceptions of normative pressure when considering mammography screening. In other words, most women perceive mammography screening as something that most of their important others would want them to do and hence there is limited variability in their responses, reducing the prediction of intentions.

PBC had large-sized relationships with intentions for colorectal and cervical screening. As colorectal screening was not part of any national screening programme in the studies examined, and is down to the individual to perform, individuals may perceive more barriers relative to other tests and this may explain the strong effect for PBC. Although cervical screening is offered free of charge to women in many countries, there appear to be other barriers to attendance. Fernbach (2002) found that under-screened women perceived barriers to screening, such as remembering to attend, as more difficult to overcome than well-screened women.

PBC appears a weaker predictor of prenatal screening intentions. Michie et al. (2004) comment in their study that PBC may have been a poor predictor of intentions because the women completed the questionnaire whilst in a hospital context and may have been unaware of barriers which may prevent them attending screening at a later date. In other words, their PBC may have been a less accurate reflection of the actual control they had over attendance.

Intentions were better predictors of prenatal screening attendance compared to other screening tests. One explanation for this difference is that attending prenatal screening is a one-off behaviour performed when a woman is pregnant. In contrast,

Meta-Analysis of TPB/TRA studies predicting screening attendance

mammography screening occurs as part of a screening cycle every few years, and frequently performed behaviours are often associated with weaker intention-behaviour relations compared to infrequently performed behaviours (Ouellette & Wood, 1998).

PBC was a relatively unimportant predictor of attendance behaviour, regardless of test. Screening attendance may be an example of a behaviour where perceptions of control do not reflect actual control and as a result are ineffective predictors of behaviour (cf. Ajzen, 1988). It is also likely that PBC may be very high and, because of limited variability in PBC responses, prediction of behaviour may be low.

Location of Recruitment

Prediction of intentions by attitudes and subjective norms was strongest in studies that recruited via GP practices. The large attitude-intention relationship for GP settings suggests that individuals may value visits to their GP more than visits to more remote, less familiar, settings such as hospitals. The large subjective norm-intention relation for GP settings may be a reflection of participants having family who attend the same practice and/or having a good relationship with the GPs at their local practice.

PBC-intention relationships were highest in studies where participants were recruited in a setting where screening does not take place, (e.g., university, community centre). Ajzen (1991) argues that as control over behaviour becomes harder, then PBC becomes a stronger predictor of intentions and behaviour. When recruitment occurs in settings where screening does not take place, this may increase the difficulty of attendance, and increase the strength of the PBC-intention relation. In contrast, when participants are recruited through hospitals, there are likely to be fewer control issues and

Meta-Analysis of TPB/TRA studies predicting screening attendance

the PBC-intention relation is reduced. This is consistent with our finding that the PBC-intention relation for hospitals was the smallest effect for any of the locations.

Prediction of attendance is also affected by location of recruitment: The large-sized intention-behaviour relationship reported for hospital studies was significantly greater than small-to-medium-sized relations found in health authorities and GP settings. One explanation of this finding is that participants may not treat invitations to screen in GP settings as seriously as invitations from hospitals. Participants asked about their intentions to attend a health check at the local GP might respond that they intend to attend screening, but fail to do so because this screening is viewed as unimportant or unnecessary. This may be very different to participants' response to invitations from hospitals, which may be treated with greater respect particularly if the individual has personal (or family) experience of attending hospital for treatment for serious conditions.

Screening Cost

Attitudes and PBC were significantly better predictors of intentions in free versus paid contexts whereas intentions and PBC were significantly better predictors of attendance in paid versus free contexts. The subjective norm-intention relationship was not affected by screening cost. Thus, there is better prediction of intention (the motivational part of the TPB) in free contexts, but better prediction of behaviour (the volitional part of the TPB) in contexts where screening is not free.

Attitude and PBC may be more predictive of intentions in free contexts compared to paid contexts because screening is not limited to those who can afford it in free contexts. So, other factors such as how easy it is to access screening (in terms of proximity of testing centre), time constraints, and the value of screening, have a greater

Meta-Analysis of TPB/TRA studies predicting screening attendance

impact on decisions to attend screening. On the other hand, there may be lower attitude-intention relations in paid contexts because individuals are well disposed to screening, but do not intend to screen as they cannot afford to attend.

Intentions and PBC were better predictors of attendance in paid contexts than in free contexts. These differences may reflect greater variation in PBC and intention where there is not a free national healthcare system, due to the perception that paying for screening is a significant barrier and hence many people will not intend to attend. It is now well established that people who do not intend to attend for screening are quite successful at not doing so (Orbell & Sheeran, 1998).

In paid contexts, people may possess more accurate perceptions of control which, according to Ajzen (1988), should lead to better prediction of behaviour. For example, people will be aware of their ability to pay, which is likely to be an important control factor, thus producing more accurate PBC. In free contexts, cost is not an important control factor, and it may be that people are less aware of other control factors that impact their ability to attend screening.

Invitation to Screen

Attitudes were significantly better predictors of intentions in invitation versus no invitation contexts whereas subjective norms and PBC were better predictors of intention in the no invitation studies. It is possible that attitudes were a better predictor of intentions in the invitation studies because participants feel more informed about the screening procedure prior to completing a questionnaire and hence have more stable attitudes. The contrast between the large PBC-intention relations found for no invitation studies and the medium relationship for the invitation studies is noteworthy. One

Meta-Analysis of TPB/TRA studies predicting screening attendance

explanation is that participants in some of the no invitation studies were being asked about their perceptions of control over screening attendance for a test that was not yet available. For example, Orbell and Hagger (2006a) asked participants about their perceptions of control over attending diabetes screening if it was offered to them in the next few years. So, PBC responses in some of the no invitation studies were hypothetical. None of the invitation studies examined tests that were not yet available.

We also examined the impact of the timing of the invitation, to see if this affected TPB relationships. For every relationship, except the subjective norm-intention relationship, completing TPB measures after receiving an invitation to screen led to stronger relationships, relative to completing measures before receiving an invitation. These findings support Ajzen's (1991) claim that the TPB provides best prediction when constructs are measured near to behaviour, because there is less chance that beliefs will change between measurement and behavioural performance. Participants who complete questionnaires after they have received an invitation are likely to be nearer to attendance compared to participants who completed measures before receiving an invitation.

Limitations

The present meta-analysis is limited by the studies selected for inclusion; we decided to focus on the published literature, which means we may have missed relevant papers from the 'grey literature'. Glass, McGraw and Smith (1981) found that published papers tend to report stronger effect sizes compared to unpublished papers, so one minor limitation of this review is that it may overestimate the size of TRA/TPB relationships found in screening contexts. Second, it is possible to be critical of some of the moderator analyses conducted, due to few studies being available for some comparisons. We feel

Meta-Analysis of TPB/TRA studies predicting screening attendance

that these analyses are useful, however, in generating hypotheses for future research, rather than as providing definitive conclusions. In addition, comparing screening programmes is problematic because programmes target populations that differ on key characteristics such as age and gender.

Implications

Attitudes appear to be the best predictor of intentions to attend for screening. Consequently, to increase attendance, screening organisations would be best advised to send people information designed to generate positive attitudes, rather than alter subjective norms or PBC. An alternative aim of screening programmes is to promote informed choices. The large-sized attitude-intention relationship suggests that many individuals are making informed choices regarding screening attendance, based on the definition of “informed choice” as consistency between attitudes towards screening attendance and actual attendance, as well as high levels of knowledge about screening (Marteau, Dormandy & Michie, 2001). Research by Dormandy, Hankins and Marteau (2006) suggests that individuals are ambivalent about screening attendance and possess both negative and positive attitudes regarding screening. Given the importance of attitudes as a predictor of screening intentions, it is important for future research to assess the impact of properties of attitude, such as ambivalence, on attitude-intention consistency (see Cooke & Sheeran, 2004 for a review of this literature).

Receiving an invitation to screen does not necessarily make people think about how they will attend. One way to reduce the ‘gap’ between intentions and screening attendance is to use implementation intentions (Gollwitzer, 1993), where the individual outlines when, where and how they will perform a behaviour. Sheeran and Orbell (2000)

Meta-Analysis of TPB/TRA studies predicting screening attendance

found that cervical screening attendance was significantly higher for women who formed implementation intentions (see also Walsh, 2005). Asking people to form implementation intentions is another way to increase attendance at screening programmes.

More research is required to explain why there are differences in predicting attendance in free versus paid contexts. It is possible that there are differences between participants attending screening in paid contexts versus free contexts. The studies conducted outside of the USA tend to concern the whole of a defined age group who are eligible for that screening test, whereas the studies in the USA tend to employ more restricted samples, (e.g., siblings of patients, partners of servicemen). Variation in sampling may explain the differences in the relationships found. To rigorously test the impact on TPB relationships of free versus paid systems of screening, a quasi-experimental test is required, where a country or region shifts from one system of organising screening to the other.

We should investigate how individuals react to invitations to screen. Results for test context and location of recruitment suggest that invitations may be viewed differently depending on where they come from and the nature of the test. For example, an invitation to a health check may be treated as an activity that one should do, but may not, with the consequences of non-attendance viewed as limited. In contrast, receiving an invitation from a hospital to attend for colorectal cancer screening will likely be treated as something one must do, particularly if there is a family history of cancer.

The present study found that the TRA/TPB was an effective framework for predicting screening intentions and attendance. The next step is to perform experimental research that builds on these findings to improve screening attendance. Interventions

Meta-Analysis of TPB/TRA studies predicting screening attendance

designed to make attitudes more positive may increase attendance, because more positive attitudes should create more positive intentions, and more positive intentions increase the likelihood of attendance. Implementation intentions should also increase screening attendance, so research is needed to test this proposition. We also hypothesise that invitations to screen from GP practices are seen as less serious compared to invitations to screen at hospital. Research is needed to test these hypotheses and thereby increase our understanding of screening attendance.

Meta-Analysis of TPB/TRA studies predicting screening attendance

References

An asterisk precedes references included in the meta-analysis

Ajzen, I. (1988). Attitudes, personality, and behavior. Milton Keynes: Open University Press.

Ajzen, I. (1991). The theory of planned behavior. Organizational Behavior and Human Decision Processes, 50, 179-211.

Ajzen, I., & Fishbein, M. (1980). Understanding attitudes and predicting social behavior. Englewood Cliffs, NJ: Prentice-Hall.

Armitage, C.J., & Conner, M. (2001). Efficacy of the theory of planned behaviour: A meta-analytic review. British Journal of Social Psychology, 40, 471-499.

*Armitage, C.J., Norman, P., & Conner, M. (2002). Can the Theory of Planned Behaviour mediate the effects of age, gender and multidimensional health locus of control? British Journal of Health Psychology, 7, 299-316.

*Barling, N.R., & Moore, S.M. (1996) Prediction of cervical cancer screening using the theory of reasoned action. Psychological Reports, 79, 77-78.

*Bish, A., Sutton, S., & Golombok, S. (2000). Predicting uptake of a routine cervical smear test: A comparison of the Health Belief Model and the Theory of Planned Behaviour. Psychology & Health, 15, 35-50.

*Braithwaite, D., Sutton, S., & Steggle, N. (2002). Intention to participate in predictive genetic testing for hereditary cancer: The role of attitude toward uncertainty. Psychology and Health, 17, 761-772.

Cohen, J. (1992). A power primer. Psychological Bulletin, 112, 155-159.

Meta-Analysis of TPB/TRA studies predicting screening attendance

Conner, M., & Norman (2005). Predicting Health Behaviour. Maidenhead: Open University Press.

*Conner, M., Sheeran, P., Norman, P., & Armitage, C.J. (2000). Temporal stability as a moderator of relationships in the Theory of Planned Behaviour. British Journal of Social Psychology, *39*, 469-493.

Cooke, R., & Sheeran, P. (2004). Moderation of cognition-intention and cognition-behaviour relations: A meta-analysis of properties of variables from the theory of planned behaviour. British Journal of Social Psychology, *43*, 159-186.

Department of Health (February 2005). Breast Screening Programme, England: 2003-04. Bulletin 2005/06. London, England: Department of health. ISBN 1 84182 949 8.

*DeVellis, B.M., Blalock, S.J., & Sandler, R.S. (1990). Predicting participation in cancer screening: The role of perceived behavioral control. Journal of Applied Social Psychology, *20*, 639-660.

Dormandy, E., Hankins, M., & Marteau, T.M. (2006). Attitudes and uptake of a screening test: The moderating role of ambivalence. Psychology & Health, *21*, 499-511.

*Drossaert, C.H.C., Boer, H., & Seydel, E.R. (2003). Prospective study of the determinants of repeat attending and attendance patterns in breast cancer screening using the theory of planned behaviour. Psychology & Health, *18*, 551-565.

Fishbein, M, & Ajzen, I. (1975). Beliefs, attitude, intention and behavior. Reading, MA: Addison-Wesley.

Fernbach, M. (2002). The impact of a media campaign on cervical screening knowledge and self-efficacy. Journal of Health Psychology, *7*, 85-97.

Meta-Analysis of TPB/TRA studies predicting screening attendance

*Frost, S., Myers, L.B., & Newman, S.P. (2001). Genetic screening for Alzheimer's disease: What factors predict intentions to take a test? Behavioral Medicine, 27, 101-109.

Glass, G.V., McGraw, B., & Smith, M.L. (1981). Meta-analysis in social research. Beverly Hills, CA: Sage.

Godin, G., Boyer, R., Duval, B., Fortin, C., & Nadeau, D. (1992). Understanding physicians' decisions to perform a clinical examination on an HIV seropositive patient. Medical Care, 30, 199-207.

*Godin, G., Gagné, C., Maziade, J., Moreault, L., Beaulieu, D., & Morel, S. (2001). Breast cancer: The intention to have a mammography and a clinical breast examination-- Application of the theory of planned behavior. Psychology & Health, 16, 423-441.

Godin, G., & Kok, G. (1996). The theory of planned behavior: A review of its applications to health-related behaviors. American Journal of Health Promotion, 11, 87-98.

Gollwitzer, P.M. (1993). Implementation intentions: Strong effects of simple plans. American Psychologist, 54, 493-503.

*Hill, D., Gardner, G., & Rassaby, J. (1985). Factors predisposing women to take precautions against breast and cervix cancer. Journal of Applied Social Psychology, 15, 59-79.

Hedges, L., & Olkin, I. (1985). Statistical methods for meta-analysis. New York: Academic Press.

Hounsa, A., Godin, G., Alihonou, E., Valois, P., & Girard, J. (1993). An application of Ajzen's theory of planned behaviour to predict mother's intention to use oral rehydration therapy in a rural area of Benin. Social Science & Medicine, 37, 253-261.

Meta-Analysis of TPB/TRA studies predicting screening attendance

Jepson, C., Clegg, A., Forbes, C., Lewis, R., Sowden, A., & Kleijnen J. (2000). The determinants of screening uptake and interventions for increasing uptake: a systematic review. Health Technology Assessment, 4(14).

*Jennings-Dozier, K. (1999) Predicting intentions to obtain a pap smear among African-American and Latina women: Testing the theory of planned behavior. Nursing Research, 48, 198-205.

Johnston, M., French, D.P., Bonetti, D., & Johnston, D. (2004). Assessment and measurement in health psychology. In S Sutton, A Baum, & M Johnston (eds.), *The Sage Handbook of Health Psychology*. London: Sage. (pp 288-323)

Marteau, T.M., Dormandy, E, & Michie, S. (2001). A measure of informed choice. Health Expectations, 4, 99-108.

McCaffery, K., Wardle, J., & Waller, J. (2003). Knowledge, attitudes and behavioral intentions in relation to the early detection of colorectal cancer in the United Kingdom. Preventive Medicine, 36, 525-535.

McCaul, K., O'Neill, K., & Glasgow, R. (1988). Predicting the performance of dental hygiene behaviors: An examination of the Fishbein and Ajzen model and self-efficacy expectations. Journal of Applied Social Psychology, 18, 114-128.

*Michels, T.C., Taplin, S.M., Carter, W.B., & Kugler, J, P. (1995). Barriers to screening: The theory of reasoned action applied to mammography use in a military beneficiary population. Military Medicine, 160, 431-437.

*Michie, S., Dormandy, E., French, D.P., & Marteau, T.M. (2004). Using the theory of planned behaviour to predict screening uptake in two contexts. Psychology and Health, 19, 705-718.

Meta-Analysis of TPB/TRA studies predicting screening attendance

*Montaño, D.E., & Taplin, S.H. (1991). A test of an expanded theory of reasoned action to predict mammography participation. Social Science & Medicine, *32*, 733-741.

*Montaño, D.E., Thompson, B., Taylor, V.M., & Mahloch J. (1997). Understanding mammography intention and utilization among women in an inner city public hospital clinic. Preventive Medicine, *26*, 817-824.

*Norman, P., & Conner, M. (1996). Predicting health-check attendance among prior attenders and nonattenders: The role of prior behavior in the theory of planned behavior. Journal of Applied Social Psychology, *26*, 1010-1026.

*Orbell, S., & Hagger, M. (2006a). Temporal framing and the decision to take part in Type 2 diabetes screening: Effects of individual differences in consideration of future consequences on persuasion. Health Psychology, *25*, 537-548.

*Orbell, S., & Hagger, M. (2006b). “When no means no”: Can reactance augment the theory of planned behavior? Health Psychology, *25*, 586-594.

*Orbell, S., Perugini, M., & Rakow, T. (2004). Individual differences in sensitivity to health communications: Consideration of future consequences. Health Psychology, *23*, 388-396.

Orbell, S., & Sheeran, P. (1998). ‘Inclined abstainers’. A problem for predicting health-related behaviour. British Journal of Social Psychology, *37*, 151-165.

Ouellette, J., & Wood, W. (1998). Habit and intention in everyday life: The multiple processes by which past behavior predicts future behavior. *Psychological Bulletin*, *124*, 54-74

*Poss, J.E. (2000). Factors associated with participation by Mexican migrant farmworkers in a tuberculosis screening program. Nursing Research, *49*, 20-28.

Meta-Analysis of TPB/TRA studies predicting screening attendance

Rosenthal, R.R. (1984). Meta-analytic procedures for social research. Beverly Hills, CA: Sage.

*Rutter, D.R. (2000). Attendance and reattendance for breast cancer screening: A prospective 3-year test of the Theory of Planned Behaviour. British Journal of Health Psychology, *5*, 1-13.

Schwarzer, R. (1988). Meta: Programs for secondary data analysis. Berlin, Germany: Free University of Berlin.

*Sheeran, P., Conner, M., & Norman, P. (2001). Can the theory of planned behavior explain patterns of health behavior change? Health Psychology, *20*, 12-19.

Sheeran, P., & Orbell, S. (2000) Using implementation intentions to increase attendance for cervical cancer screening. Health Psychology, *19*, 283-289.

Shepherd, B.H., Hartwick, J., & Warshaw, P.R. (1988). The theory of reasoned action: A meta-analysis of past research with recommendations for future research. Journal of Consumer Research, *15*, 325-343.

*Steadman, L., & Rutter, D.R. (2004) Belief importance and the theory of planned behaviour: Comparing modal and ranked modal beliefs in predicting attendance at breast screening. British Journal of Health Psychology, *9*, 447-464.

*Steadman, L., Rutter, D.R., & Field, S. (2002). Individually elicited versus modal normative beliefs in predicting attendance at breast screening: Examining the role of belief salience in the Theory of Planned Behaviour. British Journal of Health Psychology, *7*, 317-330.

*Tolma, E.L., Reininger, B.M., Ureda, J., & Evans, A. (2003). Cognitive motivations associated with screening mammography in Cyprus. Preventive Medicine, *36*, 363-373.

Meta-Analysis of TPB/TRA studies predicting screening attendance

*Walsh, J.C. (2005). Increasing screening uptake for a cervical smear test: Predictors of attendance and the use of action plans in prior non-attenders. Irish Journal of Psychology, 26, 65-73.

Meta-Analysis of TPB/TRA studies predicting screening attendance

Acknowledgments

We thank Constance Drossaert, Susan Moore, Sheina Orbell, Paschal Sheeran and Jane Walsh for providing additional data, and Elizabeth Dormandy and Ann Louise Kinmonth for helpful discussions of our results.

Meta-Analysis of TPB/TRA studies predicting screening attendance

Table 1. Studies included in the meta-analysis, with correlation coefficients between Motivational variables (Attitude, Subjective Norm, and Perceived Behavioural Control) and Intention

Authors	Sample	Country	Behaviour	N	A	SN	PBC
Armitage, Norman, & Conner (2002)	Men/Women (31-42)	UK	Health Check	201	.46	.26	.33
Barling & Moore (1996)	Women (18-63)	Australia	Cervical Smear	72	.49	.57	
Bish, Sutton, & Golombok (2000)	Women ^a	UK	Cervical Smear	142	.67	.54	.13
Braithwaite, Sutton, & Steggle (2002)							
Breast Cancer	Women (18-60)	UK	Genetic Test	124	.71	.48	
Colon Cancer	Men/Women (18-60)	UK	Genetic Test	168	.67	.55	
Conner, Sheeran, Norman, & Armitage (2000)	Men/Women (31-42)	UK	Health Check	201	.60	.67	.33
DeVellis, Blalock, & Sandler (1990)							
High risk sample	Men/Women (40-75)	USA	Colorectal Screen	96	.36	.18	.35
Low risk sample	Men/Women (40-75)	USA	Colorectal Screen	144	.42	.36	.43
Drossaert, Boer, & Seydel (2003)	Women (50-69)	Netherlands	Mammogram	2657	.60	.20	.49

Meta-Analysis of TPB/TRA studies predicting screening attendance

Frost, Myers, & Newman (2001)	Men/Women (18-25)	UK	Genetic test	449	.53	.54	
Godin, Gagné, Maziade, Moreault, Beaulieu, & Morel (2001)	Women (40-69)	Canada	Mammogram	354	.10	.31	.59
	Women (40-69)	Canada	CBE	344	.35	.05	.54
Hill, Gardner, & Rassaby (1985)	Women (18-70)	Australia	Cervical Smear	123	.47	.29	
Jennings-Dozier (1999)							
African American Sample	Women (18-83)	USA	Cervical Smear	108	.58		.30
Latina Sample	Women (18-83)	USA	Cervical Smear	96	.40		.35
Michels, Taplin, Carter, & Kugler (1995)	Women (41-89)	USA	Mammogram	309	.23	.36	
Michie, Dormandy, French, & Marteau (2004) (Routine Testing)	Women (under 37)	UK	Prenatal Screening	446	.83	.70	.17
(Separate Testing)	Women (under 37)	UK	Prenatal Screening	597	.76	.68	.17
Montano & Taplin (1996)	Women (40 and over)	USA	Mammogram	665	.47	.43	.39
Montano, Thompson, Taylor, & Mahloch (1997)	Women (50-69)	USA	Mammogram	361	.39	.41	
Norman & Conner (1996)	Men/Women (31-42)	UK	Health Check	262	.64	.56	.48

Meta-Analysis of TPB/TRA studies predicting screening attendance

Orbell & Hagger (2006a)	Men/Women (40 and over)	UK	Diabetes	210	.54	.66	.77
Orbell & Hagger (2006b)	Women (20-64)	UK	Cervical	660	.35	.35	.67
Orbell, Perugini, & Rakow (2004)	Women (50-69)	UK	Colorectal	220	.47	.71	.78
Poss (2000)	Men/Women (18 and over)	USA	TB Screening	206	.50	.65	
Rutter (2000)	Women (50-64)	UK	Mammogram	1108	.45	.36	.40
Sheeran, Conner, & Norman (2000)	Men/Women (30-41)	UK	Health Check	389	.61	.61	.53
Sheeran & Orbell (2000)	Women (20-67)	UK	Cervical Smear	114	.38	.44	.69
Steadman & Rutter (2004)	Women (50-64)	UK	Mammogram	785	.21	.34	.43
Steadman, Rutter, & Field (2002)							
(Individual condition)	Women (50-64)	UK	Mammogram	228	.49	.27	.29
(Modal condition)	Women (50-64)	UK	Mammogram	270	.44	.28	.45
Tolma, Reininger, Ureda & Evans (2003)	Women (40-65)	Cyprus	Mammogram	293	.33	.39	.33
Walsh (2005)	Women (25-60)	Ireland	Cervical Smear	156	.41	.50	.67

^aNo mention of sample age range in paper; CBE = Clinical Breast Exam; TB = Tuberculosis

Meta-Analysis of TPB/TRA studies predicting screening attendance

Table 2. Studies included in the meta-analysis, including correlation coefficients between intentions and behaviour and perceived behavioural control and behaviour.

Authors	Sample	Country	Behaviour	N	I	PBC
Armitage, Norman & Conner (2002)	Men/Women (31-42)	UK	Health Check	201	.22	.19
Bish, Sutton, & Golombok (2000)	Women	UK	Cervical Smear	142	.17	.09
Conner, Sheeran, Norman, & Armitage (2000)	Men/Women (31-42)	UK	Health Check	201	.34	.26
DeVellis, Blalock, & Sandler (1990)						
High risk sample	Men/Women (40-75)	USA	Colorectal Screen	96	.33	.34
Low risk sample	Men/Women (40-75)	USA	Colorectal Screen	144	.50	.19
Drossaert, Boer, & Seydel (2003)	Women (50-69)	Netherlands	Mammogram	2039	.32 ^a	.23 ^a
Michie, Dormandy, French, & Marteau (2004) (Routine Testing)	Women (under 37)	UK	Prenatal Screening	446	.86	.14
(Separate Testing)	Women (under 37)	UK	Prenatal Screening	597	.69	.13
Montano & Taplin (1991)	Women (40 or older)	USA	Mammogram	665	.50	.37

Meta-Analysis of TPB/TRA studies predicting screening attendance

Norman & Conner (1996)	Men/Women (31-42)	UK	Health Check	268	.16	.13
Orbell & Hagger (2006b)	Women (20-64)	UK	Cervical	660	.19	.12
Poss (2000)	Men/Women (18 or older)	USA	TB Screening	206	.84	
Rutter (2000)	Women (50-64)	UK	Mammogram	600	.32 ^a	.16 ^a
Sheeran, Conner, & Norman (2000)	Men/Women (30-41)	UK	Health Check	389	.20 ^a	.19 ^a
Sheeran & Orbell (2000)						
(Control group)	Women (20-67)	UK	Cervical Smear	55	.57	.41
Steadman & Rutter (2004)	Women (50-64)	UK	Mammogram	785	.34	.15
Steadman, Rutter, & Field (2002)						
(Individual condition)	Women (50-64)	UK	Mammogram	228	.49	.04
(Modal condition)	Women (50-64)	UK	Mammogram	270	.44	.03
Walsh (2005)	Women (25-60)	Ireland	Cervical Smear	156	.18	.25

Notes: ^a Average of measures taken at two timepoints

Meta-Analysis of TPB/TRA studies predicting screening attendance

Table 3. Summary of TRA/TPB relationships estimated by meta-analysis

Relation	n	k	CI	χ^2	r_+	FSN
Attitude-Intention	12558	33	.49 to .53	737.96***	.51	305
Subjective Norm-Intention	12354	31	.39 to .42	560.31***	.41	221
PBC-Intention	10746	25	.45 to .48	415.74***	.46	205
Intention-Behaviour	8148	19	.40 to .44	483.58***	.42	141
PBC-Behaviour	7942	18	.17 to .21	58.13***	.19	50

Note. n = number of participants, k = number of tests of the relationship, CI = 95% confidence interval, χ^2 = chi-square test of homogeneity, r_+ = sample-weighted average correlation, FSN = Rosenthal's (1984) Fail Safe N. * $p < .05$; ** $p < .01$; *** $p < .001$.

Table 4. Estimated effect sizes for Type of Screening Test for Attitude-Intention, Subjective Norm-Intention and Intention-Behaviour Relationships

Moderator	Attitude-Intention			Subjective Norm-Intention			PBC-Intention			Intention-Behaviour		
	<u>n</u>	<u>k</u>	<u>r</u> ₊	<u>n</u>	<u>k</u>	<u>r</u> ₊	<u>n</u>	<u>k</u>	<u>r</u> ₊	<u>n</u>	<u>k</u>	<u>r</u> ₊
Cervical	1471	8	.43 _c	1267	6	.43 _c	1276	6	.58 _a	1013	4	.21 _c
Colorectal	460	3	.43 _c	460	3	.52 _b	460	3	.62 _a	240	2	.44 _b
Genetic Test	741	3	.60 _b	741	3	.53 _b	-	-	-	-	-	-
Health Check	1053	4	.59 _b	1053	4	.55 _b	1053	4	.45 _b	1059	4	.22 _c
Mammography	7030	10	.46 _c	7030	10	.30 _d	6360	8	.45 _b	4587	6	.37 _b
Prenatal	1043	2	.79 _a	1043	2	.69 _a	1043	2	.17 _c	1043	2	.78 _a

Note. Correlations within columns not sharing the same subscript are significantly different from one other ($p < .05$). n = number of participants, k = number of tests of the relationship, r = sample-weighted average correlation.

Table 5. Estimated effect sizes for Location of Recruitment for Attitude-Intention, Subjective Norm-Intention, PBC-Intention and Intention-Behaviour Relations

Moderator	Attitude-Intention			Subjective Norm-Intention			PBC-Intention			Intention-Behaviour		
	n	k	r ₊	n	k	r ₊	n	k	r ₊	n	k	r ₊
General Practices	1601	8	.60 _a	1601	8	.54 _a	1309	6	.44 _b	1256	6	.23 _c
Hospitals	3571	9	.53 _b	3571	9	.49 _b	2901	7	.39 _c	2608	6	.57 _a
Health Authorities	5204	6	.50 _c	5204	6	.27 _d	5204	6	.46 _b	4078	6	.34 _b
Other Settings	2622	10	.40 _d	2418	8	.41 _c	1772	6	.60 _a			

Note. Correlations within columns not sharing the same subscript are significantly different from one other ($p < .05$). n = number of participants, k = number of tests of the relationship, r = sample-weighted average correlation.

Table 6. Comparison between TRA/TPB relationships reported where screening is paid for versus where screening is free

Relation	Free					Paid For					Z
	n	k	CI	χ^2	r ₊	n	k	CI	χ^2	r ₊	
Attitude-Intention	10573	25	.51 to .54	715.72***	.53	1985	8	.38 to .45	28.75***	.42	5.89***
Subjective Norm-Intention	10573	25	.39 to .42	524.98***	.40	1781	6	.39 to .46	30.43***	.43	1.41
PBC-Intention	9637	20	.45 to .48	414.32***	.47	1109	5	.33 to .43	2.40	.38	3.47***
Intention-Behaviour	7037	15	.38 to .41	366.59***	.40	1111	4	.53 to .61	82.50***	.58	7.39***
PBC-Behaviour	7037	15	.15 to .19	28.92***	.17	905	3	.28 to .40	5.17	.34	5.16***

Note. n = number of participants, k = number of tests of the relationship, CI = 95% confidence interval, χ^2 = chi-square test of homogeneity, r₊ = sample-weighted average correlation. *** $p < .001$.

Meta-Analysis of TPB/TRA studies predicting screening attendance

Table 7. Comparison between TPB relationships reported in Invitation versus No Invitation studies

Relation	Invitation					No Invitation					Z
	n	k	CI	χ^2	r ₊	n	k	CI	χ^2	r ₊	
Attitude-Intention	9413	20	.53 to .56	605.54***	.55	3145	13	.36 to .42	91.39***	.39	10.03***
Subjective Norm-Intention	9413	20	.38 to .41	384.28***	.39	2941	11	.41 to .47	174.00***	.44	2.85**
PBC-Intention	9121	18	.42 to .45	274.37***	.44	1625	7	.53 to .61	144.42***	.57	6.50***

Note. n = number of participants, k = number of tests of the relationship, CI = 95% confidence interval, χ^2 = chi-square test of homogeneity, r₊ = sample-weighted average correlation. * p < .05; ** p < .01; *** p < .001.

Table 8. Estimated effect sizes depending on Timing of Questionnaire completion

Moderator	Before Invitation			Same Time as Invitation			After Invitation		
	n	k	r ₊	n	k	r ₊	n	k	r ₊
Attitude-Intention	3041	10	.51 _b	448	3	.61 _a	5139	6	.60 _a
Subjective Norm-Intention	3041	10	.42 _b	448	3	.51 _a	5139	6	.37 _c
PBC-Intention	3041	10	.40 _b				5139	6	.45 _a
Intention-Behaviour	2539	10	.31 _b				4462	6	.47 _a
PBC-Behaviour	2539	10	.15 _b				4462	6	.22 _a

Note. Correlations within columns not sharing the same subscript are significantly different from one other ($p < .05$). n = number of participants, k = number of tests of the relationship, r = sample-weighted average correlation.