

## **Highlights**

- traditional weed control strategies tend to impact negatively the environment
- cover crop cultivation prior to a maize crop can be efficient to control weeds
- niger, sunflower, pea and phacelia were the most successful species in this respect
- some cover crops also improved maize yield compared to the control
- this study shows that cover crops are efficient to reduce weed control intensity

1 **Title**

2 Cover crops to secure weed control strategies in a maize crop with reduced tillage

3

4 **Authors**

5 Lucie Büchi <sup>1,2,\*</sup>, Marina Wendling <sup>1,3</sup>, Camille Amossé <sup>1</sup>, Bernard Jeangros <sup>1</sup>, Raphaël

6 Charles <sup>1,3</sup>

7

8 **Affiliations**

9 <sup>1</sup> Agroscope, Institute for Plant Production Sciences, Nyon, Switzerland

10 <sup>2</sup> Natural Resources Institute, University of Greenwich, Chatham, United Kingdom

11 <sup>3</sup> Research Institute of Organic Agriculture FiBL, Lausanne, Switzerland

12

13 **\*Corresponding author**

14 L.A.Buchi@greenwich.ac.uk

15 Natural Resources Institute

16 University of Greenwich

17 Chatham

18 ME4 4TB

19 United Kingdom

20

21 **Declarations of interest: none**

22 **Abstract**

23 To better understand the ability of cover crops to control weeds in a maize crop (*Zea mays*, L.)  
24 grown with reduced tillage, four field experiments were set up from 2009 to 2014 in the  
25 western part of Switzerland. Ten non-wintering cover crop species were compared to a no  
26 cover crop control in strip plot experiments including different weeding strategies. The  
27 weeding strategies included no or minimum tillage before maize seeding. Soil coverage by  
28 weeds at early maize stage (2-4 leaf stage) varied drastically between weeding strategies and  
29 years. In most cases, cover crops allowed to reduce the weed pressure compared to the no  
30 cover crop control. The most efficient cover crop species varied from year to year, but niger  
31 (*Guizotia abyssinica*, (L.f.) Cass.), sunflower (*Helianthus annuus*, L.), field pea (*Pisum*  
32 *sativum*, L.) and phacelia (*Phacelia tanacetifolia*, Benth.) gave the best overall results. Maize  
33 yield differed significantly between weeding strategies only one year, with higher yield  
34 observed with minimum tillage. In some situation, cover crops cultivated in autumn still  
35 showed a significant impact on maize yield, with common vetch (*Vicia sativa*, L.) as the most  
36 successful species. Interestingly, the effect of cover crop on weed cover and maize yield was  
37 not limited to the less intense strategy (no tillage). These results show that cultivating cover  
38 crops before maize in this type of conditions is a promising method to help controlling weeds.  
39 In addition, cover crops are known for providing multiple ecosystem services which could  
40 altogether improve the sustainability of cropping systems on the long term.

41

42

43 **Keywords**

44 no till, cover crop biomass, residue cover, integrated weed management

45

46

47

## 48 **1 Introduction**

49 Reduced tillage has gained popularity over the last thirty years among farmers. The occasional  
50 or systematic suppression of full-inversion ploughing implies an adaptation of the cropping  
51 system, notably in the control of weeds as well as of soil-borne organisms and pests (Lahmar,  
52 2010; Peigné et al., 2007). Consequently, farmers using reduced tillage may choose to rely  
53 increasingly on herbicides and pesticides to deal with these threats (Lahmar, 2010). Weed  
54 control is currently identified as the main issue associated with reduced tillage and the key to  
55 sustainability (Bajwa, 2014; Eslami, 2014; Ramesh, 2015), especially in the context of  
56 conservation agriculture (Giller et al., 2009). The recent preoccupation concerning the toxicity  
57 of glyphosate (Braz-Mota et al., 2015; Gaupp-Berghausen et al., 2015; Sihtmäe et al., 2013),  
58 the main herbicide associated with reduced tillage (Nalewaja, 2003), requires new solutions to  
59 reduce its use and the appearance of resistances (Beckie, 2014; Nauen and Denholm, 2005;  
60 Walsh and Powles, 2014). In addition, the extensive reliance on herbicides in reduced tillage  
61 systems is particularly problematic in context where the cost of herbicides renders their access  
62 and use difficult or even impossible (Giller et al., 2009). Melander et al. (2013) and Nichols et  
63 al. (2015) discussed the perspective of adopting nonchemical weed management in reduced  
64 tillage systems by optimising crop rotation, cover crop use, stubble management, enhancement  
65 of crop growth and direct non-chemical methods.

66 The use of a cover crop between two cash crops brings together (i) soil cover as a living crop  
67 or a dead mulch and (ii) diversification of the crop rotation. Cover crops are expected to  
68 control weeds and may reduce the need to use herbicides (Alonso-Ayuso et al., 2018; Brust et  
69 al., 2014; Teasdale and Mohler, 2000). However, the beneficial effect of cover crops on weeds  
70 is generally linked to a high cover crop biomass or to a fast soil cover. If biomass and residues  
71 are scarce or rapidly decomposed, herbicides can then be needed, depending on weed pressure.  
72 Consequently, the choice of the best adapted cover crop species is crucial.

73 Cover crop species previously tested in reduced tillage systems were mainly wintering cover  
74 crop species (Carrera et al., 2004; Hayden et al., 2014; Mirsky et al., 2013; Sainju et al., 2002;  
75 Teasdale and Mohler, 1993; Williams et al., 1998). Non-wintering (i.e. frost-killed) species  
76 could be useful when the aim is to reduce simultaneously ploughing and herbicide use. Yet the  
77 practical issues are currently in the optimal combination of soil tillage intensity, herbicide  
78 timing/use and cover crop species choice. Consistent weeding practices based on annual weed  
79 pressure need to be studied to develop an integrated approach to control weeds.

80 The objective of this multi-year study is to highlight the contribution of a set of ten cover crop  
81 species to weed control and maize yield, in combination with two weeding strategies including  
82 no or minimum soil tillage, before a silage maize crop (*Zea mays*, L.). For this purpose, on-  
83 station field experiments were conducted during the period 2009-2014 in Switzerland, to  
84 investigate 1. the performance of cover crops in terms of biomass production in autumn and  
85 residue cover in early spring, 2. the influence of weeding strategies on weed cover in spring, 3.  
86 the effect of cover crops on weed cover within each weeding strategy, and 4. the effect of  
87 weeding strategies and cover crops on maize yield.

88

89

## 90 **2 Materials and methods**

### 91 **2.1 Experimental setup**

92 The experiments were set up at the research station Agroscope Changins (46° 24' N, 06° 14' E,  
93 430 m above sea level), Switzerland, on a Cambisol (FAO classification). At this location, the  
94 average total annual precipitation is 999 mm and the mean air temperature 10.2°C (30-year  
95 averages, 1981-2010).

96 Four experiments were implemented, in 2009-2010 (hereafter named 'year 0'), 2010-11 ('year  
97 1'), 2011-12 ('year 2') and 2013-14 ('year 3') on different fields at the same experimental  
98 station to study the influence of cover crops and weeding strategies on weed infestation and

99 maize yield. Soil characteristics, crop management and observation dates for each experiment  
100 are presented in Supplementary Material Table S1. Each year, the experimental design  
101 corresponded to a strip plot with three replicates. Cover crop treatments represented the  
102 ‘horizontal’ factor of the strip plot, whereas weeding strategies represented the ‘vertical’ factor.  
103 Each unit plot (cover crop species x weeding strategy) had a size of 3 x 8 m.  
104 Ten cover crop species (Table 1) and a no cover crop control were compared. The species list  
105 included three Brassicaceae species (Indian mustard, *Brassica juncea*, species code b1; turnip  
106 rape, *Brassica rapa campestris*, b2 and daikon radish, *Raphanus sativus longipinnatus*, b3),  
107 three Fabaceae species (field pea, *Pisum sativum*, f1; berseem clover, *Trifolium alexandrinum*,  
108 f2 and common vetch, *Vicia sativa*, f3), one Poaceae species (black oat, *Avena strigosa*, p1),  
109 two Asteraceae species (niger, *Guizotia abyssinica*, a1 and sunflower, *Helianthus annuus*, a2)  
110 and one Hydrophyllaceae species (phacelia, *Phacelia tanacetifolia*, h1). The cover crops were  
111 non-wintering species, known to grow well at this location. The control treatment was left non-  
112 seeded. In year 0, only seven of these ten cover crop species were tested (all species except  
113 daikon radish, common vetch and black oat, Table 1).  
114 The second factor of the strip plot consisted in two different weeding strategies. The first, less  
115 intense one, involved no weeding (herbicide or tillage) interventions from cover crop seeding  
116 to maize seeding (1.NoTill). The second strategy involved minimum tillage (rotary harrow)  
117 prior to maize seeding. Herbicides (Dicamba, Terbutylazine, Mesotrione, Nicosulfuron) were  
118 applied in all treatments at the end of May – beginning of June just after weed cover evaluation  
119 (Table S1), except in 2.MinTill in year 3 where a global very low weed infestation decided for  
120 the use of mechanical weeding instead of using herbicides. Figure 1 shows the timing of the  
121 weeding interventions.  
122

## 123 2.2 Experiment management

124 All the experiments started in summer, after the harvest of the preceding crop. Deep inversion  
125 tillage (mouldboard plough, 20-30 cm depth) followed by rotary harrow (8 cm depth) was  
126 applied before cover crop seeding, except in year 3 where ploughing was not necessary, as a  
127 legume species and not a cereal was cultivated as preceding crop (rotary harrow only, 8 cm  
128 depth). Cover crops were seeded in mid-summer (end of July-beginning of August, Table S1),  
129 with an experimental seeder, at 2-3 cm depth. Cover crops were fertilised just after seeding  
130 with 30 kg N/ha, except in year 3 where the preceding crop was *Medicago sativa*, a legume  
131 species. The cover crops were shredded before maize seeding (horizontal axis shredder with  
132 hammer knives). Silage maize (cv. Ricardinio) was seeded in May - end of April in all plots  
133 with a pneumatic seeder, at a density of 10 plants/m<sup>2</sup>, with an inter-row spacing of 0.75 m.  
134 Nitrogen fertilisation was applied twice, the first time in early May and then in early June  
135 (~120 kg N/ha in total).

136

## 137 2.3 Data collection

138 The aboveground biomass of the cover crops was evaluated each year at the end of the growing  
139 period (beginning of November, 96-97 days after seeding, Table S1), except for the first  
140 experiment in year 0, where no biomass sampling was done. Plants were collected at ground  
141 level in one 0.5 x 0.5 m quadrat from each plot. Biomass was then oven-dried at 60°C during  
142 72 h and weighed to determine the aboveground dry matter of each cover crop species.  
143 In early spring, the proportion of soil covered by the cover crops or their residues (when the  
144 cover crops were killed by frost) (abbreviated as ‘residue cover’ hereafter) was estimated  
145 visually in each plot, using a soil cover scale (example in Figure S2 in Büchi et al., 2018).  
146 Soil coverage by weeds (abbreviated as ‘weed cover’ hereafter) was evaluated at the end of  
147 May, at the highly sensitive 2-4 leaf stage of maize (BBCH 12-14, 20-29 days after seeding),  
148 using the same method as for the cover crops.

149 Maize was harvested as whole plants with a combine harvester at the end of August –  
150 beginning of September (Table S1), except for the first experiment in year 0, where maize  
151 harvest was not conducted separately for each treatment. Fresh maize biomass was weighed  
152 and a subsample dried (72 h, 60°C) to determine its water content. The maize shoot dry yield  
153 (t/ha) was then calculated. Figure 1 shows the timing of these measurements in relation to crop  
154 management interventions.

155

## 156 **2.4 Data analyses**

157 The influence of cover crop species on biomass production and residue cover was tested  
158 independently each year using analyses of variance, followed by post-hoc Tukey HSD tests.  
159 Correlation between cover crop biomass and residue cover was tested using Kendall non-  
160 parametric correlation test.

161 The influence of the weeding strategies on weed cover and maize yield in the control plots  
162 without cover crop was tested independently each year using analyses of variance.

163 Independently for each year and each weeding strategy, the effect of cover crops on weed  
164 cover and maize yield, compared to the no cover crop control, was tested by analyses of  
165 variance followed by pairwise comparisons with least-square significant difference tests (R  
166 package "agricolae", de Mendiburu, 2017).

167 The contribution of cover crop biomass and residue cover to weed cover was analysed with  
168 multiple linear regressions. Correlation between weed cover and maize yield was tested using  
169 Kendall non-parametric correlation test. R 3.6.0 (R Core Team, 2019) was used to perform all  
170 statistical analyses.

171

172



## 173 3 Results

### 174 3.1 Cover crop biomass in autumn and residue cover in spring

175 For each year, average daily temperature was around 20°C at the beginning of the cover crop  
176 growth and decreased progressively to reach about 10°C at the beginning of November, when  
177 cover crop biomass was evaluated. During the cover crop growth period, between seeding and  
178 biomass sampling, the mean temperature was similar each year, around 15°C (14.4°C, 16.1°C,  
179 14.9°C in year 1, year 2 and year 3). Growing degree days (GDD, with a base temperature of  
180 0°C) were also quite similar between years, 1410, 1561 and 1461 GDD respectively. In  
181 contrast, the amount of rainfall between cover crop seeding and biomass sampling changed  
182 drastically between years, from 179 mm in year 1 and 209 in year 2 to 415 mm in year 3.  
183 Overall, cover crops produced about 4.7 t/ha aboveground biomass at the beginning of  
184 November (Table 2). Mean cover crop biomass was, on average, higher in year 3 compared to  
185 year 1 and year 2 (6.2 t/ha vs 3.7 and 4.2 t/ha, respectively). There was a significant interaction  
186 between year and cover crop species ( $p < 0.001$ , Supplementary Material Figure S1) and so  
187 differences between cover crop biomass were tested independently each year (Table 2).  
188 General patterns of biomass production could nevertheless be identified (Table 2, Figure 2). At  
189 the beginning of November, sunflower (species code a2) showed, on average over the three  
190 years, the highest shoot dry matter among the tested species (8.7 t/ha). Niger (a1), the other  
191 Asteraceae, also presented high biomass (6.4 t/ha, 3-year average). Oat (p1), Indian mustard  
192 (b1) and phacelia (h1) performed rather well, with a 3-year average shoot dry matter between  
193 4.8 and 5.6 t/ha. The three legume species (pea, f1; clover, f2 and vetch, f3), and the two  
194 Brassicaceae turnip rape (b2) and daikon radish (b3) presented the lowest 3-year average shoot  
195 dry matter ( $\leq 4.0$  t/ha) (Table 2, Figure 2).  
196 During winter, all cover crops were killed by frost except daikon radish and clover in year 3  
197 due to insufficiently low temperatures. Overall, cover crop residue cover was around 44% in  
198 spring, but it varied between years and species (Figure 2, Table 2). The three legume species

199 (f1, f2, f3) and oat (p1) showed the highest residue cover in early spring ( $\geq 55\%$ , 3-year  
200 average, Table 2). Both Asteraceae (a1, a2) and phacelia (h1) showed intermediate soil cover  
201 (between 28 and 42%, 3-year average). The species with the lowest soil cover were the  
202 Brassicaceae (b1, b2, b3) species ( $\leq 13\%$ , 3-year average).  
203 Cover crop residue cover in early spring was not correlated with cover crop biomass in autumn  
204 (Kendall correlation coefficient  $\tau = -0.04$ , p-value = 0.748). Depending on the species  
205 characteristics, high residue cover in early spring could be achieved with low biomass in  
206 autumn, like for the three Fabaceae species (Figure 2). In contrast, sunflower produced high  
207 amount of biomass but resulting in low cover in spring, and the Brassicaceae species tended to  
208 produce low biomass and low residue cover (Figure 2).

209

### 210 **3.2 Influence of weeding strategy on weed cover at early maize stage**

211 Among the eight cases tested over the four years, only one (2.MinTill in year 3) led to a mean  
212 weed cover lower than the 5% threshold, and one had a mean weed cover around 10%  
213 (2.MinTill in year 1) (Figure 3A). The six other strategies showed higher weed cover ( $>40\%$ ,  
214 Figure 3A and Supplementary Material Table S2). The less intense weeding strategy  
215 (1.NoTill), involving no tillage prior to maize seeding can be used as an indicator of overall  
216 weed pressure. Without any weeding intervention, weed cover after maize seeding ranged from  
217 43% in year 2 up to 90% in year 0. Compared to this low intensive weeding strategy, the other  
218 weeding strategy gave contrasting results for each year (Figure 3A). In years 0 and 2, no  
219 significant effect of the tillage before maize seeding on the weed cover was observed, whereas  
220 in years 1 and 3 a highly significant reduction of weed cover was observed, allowing to  
221 decrease weed pressure to around or less than 10%.

222

### 223 **3.3 Influence of cover crops on weed cover at early maize stage**

224 Comparing the weed cover in the cover crop treatments and in the control with no cover crop  
225 allows to highlight the potential of cover crops to reduce weed pressure at early maize stage (2-

226 4 leaf stage) (Figure 3B and Supplementary Material Table S2). It was expected that the more  
227 pronounced cover crop impact would be observed in the less intense weeding strategy  
228 (1.NoTill), where no tillage was applied. This was however not systematically the case. No  
229 effect of cover crops compared to the control could be observed in years 0 and 1 (Figure 3B).  
230 In year 2, all species induced a significant reduction of weed cover except clover f2, vetch f3  
231 and oat p1. In year 3, niger a1 and oat p1 allowed a reduction of weed cover, from 62%  
232 (control) to 11% for oat.

233 In the strategy involving minimum tillage before maize seeding (2.MinTill), a significant effect  
234 of cover crops could still be observed despite this late tillage intervention, except in year 3  
235 where the no cover crop control was at only 3% weed cover (Figure 3B). Four cover crop  
236 species always induced a significant reduction of weed cover at early maize stage in this  
237 strategy: field pea f1, niger a1, sunflower a2 and phacelia h1.

238 When looking at the cover crop species which systematically allowed to reducing weed cover  
239 across years and strategies, the most successful species was niger a1, followed by field pea f1,  
240 sunflower a2 and phacelia h1 (Figure 3B).

241 Cover crop biomass production in autumn and residue cover in spring did not, in general,  
242 explain the weed cover observed in early spring for most of the strategies. These two variables  
243 accounted for a significant part of the variance in weed cover, but with really low  $R^2$ , only in  
244 year 2 (1.NoTill:  $R^2 = 24\%$  and 2.MinTill:  $R^2 = 34\%$ ,  $p < 0.05$ ). Residue cover alone appeared to  
245 have a significant effect in year 3 for the less intense strategy (1.NoTill:  $R^2 = 16\%$ ,  $p < 0.05$ ).

246

### 247 **3.4 Influence of tillage and cover crops on maize yield**

248 Across years 1, 2 and 3, the mean maize yield in the no cover crop control ranged from 11.3  
249 t/ha to 16.9 t/ha (Figure 4A). A significant difference between the two weeding strategies was  
250 observed only in year 1, with 13.5 t/ha in 1.NoTill and 16.9 t/ha in 2.MinTill. Significant  
251 differences between some cover crop species and the no cover crop control could be observed

252 for maize yield in four out of the six cases (Figure 4B and Supplementary Material Table S3),  
253 but generally cover crop treatments show yield similar to the control. In year 1, where a  
254 significant reduction of maize yield in the 1.NoTill strategy could be observed in the absence  
255 of cover crops, some cover crop species allow to reach a yield similar to that observed in the  
256 2.MinTill strategy (Figure 4B).

257 Vetch f3 was the species appearing most often (three times) as allowing to improve yield  
258 compare to the no cover crop control, followed by field pea f1, oat p1 and niger a1 (twice each)  
259 (Figure 4B). Significant negative correlations between maize yield and weed cover at early  
260 maize stage was observed in three out of six cases (1.NoTill in year 1 and 3, 2.MinTill in year  
261 3), despite all treatments having been weeded in between weed cover estimation and maize  
262 harvest.

263

264

## 265 **4 Discussion**

### 266 **4.1 Influence of weeding strategy on weed infestation and maize yield**

267 Comparing the weed cover observed in the weeding strategies involving minimum tillage to  
268 that without tillage highlighted a high inconsistency in the efficiency of this method. Only in  
269 50% of the cases (two out of four cases), the use of tillage allowed to reduce significantly the  
270 weed cover in the no cover crop controls, compared to the strategy with no intervention.

271 Tillage before maize seeding can disrupt standing weeds and thus control their proliferation,  
272 but at the same time the soil loosening may also recruit weed seeds from the soil seedbank (and  
273 in particular *Echinochloa crus-galli* in these experiments) and give them favourable emergence  
274 conditions (Nichols et al., 2015, Sadeghpour et al., 2014). Therefore, depending on the weather  
275 conditions, minimum tillage before maize seeding is not sufficient to prevent weed emergence.

276 Despite the herbicide application in all treatments just after weed cover evaluation, maize yield  
277 in the control with no cover crop still shows a pattern mirroring the pattern observed for weed

278 cover, with higher yield in the minimum tillage compared to the no till strategy when the weed  
279 cover at early maize stage was significantly reduced. This confirms the observation that maize  
280 performance is highly sensitive to weed infestation in early stages (Page et al., 2012).

281

## 282 **4.2 Cover crop performance and effect on weed infestation and maize yield**

283 Cover crop biomass production varied between years, partly as a response to rainfall amount  
284 during cover crop growth. For most species, the biomass produced in three months of growth  
285 was higher than 3 t/ha, previously identified as a threshold for good cover crop performance  
286 (Gfeller et al., 2018; Wendling et al., 2019). Only turnip rape and daikon radish in years 1 and  
287 2 failed to reach this threshold, with a mean biomass lower than 3 t/ha (between 1.2 and 1.6  
288 t/ha).

289 Cover crops induced a reduction of weed infestation at the early stage of maize in most of the  
290 situations, even when tillage was applied after cover crop cultivation. However, the presence or  
291 magnitude of infestation reduction was not easily predicted based on the data collected in this  
292 study. It is generally admitted that high biomass production or good soil cover are the key  
293 characteristics allowing an efficient weed control by cover crops (Brust et al., 2014, Buchanan  
294 et al., 2016). However, an influence of species identity, and of the use of mixtures, has also  
295 been demonstrated (Baraibar et al., 2018).

296 Different factors could explain why cover crops are generally associated with weed control  
297 (Dorn et al., 2015; Kruidhof et al., 2008; Weber et al., 2017). As living crops, they modify  
298 environmental conditions to the detriment of weeds or directly compete with them for  
299 resources (Rueda-Ayala et al., 2015; Smith et al., 2015). As dead mulches, they physically  
300 constrain weed seedling emergence (Teasdale and Mohler, 2000), by reducing light  
301 transmittance to the soil, soil maximum temperature and daily soil temperature amplitude  
302 (Teasdale and Mohler, 1993). Additionally, cover crop residues can improve the environmental  
303 conditions for the growth of seed predators or soil fauna which may destroy or degrade seeds,  
304 but field studies have reported contradictory results (Nichols et al., 2015). Living or dead,

305 cover crops may express allelopathy (Farooq et al., 2013; Gfeller et al., 2018). The diversity of  
306 the cover crop species tested and the variability of weed responses illustrates the many factors  
307 involved and the uncertain outcome according to the context.

308 In this study, despite high variability between years, some cover crop species systematically  
309 allowed to reduce weed infestation. The Asteraceae species (niger and sunflower) produced  
310 large amounts of biomass during each year of experiment and generally showed good weed  
311 control, and appear in some cases to improve maize yield. This performance may rely on their  
312 competitiveness as living plants against autumn germinating weeds and as dead mulch against  
313 spring germinating weeds, especially when shredded residues were left on the soil surface. In  
314 case of soil tillage before maize seeding, the incorporation of high amounts of carbon-rich and  
315 slowly mineralizable residues may have disturbed weed emergence through nitrogen  
316 immobilization (Justes et al., 2009) or the creation of a physical barrier (Kruidhof et al., 2008).  
317 Sunflower is also known to express allelopathy against weeds and to improve subsequent  
318 wheat growth (Alsaadawi et al., 2012).

319 Among the legume species, field pea and common vetch gave good soil cover at the end of  
320 winter despite medium to low shoot growth during autumn. These legumes were the only cover  
321 crop species with a creeping growing habit compared to the other tested species, which had a  
322 more erected architecture. A long-lasting good soil cover could partly explain the successful  
323 weed control of these species. However, common vetch failed to control weeds and was often  
324 among the less efficient species, together with berseem clover, which had a more erected  
325 architecture. Isik et al. (2009) also observed that berseem clover was the worst species in the  
326 control of weeds when used as cover crop before spring planted sweet pepper. Despite the  
327 deficient weed control by common vetch, this species shows a positive effect on maize yield in  
328 three cases, which can be probably explained by a beneficial input of nitrogen from this legume  
329 species. No assessment of nitrogen concentration in the cover crops were made in this study,  
330 but an experiment on biological nitrogen fixation conducted in a neighbouring field in the year

331 2011-2012 allowed to highlight the huge amount that legume cover crops can accumulate in  
332 only three months of growth (Büchi et al., 2015). Using the values of nitrogen fixation from  
333 this previous study combined with the biomass observed in the current study, it can be  
334 estimated that common vetch could have accumulated about 121 kgN/ha of nitrogen on  
335 average. Meanwhile, the estimates for field pea would be only 67 kgN/ha and 42 kgN/ha for  
336 berseem clover.

337 The three Brassicaceae species were characterized by intermediate to low biomass in autumn  
338 and really low residue cover in spring, and consequently did not achieved good weed control,  
339 despite potential allelopathic effects (Bangarwa and Norsworthy, 2014; Haramoto and  
340 Gallandt, 2005).

341 Black oat froze and lodged during winter, providing a thick soil cover despite intermediate  
342 growth compared to other cover crop species. Grimmer and Masiunas (2004) made a similar  
343 observation with *Avena sativa*. The good weed control of this grass in years 1 and 3 can thus be  
344 linked to the good soil cover during winter. In year 3, oat was the most efficient species against  
345 weeds in the less intensive weeding strategy (1.NoTill), and induced a really high maize yield  
346 afterwards.

347 Finally, phacelia presented an intermediate performance both in biomass production and  
348 residue cover, and was generally efficient in terms of weed control but did not appear to  
349 improve maize yield.

350

351 Overall, an increase of maize yield associated with the overall improvement of soil quality and  
352 fertility induced by the use of cover crops could have been expected (Fageria et al., 2005).

353 However, in these experiments, the maize fertilisation level was as recommended and could  
354 have thus reduced the potential differences between cover crop species, and in particular the  
355 distinction of legume versus non legume species. In addition, the weeding intervention in the  
356 early maize could have cancelled some of the residual effect of cover crop species on weed

357 infestation. Future experiments should thus be conducted in less optimal conditions, poorer  
358 soil, less fertilisers, challenging weather conditions, to reveal further the potential of cover  
359 crops to maintain maize yield. This would be of particular relevance in contexts where the  
360 access and price of herbicides do not allow to rely on these inputs, and thus where weed control  
361 is crucial to insure sufficient yield and sustainability of the system, such as in Sub-Saharan  
362 Africa.

363

## 364 **5 Conclusions**

365 Overall, the application of tillage prior to maize seeding did not always guarantee a low weed  
366 cover in the early stage of maize growth. In most cases, the cultivation of non-wintering cover  
367 crops species before maize seeding allowed to reduce weed infestation, down to only 15%  
368 weed cover in a lot of cases. The most efficient cover crop species for weed control varied  
369 from year to year, but niger, sunflower, field pea and phacelia gave the best results throughout  
370 the experiments. An effect of cover crops on maize yield could still be observed in some  
371 situations. Therefore, the use of cover crops is recommended to limit weed incidence and  
372 improve yield. Besides weed control, cover crops also provide other ecosystem services, such  
373 as soil protection during winter, nitrogen recycling or auxiliary insect promotion. However, as  
374 trade-offs between these services exist, cultivation of cover crop species mixtures may offer a  
375 solution for accumulating multiple and complementary services.

376

## 377 **Acknowledgements**

378 The authors wish to thank Cindy Bally, Nicolas Widmer and all the persons who helped with  
379 the management of experiments and data collection, and Jonne Rodenburg for help and  
380 comments on previous versions of the manuscript. We also thank the two anonymous  
381 reviewers who provided valuable comments on a previous version of this manuscript. This  
382 study was partly funded by the Swiss National Science Foundation in the framework of the



383 National Research Program NRP 68 'Sustainable Use of Soil as a Resource' [grant 406840-  
384 143063].

385 **References**

- 386 Alonso-Ayuso, M., Gabriel, J.L., Garcia-Gonzales, I., Del Monte, J.P., Quemada, M., 2018.  
387 Weed density and diversity in a long-term cover crop experiment background. *Crop*  
388 *Prot.* 112, 103-111.
- 389 Alsaadawi, I.S., Sarbout, A.K., Al-Shamma, L.M., 2012. Differential allelopathic potential of  
390 sunflower (*Sunflower*L.) genotypes on weeds and wheat (*Triticum aestivum* L.) crop.  
391 *Arch. Acker. Pfl. Boden.* 58, 1139–1148.
- 392 Bajwa, A.A., 2014. Sustainable weed management in conservation agriculture. *Crop Prot.* 65,  
393 105–113.
- 394 Bangarwa, S.K., Norsworthy, J.K., 2014. Brassicaceae cover-crop effects on weed  
395 management in plasticulture tomato. *J. Crop Improv.* 28, 145–158.
- 396 Baraibar, B., Mortensen, D.A., Hunter, M.C., Barbercheck, M.E., Kaye, J.P., Finney, D.M.,  
397 Curran, W.S., Bunchek, J., White, C.M., 2018. Growing degree days and cover crop  
398 type explain weed biomass in winter cover crops. *Agron. Sustain. Dev.* 38, 65.
- 399 Beckie, H.J., 2014. Herbicide resistance in weeds and crops: challenges and opportunities, in:  
400 Chauhan, B.S., Mahajan, G. (Eds.), *Recent Advances in Weed Management*. Springer  
401 New York, pp. 347–364.
- 402 Braz-Mota, S., Sadauskas-Henrique, H., Duarte, R.M., Val, A.L., Almeida-Val, V.M.F., 2015.  
403 Roundup (R) exposure promotes gills and liver impairments, DNA damage and  
404 inhibition of brain cholinergic activity in the Amazon teleost fish *Collossoma*  
405 *macropomum*. *Chemosphere* 135, 53–60.
- 406 Brust, J., Claupein, W., Gerhards, R., 2014. Growth and weed suppression ability of common  
407 and new cover crops in Germany. *Crop Prot.* 63, 1–8.
- 408 Buchanan, A.L., Kolb, L.N., Hooks, C.R.R., 2016. Can winter cover crops influence weed  
409 density and diversity in a reduced tillage vegetable system? *Crop Prot.* 90, 9-16.

410 Büchi, L., Gebhard, C.-A., Liebisch, F., Sinaj, S., Ramseier, H., Charles, R., 2015.  
411 Accumulation of biologically fixed nitrogen by legumes cultivated as cover crops in  
412 Switzerland. *Plant Soil* 393, 163–175.

413 Büchi, L., Wendling, M., Mouly, P., Charles, R., 2018. Comparison of Visual Assessment and  
414 Digital Image Analysis for Canopy Cover Estimation. *Agron. J.* 110, 1-7.

415 Carrera, L.M., Abdul-Baki, A.A., Teasdale, J.R., 2004. Cover crop management and weed  
416 suppression in no-tillage sweet corn production. *Hort. Science* 39, 1262–1266.

417 de Mendiburu, F., 2017. agricolae: Statistical Procedures for Agricultural Research. R package  
418 version 1.2-8.

419 Dorn, B., Jossi, W., van der Heijden, M.G.A., 2015. Weed suppression by cover crops:  
420 comparative on-farm experiments under integrated and organic conservation tillage.  
421 *Weed Res.* 55, 586–597.

422 Eslami, S.V., 2014. Weed management in conservation agriculture systems, in: Chauhan, B.S.,  
423 Mahajan, G. (Eds.), *Recent Advances in Weed Management*. Springer New York, pp.  
424 87–124.

425 Fageria, N.K., Baligar, V.C., Bailey, B.A., 2005. Role of cover crops in improving soil and  
426 row crop productivity. *Commun. Soil Sci. Plant* 36, 2733–2757.

427 Farooq, M., Bajwa, A.A., Cheema, S.A., Cheema, Z.A., 2013. Application of allelopathy in  
428 crop production. *Intern. J. Agric. Biol.* 15, 1367–1378.

429 Gaupp-Berghausen, M., Hofer, M., Rewald, B., Zaller, J.G., 2015. Glyphosate-based  
430 herbicides reduce the activity and reproduction of earthworms and lead to increased soil  
431 nutrient concentrations. *Sci. Rep.* 5, 12886–12886.

432 Gfeller, A., Herrera, J.M., Tschuy, F., Wirth, J., 2018. Explanations for *Amaranthus retroflexus*  
433 growth suppression by cover crops. *Crop Prot.* 104, 11-20.

434 Giller, K., Witter, E., Corbeels, M., Tittonell, P., 2009. Conservation agriculture and  
435 smallholder farming in Africa: The heretics' view. *Field Crops Res.* 114, 23-34.

436 Grimmer, O.P., Masiunas, J.B., 2004. Evaluation of winter-killed cover crops preceding snap  
437 pea. *Hort. Technology* 14, 349–355.

438 Haramoto, E.R., Gallandt, E.R., 2005. Brassica cover cropping: I. Effects on weed and crop  
439 establishment. *Weed Sci.* 53, 695–701.

440 Hayden, Z.D., Ngouajio, M., Brainard, D.C., 2014. Rye–vetch mixture proportion tradeoffs:  
441 cover crop productivity, nitrogen accumulation, and weed suppression. *Agron. J.* 106,  
442 904.

443 Isik, D., Kaya, E., Ngouajio, M., Mennan, H., 2009. Weed suppression in organic pepper  
444 (*Capsicum annuum* L.) with winter cover crops. *Crop Prot.* 28, 356–363.

445 Justes, E., Mary, B., Nicolardot, B., 2009. Quantifying and modelling C and N mineralization  
446 kinetics of catch crop residues in soil: parameterization of the residue decomposition  
447 module of STICS model for mature and non mature residues. *Plant Soil* 325, 171–185.

448 Kruidhof, H.M., Bastiaans, L., Kropff, M.J., 2008. Ecological weed management by cover  
449 cropping: effects on weed growth in autumn and weed establishment in spring. *Weed*  
450 *Res.* 48, 492–502.

451 Lahmar, R., 2010. Adoption of conservation agriculture in Europe: Lessons of the KASSA  
452 project. *Land Use Policy* 27, 4–10.

453 Melander, B., Munier-Jolain, N., Charles, R., Wirth, J., Schwarz, J., van der Weide, R., Bonin,  
454 L., Jensen, P.K., Kudsk, P., 2013. European perspectives on the adoption of  
455 nonchemical weed management in reduced-tillage systems for arable crops. *Weed*  
456 *Technol.* 27, 231–240.

457 Mirsky, S.B., Ryan, M.R., Teasdale, J.R., Curran, W.S., Reberg-Horton, C.S., Spargo, J.T.,  
458 Wells, M.S., Keene, C.L., Moyer, J.W., 2013. Overcoming weed management  
459 challenges in cover crop–based organic rotational no-till soybean production in the  
460 eastern United States. *Weed Technol.* 27, 193–203.

461 Nalewaja, J.D., 2003. Weeds and Conservation Agriculture, in: García-Torres, L., Benites, J.,  
462 Martínez-Vilela, A., Holgado-Cabrera, A. (Eds.), Conservation Agriculture. Springer  
463 Netherlands, pp. 201–210.

464 Nauen, R., Denholm, I., 2005. Resistance of insect pests to neonicotinoid insecticides: Current  
465 status and future prospects. *Arch. Insect Biochem. Physiol.* 58, 200–215.

466 Nichols, V., Verhulst, N., Cox, R., Govaerts, B., 2015. Weed dynamics and conservation  
467 agriculture principles: A review. *Field Crop. Res.* 183, 56–68.

468 Page, E.R., Cerrudo, D., Westra, P., Loux, M., Smith, K., Foresman, C., Wright, H., Swanton,  
469 C.J., 2012. Why early season weed control is important in maize? *Weed Sci.* 60, 423–  
470 430.

471 Peigné, J., Ball, B.C., Roger-Estrade, J., David, C., 2007. Is conservation tillage suitable for  
472 organic farming? A review. *Soil Use Manage.* 23, 129–144.

473 R Core Team, 2019. R: A language and environment for statistical computing. R Foundation  
474 for Statistical Computing, Vienna, Austria.

475 Ramesh, K., 2015. Weed problems, ecology, and management options in conservation  
476 agriculture: issues and perspectives. *Adv. Agron.* 251–303.

477 Rueda-Ayala, V., Jaeck, O., Gerhards, R., 2015. Investigation of biochemical and competitive  
478 effects of cover crops on crops and weeds. *Crop Prot.* 71, 79–87.

479 Sadeghpour, A., Hashemi, M., DaCosta, M., Jahanzad, E., Herbert, S.J., 2014. Switchgrass  
480 establishment influenced by cover crop, tillage systems, and weed control. *Bioenerg.*  
481 *Res.* 7, 1402–1410.

482 Sainju, U.M., Singh, B.P., Whitehead, W.F., 2002. Long-term effects of tillage, cover crops,  
483 and nitrogen fertilization on organic carbon and nitrogen concentrations in sandy loam  
484 soils in Georgia, USA. *Soil Till. Res.* 63, 167–179.

485 Sihtmäe, M., Blinova, I., Künnis-Beres, K., Kanarbik, L., Heinlaan, M., Kahru, A., 2013.  
486 Ecotoxicological effects of different glyphosate formulations. *Appl. Soil Ecol.* 72, 215–  
487 224.

488 Smith, R.G., Atwood, L.W., Pollnac, F.W., Warren, N.D., 2015. Cover-crop species as distinct  
489 biotic filters in weed community assembly. *Weed Sci.* 63, 282–295.

490 Teasdale, J.R., Mohler, C.L., 1993. Light transmittance, soil temperature, and soil moisture  
491 under residue of hairy vetch and rye. *Agron. J.* 85, 673.

492 Teasdale, J.R., Mohler, C.L., 2000. The quantitative relationship between weed emergence and  
493 the physical properties of mulches. *Weed Sci.* 48, 385–392.

494 Walsh, M.J., Powles, S.B., 2014. Management of herbicide resistance in wheat cropping  
495 systems: learning from the Australian experience. *Pest Manag. Sci.* 70, 1324–1328.

496 Weber, F.J., Kunz, C., Peteinatos, G.G., Zikeli, S., Gerhards, R., 2017. Weed Control Using  
497 Conventional Tillage, Reduced Tillage, No-Tillage, and Cover Crops in Organic  
498 Soybean. *Agriculture* 7.

499 Wendling, M., Charles, R., Herrera, J.M., Amossé, C., Jeangros, B., Walter, A., Büchi L.  
500 Effect of species identity and diversity on biomass production and its stability in cover  
501 crop mixtures. *Agr. Ecosyst. Environ.* *In press*.

502 Williams, M.M., Mortensen, D.A., Doran, J.W., 1998. Assessment of weed and crop fitness in  
503 cover crop residues for integrated weed management. *Weed Sci.* 46, 595–603.

504

505 **Table and figure captions**

506 **Table 1** Name, botanical family, code and seeding rate for the ten cover crop species.

507 **Table 2** Mean dry biomass of cover crops (t/ha) in autumn, and residue cover (%) in early  
508 spring for the three year of experiment with biomass/cover estimation. Within each column,  
509 values with the same letters are not significantly different (Tukey HSD test,  $p < 0.05$ ).

510 **Figure 1** Schematic representation of the implemented weeding strategies and observations for  
511 the four years of experiment.

512 **Figure 2** Relative values (to the yearly average of all ten cover crop species values) of biomass  
513 production in autumn and residue cover in early spring for the ten cover crop species, for the  
514 three years of experiment with biomass/cover estimation.

515 **Figure 3** Weed cover [%] at early maize stage. A. No cover crop control plots. Different letters  
516 indicate significant differences ( $p < 0.05$ ) between weeding strategies, within each year. B.  
517 Cover crop treatment plots. The solid horizontal black lines show the mean weed cover in the  
518 respective no cover crop controls (i.e. mean values of the boxplots shown in panel A). Cover  
519 crop species showing significantly different weed cover than the control are indicated with the  
520 code of the species name, non significant ones are shown with a circle. The horizontal dotted  
521 line shows the 5% threshold.

522 **Figure 4** Maize yield (dry silage) [t/ha]. A. No cover crop control plots. Different letters  
523 indicate significant differences ( $p < 0.05$ ) between weeding strategies, within each year. B.  
524 Cover crop treatment plots. The solid horizontal black lines show the mean maize yield in the  
525 respective no cover crop controls (i.e. mean values of the boxplots shown in panel A). Cover  
526 crop species showing significantly different weed cover than the control are indicated with the  
527 code of the species name, non significant ones are shown with a circle.

528

529

530

531

532 **Supplementary Material**

533 **Table S1** Soil characteristics, crop management and observation dates for each year of  
534 experiment

535 **Table S2** Mean and standard error of weed cover at early maize stage (2-4 leaf stage) for each  
536 cover crop treatment, weeding strategy and year of experiment.

537 **Table S3** Mean and standard error of maize yield for each cover crop treatment, weeding  
538 strategy and year of experiment.

539 **Figure S1** Cover crop biomass in autumn (A.) and residue cover in spring (B.) for each year of  
540 experiment.

541



542 **Table 1**

Latin name	Common name	Family	Code	Mean seeding rate grain/m <sup>2</sup>
<i>Brassica juncea</i> , (L.) Czern.	Indian mustard	Brassicaceae	b1	515
<i>Brassica rapa</i> L., var <i>campestris</i>	Turnip rape	Brassicaceae	b2	630
<i>Raphanus sativus</i> L., var <i>longipinnatus</i>	Daikon radish*	Brassicaceae	b3	110
<i>Pisum sativum</i> , L.	Field pea	Fabaceae	f1	135
<i>Trifolium alexandrinum</i> , L.	Berseem clover	Fabaceae	f2	605
<i>Vicia sativa</i> , L.	Common vetch*	Fabaceae	f3	225
<i>Avena strigosa</i> , Schreb.	Black oat*	Poaceae	p1	490
<i>Guizotia abyssinica</i> , (L.f.) Cass.	Niger	Asteraceae	a1	270
<i>Helianthus annuus</i> , L.	Sunflower	Asteraceae	a2	75
<i>Phacelia tanacetifolia</i> , Benth.	Phacelia	Hydrophyllaceae	h1	450

543 \* these three species were not tested in year 0


544


545

Table 2

Species	Dry biomass [t/ha]				Residue cover [%]									
	year 1	year 2	year 3	mean	year 1	year 2	year 3	mean						
<b>b1</b> Indian mustard	3.5	bc	5.1	bc	7.3	bc	5.3	10	de	15	f	13	de	13
<b>b2</b> Turnip rape	1.6	cd	1.3	d	5.2	cd	2.7	7	e	15	f	18	cde	13
<b>b3</b> Daikon radish	1.2	d	1.6	d	4.9	cd	2.6	5	e	22	ef	6	e	11
<b>f1</b> Field pea	3.2	bcd	3.1	cd	3.4	d	3.2	87	a	87	ab	46	b	73
<b>f2</b> Berseem clover	3.6	bc	3.2	cd	5.0	cd	3.9	70	ab	58	cd	37	bc	55
<b>f3</b> Common vetch	5.2	ab	3.4	cd	3.4	d	4.0	90	a	92	a	82	a	88
<b>p1</b> Black oat	3.6	bc	4.1	cd	9.2	ab	5.6	70	ab	73	bc	86	a	77
<b>a1</b> Niger	4.2	b	7.4	ab	7.6	bc	6.4	43	bc	52	d	31	bcd	42
<b>a2</b> Sunflower	6.5	a	8.4	a	##	a	8.7	27	cde	33	e	24	bcde	28
<b>h1</b> Phacelia	4.5	ab	4.7	bc	5.3	cd	4.8	40	bcd	60	cd	27	bcde	42
<b>mean</b>	3.7		4.2		6.2		4.7	45		51		37		44

546









**Table S1** Soil characteristics, crop management and observation dates for each year of experiment.

<b>Experiment</b>	<b>year 0 2009-2010</b>	<b>year 1 2010-2011</b>	<b>year 2 2011-2012</b>	<b>year 3 2013-2014</b>
<i>Soil characteristics</i>				
Clay (%)	26.2	24.6	32.1	26.1
Sand (%)	31.6	29.7	34.1	28.5
USDA texture	Loam	Loam	Clay loam	Loam
pH	8.0	7.7	7.3	7.4
Organic matter (%)	2.1	1.9	3.2	2.4
<i>Cover crop and maize management</i>				
Cover crop seeding	31.07.2009	29.07.2010	29.07.2011	06.08.2013
Cover crop residue shredding	25.05.2010	04.05.2011	03.04.2012	05.05.2014
Cover crop residue harrowing*	25.05.2010	04.05.2011	30.04.2012	05.05.2014
Maize seeding	26.05.2010	04.05.2011	30.04.2012	06.05.2014
Herbicide	-	28.05.2011	26.05.2012	03.06.2014
Maize harvest	-	06.09.2011	23.08.2012	12.09.2014
<i>Observation dates</i>				
Cover crop biomass [DAS]	-	03.11.2010 [97]	02.11.2011 [96]	11.11.2013 [97]
Cover crop residue cover	-	04.05.2011	03.04.2012	19.03.2014
<b>Weed cover (2-4 leaf stage of maize) [DAS]</b>	<b>24.06.2010 [29]</b>	<b>27.05.2011 [23]</b>	<b>25.05.2012 [25]</b>	<b>26.05.2014 [20]</b>

\* Only on the respective treatments (see Figure 1)

**Table S2** Mean and standard error of weed cover at early maize stage (2-4 leaf stage) for each cover crop treatment, weeding strategy and year of experiment.


Species		Mean weed cover [%] +- 1*se															
		year 0				year 1				year 2				year 3			
		1. NoTill		2. MinTill		1. NoTill		2. MinTill		1. NoTill		2. MinTill		1. NoTill		2. MinTill	
b1	Indian mustard	87	4.4	77	3.3	43	8.8	2	1.5	16	4.0	13	4.4	58	13	4	3.0
b2	Turnip rape	87	3.3	73	6.0	40	5.8	1	0.3	19	0.1	12	4.4	45	16	3	1.9
b3	Daikon radish					40	10.0	1	0.0	22	1.5	30	10.0	28	4	1	0.3
f1	Field pea	75	14.1	58	12.0	33	6.7	2	1.3	13	6.2	18	6.0	42	16	2	0.3
f2	Berseem clover	80	10.1	78	1.7	47	12.0	1	0.0	41	2.2	50	11.5	52	10	4	1.5
f3	Common vetch					30	10.0	1	0.0	44	10.6	53	13.3	30	10	1	0.0
p1	Black oat					43	13.3	2	1.3	29	4.7	27	3.3	11	5	1	0.0
a1	Niger	73	1.7	58	4.4	27	3.3	2	1.3	12	3.9	12	3.3	22	7	1	0.3
a2	Sunflower	80	2.9	62	1.7	43	20.3	1	0.0	16	6.1	4	1.3	47	9	2	0.6
h1	Phacelia	77	8.8	68	1.7	53	13.3	1	0.0	22	1.5	27	7.3	62	14	4	1.8
	no cover crop control	90	2.6	83	1.7	53	3.3	12	9.1	43	4.8	49	6.0	62	11	3	1.2
	mean	81		70		41		2		25		27		42		3	



**Table S3** Mean and standard error of maize yield for each cover crop treatment, weeding strategy and year of experiment.

**Mean maize yield [t/ha] +- 1\*se**

Species		year 1				year 2				year 3			
		1. NoTill		2. MinTill		1. NoTill		2. MinTill		1. NoTill		2. MinTill	
b1	Indian mustard	16.9	1.2	17.2	0.5	13.3	0.5	14.3	0.2	12.3	0.7	16.2	0.6
b2	Turnip rape	16.1	1.6	15	0.4	13.2	0.4	12.3	0.4	13.8	0.9	17.2	1.5
b3	Daikon radish	16.5	0.3	16.1	0.6	14.3	0.6	15.2	1.3	12.6	0.5	17.7	0.4
f1	Field pea	17.9	1.1	18.1	0.0	13.9	0.2	13.7	0.6	12.9	0.6	17.4	0.5
f2	Berseem clover	15.3	1.0	18.1	0.6	14	0.8	16	1.6	9.27	3.1	16.7	1.0
f3	Common vetch	18.1	0.8	19.7	0.3	13.5	0.6	14.6	0.8	13.7	0.3	17.8	0.7
p1	Black oat	17.1	1.5	16.7	0.0	13.3	0.6	14.4	0.5	15.6	0.1	17.4	0.4
a1	Niger	17.4	1.1	16.7	0.1	13.6	1.0	14	0.2	13.1	1.4	18.4	1.2
a2	Sunflower	16.3	0.5	17.4	0.7	14.5	0.7	14.8	1.1	11.8	0.9	18.1	0.6
h1	Phacelia	15.2	0.6	17.5	0.4	14.8	1.0	13.2	0.5	8.77	3.3	16.5	0.5
	no cover crop control	13.5	0.2	16.9	0.3	14.1	0.4	13.3	0.5	11.3	1.4	14.6	1.0
	mean	16.4		17.2		13.9		14.2		12.3		17.1	



**Figure S1** Cover crop biomass in autumn (A.) and residue cover in spring (B.) for each year of experiment.