

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17

This is the final peer-reviewed accepted manuscript of:  
Merlo B, Zambelli D, Cunto M, Iacono E, Nasi L, Giaretta E, Galeati G, Bucci D, Spinaci M. Sex-sorted canine sperm cryopreservation: limits and procedural considerations. Theriogenology Volume 83, Issue 7, 15 April 2015, Pages 1121-1127  
The final published version is available online at:  
[10.1016/j.theriogenology.2014.12.018](http://dx.doi.org/10.1016/j.theriogenology.2014.12.018)

©2015. This manuscript version is made available under the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) 4.0 International License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

18 Sex-sorted canine sperm cryopreservation: limits and procedural considerations

19

20 Barbara Merlo<sup>a\*</sup>, DVM, PhD, ECAR Diplomate, Associate Professor

21 Daniele Zambelli<sup>a</sup>, DVM, PhD, ECAR Diplomate, Associate Professor

22 Marco Cunto<sup>a</sup>, DVM, PhD, Researcher

23 Iacono Eleonora<sup>a</sup>, DVM, PhD, Associate Professor

24 Ludovica Nasi<sup>a</sup>, DVM

25 Elisa Giaretta<sup>a</sup>, PhD student

26 Giovanna Galeati<sup>a</sup>, DVM, PhD, Associate Professor

27 Diego Bucci<sup>a</sup>, DVM, PhD, Research grant student

28 Marcella Spinaci<sup>a</sup>, DVM, PhD, Associate Professor

29

30 <sup>a</sup>Department of Veterinary Medical Sciences, University of Bologna, via Tolara di Sopra 50, 40064

31 Ozzano Emilia (BO), Italy.

32

33 \*Corresponding Author at: Barbara Merlo, DVM, PhD, ECAR Diplomate, Researcher, Department

34 of Veterinary Medical Sciences, University of Bologna, via Tolara di Sopra 50, 40064 Ozzano

35 Emilia (BO), Italy.

36 Tel.: +39-051-2097567

37 E-mail: [barbara.merlo@unibo.it](mailto:barbara.merlo@unibo.it)

38

39

40 **Abstract**

41

42 The aim of this study was to define a protocol to store dog sperm pre- and post-sorting in order to  
43 obtain an insemination dose sufficient to allow the conception by artificial insemination.

44 Experiment 1 and 2 were performed to evaluate the more appropriate extender for preserving at  
45 room temperature dog sperm pre and post sorting. Four extenders were tested: a) Tris-fructose-  
46 citrate (TFC), b) Tris-glucose-citrate (TGC), c) modified Tyrode's albumin-lactate-pyruvate  
47 medium (mTALP), d) third fraction of the ejaculate (after centrifugation at 5000 x g for 10 min) (III  
48 FRAC). Experiment 3 and 4 were performed to evaluate the ability of dog semen to withstand sex-  
49 sorting and freezing/thawing. mTALP was the best extender for canine sperm storage at room  
50 temperature (20-25°C) pre- (total motility: TFC  $8.3 \pm 1.7$ , TGC  $50.0 \pm 11.5$ , mTALP  $70.0 \pm 0.1$ , III  
51 FRAC  $25.0 \pm 1.0$ ,  $P < 0.05$ ) and post-sorting (total motility: TFC  $7.3 \pm 1.5$ , TGC  $10.3 \pm 1.5$ ,  
52 mTALP  $33.3 \pm 6.7$ , III FRAC  $8.7 \pm 5.8$ ,  $P < 0.05$ ), even if at 24h sorted sperm quality was impaired  
53 in all extenders tested herein. Sperm quality decreased after sorting (total motility: control  $92.5 \pm$   
54  $0.9$ , sorted  $52.9 \pm 6.0$ ,  $P < 0,05$ ) and, especially, after freezing/thawing (total motility: frozen-  
55 control  $25.7 \pm 4.1$ , frozen-sorted  $2.4 \pm 1.2$ ,  $P < 0,05$ ). In conclusion mTALP is an appropriate  
56 medium for canine sperm storage pre- and soon after sorting (hours), but a long storage period of  
57 sexed sperm at room temperature is not adequate. Cryopreservation greatly impaired sperm quality  
58 and further studies are needed to optimize the freezing protocol for sexed dog sperm.

59

60 **Keywords:** dog, spermatozoa, sex-sorting, extender, cryopreservation

61

## 62 **1. Introduction**

63 Flow cytometric sperm sorting based on X and Y sperm DNA difference has been established as the  
64 only effective method to pre-determinate the sex of mammalian offspring before fertilization.

65 While sexing technique has already reached a commercial level in the bovine species, sorting  
66 efficacy has been demonstrated suggesting future applications in a variety of mammals (pig, horse,  
67 sheep, goat, cat, endangered species) [1].

68 The production of animals of a predetermined sex by sperm sexing is an interesting target also in  
69 dog reproduction as, according to the different fields of breeding and use of animals (purebred pet  
70 dog, working dogs such as guide dogs, rescue dogs, farm dogs), male or female are preferably  
71 sought by dog breeders and owners [2-5].

72 The possible association of reproductive technology, such as cryopreservation and artificial  
73 insemination (AI), with sex sorting of sperm could optimize the profitability of the breeding and  
74 should have a great potential application also for the preservation of endangered canids maintaining  
75 the best male-female ratio for the animal repopulation [6].

76 On the best of our knowledge only a couple of studies have been performed until now on dog sperm  
77 sexing [3,5].

78 One of the major limitation for the use of sperm sexing in the dog, as well as in other species, is the  
79 long sorting time necessary in order to obtain the number of sexed spermatozoa necessary for  
80 artificial insemination (AI) (approximately 150-200 million motile spermatozoa) [7]. Rodenas et  
81 al.,[5] reported that is possible to collect  $14-16 \times 10^6$  dog sexed spermatozoa per hour in the best  
82 samples; therefore conventional AI is not feasible in this species. Moreover it has been reported that  
83 spermatozoa can be damaged during the sorting process [1]. The possibility to overcome this  
84 limitation by reducing both the number of spermatozoa and the distance from the site of fertilization  
85 has been demonstrated by Meyers et al. [3] producing offspring of the desired sex in one bitch by  
86 intrauterine low-dose insemination using sexed spermatozoa.

87 Furthermore, the number of spermatozoa requested in dog for a intrauterine low-dose insemination  
88 (at least 50 million) [7] is very high considering the sorting time and, therefore, sperm cells have to  
89 be stored properly at room temperature waiting to be sexed.

90 Another problem to solve is the storage of sexed semen, as the bitch can be far from the sorting  
91 facilities or more than one insemination could be required. Until now no data about storage of  
92 canine sexed semen are available in the literature.

93 Cryopreservation permits long-term sperm storage, but it has to be taken in mind that sexed  
94 spermatozoa are cells already stressed by the sorting procedure and usually present a greater  
95 susceptibility to cryopreservation procedures compared to unsexed semen.

96 The aim of this study was to define: 1) an appropriate extender in order to liquid store dog sperm  
97 pre- and post-sorting; 2) evaluate if sexed canine spermatozoa are able to withstand  
98 freezing/thawing procedure. For this purpose motility, viability and acrosome integrity were  
99 assessed in unsorted and sorted sperm either maintained at room temperature or cryopreserved.

100

## 101 **2. Materials and methods**

102

### 103 ***2.1 Experimental design***

104

105 This study was divided into 4 experiments.

106 As the sorting process is slow, the aim of the experiment 1 was to evaluate which of four extenders  
107 is the more appropriate for storing at room temperature dog spermatozoa waiting to be sexed. Total  
108 motile spermatozoa, movement, viability and acrosome integrity were evaluated at 0, 2, 4, 6, 8, and  
109 24h.

110 In the experiment 2, the best extender from experiment 1 (mTALP, see below) was used for diluting  
111 pre-sorted spermatozoa while the four extenders evaluated in the experiment 1 were tested for  
112 sperm storage at room temperature (20-25°C) after sorting. The above mentioned parameters were  
113 evaluated at 0, 2, 4, 24h after sorting.

114 The aim of the experiment 3 was to evaluate if dog spermatozoa are able to withstand sexing  
115 procedure followed by freezing. For that purpose ejaculates from purebred dogs (Australian  
116 Shepherd) were analyzed: immediately after collection (CTR), after the sorting (SORT) and after  
117 freezing and thawing of sperm for all groups (CTR-FR, SORT-FR).

118 In the experiment 4 the same protocol of the experiment 3 was performed on semen from mixed  
119 breed dogs.

120

## 121 ***2.2 Extenders***

122

123 All reagents were purchased from Sigma-Aldrich (Milan, Italy) unless otherwise stated.

124 Extenders tested in experiment 1 were Tris-fructose-citrate (TFC) (3% wt/vol Tris, 1.7% wt/vol  
125 citric acid, 1.25% wt/vol fructose, 0.06% wt/vol Na benzyl-penicillin, 0.1% wt/vol streptomycin  
126 sulphate in distilled water), Tris-glucose-citrate (TGC) (2.4% wt/vol Tris, 1.4% wt/vol citric acid,  
127 0.8% wt/vol glucose, 0.06% wt/vol sodium benzyl-penicillin, 0.1% wt/vol streptomycin sulphate in  
128 distilled water), modified Tyrode's solution (mTALP) (3.3 mM MgCl<sub>2</sub>, 99 mM NaCl, 3.1 mM KCl,  
129 25 mM NaHCO<sub>3</sub>, 0.35 Mm NaH<sub>2</sub>PO<sub>4</sub>, 1 Mm sodium pyruvate, 21.6 mM DL lactic acid, 0.125 mM  
130 Kanamycin in distilled water), third fraction (III FRAC) of the ejaculated after centrifugation at  
131 5000 x g for 10 min.

132 Freezing extender for the first step was TGC supplemented with 20% (v/v) egg yolk, 3% (v/v)  
133 glycerol (EYTGCl), while for the second step TGC was supplemented with 20% (v/v) egg yolk,

134 7% (v/v) glycerol and 1 % (v/v) Equex STM Paste (Nova Chemical Sales Inc., Scituate, Ma, USA)  
135 (EYTGC2).

136

### 137 ***2.3 Semen collection and evaluation***

138

139 A total of 14 adult mixed breed and purebred male dogs between 1 and 7 years of age were enrolled  
140 in these experiments, between January 2008 and March 2013. The three fractions of the ejaculates  
141 were collected in calibrated plastic vials by digital manipulation. The experiment was approved by  
142 the Ethic-scientific Committee of Alma Mater Studiorum, University of Bologna.

143 The sperm-rich fraction was evaluated for volume, concentration, total sperm motility, movement,  
144 sperm morphology and membrane integrity to estimate sperm viability.

145 The volume was measured by a calibrated micro-pipette and sperm concentration was determined  
146 with a Bürker chamber, after dilution of the sperm suspension 1:40 with buffered formol saline  
147 (BFS) to immobilize spermatozoa.

148 The percentage of total motile spermatozoa and movement score, based on the type of the forward  
149 movement of sperm (scale of 0–5; 0 = no forward movement, 5 = steady, rapid forward  
150 progression) [8] were subjectively estimated at a phase contrast microscope (x400) (AxioLab, Zeiss,  
151 Italy) equipped with a warming plate (37°C) (Thermo Plate, Tokai Hit, Japan). Percentages of  
152 morphologically normal spermatozoa were determined at the same microscope (x1000) after  
153 dilution of semen 1:1 with BFS, and at least 200 spermatozoa per sample were examined.

154 In order to evaluate plasma membrane integrity, 25 µL of semen were incubated with 2 µL of a 300  
155 µM propidium iodide (PI) stock solution, 2 µL of a 10 µM SYBR-14 (green) stock solution, both  
156 obtained from the live/dead sperm viability kit (Life Technologies, Molecular Probes®, Monza,  
157 Italy) for 5 min at 37°C in the darkness. Aliquots of the stained suspensions were placed on clean  
158 microscope slides, overlaid carefully with coverslips and at least 200 spermatozoa per sample were

159 scored with Nikon Eclipse E 600 epifluorescence microscope (Nikon Europe BV, Badhoevedop,  
160 The Netherlands). Spermatozoa stained with SYBR-14 and not stained with PI were considered as  
161 viable. Spermatozoa SYBR positive and PI positive and those SYBR negative / PI positive were  
162 considered as cells with non-intact membranes or dead as an estimate of sperm viability.

163 Acrosome integrity was measured with a FITC-coniugated lectin from *Pisum sativum* (FITC-PSA)  
164 which labelled acrosomal matrix glycoproteins. Briefly, spermatozoa were washed twice in PBS,  
165 resuspended with ethanol 95% and fixed at 4°C for 30 min. Samples were dried in heated slides and  
166 incubated with FITC-PSA solution (5 µg PSA-FITC/1 mL H<sub>2</sub>O) for 15 min in darkness. After  
167 staining samples were washed in PBS and mounted with Vectashield mounting medium with PI  
168 (Vector Laboratories, Burlingame, CA, USA). The slides were then observed with the above  
169 mentioned microscope and at least 200 spermatozoa per sample were scored.

170 The presence of a green acrosomal fluorescence was considered indicative for an intact acrosome,  
171 while a partial or total absence of fluorescence is indicative of acrosome disruption or acrosome  
172 reaction.

173

#### 174 ***2.4 Sperm sorting***

175

176 Sperm samples were diluted with mTALP to 100 x 10<sup>6</sup> spermatozoa/mL. Aliquots of 1 mL of  
177 diluted semen were then transferred into Falcon tubes and stained with 10 µl of 5 mg/mL Hoechst  
178 33342 stock solution for 1 h at 35° C in the dark. Just prior to sorting, 1 µL of food dye (FD&C#40,  
179 Warner Jenkinson, St. Louis, MO, USA) stock solution (25 mg/mL) was added to each sample. The  
180 samples were then filtered through a 60 µm nylon mesh filter to remove debris or clumped  
181 spermatozoa.

182 A MoFlo SX® flow cytometer/sperm sorter (DakoCytomation Inc., Fort Collins, CO, USA)  
183 equipped with an argon laser (wavelength 351 at 150 mW) was used. Dulbecco's phosphate



184 buffered saline (DPBS) served as sheath fluid and the instrument sheath pressure was 40 psi. Sorted  
185 spermatozoa were deflected into 20 mL polypropylene tubes containing 500  $\mu$ L of 2 % Tes-Tris-  
186 egg yolk buffer [9]. After collection of a total of  $20 \times 10^6$  sperm/sex (around  $7 \times 10^6$  sperm/ tube),  
187 the two populations were pooled (since sex predetermination was not an objective of the  
188 experiment). The samples were divided in four aliquots in experiment 2 and after a centrifugation at  
189  $800 \times g$  for 20 min the pellets were resuspended in the different extenders tested (TFC, TGC,  
190 mTALP, III FRAC). In experiments 3 and 4, after centrifugation, the pellets were resuspended in  
191 EYTGC1 .

192

### 193 ***2.5 Freezing and thawing***

194

195 After centrifugation at  $300 g$  for 10 min (CTR group) and after the centrifugation performed in  
196 order to concentrate sorted sperm ( $800 \times g$  for 20 min) (SORT group), the supernatant was removed  
197 and the sperm pellets (about of  $40 \times 10^6$  spermatozoa/pellet) were resuspended in two steps in  
198 freezing extender into a 1.5 mL conical centrifuge tube. Firstly, spermatozoa were diluted slowly at  
199 room temperature ( $20\text{-}25^\circ\text{C}$ ) by dropwise addition of  $250 \mu\text{L}$  of EYTGC1. After that, semen was  
200 cooled to  $4^\circ\text{C}$  for 60 min in a controlled temperature refrigerator (about  $-0.3^\circ\text{C}/\text{min}$ ). Then the  
201 second dilution with  $250 \mu\text{L}$  of pre-cooled EYTGC2 was made at  $4^\circ\text{C}$ , reaching a final  
202 concentration of  $80 \times 10^6$  spermatozoa/mL. After an additional equilibration period of 15 min at  
203  $4^\circ\text{C}$ , semen samples were loaded in plastic straws ( $0.5 \text{ mL}$ ) and the straws were heat sealed. Sperm  
204 were frozen on liquid nitrogen vapors into a polystyrene box ( $36 \times 33 \times 30 \text{ cm}$ ) keeping straws at  
205 distance of 7 cm from liquid nitrogen (4 cm) for 10 min, before plunging straws directly into liquid  
206 nitrogen. Finally, the straws were transferred into a Dewar flask at  $-196^\circ\text{C}$ .

207 After at least 4 d of storage, one straw of each sample was thawed by immersion in a  $37^\circ\text{C}$  water  
208 bath for 30 s and the content emptied into an 1.5 mL conical centrifuge tube for evaluations.

209

## 210 **2.6 Statistical analysis**

211

212 Values are reported as mean  $\pm$  standard error of the mean (SEM). Data were checked for normality  
213 using a Shapiro-Wilk test and analyzed using a one-way ANOVA or a Kruskal-Wallis H test  
214 depending on distribution (IBM SPSS Statistics 21, IBM Corporation, Milan, Italy). When  
215 significant differences were found Tukey post-hoc test was performed. The level of significance  
216 was set at  $P < 0.05$ .

217

## 218 **3. Results**

219

### 220 **3.1 Experiment 1**

221

222 Mean data from the evaluation of 4 ejaculates from 4 mixed breed dogs after collection were:  
223 volume  $0.7 \pm 0.3$  mL, concentration  $306 \pm 108 \times 10^6$  spermatozoa/mL, total motility  $86.7 \pm 3.3\%$ ,  
224 movement  $5.0 \pm 0.0$ , and rate of morphologically normal spermatozoa greater than 90%.

225 The results of experiment 1 are reported in Figure 1.

226 After 24h storage at room temperature total sperm motility was higher in mTALP compared to the  
227 TFC and III FRAC (Fig.1A) ( $P < 0.05$ ). Sperm movement was significantly higher in mTALP  
228 compared to III FRAC at 6h (Fig.1B) ( $P < 0.05$ ). No differences were observed in the percentage of  
229 sperm with intact plasma membrane and sperm with intact acrosome in the different media (Fig.1C  
230 and D).

231

### 232 **3.2 Experiment 2**

233

234 Mean data from the evaluation of 4 ejaculates from 4 mixed breed dogs after collection were:  
235 volume  $1.6 \pm 0.6$  mL, concentration  $154.3 \pm 2.2 \times 10^6$  spermatozoa/mL, total motility  $91.7 \pm 1.7\%$ ,  
236 movement  $5.0 \pm 0.0$ , and rate of morphologically normal spermatozoa greater than 90%.

237 Four h after sorting, total motility and movement were significantly higher in semen resuspended in  
238 mTALP (Fig.2 A e B) ( $P < 0.05$ ). The percentage of sperm with intact membranes in sorted group  
239 resuspended in mTALP compared to TFC and TGC resulted higher at 0 and 4h (Fig. 2 C) ( $P < 0.05$ ).  
240 The difference between mTALP and TGC was significant also after 24h ( $P < 0.05$ ). Regarding semen  
241 resuspended in III FRAC, sorted spermatozoa presented a percentage of viable cells similar to  
242 mTALP at 0h; this percentage, however, rapidly declined resulting the lowest already at 4h  
243 ( $P < 0.05$ ). The resuspension of sorted semen in mTALP resulted in a percentage of acrosome intact  
244 sperm higher than the other extenders at 2 and 4h (Fig. 2 D) ( $P < 0.05$ ).

245

### 246 **3.3 Experiment 3**

247

248 Mean data from the evaluation of 8 ejaculates from 8 purebred dogs after collection were: volume  
249  $2.3 \pm 0.5$  mL, concentration  $159 \pm 22 \times 10^6$  spermatozoa/mL, total motility  $92.5 \pm 0.9\%$ , movement  
250  $5.0 \pm 0.0$ , and rate of morphologically normal spermatozoa greater than 90%.

251 The sorting procedure induced a significant reduction of total motility and movement compared to  
252 control semen evaluated immediately after collection (CTR) ( $P < 0.05$ ). Freezing caused a significant  
253 decrease ( $P < 0.05$ ) of total motile sperm and movement in both CTR-FR and SORT-FR samples  
254 (Tab. 1).

255 The percentage of sperm with intact membranes and that of acrosome intact cells did not differ  
256 between CTR and SORT but freezing induced a significant reduction of these parameters in both  
257 CTR-FR and SORT-FR samples ( $P < 0.05$ ).

258

### 259 **3.4 Experiment 4**

260

261 Mean data from evaluation of 8 ejaculates from 2 mixed breed dogs after collection were: volume  
262  $3.6 \pm 0.2$  mL, concentration  $282 \pm 21 \times 10^6$  spermatozoa/mL, total motility  $94.4 \pm 0.6\%$ , movement  
263  $5.0 \pm 0.0$ , and morphologically normal spermatozoa greater than 90%.

264 A decrease of total motile sperm and sperm with intact membranes after sorting compared to  
265 control semen evaluated immediately after collection was observed ( $P < 0.05$ ). Freezing induced a  
266 reduction of total motile sperm, sperm with intact membranes and acrosome integrity in both  
267 unsorted and sorted sperm ( $P < 0.05$ ) (Tab. 1).

268

### 269 **4. Discussion**

270

271 This study demonstrates that the use of mTALP can be considered a suitable medium for the storage  
272 of dog semen at room temperature (20-25°C) before sorting. Even if all the parameters evaluated  
273 underwent a decrease after 24h storage, usually spermatozoa are sexed within the first hours after  
274 collection when sperm in mTALP maintained parameters similar to those of semen analysed  
275 immediately after collection. For this reason mTALP was used in the subsequent experiments for  
276 diluting dog semen pre-sorting.

277 Concerning the sex sorting procedure, even if it caused a decrease of sperm motility parameters, it  
278 did not seem to severely affect dog sperm membranes as viability and acrosome integrity were  
279 similar to that of control semen except for viability in experiment 4 that anyway resulted higher  
280 than 70%. Rodenas et al. [5] reported that, while the quality of spermatozoa immediately after  
281 sorting before centrifugation step, performed in order to concentrate the highly diluted sorted cells,

282 was similar to that of fresh semen samples, the high speed centrifugation of sexed sperm at 3000 x g  
283 for 4 min resulted in a significant reduction of total motility, viability and sperm with intact  
284 acrosome. In our study we performed a low speed centrifugation over a long period (800 x g for 20  
285 min), basing on the protocol routinely used in our lab for horse and boar semen [10,11]. We did not  
286 check viability and acrosome integrity before centrifugation step but the maintenance of a good  
287 viability and acrosome integrity suggests that the centrifugation protocol after sorting used in this  
288 study could be adequate for concentrating sexed spermatozoa in this species.

289 The bitch that has to be inseminated with sexed sperm can be far from the sorting facilities or  
290 require multiple inseminations; for this reason it is very important to define a proper method to store  
291 sexed sperm. As no data about storage of canine sexed sperm are available in literature, we decided  
292 to evaluate the possibility of either liquid store sexed sperm at room temperature (20-25°C) or  
293 freeze it.

294 In the experiment 2, among the four extender tested for liquid storage of dog sexed semen, mTALP  
295 resulted in the maintenance of better sperm quality; this positive effect was particularly evident for  
296 total motility. Anyway the low quality of sorted semen after 24h suggests the necessity of  
297 optimizing an alternative storage method for long periods.

298 In experiment 3 the ability of sperm from 8 purebred dogs to withstand sexing procedure followed  
299 by freezing was tested and freezing caused a significant reduction of all the parameters evaluated in  
300 both unsorted and sorted sperm. SORT-FR semen was characterized by lower motility  
301 characteristics but similar viability and acrosome integrity compared to CTR-FR. These results  
302 suggest that the semen used in this experiment was mainly injured by freezing even if semen  
303 susceptibility to freezing seems to be, in a certain manner increased, by sorting.

304 To exclude the hypothesis that the unsatisfactory quality of both unsorted and sorted frozen sperm  
305 could be due to the fact that dogs of a single breed were enrolled in the experiment, the same  
306 protocol was used for ejaculates from mixed breed dogs (experiment 4). The results evidenced that  
307 sorting and subsequent freezing caused severe damage to spermatozoa. However the freezing

308 injuries to sperm observed could be due, at least in part, to the cooling and the second dilution with  
309 EYTGC2 made at 4°C performed in small volumes (500µl) that can increase the sperm  
310 susceptibility to cooling/freezing stress as performing the same procedure with higher volumes of  
311 unsorted semen (3 mL) significantly improved sperm quality (data not shown). This information  
312 must be considered in the future in order to optimize the procedure of freezing dog sorted semen;  
313 sex-sorted sperm in fact require modification of the basic protocol used for cryopreserving unsorted  
314 sperm due to the small volume of sex-sorted sperm at a low concentration compared to unsorted  
315 sperm [12].

316

#### 317 ***4.1 Conclusions***

318

319 In conclusion mTALP is an appropriate medium for canine sperm storage pre- and soon after  
320 sorting (hours), but a long storage period at room temperature is not tolerated by sex-sorted  
321 spermatozoa. Cryopreservation resulted in a great loss of sexed spermatozoa and further studies are  
322 needed to optimize the freezing protocol.

323

#### 324 **Acknowledgement**

325

326 The authors wish to thank Società Italiana Produzione Sementi and Fondazione Sfameni for  
327 financially supporting Elisa Giaretta. Work funded by a Bologna University grant (RFO, Ricerca  
328 Fondamentale Orientata).

#### 329 **References**

330 [1] Garner DL. Flow cytometric sexing of mammalian sperm. *Theriogenology* 2006;65:943-57.

331 [2] Garner D., Seidel GE Jr. Past, present and future perspectives on sexing sperm. *Can J Anim Sci*  
332 2003;83:375–84.

- 333 [3] Meyers MA, Burns G, Arn D, Schenk JL. Birth of canine offspring following insemination of a  
334 bitch with flow-sorted spermatozoa. *Rep Fertil Dev* 2008;20: 213.
- 335 [4] Oi M, Yamada K, Hayakawa H, Suzuki H. Sexing of dog sperm by fluorescence in situ  
336 hybridization. *J Reprod Dev* 2013;59:92–6.
- 337 [5] Rodenas C, Lucas X, Tarantini T, Del Olmo D, Roca J, Vazquez JM, Martinez EA, Parrilla I.  
338 The effects of hoechst 33342 staining and the male sample donor on the sorting efficiency of canine  
339 spermatozoa. *Reprod Domest Anim* 2014;49:115-21.
- 340 [6] Johnston SD, Ward D, Lemon J, Gunn I, MacCallum CA, Keeley T, Blyde D. Studies of male  
341 reproduction in captive African wild dogs (*Lycaon pictus*). *Anim Reprod Sci* 2007;100:338–55.
- 342 [7] Payan-Carreira R, Miranda S, Nizański W. Artificial Insemination in dogs. In: Manafi M editor.  
343 Artificial insemination in farm animals, InTech; 2011, p. 51-78.
- 344 [8] Howard JG, Bush M, Wildt DE. Semen collection, analysis and cryopreservation in  
345 nondomestic mammals. In: Morrow DA editor. *Current therapy in theriogenology* 2<sup>nd</sup> edn.,  
346 Philadelphia: Saunders Company, 1986; p. 1047-53.
- 347 [9] Johnson LA. Sex preselection in swine: altered sex ratios in offspring following surgical  
348 insemination of flow sorted X- and Y-bearing sperm. *Reprod Domest Anim* 1991;26:309–314.
- 349 [10] Spinaci M, Chlapanidas T, Bucci D, Vallorani C, Perteghella S, Lucconi G, Communod R,  
350 Vigo D, Galeati G, Faustini M, Torre ML. Encapsulation of sex sorted boar semen: sperm  
351 membrane status and oocyte penetration parameters. *Theriogenology* 2013;79:575-81.
- 352 [11] Balao da Silva CM, Spinaci M, Bucci D, Giaretta E, Peña FJ, Mari G, Galeati G. Effect of sex  
353 sorting on stallion spermatozoa: heterologous oocyte binding, tyrosine phosphorylation and  
354 acrosome reaction assay. *Anim Reprod Sci* 2013;141:68-74.
- 355 [12] Bathgate R. Functional integrity of sex-sorted, frozen-thawed boar sperm and its potential for  
356 artificial insemination. *Theriogenology* 2008;70:1234–41.

357

358

359 **Table 1.** Seminal quality evaluations of sperm of purebred (n=8) and mixed breed dogs (n=8) after  
 360 collection (CTR), after flow cytometric sex-sorting (SORT), and after thawing of non-sorted and  
 361 sorted cryopreserved samples (CTR-FR, SORT-FR).

	<b>Total motility (%)</b>	<b>Movement (0-5)</b>	<b>Viability (%)</b>	<b>Acrosome intact sperm (%)</b>
<b>Purebred</b>				
<b>CTR</b>	92.5 ± 0.9 <sup>a</sup>	5.0 ± 0.0 <sup>a</sup>	83.9 ± 1.8 <sup>a</sup>	94.5 ± 1.2 <sup>a</sup>
<b>SORT</b>	52.9 ± 6.0 <sup>b</sup>	2.1 ± 0.6 <sup>b</sup>	85.2 ± 6.8 <sup>a</sup>	90.6 ± 1.3 <sup>a</sup>
<b>CTR-FR</b>	25.7 ± 4.1 <sup>c</sup>	3.1 ± 0.4 <sup>b</sup>	31.3 ± 4.0 <sup>b</sup>	76.5 ± 2.8 <sup>b</sup>
<b>SORT-FR</b>	2.4 ± 1.2 <sup>d</sup>	0.6 ± 0.4 <sup>c</sup>	22.8 ± 4.5 <sup>b</sup>	78.0 ± 2.8 <sup>b</sup>
<b>Mixed breed</b>				
<b>CTR</b>	94.4 ± 0.6 <sup>a</sup>	5.0 ± 0.0 <sup>a</sup>	91.4 ± 1.1 <sup>a</sup>	97.9 ± 0.5 <sup>a</sup>
<b>SORT</b>	54.4 ± 9.1 <sup>b</sup>	4.6 ± 0.2 <sup>a</sup>	71.3 ± 5.2 <sup>b</sup>	89.9 ± 3.9 <sup>a</sup>
<b>CTR-FR</b>	37.5 ± 4.1 <sup>b</sup>	5.0 ± 0.0 <sup>a</sup>	33.2 ± 2.8 <sup>c</sup>	33.9 ± 4.0 <sup>b</sup>
<b>SORT-FR</b>	9.4 ± 4.6 <sup>c</sup>	2.7 ± 0.5 <sup>b</sup>	12.2 ± 2.5 <sup>d</sup>	43.7 ± 3.4 <sup>b</sup>

362 <sup>ab</sup> Values with different superscripts within a column differ significantly (P<0.05)

363



364 **Figure legends**

365

366 **Figure 1.** (A) Total sperm motility, (B) movement, (C) membrane integrity as an estimate of sperm  
367 viability, and (D) acrosome integrity evaluated in dog semen stored at room temperature (20-25°C)  
368 in different extenders. (n=4)

369 TFC - Tris-fructose-citrate; TGC - Tris-glucose-citrate; mTALP - modified Tyrode's solution, III

370 FRAC - third fraction of the ejaculate.

371 Different letters on the same type bars indicate a significant ( $P<0.05$ ) difference. Values are  
372 expressed as mean  $\pm$ SEM.

373

374 **Figure 2.** (A) total sperm motility, (B) movement, (C) membrane integrity as an estimate of sperm  
375 viability and (D) acrosome integrity evaluated in sorted dog sperm resuspended and stored with  
376 different extender at room temperature (20-25°C). (n=4)

377 TFC - Tris-fructose-citrate; TGC - Tris-glucose-citrate; mTALP - modified Tyrode's solution, III

378 FRAC - third fraction of the ejaculated.

379 Different letters on the same type bars indicate a significant ( $P<0.05$ ) difference. Values are  
380 expressed as mean  $\pm$ SEM.

381