

This is the final peer-reviewed accepted manuscript of:

**Mercury and protein thiols: Stimulation of mitochondrial F(1)F(O)-ATPase and inhibition of respiration.**

**Nesci S, Trombetti F, Pirini M, Ventrella V, Pagliarani A. Chem Biol Interact. 2016 Dec 25;260:42-49.**

The final published version is available online at:  
**<https://doi.org/10.1016/j.cbi.2016.10.018>**

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

*This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)*

***When citing, please refer to the published version.***

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

# Mercury and protein thiols: stimulation of mitochondrial $F_1F_0$ -ATPase and inhibition of respiration

*Salvatore Nesci, Fabiana Trombetti, Maurizio Pirini, Vittoria Ventrella, Alessandra Pagliarani\**

Department of Veterinary Medical Sciences, University of Bologna, Italy

\*Corresponding author: Alessandra Pagliarani [alessandra.pagliarani@unibo.it](mailto:alessandra.pagliarani@unibo.it) Tel +390512097017

Fax +390512097037, Department of Veterinary Medical Sciences (DIMEVET), University  
of Bologna, via Tolara di Sopra, 50, 40064, Ozzano Emilia, Bologna, Italy

## Abstract

1  
2  
3 In spite of the known widespread toxicity of mercury, its impact on mitochondrial bioenergetics is a  
4  
5 still poorly explored topic. Even if many studies have dealt with mercury poisoning of  
6  
7 mitochondrial respiration, as far as we are aware  $\text{Hg}^{2+}$  effects on individual complexes are not so  
8  
9 clear. In the present study changes in swine heart mitochondrial respiration and  $\text{F}_1\text{F}_0$ -ATPase (F-  
10  
11 ATPase) activity promoted by micromolar  $\text{Hg}^{2+}$  concentrations were investigated.  $\text{Hg}^{2+}$  was found  
12  
13 to inhibit the respiration of NADH-energized mitochondria, whereas it was ineffective when the  
14  
15 substrate was succinate. Interestingly, the same micromolar  $\text{Hg}^{2+}$  doses which inhibited the NADH-  
16  
17  $\text{O}_2$  activity stimulated the F-ATPase, most likely by interacting with adjacent thiol residues.  
18  
19 Accordingly,  $\text{Hg}^{2+}$  dose-dependently decreased protein thiols and all the elicited effects on  
20  
21 mitochondrial complexes were reversed by the thiol reducing agent DTE. These findings clearly  
22  
23 indicate that  $\text{Hg}^{2+}$  interacts with Cys residues of these complexes and differently modulate their  
24  
25 functionality by modifying the redox state of thiol groups. The results, which cast light on some  
26  
27 implications of metal-thiol interactions up to now not fully explored, may contribute to clarify the  
28  
29 molecular mechanisms of mercury toxicity to mitochondria.  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39

## Keywords

40  
41  
42  $\text{F}_1\text{F}_0$ -ATPase; Mitochondria; Oxidative phosphorylation; Mercury; Thiol groups.  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## Abbreviations

$\Delta\mu_{H^+}$ , Transmembrane electrochemical gradient of protons; OXPHOS, oxidative phosphorylation; DTE, 1,4-Dithioerythritol; DNP, 2,4-Dinitrophenol; PAO, Phenylarsine oxide; NEM, *N*-Ethylmaleimide; MBBr, Monobromobimane; DTNB, 5,5'-Dithiobis(2-nitrobenzoic acid); MTRs, monothiol reagents

**Funding:** This work was supported by the University of Bologna (RFO grant).

## 1. Introduction

Mercury is a toxic heavy metal and mercury contamination causes pathological symptoms primarily in brain but also in lungs, kidney and heart [1-4]. The different chemical forms of mercury have been related to their different toxicity. Inorganic mercury comprises the elemental forms metallic mercury and mercury vapor ( $\text{Hg}^0$ ), and the ionic forms mercurous ( $\text{Hg}^+$ ) and mercuric ( $\text{Hg}^{2+}$ ) ions. Organic mercury, in which the metal is covalently bound to an organic moiety, mainly stems from biological sources. The organic forms have long been considered highly dangerous, being more easily absorbed and life-long accumulated especially in animals at the top of the food chain. Organic mercury can be also converted into inorganic salts, and especially into mercuric chloride ( $\text{HgCl}_2$ ) [4]. This inorganic salt has raised concern as a significant source of acute and chronic poisoning by mercury [5]. From a molecular insight, the chemical interaction between  $\text{Hg}^{2+}$  and biomolecules, only partially known, may substantially contribute to the highly varied mercury toxicity [3].

Due to their biological features, mitochondria are extremely vulnerable to pollutants [6] including highly toxic heavy metals [7]. Exposure to mercury, mainly through contaminated seafood and dental amalgam [8], has been associated with mitochondrial dysfunctions in aging [6], cardiovascular [2,4,9] and neurodegenerative diseases [7,10] and even with the development of autoimmune diseases [11] and autism [12].

All mercury compounds have a great affinity for sulphhydrylic groups of proteins and other biomolecules [3,5]. The inorganic  $\text{Hg}^{2+}$  [8] enters mitochondria by exploiting the negative charge in the mitochondrial matrix built by the respiratory chain [6] and uncouples oxidative phosphorylation (OXPHOS) [7,13,14]. Mercury mainly localizes in the protein fraction of mitochondria [15].  $\text{Hg}^{2+}$  covalently binds to thiol ( $-\text{SH}$ ) groups of Cys residues of proteins and forms adducts which change the protein conformation [3,16,17]. Moreover, mercury may bind to intracellular reduced glutathione (GSH) and deplete GSH stores, thus contributing to an increase in reactive oxygen species (ROS) [18]. Furthermore  $\text{Hg}^{2+}$  formation from organomercury compounds in the

1 mitochondrial matrix is stimulated by superoxide production. Additionally, oxidative stress greatly  
2 amplifies mercury capability to bind Cys thiols [19]. As a response to cellular stress, mitochondria  
3  
4 can dictate the cell fate by increasing the mitochondrial membrane permeability, an event that  
5  
6 triggers apoptosis [20], another process in which mercury has been involved [4,7,11]. So, a tight  
7  
8 link exists between mercury, oxidative stress and cell death. Being among mercury targets and ROS  
9  
10 producers, the mitochondrion plays a key role in this interaction network.  
11  
12

13  
14 The primary role of mitochondria is ATP production through OXPHOS. Basically OXPHOS  
15  
16 involves four respiratory complexes in the inner mitochondrial membrane (IMM) which transfer the  
17  
18 electrons withdrawn from the reduced substrates  $\text{NADH} + \text{H}^+$  and  $\text{FADH}_2$ , in turn generated by  
19  
20 nutrient oxidation, to molecular oxygen [21]. The energy released by this electron cascade through  
21  
22 the so-called electron transport chain is converted into an electrochemical proton gradient ( $\Delta\mu_{\text{H}^+}$ )  
23  
24 across the IMM by three of these complexes (complex I, III and IV, respectively), which pump  
25  
26 protons. The ATP synthase or  $\text{F}_1\text{F}_0$ -ATPase is able to dissipate  $\Delta\mu_{\text{H}^+}$  to synthesize ATP; under  
27  
28 pathological conditions when IMM is depolarized, the same enzyme complex can work in reverse,  
29  
30 namely it hydrolyzes ATP. In this case the enzyme complex exploits ATP hydrolysis to pump  
31  
32 protons and re-energize the membrane [22]. The two interlocked domains of the  $\text{F}_1\text{F}_0$ -ATPase,  $\text{F}_1$   
33  
34 and  $\text{F}_0$ , work together matched by a torque generation mechanism [23]: the hydrophilic  $\text{F}_1$  builds  
35  
36 ATP, while the membrane-embedded domain  $\text{F}_0$  rotates as a  $\text{H}^+$  turbine. The rotation direction of  
37  
38 these engines and their catalytic task (synthesis/hydrolysis) depend on the thermodynamic balance  
39  
40 between  $\Delta\mu_{\text{H}^+}$  and  $\Delta G_p$  (the Gibbs free energy for ATP synthesis).  $\Delta\mu_{\text{H}^+}$  drives ATP synthesis  
41  
42 (counterclockwise rotation, viewed from the matrix side), and *vice versa* an overwhelming  $\Delta G_p$   
43  
44 leads to ATP hydrolysis (clockwise rotation) [24].  
45  
46  
47  
48  
49  
50  
51  
52

53 Post-translational modifications altering the redox state of thiols concur to physiologically rule the  
54  
55 mitochondrial functions associated with signal transduction pathways triggered by ROS production  
56  
57 [25]. Since antioxidant defenses, mainly GSH pool, the related regenerating systems and ROS  
58  
59 sources, are differently distributed, mitochondria contain different redox compartments.  
60  
61  
62  
63  
64  
65

1 Interestingly, OXPHOS complexes lie on the boundary line of two redox environments: the  
2 reducing matrix and the intermembrane space with cristae lumen which constitutes a relatively  
3 oxidizing environment [26]. Within this boundary line, protein Cys residues may act as a “sulfur  
4 switch” [25], thus behaving as chemical interface between the mitochondrial environment and the  
5 enzyme function. The protein thiol redox state, already involved in the conformational and/or  
6 activity changes of Complex I [27] and F<sub>1</sub>F<sub>0</sub>-ATPase [28] under pathological conditions, may also  
7 play a role in the mercury-driven mitochondrial dysfunction. The latter leads to a decrease in ATP  
8 production and to an increase in oxidative stress [4,7]. In spite of the wealth of studies on mercury  
9 effects on the respiratory chain, the modulation of the enzyme complexes, with the exclusion of any  
10 interference of membrane potential driven by mercury cations, has not been investigated yet. The  
11 molecular mechanisms of metal interaction with protein thiols, here investigated in uncoupled  
12 mitochondria, reveal that the same basic chemical mechanism differently modulates the F<sub>1</sub>F<sub>0</sub>-  
13 ATPase (henceforth defined as F-ATPase) and other OXPHOS complexes. The findings may  
14 contribute to understand the complex and varied pattern of mercury toxicity by casting light on  
15 some implications of metal-thiol interactions which up to now have not been fully explored.  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38

## 39 2. Materials and methods

### 40 2.1. Chemicals

41 Na<sub>2</sub>ATP, oligomycin mixture (A:B:C 64:15:17%), 1,4-Dithioerythritol (DTE), 2,4-Dinitrophenol  
42 (DNP), Phenylarsine oxide (PAO), N-Ethylmaleimide (NEM), NADH, sodium succinate, rotenone,  
43 mercury(II) chloride and 5,5'-Dithiobis(2-nitrobenzoic acid) (DTNB) were obtained from Sigma-  
44 Aldrich (Milan, Italy). Monobromobimane (MBr) was purchased by Vinci-Biochem (Vinci, Italy).  
45  
46 All other chemicals were reagent grade. Quartz double distilled water was used for all reagent  
47 solutions except when differently stated.  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

### 61 2.2. Preparation of the mitochondrial fraction

62  
63  
64  
65

1 Swine hearts (*Sus scrofa domesticus*) were collected at a local abattoir and transported to the lab  
2 within 2 h in ice buckets at 0-4°C. After removal of fat and blood clots as much as possible,  
3  
4 approximately 30-40 g of heart tissue from the left ventricle were rinsed in ice-cold washing Tris-  
5  
6 HCl buffer consisting of 0.25 M sucrose, 10 mM Tris(hydroxymethyl)-aminomethane (Tris), pH  
7  
8 7.4 and finely chopped into fine pieces with scissors. Each preparation was made from one heart.  
9  
10 Mitochondria were isolated by stepwise centrifugation as described previously [29]. The protein  
11  
12 concentration was determined according to the spectrophotometric method of Bradford [30] by Bio-  
13  
14 Rad Protein Assay kit II. The mitochondrial preparations were then stored in liquid nitrogen until  
15  
16 analyses.  
17  
18  
19  
20  
21  
22  
23

### 24 2.3. Assay of the mitochondrial F<sub>1</sub>F<sub>0</sub>-ATPase (F-ATPase) activity

25  
26 Immediately after thawing, the mitochondrial fractions were used for the F-ATPase activity assays.  
27  
28 The capability of ATP hydrolysis was assayed in a reaction medium (1 mL) containing 75 mM  
29  
30 ethanolamine-HCl buffer pH 9.3, 0.15 mg mitochondrial protein and 2.0 mM MgCl<sub>2</sub> for F<sub>1</sub>F<sub>0</sub>-  
31  
32 ATPase assay. After 5 min preincubation at 37 °C, the reaction, carried out at the same temperature,  
33  
34 was started by the addition of 6.0 mM Na<sub>2</sub>ATP as substrate and stopped after 5 min by the addition  
35  
36 of 1 mL of ice-cold 15% (w/w) aqueous solution trichloroacetic acid. Once the reaction was  
37  
38 stopped, vials were centrifuged for 15 min at 5000 rpm (ALC 4225 Centrifuge). In the supernatant,  
39  
40 the concentration of inorganic phosphate (P<sub>i</sub>) hydrolyzed by known amounts of mitochondrial  
41  
42 protein, which is an indirect measure of ATPase activity, was colorimetrically evaluated [31].  
43  
44  
45  
46  
47

48 The ATPase activity was routinely measured by subtracting, from the P<sub>i</sub> hydrolyzed by known  
49  
50 amounts of mitochondrial protein (which indirectly indicates the total ATPase activity); the P<sub>i</sub>  
51  
52 hydrolyzed in the presence of 3 µg/mL oligomycin. To this aim, in vials run in parallel, 1 µL from a  
53  
54 mother solution of 3 mg/mL oligomycin in dimethylsulfoxide (DMSO) was directly added to the  
55  
56 reaction mixture before starting the reaction. The employed dose of oligomycin, a specific inhibitor  
57  
58 of F-ATPases which selectively blocks the F<sub>0</sub> subunit [32-34], ensured maximal enzyme activity  
59  
60  
61  
62  
63  
64  
65



1 inhibition and was currently used in ATPase assays [29,32-34]. The total ATPase activity was  
2 calculated by detecting the  $P_i$  in control tubes run in parallel and containing 1  $\mu$ L DMSO per mL  
3 reaction system. In each experimental set, control tubes were alternated to the condition to be  
4 tested.. Small volumes of the effectors in aqueous solutions were directly added to the reaction  
5 system so as to obtain the required concentrations. In all experiments the enzyme activity was  
6 calculated as  $\mu$ mol  $P_i$ ·mg protein<sup>-1</sup>·min<sup>-1</sup>.  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16

#### 17 2.4. Preincubation and treatment of mitochondria

18 When combined effects of different effectors, *i.e.* HgCl<sub>2</sub> and mono- or di-thiol reagents were tested,  
19 mitochondrial suspensions were preincubated for 30 min with appropriate concentrations of these  
20 effectors in Eppendorf vials on ice. This procedure was designed to ensure the interaction of the  
21 effectors to be tested with the mitochondrial proteins [34]. After this preincubation time, the  
22 ATPase reaction was carried out as described in Section 2.3.  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33

#### 34 2.5. Assay of the mitochondrial respiration

35 Immediately after thawing, the mitochondrial fractions were used to evaluate the mitochondrial  
36 respiration. The oxygen consumption rates were polarographically evaluated by Clark-type  
37 electrode using a thermostated Oxytherm System (Hansatech Instruments) equipped with a 1 mL  
38 polarographic chamber. The reaction mixture (120 mM KCl, 10 mM Tris-HCl buffer pH 7.2),  
39 maintained under Peltier thermostatisation at 37 °C and continuous stirring, contained 0.25 mg  
40 mitochondrial protein.  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50

51 To evaluate the NADH-O<sub>2</sub> activity, the mitochondrial oxidation was run under saturating substrate  
52 conditions (75  $\mu$ M NADH) after 2 min of stabilization of oxygen signal. Preliminary tests assessed  
53 that under these conditions the O<sub>2</sub> consumption was suppressed by the presence of 2.5  $\mu$ M rotenone,  
54 known inhibitor of Complex I [21]. The succinate-O<sub>2</sub> activity by Complex II was determined by the  
55 oxidation of succinate in the presence of 2.5  $\mu$ M rotenone. The reaction was started by the addition  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 of 10 mM succinate after 2 min of stabilization of oxygen signal. Also in this case preliminary tests  
2 assessed that, under the conditions applied, succinate oxidation was suppressed by 1µg/mL of  
3 antimycin A, a selective inhibitor of complex III [21].  
4

5 To evaluate Hg<sup>2+</sup>-effects, HgCl<sub>2</sub> solutions of adequate concentrations, prepared immediately before  
6 the experiments by dissolving the contaminant as chloride salt in water, were added 5 min before to  
7 start the reaction at 37 °C at the same time of mitochondria to the polarographic chamber. The rate  
8 of mitochondrial respiratory activities were evaluated automatically by O<sub>2</sub>view software as nmoles  
9 O<sub>2</sub>·mg protein<sup>-1</sup>·min<sup>-1</sup>. In routine experiments, reagents were added by a syringe to the  
10 polarographic cell containing the reaction mixture and the mitochondrial protein suspension in the  
11 following order: HgCl<sub>2</sub> when required, rotenone for succinate-O<sub>2</sub> activity, substrate (NADH or  
12 succinate) and DTE. Polarographic assays were run at least in triplicate on mitochondrial fractions  
13 from distinct animal pools.  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30

## 31 2.6. Quantitative evaluation of free thiols

32 Free thiols in mitochondrial suspensions in the absence and in the presence of 1.0, 2.0, 3.0, 5.0, 10.0  
33 µM HgCl<sub>2</sub> were colorimetrically quantified by Ellman's reagent [35]. The selected HgCl<sub>2</sub>  
34 concentrations were added to the mitochondrial suspensions immediately before the colorimetric  
35 analysis. The widely used Ellman's method is based on the capability of 5,5'-Dithiobis(2-  
36 nitrobenzoic acid) (DTNB) to react with free thiol groups by disulphide bonds with thionitrobenzoic  
37 acid (TNB). As the ratio of protein thiols to TNB formed is 1:1, TNB formation is currently taken  
38 as a measure of the number of free thiol groups in mitochondrial suspensions. After the addition of  
39 15% w/v trichloroacetic acid solution (250 µL/0.15 mg protein) to precipitate proteins, the  
40 mitochondrial suspensions were centrifuged at 12,000 g for 5 min at 4 °C. After removal of the  
41 supernatant, the mitochondrial pellet was carefully resuspended with potter Eppendorf pestle. Then,  
42 400 µL of reagent solution containing 0.5 M phosphate buffer (KH<sub>2</sub>PO<sub>4</sub>/K<sub>2</sub>HPO<sub>4</sub>, pH 7.4), 0.2 mM  
43 DTNB, were added to the suspensions and incubated for 20 min at 4 °C. The absorbance at 412 nm  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 (maximum TNB absorption wavelength) of the supernatant from a second centrifugation at 12,000  
2 g for 5 min at 4 °C was read on a Perkin–Elmer lambda 45 spectrophotometer. Mitochondrial thiol  
3  
4 groups were quantitatively evaluated by interpolating the absorbance values in a calibration curve  
5  
6 built by employing known cysteine concentrations as -SH standard. In each experiment set, data  
7  
8 were expressed as % free thiol/mg protein in mitochondria detected in HgCl<sub>2</sub>-free medium taken as  
9  
10 control.  
11  
12

### 13 2.7. Kinetic analyses

14  
15  
16 The mechanism of the enzyme activation by HgCl<sub>2</sub> on the F-ATPase was explored by the aid of the  
17  
18 graphical method of Lineweaver-Burk (double reciprocal plots). In all kinetic analyses the enzyme  
19  
20 specific activity was taken as the expression of the reaction rate ( $v$ ). Briefly, to build double  
21  
22 reciprocal plots the reciprocal of the reaction rate  $1/v$  ( $y$  axis) was plotted as a function of the  
23  
24 reciprocal of concentration of ATP substrate or Mg<sup>2+</sup> cofactor ( $x$  axis) [36]. By plotting the enzyme  
25  
26 activity data, distinct straight lines, each of which corresponded to a fixed concentration of HgCl<sub>2</sub>  
27  
28 were obtained by linear regression. The R<sup>2</sup> value was never lower than 0.98, thus confirming the  
29  
30 linearity of these plots.  
31  
32  
33

34 At least three independent experiments were carried out to build each plot.  
35  
36  
37  
38

### 39 2.8. Statistical Analysis

40  
41  
42 The data represent the mean  $\pm$  SD (shown as vertical bars in the figures) of the number of  
43  
44 experiments reported in the figure captions and table legends. In each experimental set, the analyses  
45  
46 were carried out on different mitochondrial preparations. The differences between data were  
47  
48 evaluated by one way ANOVA followed by Students–Newman–Keuls' test when F values indicated  
49  
50 significance ( $P \leq 0.05$ ).  
51  
52  
53  
54  
55  
56  
57  
58  
59

## 60 3. Results

61  
62  
63  
64  
65

### 3.1. Mercury effects on selected mitochondrial enzyme activities

The present work highlights a differential effect of HgCl<sub>2</sub> on some enzyme activities involved in OXPHOS. Accordingly, as shown in Figure 1, micromolar concentrations of HgCl<sub>2</sub> up to 10 μM, evaluated by detecting the oxygen consumption in the presence of two substrates which elicit the activity of Complex I (NADH) and Complex II (succinate), respectively, yielded substrate-dependent effects. The NADH-O<sub>2</sub> activity was progressively inhibited by increasing HgCl<sub>2</sub> concentrations up to attain 10% of the activity in a mercury-free medium at 10 μM HgCl<sub>2</sub>. The calculated IC<sub>50</sub> value was 1.7±0.5 μM. Conversely, the succinate-O<sub>2</sub> activity was unaffected by the HgCl<sub>2</sub> concentrations tested up to 10 μM HgCl<sub>2</sub>.

A biphasic effect was shown on the F-ATPase (Fig. 2). The enzyme activity was stimulated by low micromolar HgCl<sub>2</sub> concentrations: at 2 μM HgCl<sub>2</sub> the enzyme activity was 42% higher than the control (no HgCl<sub>2</sub>). Only at HgCl<sub>2</sub> concentrations ≥ 10 μM was the F-ATPase activity decreased: 50% enzyme inhibition occurred at 22.6±4.0 μM HgCl and 100% inhibition was detected in the presence of 100 μM HgCl<sub>2</sub> (Fig. S1).

### 3.2. Hg<sup>2+</sup> and thiols

On considering the ability of mercury to form complexes with Cys thiols, its putative binding to mitochondrial proteins was investigated by evaluating the thiol content in swine heart mitochondria incubated in the absence or in the presence of increasing concentrations of HgCl<sub>2</sub>. The amount of free thiol groups in mitochondria significantly decreased at concentrations of HgCl<sub>2</sub> ≥ 3 μM up to plateau to 20% in the range 5-10 μM HgCl<sub>2</sub>. Therefore, at least under the conditions applied, only 20% of the free mitochondrial thiols were targeted by mercury (Fig. 3).

In order to check whether HgCl<sub>2</sub> effects on mitochondrial enzyme activities could be reversed, the thiol reagent DTE, known to reduce thiols, was tested (Fig. 4). The NADH-O<sub>2</sub> activity *per se* was unaffected by 50 μM DTE. Conversely, when the enzyme activity was inhibited by 2 μM HgCl<sub>2</sub>,


1 DTE removed the enzyme inhibition and restored the mitochondrial respiration to control values  
2 (Fig. 4A). Likewise, the F-ATPase activity was *per se* unaffected by 50  $\mu$ M DTE, a dose which  
3  
4 suppressed the enzyme activation by 2  $\mu$ M HgCl<sub>2</sub> (Fig. 4B).  
5  
6

7 Interestingly, the mitochondrial F-ATPase was also activated by the presence of micromolar  
8  
9 concentrations of phenylarsine oxide (PAO), a vicinal dithiol-binding agent. Conversely binary  
10  
11 mixtures of increasing micromolar concentrations of PAO plus 50  $\mu$ M DTE were ineffective (Fig.  
12  
13 S2). To check if the F<sub>1</sub>F<sub>0</sub>-ATPase activation was due to an uncoupler-like effect of mercury, the  
14  
15 known uncoupler DNP was tested. However, no significant change in the enzyme activity was  
16  
17 detected (Fig. S3).  
18  
19  
20

21 The mechanisms involved in the F-ATPase activation by HgCl<sub>2</sub> were deepened by assaying the  
22  
23 effects of PAO and of the thiol reagents (MTRs) *N*-Ethylmaleimide (NEM) and monobromobimane  
24  
25 (MBBr), already used in a previous work [34] (Table 1). In these sets of experiments two different  
26  
27 protocols were applied, namely mitochondria were either preincubated with HgCl<sub>2</sub> or with  
28  
29 individual MTRs in the absence of other effectors. When mitochondria were preincubated with  
30  
31 HgCl<sub>2</sub>, individual MTRs were added to the reaction system immediately after the addition of  
32  
33 HgCl<sub>2</sub>-preincubated mitochondrial suspensions. In the alternative mode, HgCl<sub>2</sub> was added to the  
34  
35 reaction system immediately after the addition of MTR-preincubated mitochondrial suspensions.  
36  
37 The two procedures resulted in the same F-ATPase activation by HgCl<sub>2</sub> which was suppressed by  
38  
39 DTE addition. In the absence of mercury, all the MTRs, both in the presence and in the absence of  
40  
41 DTE, did not affect the F-ATPase activity. Conversely, in HgCl<sub>2</sub>-preincubated mitochondria the  
42  
43 subsequent addition of PAO to the reaction system did not result in any enzyme activity  
44  
45 stimulation. In PAO-preincubated mitochondria, HgCl<sub>2</sub> added to the reaction system did not  
46  
47 stimulate the F-ATPase activity. In all cases, DTE restored the enzyme activity detected in the  
48  
49 control.  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

### 61 3.3. Kinetics 62 63 64 65

Finally, the F-ATPase activation kinetics by  $\text{Hg}^{2+}$  was studied by evaluating the F-ATPase activities raised by different concentrations of ATP substrate at fixed cofactor  $\text{Mg}^{2+}$  concentration, or by different concentrations of  $\text{Mg}^{2+}$  at fixed ATP concentration, in the presence and in the absence of  $2 \mu\text{M HgCl}_2$ . This approach, adequately adapted from kinetic analysis methods [36], aimed at evaluating the kinetic activation by  $\text{HgCl}_2$  of the F-ATPase with respect to ATP substrate and  $\text{Mg}^{2+}$  cofactor, respectively. The unparallel lines of the obtained double reciprocal plots (Figure 5), showed that in the presence of  $\text{Hg}^{2+}$  the line was below the control line in both cases. Therefore  $\text{Hg}^{2+}$  increases both the  $V_{\max}$  and  $K_m$  values for the ATP substrate and the  $\text{Mg}^{2+}$  cofactor (Table 2). On these bases mixed-type or non-competitive activation are suggested. Mixed-type activation indicates that the activator ( $A$ ) binds both to the free enzyme, yielding the binary complex  $EA$  and to  $ES$ , forming the ternary complex  $ESA$ . Non-competitive activation means that these two possibilities are equal. The whole pattern of mixed-type enzyme activation by the modulator  $A$  is illustrated below:



To explore this point, once obtained the modified  $K_m$  and  $V_{\max}$  values from the double reciprocal plot in the presence of  $\text{Hg}^{2+}$ , the factor  $\alpha$ , which represents the proportionality constant which modifies the slope in the presence of a modulator, was calculated from the relationship  $\alpha = \text{slope} \cdot V_{\max} / K_m$ . Since in mixed-type activation the intercept on  $y$  axis is modified by the constant  $\alpha'$ , namely  $y \text{ intercept} = \alpha' / V_{\max}$ ,  $\alpha'$  was obtained by multiplying the  $y$  intercept per  $V_{\max}$ . In other words,  $\alpha$  refers to the formation of  $EA$  complex, while  $\alpha'$  refers to the formation of  $ESA$  complex.

1  
2 Therefore, the activation constants  $K_a$  and  $K'_a$ , which represent the equilibrium constants of the  
3 dissociation of the *EA* and *ESA* complexes respectively, were obtained from the relationships:  
4  $\alpha=1+[A]/K_a$  and  $\alpha'=1+[A]/K'_a$ . Since  $K_a$  is higher than  $K'_a$  (Table 3), *ESA* is more stable than *EA*.  
5  
6 This means that  $Hg^{2+}$  mainly forms the ternary enzyme-substrate-activator complex.  
7  
8  
9

#### 10 11 4. Discussion

12  
13 Mercury toxicity renewed concern, especially for the cardiovascular consequences of the chronic  
14 human exposure [2-4]. The cationic form  $Hg^{2+}$  which alters the mitochondrial integrity and  
15 metabolism [4,37,39] is long known to perturb OXPHOS in mammalian [4,7,15] and non  
16 mammalian [39] species. Given the documented wide range of action of the toxicant [3,5] and the  
17 interconnection between membrane and enzyme perturbations, it is quite difficult to elucidate the  
18 deleterious mechanisms triggered by the toxicant. As far as we are aware, up to now no study has  
19 considered mercury effects on the mitochondrial respiratory complexes and  $F_1F_0$ -ATPase activity  
20 independently of IMM potential and structural perturbations. So, the present work on freeze-thawed  
21 mitochondria had the main goal of clarifying mercury effects on selected enzyme activities of the  
22 IMM, under conditions which rule out any possible concomitant effect of  $\Delta\mu_{H^+}$  changes. With the  
23 aim of casting light on the molecular mechanisms involved, we primarily focused on the metal  
24 interaction with protein thiols. Recent advances hint that this is probably the main molecular  
25 mechanism of mercury toxicity [3,4,17], now recognized as connected with oxidative damage [7,  
26 9]. Accordingly, thiol-based redox changes in OXPHOS complexes are accompanied by ROS  
27 production [7] and/or come from a reduction potential of a critical redox couple [40]. Moreover,  
28 independently of major redox changes within the membrane, under appropriate conditions, post-  
29 translational changes lead to a variety of thiol modifications such as sulfenic acid formation,  
30 glutathionylation, sulfenyl amides, *S*-nitrosothiols and to the onset of intra- and inter-protein  
31 disulfide bridges [41]. Thiol modifications rule Complex I functions in heart failure: inactive  
32 Complex I is likely to be *S*-nitrosylated, while the active Complex I is unaffected. This reversible  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 manipulation would inhibit superoxide production by reverse electron transfer [42]. Redox changes  
2 in some thiol groups in the F<sub>O</sub> portion of the mitochondrial F-ATPase result in uncoupling [43], loss  
3 of sensitivity to oligomycin and other macrolide antibiotics [31]. The reversible thiol modification  
4 by mercurials prevents the F-ATPase from being irreversibly blocked by DCCD [44], thus hinting  
5 that some susceptible thiols occur in critical proton binding points. Past *in vitro* research pointed out  
6 a widespread HgCl<sub>2</sub> toxicity on OXPHOS. Various effects were recorded: Hg<sup>2+</sup> mainly acted as  
7 uncoupler by dissipating  $\Delta\mu_{H^+}$ , enhanced basal respiration [7,45], while state 3 respiration was  
8 unaffected or depressed [7], inhibited the succinate dehydrogenase and ATP synthase activities  
9 [39], suppressed the membrane cation selectivity [46] and prevented the membrane re-energization  
10 [14]. Concomitantly, Hg<sup>2+</sup> promoted oxidative stress [7], leading to lipid peroxidation [47], in turn  
11 compromising the IMM structural integrity [48]. In heart mitochondria the ATP synthase reversible  
12 inhibition by nanomolar Hg<sup>2+</sup> concentrations was already reported, without elucidating the  
13 mechanism(s) involved [49]. Conversely, methylmercury stimulated the F-ATPase activity in the  
14 presence of K<sup>+</sup>, which by entering mitochondria would collapse the membrane potential [15].

15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34 In the present study the NADH-O<sub>2</sub> inhibition by micromolar HgCl<sub>2</sub> concentrations, mirrors the  
35 activities of three respiratory complexes (I, III and IV) which act as proton pumps [21]. Since the  
36 mercury-driven inhibition is confined to the NADH-O<sub>2</sub> activity, being the succinate-O<sub>2</sub> activity  
37 unaffected (Fig. 1), Complex I is confirmed as main target of mercury among OXPHOS complexes  
38 and its inhibition may *per se* lead to mitochondrial dysfunction [4]. In contrast, the Complex II  
39 features may explain the succinate oxidase activity refractoriness to the tested Hg<sup>2+</sup> concentrations.  
40 Accordingly, in spite of its content of thiols and dithiols [50], Complex II contains less reactive or  
41 less accessible thiol groups than Complex I [51].

42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000



1 presence of  $\mu\text{molar HgCl}_2$ , we can infer that only the succinate- $\text{O}_2$  activity contributes to  $\Delta\mu_{\text{H}^+}$   
2 generation since the electron transfer from Complex I to Complex 3 is prevented by mercury, which  
3 blocks Complex I by thiol complexation. By this chemical mechanism, mercury would poison the  
4 respiratory chain in a different way from other mechanisms already described in mitochondria such  
5 as the displacement of  $\text{Fe}^{2+}$  and  $\text{Cu}^+$  ions from redox centers and membrane lipid peroxidation [2, 4,  
6 7]. To sum up, it seems likely that the direct binding of  $\text{Hg}^{2+}$  to critical thiols of Complex I may  
7 lessen Complex I functionality and significantly concur to the mercury-driven mitochondrial  
8 dysfunction.

9 An opposite effect, namely the enzyme activity stimulation, was shown on the mitochondrial F-  
10 ATPase. As bifunctional enzyme [53,54], the F-ATPase not only synthesizes ATP but also can  
11 exploit ATP hydrolysis to pump protons when the IMM is insufficiently polarized, to overcome the  
12 reversal potential threshold [54] and prevent the depolarization of mitochondria. In this context  
13  $\text{HgCl}_2$ , by stimulating the ATPase activity, may improve the coupling between glycolytic and  $\Delta\mu_{\text{H}^+}$   
14 oscillations in non-respiring mitochondria [55]. Interestingly, mercury complexation to thiols can  
15 produce either inhibition ( $\text{NADH-O}_2$ ) or activation (F-ATPase), depending on the localization and  
16 the role of the susceptible thiols in the mitochondrial complexes. Consistently, even if mercury  
17 binding to enzyme thiols is believed a common motif for enzyme inactivation [3], it activates some  
18 phospholipases [9,56].

19 Finally, the failed effect of the protonophore DNP on the F-ATPase activation rules out any  
20 uncoupling effect of  $\text{Hg}^{2+}$  under our experimental conditions, even if  $\text{Hg}^{2+}$  is a recognized ionophore  
21 [46].

22 The reversibility of these modifications of thiols shoulders the thiol role as sulfur switch to  
23 modulate enzyme function [3]. Since the F-ATPase activation by mercury is not sensitive to MTRs  
24 but is prevented by PAO, which specifically binds to dithiols, two adjacent thiol groups, known to  
25 play a significant role in mitochondrial proteins [57] are required for mercury binding to the F-  
26 ATPase. Mercury complexation to the F-ATPase thiols is facilitated when ATP and  $\text{Mg}^{2+}$  are bound  
27

1 to the enzyme, since mercury preferentially binds to the enzyme-substrate complex, namely it  
2 shows mixed-type activation kinetics (Figure 5). It seems likely that the MgATP-complex  
3 formation may induce conformational changes which somehow make dithiols more accessible to  
4 mercury. Consistently, with respect to controls, the reaction rate increases in spite of a decreased  
5 affinity for ATP and Mg<sup>2+</sup> (Table 2).  
6  
7  
8  
9

10 This work shoulders the concept that enzyme alterations play a great role in mercury toxicity [3].  
11 From a literature overview, this is probably the first report which shows that Hg<sup>2+</sup>, by targeting  
12 crucial thiols, acts differently on the mitochondrial complexes, producing either inhibition or  
13 stimulation of IMM enzyme activities or even no effect, as on Complex II. Even if we must be  
14 cautious to transpose *in vitro* to *in vivo* responses, it is intriguing to speculate that the shown duality  
15 of mercury yielding Complex I inhibition and F-ATPase stimulation could make mitochondria  
16 resilient to mercury contamination. Accordingly, the stimulation of the reverse functioning mode of  
17 the ATP synthase (F-ATPase) would preserve, at least to a certain extent, the IMM polarization and  
18 ionic homeostasis under conditions of respiratory chain inhibition. If this mechanism also works *in*  
19 *vivo*, mercury-contaminated mitochondria would maintain at least partially their functionality.  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35

36 Even if vulnerable to pollutants, mitochondria are at the same time robust [6], being endowed with  
37 multiple mechanisms to ensure their function. Consistently, the cysteine proportion in proteins  
38 increases with organism complexity [58] and this thiol abundance on the protein surface may be  
39 there for a reason [59]. So, among the multiple targets of mercury in mitochondria, the most  
40 reactive thiols of Complex I and F-ATPase could work together to minimize the mitochondrial  
41 damage by mercury.  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53

## 54 5. Acknowledgment

55 Danilo Matteuzzi (Department of Veterinary Medical Sciences, University of Bologna) is gratefully  
56 acknowledged for kindly conferring pig hearts from a local abattoir to our lab.  
57  
58  
59  
60  
61  
62  
63  
64  
65

## 6. References

- 1  
2  
3 [1]World Health Organisation Geneva, Inorganic mercury. Sources of human and  
4 environmental exposure, Environmental Health Criteria 118 (1991) 29-33.  
5  
6  
7 [2]M.C. Houston , Role of mercury toxicity in hypertension, cardiovascular disease and stroke.  
8  
9 J. Clin. Hyper. 8 (2011) 621-627.  
10  
11  
12 [3]R. Ynalvez, J. Gutierrez, H. Gonzalez-Cantu, Minireview: toxicity of mercury as a  
13 consequence of enzyme alteration, Biometals (2016), in press doi; 10.1007/s10534-015-  
14 9967-8.  
15  
16  
17  
18  
19 [4] M.C. Houston, The role of mercury in cardiovascular disease, J. Cardiovasc. Dis. Diagn. 2  
20 (2014) 170.  
21  
22  
23  
24 [5]R.A. Bernhoft, Mercury toxicity and treatment: a review of the literature. J. Environ. Public  
25 Health (2012) 2012, ID 460508 <http://dx.doi.org/10.1155/2012/460508>.  
26  
27  
28  
29 [6]J.N. Meyer, M.C.K. Leung, J.P. Rooney, A. Sandoel, M.O. Hengartner, G.E. Kisby, A.S.  
30 Bess, Mitochondria as a target of environmental toxicants, Toxicol. Sci. 134 (2013) 1-17.  
31  
32  
33  
34 [7]E.A. Belyaeva, T.V. Sokolova, L.V. Emelyanova. I.O. Zakharova, Mitochondrial electron  
35 transport chain in Heavy metal-induced neurotoxicity: effects of cadmium, mercury and  
36 copper, The Scientific World J. 2012 (2012) Article ID 136063 doi: 10.1100/2012/136063.  
37  
38  
39  
40 [8]T. Syversen, P. Kaur, The toxicology of mercury and its compounds, J. Trace El. Med. Biol.  
41 26 (2012) 215-26.  
42  
43  
44  
45  
46 [9]T.J. Hagele, J.N. Mazerik, A. Gregory, B. Kaufman, U. Magalang, M.L. Kuppusamy, C.B.  
47 Marsch, P. Kuppusamy, N.L. Parinandi, Mercury activates vascular endothelial cell  
48 phospholipase D through thiols and oxidative stress, Int. J. Toxicol. 26 (2007) 57-69.  
49  
50  
51  
52  
53 [10]A. Carocci, N., Rovito, M.S. Sinicropi, G., Genchi, Mercury toxicity and neurodegenerative  
54 effects, Rev. Environ.Contam. Toxicol. 229 (2014) 1-18.  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- [11]M.P. Waalkes, D.A. Fox, J.C. States, S.R. Patierno, M.J. McCabe Jr., Metals and disorders of cell accumulation: modulation of apoptosis and cell proliferation, *Toxicol. Sci.* 56 (2000) 255-61.
- [12]S. Rose, R. Wynne, R.E. Frye, S. Melnyk, S.J. James, Increased susceptibility to ethylmercury-induced mitochondrial dysfunction in a subset of autism lymphoblastoid cell lines, *J. Toxicol.* 2015 (2015) 573701
- [13]A.C. Santos, S.A. Uyemura, N.A. Santos, F.E. Mingatto, C. Curti, Hg(II)-induced renal cytotoxicity: *in vitro* and *in vivo* implications for the bioenergetic and oxidative status of mitochondria, *Mol. Cell. Biochem.* 177 (1997) 53-9.
- [14]H. Reyes-Vivas, M. Lopez-Moreno, E. Chavez, Protective effect of dithiocarbamate on mercury-induced toxicity in kidney mitochondria, *Comp. Biochem. Physiol.* 113C (1996) 349-352.
- [15]N. Sone, M.K. Larstuvold, Y. Kagawa, Effect of methyl mercury on phosphorylation, transport and oxidation in mammalian mitochondria, *J. Biochem.* 82 (1977) 859-68.
- [16]P.R. van Iwaarden, A.J. Driessen, W.N. Konings, What we can learn from the effects of thiol reagents on transport proteins, *Biochim. Biophys. Acta* 1113 (1992) 161-70.
- [17]M. Farina, D.S. Avila, J.B. Teixeira da Rocha, M. Aschner, Metals, oxidative stress and neurodegeneration: a focus on iron, manganese and mercury. *Neurochem. Int.* 62 (2013) 575-591.
- [18]J.M. Hansen, H. Zhang, D.P. Jones, Differential oxidation of thioredoxin-1, thioredoxin-2, and glutathione by metal ions, *Free Radic. Biol. Med.* 40 (2006) 138-45.
- [19]R.J. Mailloux, E. Yumvihoze, H.M. Chan, Superoxide produced in the matrix of mitochondria enhances methylmercury toxicity in human neuroblastoma cells, *Toxicol. Appl. Pharmacol.* 289 (2015) 371-80.
- [20]D.R. Green, L. Galluzzi, G. Kroemer, Cell biology. Metabolic control of cell death, *Science* 345 (2014) 1250256.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- [21] D.G. Nicholls, S.J. Ferguson, *Bioenergetics* 4, fourth ed., Academic Press, Amsterdam, 2013.
- [22] P. Dimroth, C. von Ballmoos, T. Meier, Catalytic and mechanical cycles in F-ATP synthases. Fourth in the Cycles Review Series, *EMBO Rep.* 7 (2006) 276-82.
- [23] W. Junge, H. Sielaff, S. Engelbrecht, Torque generation and elastic power transmission in the rotary F(O)F(1)-ATPase, *Nature* 459 (2009) 364-70.
- [24] S. Nesci, F. Trombetti, V. Ventrella, A. Pagliarani, Opposite rotation directions in the synthesis and hydrolysis of ATP by the ATP synthase: hints from a subunit asymmetry, *J. Membr. Biol.* 248 (2015) 163-9.
- [25] Y.M. Go, D.P. Jones, The redox proteome, *J. Biol. Chem.* 288 (2013) 26512-20.
- [26] S. Dröse, U. Brandt, I. Wittig, Mitochondrial respiratory chain complexes as sources and targets of thiol-based redox-regulation, *Biochim. Biophys. Acta.* 1844 (2014) 1344-54.
- [27] M. Babot, A. Birch, P. Labarbuta, A. Galkin, Characterisation of the active/de-active transition of mitochondrial complex I, *Biochim. Biophys. Acta.* 1837 (2014) 1083-92.
- [28] A. Pagliarani, S. Nesci, F. Trombetti, V. Ventrella, Thiol-related regulation of the mitochondrial F<sub>1</sub>F<sub>0</sub>-ATPase activity, in: S. Chakraborti, N.S. Dhalla, (Eds.), *Regulation of Ca<sup>2+</sup>-ATPases, V-ATPases and F-ATPases Series: Advances in Biochemistry in Health and Disease* Vol. 14, Springer International Publishing, Switzerland, (2016), pp 441-458.
- [29] S. Nesci, V. Ventrella, F. Trombetti, M. Pirini, A. Pagliarani, Preferential nitrite inhibition of the mitochondrial F<sub>1</sub>F<sub>0</sub>-ATPase activities when activated by Ca<sup>2+</sup> in replacement of the natural cofactor Mg<sup>2+</sup>, *Biochim. Biophys. Acta* 1860 (2016) 345-53.
- [30] M.M. Bradford, A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding, *Anal. Biochem.* 72 (1976) 248-54.
- [31] C.G. Fiske, Y. Subbarow, The colorimetric determination of phosphorus, *J. Biol. Chem.* 66 (1925) 375-400.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- [32] S. Nesci, V. Ventrella, F. Trombetti, M. Pirini, A. Pagliarani, Thiol oxidation is crucial in the desensitization of the mitochondrial  $F_1F_0$ -ATPase to oligomycin and other macrolide antibiotics, *Biochim. Biophys. Acta* 1840 (2014) 1882-91.
- [33] S. Nesci, V. Ventrella, F. Trombetti, M. Pirini, A. Pagliarani, The mitochondrial  $F_1F_0$ -ATPase desensitization to oligomycin by tributyltin is due to thiol oxidation, *Biochimie* 97 (2014) 128-37.
- [34] S. Nesci, V. Ventrella, F. Trombetti, M. Pirini, A. Pagliarani, Tri-*n*-butyltin binding to a low-affinity site decreases the  $F_1F_0$ -ATPase sensitivity to oligomycin in mussel mitochondria, *Appl. Organomet. Chem.* 26 (2012) 593–599.
- [35] G.L. Ellman, Tissue sulfhydryl groups, *Arch. Biochem. Biophys.* 82 (1959) 70-7.
- [36] I.H. Segel, *Enzyme kinetics*, John Wiley & Sons Inc, New York, 1975.
- [37] T. Toimela, H. Tähti, Mitochondrial viability and apoptosis induced by aluminum, mercuric mercury and methylmercury in cell lines of neural origin, *Arch. Toxicol.* 78 (2004) 565-74.
- [38] P.B. Pal, S. Pal, J. Das, P.C. Sil, Modulation of mercury-induced mitochondria-dependent apoptosis by glycine in hepatocytes, *Amino Acids* 42 (2012) 1669-83.
- [39] C.L. Mieiro, M. Pardal, A. Duarte, E. Pereira, C.M. Palmeira, Impairment of mitochondrial Energy metabolism of two marine fish by in vitro mercuric chloride exposure, *Mar. Poll. Bull.* 97 (2015) 488-93.
- [40] Y.M. Janssen-Heininger, B.T. Mossman, N.H. Heintz, H.J. Forman, B. Kalyanaraman, T. Finkel, J.S. Stamler, S.G. Rhee, A. van der Vliet, Redox-based regulation of signal transduction: principles, pitfalls, and promises, *Free Radic. Biol. Med.* 45 (2008) 1-17.
- [41] M.P. Murphy, Mitochondrial thiols in antioxidant protection and redox signaling: distinct roles for glutathionylation and other thiol modifications, *Antioxid. Redox. Signal.* 16 (2012) 476-95.
- [42] E.T. Chouchani, C. Methner, S.M. Nadtochiy, A. Logan, V.R. Pell, S. Ding, A.M. James, H.M. Cochemé, J. Reinhold, K.S. Lilley, L. Partridge, I.M. Fearnley, A.J. Robinson, R.C.

- 1 Hartley, R.A. Smith, T. Krieg, P.S. Brookes, M.P. Murphy, Cardioprotection by S-  
2 nitrosation of a cysteine switch on mitochondrial complex I, *Nat. Med.* 19 (2013) 753-9.  
3  
4  
5 [43] T. Yagi, Y. Hatefi, Thiols in oxidative phosphorylation: inhibition and energy-potentiated  
6 uncoupling by monothiol and dithiol modifiers, *Biochemistry* 23 (1984) 2449-55.  
7  
8  
9 [44] T. Yagi, Y. Hatefi, Thiols in oxidative phosphorylation: thiols in the F<sub>O</sub> of ATP synthase  
10 essential for ATPase activity, *Arch. Biochem. Biophys.* 254 (1987) 102-9.  
11  
12  
13 [45] C.M. Palmeira, V.M.C. Madeira, Mercuric chloride toxicity in rat liver mitochondria and  
14 isolated hepatocytes, *Toxicol. Appl. Pharmacol.* 143 (1997) 338-47.  
15  
16  
17 [46] L.P. Karniski, Hg<sup>2+</sup> and Cu<sup>+</sup> are ionophores, mediating Cl<sup>-</sup>/OH<sup>-</sup> exchange in liposomes and  
18 rabbit renal brush border membranes, *J. Biol. Chem.* 267 (1992) 19218-25.  
19  
20  
21  
22 [47] B.O. Lund, D.M. Miller, J.S. Woods, Studies on Hg(II)-induced H<sub>2</sub>O<sub>2</sub> formation and  
23 oxidative stress *in vivo* and *in vitro* in rat kidney mitochondria, *Biochem. Pharmacol.* 45  
24 (1993) 2017-24.  
25  
26  
27 [48] J.H. Southard, J.T. Penniston, D.E. Green, Induction of transmembrane proton transfer by  
28 mercurials in mitochondria. I. Ion movements accompanying transmembrane proton  
29 transfer, *J. Biol. Chem.* 248 (1973) 3546-50.  
30  
31  
32 [49] H. Shara, Z. Drahotka, Inhibition of mitochondrial ATPase by Hg<sup>++</sup> ions, *Physiol.*  
33 *Bohemoslov.* 27 (1978) 193-8.  
34  
35  
36 [50] S. Pagani, F. Bonomo, P. Cerletti, Sulfhydryl and disulfide content of succinate  
37 dehydrogenase, *FEBS Lett.* 39 (1974) 139-43.  
38  
39  
40 [51] T.K. Lin, G. Hughes, A. Muratovska, F.H. Blaikie, P.S. Brookes, D. Darley-Usmar, R.A.J.  
41 Smith, M.P. Murphy, Specific modification of mitochondrial protein thiols in response to  
42 oxidative stress-a proteomics approach, *J. Biol. Chem.* 277 (2002) 17048-56.  
43  
44  
45 [52] P.R. Rich, A. Maréchal, Electron transfer chains: Structures, mechanisms and energy  
46 coupling, *Comp. Biophys.* 8 (2012) 73-93.  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- [53] S. Nesci, F. Trombetti, V. Ventrella, A. Pagliarani, The *c*-Ring of the F<sub>1</sub>F<sub>0</sub>-ATP Synthase: Facts and Perspectives, *J. Membr. Biol.* 249 (2016) 11-21.
- [54] J.E. Walker, The ATP synthase: the understood, the uncertain and the unknown, *Biochem. Soc. Trans.* 41 (2013) 1-16.
- [55] L.F. Olsen, A.Z. Andersen, A. Lunding, J.C. Brasen, A.K. Poulsen, Regulation of glycolytic oscillations by mitochondrial and plasma membrane H<sup>+</sup>-ATPases, *Biophys. J.* 96 (2009) 3850-61.
- [56] J.N. Mazerik, H. Mikkilineni, V.A. Kuppusamy, E. Steinhour, A. Peltz, C.B. Marsch, P. Kuppusamy, N.L. Parinandi, Mercury activates phospholipase A<sub>2</sub> and induces formation of arachidonic acid metabolites in vascular endothelial cells, *Toxicol. Mechan. Methods* 17 (2007) 1-17.
- [57] R. Requejo, E.T. Chouchani, A.M. James, T.A. Prime, K.S. Lilley, J.M. Fearnley, M.P. Murphy, Quantification and identification of mitochondrial proteins containing vicinal dithiols, *Arch. Biochem. Biophys.* 504 (2010) 228-235.
- [58] A. Miseta, P. Csutora, Relationship between the occurrence of cysteine in proteins and the complexity of organisms, *Mol. Biol. Evol.* 17 (2000) 1232-9.
- [59] S.M. Marino, V.N. Gladyshev, Cysteine function governs its conservation and degeneration and restricts its utilization on protein surfaces, *J. Mol. Biol.* 404 (2010) 902-16.



**Table 1. Effects of 2.0  $\mu\text{M}$   $\text{HgCl}_2$ , thiol reagents (1.5  $\mu\text{M}$  NEM, 1.0  $\mu\text{M}$  MBBr and 50  $\mu\text{M}$  PAO) and of the reducing agent DTE on the mitochondrial F-ATPase activity.**

CONDITION	no DTE	DTE
Control	2.67 $\pm$ 0.17	2.69 $\pm$ 0.09
$\text{HgCl}_2$	3.34 $\pm$ 0.23*	2.73 $\pm$ 0.14
NEM	2.52 $\pm$ 0.03	2.67 $\pm$ 0.04
MBBr	2.68 $\pm$ 0.31	2.63 $\pm$ 0.27
PAO	3.21 $\pm$ 0.29*	2.55 $\pm$ 0.18
$\text{HgCl}_2$ + (NEM)	3.22 $\pm$ 0.16*	2.74 $\pm$ 0.03
NEM + ( $\text{HgCl}_2$ )	3.31 $\pm$ 0.04*	2.79 $\pm$ 0.05
$\text{HgCl}_2$ + (MBBr)	3.30 $\pm$ 0.17*	2.58 $\pm$ 0.19
MBBr + ( $\text{HgCl}_2$ )	3.22 $\pm$ 0.34*	2.77 $\pm$ 0.29
$\text{HgCl}_2$ + (PAO)	3.45 $\pm$ 0.29*	2.80 $\pm$ 0.46
PAO + ( $\text{HgCl}_2$ )	2.62 $\pm$ 0.18	2.37 $\pm$ 0.13

Thiol reagents were either incubated with mitochondria as detailed in Section 2.4 or, if in brackets, directly added to the reaction system; 50  $\mu\text{M}$  DTE was added to the reaction system as reported in Section 2.5. ATPase activity values ( $\mu\text{mol Pi} \cdot \text{mg protein}^{-1} \cdot \text{hr}^{-1}$ ) are the mean  $\pm$  SD of three different determinations on distinct mitochondrial preparations. The asterisk (\*) indicates significantly different values from the control ( $P \leq 0.05$ ).

**Table 2. Effect of HgCl<sub>2</sub> on kinetic parameters of the mitochondrial F-ATPase activity.**

HgCl <sub>2</sub> (μM)	0.0	2.0	0.0	2.0
	$V_{\max}$		$K_m$	
	(μmol P <sub>i</sub> ·mg protein <sup>-1</sup> ·min <sup>-1</sup> )		(μM)	
Mg <sup>2+</sup>	3.4±0.2a	5.0±0.1b	0.3±0.0a	0.4±0.1a
ATP	5.9±1.5a	8.2±1.1b	4.9±0.8a	6.0±0.6a

Data are the mean ± SD of n=3 (for Mg<sup>2+</sup> parameters) and n=4 (for ATP parameters) sets of experiments carried out on different mitochondrial preparations. In each row different letters indicate significantly different values ( $P \leq 0.05$ ).


**Table 3. Kinetic parameters of the F-ATPase activation by  $\text{Mg}^{2+}$  and ATP**

	$\alpha$	$\alpha'$	$K_a$ ( $\mu\text{M}$ )	$K'_a$ ( $\mu\text{M}$ )
$\text{Mg}^{2+}$	0.9	0.7	17.6	6.3
ATP	0.9	0.7	17.3	6.6

These parameters were calculated as detailed in the text.


Figure(s)


[Click here to download high resolution image](#)



Figure(s)



[Click here to download high resolution image](#)





Figure(s)

[Click here to download high resolution image](#)



Figure(s)

[Click here to download high resolution image](#)

