

# World Journal of *Gastroenterology*

*World J Gastroenterol* 2018 September 28; 24(36): 4093-4216





### EDITORIAL

- 4093 Interleukin 12/interleukin 23 pathway: Biological basis and therapeutic effect in patients with Crohn's disease  
*Aggeletopoulou I, Assimakopoulos SF, Konstantakis C, Triantos C*

### REVIEW

- 4104 Role of microRNAs in alcohol-induced liver disorders and non-alcoholic fatty liver disease  
*Torres JL, Novo-Veleiro I, Manzanedo L, Alvela-Suárez L, Macías R, Laso FJ, Marcos M*
- 4119 Calcium-sensing receptor in colorectal inflammation and cancer: Current insights and future perspectives  
*Iamartino L, Elajnaif T, Kallay E, Schepelmann M*
- 4132 Production of extracellular lysophosphatidic acid in the regulation of adipocyte functions and liver fibrosis  
*Yang F, Chen GX*

### MINIREVIEWS

- 4152 Ten years of sorafenib in hepatocellular carcinoma: Are there any predictive and/or prognostic markers?  
*Marisi G, Cucchetti A, Ulivi P, Canale M, Cabibbo G, Solaini L, Foschi FG, De Matteis S, Ercolani G, Valgiusti M, Frassinetti GL, Scartozzi M, Casadei Gardini A*

### ORIGINAL ARTICLE

#### Basic Study

- 4164 Differential expression of mucin 1 and mucin 2 in colorectal cancer  
*Kasprzak A, Siodla E, Andrzejewska M, Szymeja J, Seraszek-Jaros A, Cofta S, Szaflarski W*
- 4178 Mechanism of combined use of vitamin D and puerarin in anti-hepatic fibrosis by regulating the Wnt/ $\beta$ -catenin signalling pathway  
*Huang GR, Wei SJ, Huang YQ, Xing W, Wang LY, Liang LL*

#### Retrospective Study

- 4186 Frequency, types and treatment of anemia in Turkish patients with inflammatory bowel disease  
*Bengi G, Keyvan H, Durmaz SB, Akpınar H*

#### Prospective Study

- 4197 Low-dose spectral insufflation computed tomography protocol preoperatively optimized for T stage esophageal cancer - preliminary research experience  
*Zhou Y, Liu D, Hou P, Zha KJ, Wang F, Zhou K, He W, Gao JB*

### CASE REPORT

- 4208 Novel methionyl-tRNA synthetase gene variants/phenotypes in interstitial lung and liver disease: A case report and review of literature  
*Abuduxikuer K, Feng JY, Lu Y, Xie XB, Chen L, Wang JS*

**ABOUT COVER**

Editorial board member of *World Journal of Gastroenterology*, Matthias Ocker, MD, Professor, Department of Oncology Translational Medicine, Bayer AG, Berlin 13353, Germany

**AIMS AND SCOPE**

*World Journal of Gastroenterology* (*World J Gastroenterol*, *WJG*, print ISSN 1007-9327, online ISSN 2219-2840, DOI: 10.3748) is a peer-reviewed open access journal. *WJG* was established on October 1, 1995. It is published weekly on the 7<sup>th</sup>, 14<sup>th</sup>, 21<sup>st</sup>, and 28<sup>th</sup> each month. The *WJG* Editorial Board consists of 642 experts in gastroenterology and hepatology from 59 countries.

The primary task of *WJG* is to rapidly publish high-quality original articles, reviews, and commentaries in the fields of gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, hepatobiliary surgery, gastrointestinal oncology, gastrointestinal radiation oncology, gastrointestinal imaging, gastrointestinal interventional therapy, gastrointestinal infectious diseases, gastrointestinal pharmacology, gastrointestinal pathophysiology, gastrointestinal pathology, evidence-based medicine in gastroenterology, pancreatology, gastrointestinal laboratory medicine, gastrointestinal molecular biology, gastrointestinal immunology, gastrointestinal microbiology, gastrointestinal genetics, gastrointestinal translational medicine, gastrointestinal diagnostics, and gastrointestinal therapeutics. *WJG* is dedicated to become an influential and prestigious journal in gastroenterology and hepatology, to promote the development of above disciplines, and to improve the diagnostic and therapeutic skill and expertise of clinicians.

**INDEXING/ABSTRACTING**

*World Journal of Gastroenterology* (*WJG*) is now indexed in Current Contents<sup>®</sup>/Clinical Medicine, Science Citation Index Expanded (also known as SciSearch<sup>®</sup>), Journal Citation Reports<sup>®</sup>, Index Medicus, MEDLINE, PubMed, PubMed Central and Directory of Open Access Journals. The 2018 edition of Journal Citation Reports<sup>®</sup> cites the 2017 impact factor for *WJG* as 3.300 (5-year impact factor: 3.387), ranking *WJG* as 35<sup>th</sup> among 80 journals in gastroenterology and hepatology (quartile in category Q2).

**EDITORS FOR THIS ISSUE**

**Responsible Assistant Editor:** *Xiang Li*  
**Responsible Electronic Editor:** *Yan Huang*  
**Proofing Editor-in-Chief:** *Lian-Sheng Ma*

**Responsible Science Editor:** *Rao-Yu Ma*  
**Proofing Editorial Office Director:** *Ze-Mao Gong*

**NAME OF JOURNAL**

*World Journal of Gastroenterology*

**ISSN**

ISSN 1007-9327 (print)  
 ISSN 2219-2840 (online)

**LAUNCH DATE**

October 1, 1995

**FREQUENCY**

Weekly

**EDITORS-IN-CHIEF**

**Andrzej S Tarnawski, MD, PhD, DSc (Med), Professor of Medicine, Chief Gastroenterology, VA Long Beach Health Care System, University of California, Irvine, CA, 5901 E. Seventh Str., Long Beach, CA 90822, United States**

**EDITORIAL BOARD MEMBERS**

All editorial board members resources online at <http://www.wjgnet.com/1007-9327/editorialboard.htm>

**EDITORIAL OFFICE**

Ze-Mao Gong, Director  
*World Journal of Gastroenterology*  
 Baishideng Publishing Group Inc  
 7901 Stoneridge Drive, Suite 501,  
 Pleasanton, CA 94588, USA  
 Telephone: +1-925-2238242  
 Fax: +1-925-2238243  
 E-mail: [editorialoffice@wjgnet.com](mailto:editorialoffice@wjgnet.com)  
 Help Desk: <http://www.f6publishing.com/helpdesk>  
<http://www.wjgnet.com>

**PUBLISHER**

Baishideng Publishing Group Inc  
 7901 Stoneridge Drive, Suite 501,  
 Pleasanton, CA 94588, USA  
 Telephone: +1-925-2238242  
 Fax: +1-925-2238243  
 E-mail: [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)  
 Help Desk: <http://www.f6publishing.com/helpdesk>  
<http://www.wjgnet.com>

**PUBLICATION DATE**

September 28, 2018

**COPYRIGHT**

© 2018 Baishideng Publishing Group Inc. Articles published by this Open-Access journal are distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license.

**SPECIAL STATEMENT**

All articles published in journals owned by the Baishideng Publishing Group (BPG) represent the views and opinions of their authors, and not the views, opinions or policies of the BPG, except where otherwise explicitly indicated.

**INSTRUCTIONS TO AUTHORS**

Full instructions are available online at <http://www.wjgnet.com/bpg/gerinfo/204>

**ONLINE SUBMISSION**

<http://www.f6publishing.com>

## Ten years of sorafenib in hepatocellular carcinoma: Are there any predictive and/or prognostic markers?

Giorgia Marisi, Alessandro Cucchetti, Paola Ulivi, Matteo Canale, Giuseppe Cabibbo, Leonardo Solaini, Francesco G Foschi, Serena De Matteis, Giorgio Ercolani, Martina Valgiusti, Giovanni L Frassinetti, Mario Scartozzi, Andrea Casadei Gardini

Giorgia Marisi, Paola Ulivi, Matteo Canale, Serena De Matteis, Biosciences Laboratory, Istituto Scientifico Romagnolo per lo Studio e la Cura dei Tumori (IRST) IRCCS, Meldola 47014, Italy

Alessandro Cucchetti, Leonardo Solaini, Giorgio Ercolani, Department of Medical and Surgical Sciences, Alma Mater Studiorum, University of Bologna, Bologna 40126, Italy

Alessandro Cucchetti, Leonardo Solaini, Giorgio Ercolani, General and Oncologic Surgery, Morgagni-Pierantoni Hospital, Forlì 47121, Italy

Giuseppe Cabibbo, Section of Gastroenterology, D.I.B.I.M.I.S., University of Palermo, Palermo 35628, Italy

Francesco G Foschi, Department of Internal Medicine, Degli Infermi Hospital, Faenza 48018, Italy

Martina Valgiusti, Giovanni L Frassinetti, Andrea Casadei Gardini, Department of Medical Oncology, Istituto Scientifico Romagnolo per lo Studio e la Cura dei Tumori IRCCS, Meldola 47014, Italy

Mario Scartozzi, Department of Medical Oncology, University of Cagliari, Cagliari 45698, Italy

ORCID number: Giorgia Marisi (0000-0003-4012-0042); Alessandro Cucchetti (0000-0001-5269-1964); Paola Ulivi (0000-0002-4334-9662); Matteo Canale (0000-0002-9768-7149); Giuseppe Cabibbo (0000-0002-0946-3859); Leonardo Solaini (0000-0002-5031-9285); Francesco G Foschi (0000-0002-8216-0918); Serena De Matteis (0000-0001-9219-4081); Giorgio Ercolani (0000-0003-4334-5167); Martina Valgiusti (0000-0002-9843-3538); Giovanni L Frassinetti (0000-0001-9376-0057); Mario Scartozzi (0000-0001-5977-5546); Andrea Casadei Gardini (0000-0001-6289-7202).

**Author contributions:** All authors equally contributed to this paper with conception and design of the study, literature review and analysis, drafting and critical revision and editing, and final

approval of final version.

**Conflict-of-interest statement:** There is no conflict of interest associated with any of the senior author or other coauthors contributed their efforts in this manuscript.

**Open-Access:** This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

**Manuscript source:** Invited manuscript

**Correspondence to:** Alessandro Cucchetti, MD, PhD, Adjunct Professor, Surgeon, Department of Medical and Surgical Sciences, Alma Mater Studiorum, University of Bologna, Via Albertoni 15, Granarolo Dell'Emilia, Bologna 40126, Italy. [aleqko@libero.it](mailto:aleqko@libero.it)  
Telephone: +39-543-731111  
Fax: +39-543-739123

**Received:** July 3, 2018

**Peer-review started:** July 3, 2018

**First decision:** July 17, 2018

**Revised:** August 6, 2018

**Accepted:** August 24, 2018

**Article in press:** August 24, 2018

**Published online:** September 28, 2018

### Abstract

Sorafenib has been considered the standard of care for patients with advanced unresectable hepatocellular carcinoma (HCC) since 2007 and numerous studies



have investigated the role of markers involved in the angiogenesis process at both the expression and genetic level and clinical aspect. What results have ten years of research produced? Several clinical and biological markers are associated with prognosis. The most interesting clinical parameters are adverse events, Barcelona Clinic Liver Cancer stage, and macroscopic vascular invasion, while several single nucleotide polymorphisms and plasma angiopoietin-2 levels represent the most promising biological biomarkers. A recent pooled analysis of two phase III randomized trials showed that the neutrophil-to-lymphocyte ratio, etiology and extra-hepatic spread are predictive factors of response to sorafenib, but did not identify any predictive biological markers. After 10 years of research into sorafenib there are still no validated prognostic or predictive factors of response to the drug in HCC. The aim of the present review was to summarize 10 years of research into sorafenib, looking in particular at the potential of associated clinical and biological markers to predict its efficacy in patients with advanced HCC.

**Key words:** Biomarker; Angiopoietin; Neutrophil-to-lymphocyte ratio; Polymorphisms; Sorafenib; MicroRNA; Adverse events; Hepatocellular carcinoma; Vascular endothelial growth factor

© **The Author(s) 2018.** Published by Baishideng Publishing Group Inc. All rights reserved.

**Core tip:** Sorafenib has been considered the standard of care for patients with advanced unresectable hepatocellular carcinoma, but after 10 years of research into sorafenib response or resistance, there are still no validated prognostic or predictive factors of response.

Marisi G, Cucchetti A, Ulivi P, Canale M, Cabibbo G, Solaini L, Foschi FG, De Matteis S, Ercolani G, Valgiusti M, Frassinetti GL, Scartozzi M, Casadei Gardini A. Ten years of sorafenib in hepatocellular carcinoma: Are there any predictive and/or prognostic markers? *World J Gastroenterol* 2018; 24(36): 4152-4163 Available from: URL: <http://www.wjgnet.com/1007-9327/full/v24/i36/4152.htm> DOI: <http://dx.doi.org/10.3748/wjg.v24.i36.4152>

## INTRODUCTION

Sorafenib, an oral multikinase inhibitor, has been considered the standard of care for patients with advanced unresectable hepatocellular carcinoma (HCC) since 2007<sup>[1]</sup>. It works by inhibiting the activity of several tyrosine kinases involved in tumor angiogenesis and progression, including vascular endothelial growth factor receptor (VEGFR-2/3), platelet-derived growth factor receptor (PDGF-R), Flt3 and c-Kit, and also targets Raf kinases involved in the MAPK/ERK pathway<sup>[2]</sup> (Figure 1). The molecular mechanisms by

which sorafenib exerts its activity have still not been fully elucidated, and both Raf/MEK/ERK-dependent and -independent mechanisms have been observed<sup>[3]</sup>.

Sorafenib is expensive and associated with adverse events (AEs). Furthermore, a proportion of treated patients show no response to the drug. It would thus be useful to have predictive markers capable of identifying those who are more likely to benefit from therapy. The availability of more accurate predictive or prognostic factors would also help to spare potentially resistant patients from unnecessary toxicity.

Ten years have passed since sorafenib was first commercialized and about 2800 studies have been published on the kinase inhibitor. But how many associated prognostic and/or predictive markers have been identified? Numerous studies have focused on the role of markers involved in the angiogenesis process at both the expression and genetic levels. The largest biomarker study conducted to date is the SHARP trial<sup>[4]</sup>, which included an adequate number of participants and a placebo-controlled group. Smaller single-arm studies exploring predictive or prognostic markers for sorafenib have also been conducted, but the results of these have yet to be validated.

The aim of the present review was to summarize 10 years of research into sorafenib, looking in particular at the potential of associated clinical and biological markers to predict its efficacy in patients with advanced HCC (Tables 1 and 2).

## CLINICAL PARAMETERS

### *Alpha-fetoprotein*


Alpha-fetoprotein (AFP) is secreted by about 50% of all HCCs and is the main serological marker used for the diagnosis of the tumor<sup>[5]</sup>. The SHARP trial<sup>[4]</sup> showed that high baseline AFP plasma levels (> 200 ng/mL) had a negative impact on overall survival (OS), a finding recently confirmed in a pooled analysis of the SHARP trial and the Asia Pacific trial by Bruix *et al.*<sup>[6]</sup>. High baseline serum AFP levels ( $\geq$  400 ng/mL) also appear to be associated with shorter time-to-progression (TTP). Notably, in an analysis of six prospective phase II trials evaluating systemic therapies for patients with advanced HCC, no association was observed between baseline AFP levels and prognosis<sup>[7]</sup>.

Several studies<sup>[8-10]</sup> have highlighted a consistent correlation between an early decrease of > 20% in AFP levels following sorafenib and objective response and better outcome in advanced HCC patients. Shao *et al.*<sup>[8]</sup> evaluated for the first time this aspect and they observed that patients with early AFP response had an improved progression-free survival (PFS) (7.5 mo vs 1.9 mo) and OS (15.3 mo vs 4.1 mo). This data was confirmed by Personeni *et al.*<sup>[10]</sup> a few years later. They reported that early responders had a significantly better median OS and TTP than non-responders (13.8 mo vs 8.2 mo,  $P = 0.022$  and 7.9 mo vs 2.4 mo,  $P = 0.004$ ;

**Table 1 Predictive and/or prognostic value of clinical markers in hepatocellular carcinoma patients**

Clinical markers	Predictive value	Prognostic value	Ref.
Alpha-fetoprotein	No	Yes	[6]
Adverse events			
Hand-foot skin reaction	No	Yes	[13]
Hypertension	No	Uncertain	[16,19,20]
Diarrhea	No	Yes	[21]
Child-Pugh A vs B	No	Yes	[27-29]
Macroscopic vascular invasion	No	Yes	[6]
BCLC B vs C	No	Yes	[6,29,32]
Starting dose and dose reduction	No	Yes	[29,32]
Etiology HCV vs HBV	Yes	Yes	[6]
Chronic treatment with metformin	No	Yes	[35,36]
Neutrophil-to-lymphocyte ratio	Yes	Yes	[6,41,44]
Extra hepatic spread	Yes	Yes	[6]

HCC: Hepatocellular carcinoma; BCLC: Barcelona Clinic Liver Cancer; HCV: Hepatitis C virus; HBV: Hepatitis B virus.



**Figure 1 Sorafenib pathway and the main molecular factors.** Ang: Angiopoietin; Tie2: Tyrosine-protein kinase receptor; PDGFR: Platelet-derived growth factor receptors; VEGFR: Vascular endothelial growth factor receptor; SCF: Stem cell factor; PI3K: Phosphatidylinositol 3-Kinase; Akt/PKB: Protein-chinasi B; eNOS: Endothelial nitric oxide synthase; NO: Nitric oxide; P: Phospho-; MEK: Mitogen-activated protein kinase kinase; ERK: Extracellular signal-regulated kinase.

respectively). Conversely, Nakazawa *et al*<sup>[11]</sup> did not observe such an association.

**Adverse events**

The main AEs of Sorafenib are hand-foot skin reaction (HFSR), hypertension and diarrhea. Several papers have highlighted a consistent correlation between AEs and survival in patients treated with Sorafenib.

Vincenzi *et al*<sup>[12]</sup> evaluated for the first time the

correlation between HFSR and outcome. They showed, in a small series of patients treated with sorafenib, that patients with HFSR had a significantly higher disease control rate with respect to patients without HFSR. This data was confirmed in a prospective study of 147 patients by Reig *et al*<sup>[13]</sup>. They reported different OS when patients were subdivided according to the presence or not of skin toxicity during the first 60 d of treatment (18.2 mo vs 10.1 mo, respectively)<sup>[13]</sup>. A

**Table 2 Predictive and/or prognostic value of biological markers in hepatocellular carcinoma patients**

Biological markers	Predictive value	Prognostic value	Ref.
Serum and plasma proteins			
VEGF-A	No	Uncertain	[4,57]
Ang-2	No	Yes	[4]
IGF-1	No	No	[55]
Single nucleotide polymorphisms			
VEGF-A rs2010963	No	Yes	[65]
VEGF-C rs4604006	No	Yes	[65]
eNOS (eNOS-786/eNOS VNTR)	No	Yes	[66]
Ang-2 rs55633437	No	Yes	[67]
HIF-1 <i>alpha</i> rs12434438	No	Yes	[68]
Amplifications			
VEGF	No	Uncertain	[70]
FGF3/FGF4	No	Uncertain	[71]
miRNAs			
miR-425-3p	No	Yes	[74]
miR-224	No	Yes	[75]
miR-181a-5p	No	Yes	[77]
miR-339-5p	No	Yes	[77]
miR-423-5p	No	Yes	[78]
miR-10b-3p	No	Yes	[79]
miR-221	No	Uncertain	[76]
Tissue biomarker expression			
Phospho-ERK	Uncertain	Uncertain	[81,82]
PDGFR-b	No	Yes	[84]
c-Met	No	No	[84]
VEGFR	No	No	[84]
p-c-Jun	No	Yes	[85]

Ang-2: Angiopoietin-2; IGF-1: Insulin-like growth factor-1; VEGF-A: Vascular endothelial growth factor A; HIF-1: Hypoxia-inducible factor 1; FGF: Fibroblast growth factor; miRNAs: MicroRNAs; eNOS: Endothelial nitric oxide synthase; PDGFR: Platelet-derived growth factor receptors; VEGFR: Vascular endothelial growth factor receptor; ERK: Extracellular signal-regulated kinase.

recently meta-analysis confirmed that HSFR was a good indicator of outcome for OS and TTP in HCC patients receiving sorafenib<sup>[14]</sup>

Hypertension (HTN) is frequently associated with the use of angiogenesis inhibitors<sup>[15]</sup>. Casadei Gardini *et al*<sup>[16]</sup> showed that early HTN (15 d after the start of treatment) rather than later onset HTN vs patients without HTN was associated with better PFS (6.0 mo vs 2.5 mo;  $P < 0.001$ ) and OS (14.6 mo vs 3.9 mo;  $P = 0.003$ ). This finding has been confirmed in some studies<sup>[17,18]</sup> but not in others<sup>[19,20]</sup>.

Bettinger *et al*<sup>[21]</sup> reported for the first time that diarrhea was an independent positive prognostic factor (HR = 0.41;  $P = 0.001$ ) in 112 patients with advanced HCC, a finding also confirmed by Koschny *et al*<sup>[22]</sup>.

Finally, other authors showed that the number of AEs was associated with predict survival in patients treated with sorafenib. In particular, Di Costanzo *et al*<sup>[23]</sup> evaluated the potential of pretreatment clinical variables to predict survival. Three groups of patients were taken into account: patients without AEs (group 0), patients with one AE (group 1) and patients with two to three AEs (group 2). The study reported a strong correlation between this classification and disease progression at 3 mo (41.9%, 25.9% and 12.7% of patients in groups

0, 1 and 2, respectively;  $P = 0.014$ ). These data were subsequently confirmed in the validation cohort<sup>[24]</sup>. A recent meta-analysis by Abdel-Rahman *et al*<sup>[25]</sup> revealed an association between specific side-effects (hypertension, HFSR and diarrhea) and patient outcome (HR = 0.38; 95%CI: 0.30-0.48;  $P < 0.00001$ ).

### Stage, liver functionality and etiology

**Child-Pugh A vs Child-Pugh B:** In the SHARP trial<sup>[4]</sup> and the Asia Pacific trial<sup>[26]</sup>, more than 95% of patients were classified as having Child-Pugh A cirrhosis, thus preventing the investigation of the potential benefits of sorafenib in Child-Pugh B patients.

Hollebecque *et al*<sup>[27]</sup> reported for the first time the results from a prospective study on sorafenib efficacy in 120 advanced HCC patients, 20 of whom Child-Pugh B cirrhosis. OS was 11.1 mo, with a significantly longer median survival in Child-Pugh A patients than Child-Pugh B patients (13 mo vs 4.5 mo,  $P = 0.0008$ ). A few years later, Pressiani *et al*<sup>[28]</sup> studied clinical outcome in a population of 300 consecutive patients; PFS in the Child-Pugh A group was 4.3 mo vs 2.1 mo in the Child-Pugh B arm (HR = 3.23; 95%CI: 2.38-4.39;  $P < 0.001$ ), TTP was 4.2 mo vs 3.8 mo and OS was 10.0 mo vs 3.8 mo, respectively ( $P < 0.001$ ).

The most important work on the use of sorafenib in Child-Pugh subgroups was the GIDEON study published in 2016<sup>[29]</sup>. This study observed that median OS was significantly longer in patients with Child-Pugh A (13.6 mo) than in those with Child Pugh B (5.2 mo) or Child-Pugh C (2.6 mo).

**Macroscopic vascular invasion:** It is widely acknowledged that the presence of macroscopic vascular invasion leads to a poorer prognosis<sup>[26,30]</sup>. The meta-analysis by Peng *et al*<sup>[31]</sup> confirmed its prognostic value and the pooled analysis by Bruix *et al*<sup>[6]</sup> affirmed the importance of macroscopic vascular invasion as a predictor of survival but not of response to treatment.

**BCLC stage:** In the SHARP trial, patients with Barcelona Clinic Liver Cancer (BCLC) B had a median OS of 14.5 mo compared to 9.7 mo for those with BCLC C<sup>[4]</sup>. Later, SOFIA<sup>[32]</sup> and GIDEON study<sup>[29]</sup> confirmed this data. In the SOFIA trial<sup>[32]</sup> the OS was 8.4 mo in BCLC C vs 20.6 mo in BCLC B patients ( $P < 0.0001$ ), but the time to radiologic progression did not differ significantly between the 2 groups. In the GIDEON study<sup>[29]</sup>, median OS according to BCLC by Child-Pugh cross-classification followed a similar trend, *i.e.* patients with Child-Pugh A and BCLC stage B showed longer OS than those with Child-Pugh B and BCLC B (19.5 mo vs 10.0 mo); and patients with Child-Pugh A and BCLC stage C had longer OS than those with Child-Pugh B and BCLC stage C (11.2 mo vs 3.8 mo).

Recently, Bruix *et al*<sup>[6]</sup>'s pooled analysis confirmed that BCLC C patients had a poorer prognosis than those with BCLC B HCC (HR = 1.59;  $P = 0.02$ ).

**Sorafenib starting dose and dose escalation/reduction:** The two most important studies that evaluated sorafenib starting dose and dose escalation/reduction are SOFIA<sup>[32]</sup> and GIDEON trial<sup>[29]</sup>.

In the SOFIA trial<sup>[32]</sup> sorafenib was down-dosed in 161 (54%) patients because of AEs (133 patients, 83%) and a reduction in liver function (28 patients, 17%). Median OS of the 77 patients receiving a half-dose of sorafenib for 70% of the treatment period was 21.6 mo (95%CI: 13.6-29.6) compared with 9.6 mo (95%CI: 6.9-12.3) for the remaining 219 patients who had a dose reduction for < 70% of the treatment period or who maintained the full dosage.

A sub-analysis of the GIDEON study<sup>[29]</sup> evaluated the starting dose of sorafenib with respect to clinical outcome and toxicity. Patients starting on 400 mg/d were slightly older, had baseline characteristics indicative of greater disease progression and had a higher incidence of AEs than those with a starting dose of 800 mg/d (96% vs 88%). Treatment duration (18.0 wk vs 13.0 wk) and median OS (12.1 mo vs 9.4 mo) were longer in patients receiving 800 mg/d.

**Etiology:** In the subgroup analysis of the SHARP

study<sup>[4]</sup>, the HR for OS was 0.76 in HBV-positive patients (95%CI: 0.38-1.50,  $P =$  not significant) and 0.50 (95%CI: 0.32-0.77) in HCV-positive patients. Results were similar for TTP (HR = 1.03 and 0.43 for HBV-positive and HCV-positive patients, respectively). Similar data were obtained for HBV-positive HCC patients in the phase III randomized Asia Pacific trial, *i.e.* the HR for OS was 0.74 (95%CI: 0.51-1.06, not significant) with respect to patients with the other etiology, for which the HR was 0.57 (95%CI: 0.29-1.33)<sup>[26]</sup>. Bruix *et al*<sup>[6]</sup>'s pooled analysis of the SHARP/Asia Pacific trial results showed that the absence of HCV was a potential prognostic factor for poorer OS (HR = 0.7,  $P = 0.02$ ). The same authors revealed that HBV-positive patients did not show a significant difference in treatment response with respect to their HBV-negative counterparts (HR = 0.78; 95%CI: 0.57-1.06) and OS (HR = 1.128,  $P = 0.4538$ ). We believe that the 2 etiologic groups respond differently to sorafenib and that further investigation is warranted in specific studies<sup>[33]</sup>.

**Metformin treatment:** Type 2 diabetes is a significant risk factor for the development of malignancies, including HCC<sup>[34]</sup>. Casadei Gardini *et al*<sup>[35]</sup> published findings of reduced sorafenib efficacy in HCC patients treated chronically with or without metformin for type II diabetes mellitus (PFS 2.6 mo vs 5.0 mo, respectively; and OS 10.4 mo vs 15.1 mo, respectively). The same authors validated these data in a series of more than 250 cases<sup>[36]</sup>, also highlighting a possible role of sirtuin-3 in resistance to sorafenib<sup>[37]</sup>. Di Costanzo *et al*<sup>[38]</sup> recently reported an increase in TTP and OS in diabetic with respect to non-diabetic HCC patients. However, no distinction was made between the different hypoglycemic therapies administered.

#### Immune inflammation indicators

Systemic inflammatory responses have been shown to reflect the promotion of angiogenesis, DNA damage and tumor invasion through an upregulation of cytokines<sup>[39]</sup>. Previous research revealed that lymphocytes play a crucial role in tumor defense by inducing cytotoxic cell death and inhibiting tumor cell proliferation and migration<sup>[40]</sup>. Consequently, several inflammation and immune-based prognostic scores, such as lymphocyte count, neutrophil-lymphocyte ratio (NLR), and systemic immune-inflammation index (SII), have been developed to predict survival and recurrence in cancers, including HCC. Casadei Gardini *et al*<sup>[41]</sup> evaluated for the first time SII, NLR and platelet-lymphocyte ratio (PLR) in a small case series, observing that SII were independent prognostic factors for OS. Other studies showed that NLR was a significant independent risk factor for shorter survival<sup>[42,43]</sup>. NLR was also found to be an independent prognostic factor for both response and survival in Bruix *et al*<sup>[6]</sup>'s pooled analysis and Lue *et al*<sup>[44]</sup>'s retrospective



study on Spanish patients.

## IMAGING EXAMINATIONS

The response to sorafenib does not correlate with a change in lesion dimension, but it is more correlated with intralesional vascularization. For this reason, the RECIST criteria<sup>[45,46]</sup> usually used for tumor response evaluation is inappropriate to evaluate the response to sorafenib in patients with advanced HCC. The modified RECIST (mRECIST) appear more indicate for evaluation the response. They include vascularization and tumor arterial enhancement changes of the target lesion on computed tomographic (CT). Several studies have demonstrated the superiority of the mRECIST criteria with respect to the RECIST criteria in assessing the response to treatment with sorafenib<sup>[47]</sup>. Various functional imaging tools were proposed to evaluate the antiangiogenic effects, but none of these has entered normal clinical practice<sup>[48-52]</sup>. Finally, a recently study showed that texture features on pretreatment contrast material-enhanced CT images can help predict OS and TTP in these patients<sup>[53]</sup>.

## BIOLOGICAL PARAMETERS

### Serum and plasma proteins

Although plasma biomarkers are the best candidates for evaluating sorafenib efficacy, only the SHARP trial produced results with borderline significance<sup>[4]</sup>. Baseline angiopoietin-2 (Ang-2) and vascular endothelial growth factor-A (VEGF-A) plasma levels independently predicted survival in both the entire patient population and the placebo cohort. Conversely, none of the tested biomarkers significantly predicted response to sorafenib<sup>[4]</sup>. Insulin-like growth factor (IGF)-1 levels have been found to decrease in patients with cirrhosis of the liver or HCC<sup>[54]</sup>, and high pretreatment levels of IGF-1 predict better PFS and OS in advanced HCC patients receiving first-line antiangiogenic therapy<sup>[55]</sup>.

The role of serum cytokines as biomarkers for the prediction of sorafenib responses is interesting, in particular Kim *et al.*<sup>[56]</sup> developed a new prediction model for sorafenib response that combines relevant serum markers, tumor related factors, and cirrhosis-related factors in a scoring system.

**VEGF-A:** Llovet *et al.*<sup>[4]</sup> showed that, although baseline plasma VEGF-A concentrations did not exhibit a predictive value, low plasma VEGF-A was associated with improved prognosis (HR = 1.48, 95%CI: 1.08-2.03,  $P = 0.015$ ). However, other authors did not find any association between VEGF-A and prognosis in patients treated with sorafenib<sup>[57]</sup>. Tsuchiya *et al.*<sup>[58]</sup>'s analysis of plasma VEGF concentrations during sorafenib treatment revealed that a decrease in the protein 8 wk after the start of therapy predicted better overall survival

in advanced HCC patients (30.9 mo vs 14.4 mo;  $P = 0.038$ ).

**Ang-2:** In the presence of VEGF, Ang-2 destabilizes blood vessels, promotes vascular sprouting, and is associated with an invasive and metastatic cancer phenotype<sup>[59]</sup>. Llovet *et al.*<sup>[4]</sup> demonstrated that high baseline Ang-2 levels were correlated with more aggressive disease (HR = 1.58, 95%CI: 1.20-2.07,  $P = 0.001$ ). Moreover, levels of the protein increased during treatment in the placebo group, suggestive of poor outcome related to disease progression in this cohort, whereas they remained constant during treatment with sorafenib, reflecting the generally more favorable outcome of this group. Overall increased Ang-2 expression levels were associated with poorer outcome in both groups, suggesting that this marker could be useful in monitoring treatment response. In agreement with Llovet's study, Miyahara *et al.*<sup>[57]</sup> reported that high baseline Ang-2 serum levels were associated with poor outcome in advanced HCC patients receiving sorafenib (HR = 2.51, 95%CI: 1.01-6.57,  $P = 0.048$ ). Although these results indicate the potential prognostic value of Ang-2 in HCC, its role in predicting response to sorafenib remains to be verified.

**IGF-1:** Shao *et al.*<sup>[55]</sup> found that high-pretreatment serum levels of IGF-1 were associated with a better DCR and improved PFS and OS in patients undergoing antiangiogenic therapy. Although the study did not have a control arm, the substantial significant difference in DCR between patients with high and low levels of IGF-1 (71% vs 39%) denotes the potential usefulness of IGF-1 as a predictive biomarker of response to antiangiogenic therapy.

**Multiple-factor analyses:** By using baseline serum basic fibroblast growth factor (bFGF) and hepatocyte growth factor (HGF) levels as covariates together etiology (B-viral), platelet count, BCLC stage and protein induced by vitamin K absence-II, Kim *et al.*<sup>[56]</sup> reported that a total score of < 6 could be a relevant cutoff value for selecting patients who are most likely to benefit from sorafenib therapy. Moreover, Hayashi *et al.*<sup>[60]</sup> found that serum interleukin (IL)-5, IL-8, CXCL9, PDGF-BB, TGF- $\alpha$ , and VEGF-A were elevated in the long survivors group among HCC patients who received sorafenib, potentially reflecting the activation of stromal signaling in the tumor microenvironment.

### Genetic markers

Molecular and genomic analyses from tumor and non-tumor tissue have proven useful in evaluating prognosis and could open up new avenues for tailoring treatment<sup>[61]</sup>. Genetic alterations, such as single nucleotide polymorphisms (SNPs) in genes encoding for proteins involved in the angiogenic process,

have been studied as potential biomarkers for anti-angiogenic therapy. SNP evaluation would seem to be more advantageous than protein or gene expression analyses as it can be performed at any time during the course of the disease, is not substantially influenced by laboratory biases, and is relatively inexpensive. Some authors have focused on molecular profiling in formalin-fixed paraffin-embedded (FFPE) samples, comparing the mutation profiles of HCC biopsy samples and the response to sorafenib treatment<sup>[62]</sup>. Gene amplification, gene mutations and expression profiling of tumors have now become a research priority and are expected to lead to personalized treatment for HCC patients<sup>[63,64]</sup>.

**SNPs:** Specific SNPs in *VEGF* and *VEGFR* genes have been found to be correlated with PFS and OS in HCC patients treated with sorafenib. In multivariate analysis, *VEGF-A* rs2010963 and *VEGF-C* rs4604006 were found to be independent factors influencing outcome in terms of PFS (HR = 0.25, 95%CI: 0.19-1.02,  $P = 0.0376$  and HR = 0.22, 95%CI: 0.14-0.81,  $P = 0.004$ , respectively) and OS (HR = 0.28, 95%CI: 0.23-0.96,  $P = 0.02$  and HR = 0.25, 95%CI: 0.17-0.99,  $P = 0.04$ , respectively)<sup>[65]</sup>. In the Italian multicenter, retrospective ePHAS [endothelial nitric oxide synthase (*eNOS*) polymorphisms in HCC and sorafenib] study, *eNOS* polymorphisms were analyzed in relation to PFS and OS. In univariate analysis, training cohort patients homozygous for *eNOS* haplotype (HT1:T-4b at *eNOS-786/eNOS VNTR*) showed a lower median PFS (2.6 mo vs 5.8 mo, HR = 5.43, 95%CI: 2.46-11.98,  $P < 0.0001$ ) and OS (3.2 mo vs 14.6 mo, HR = 2.35, 95%CI: 1.12-4.91,  $P = 0.024$ ) than those with other haplotypes. These results were confirmed in a validation set and multivariate analysis further substantiated this haplotype as the only independent prognostic factor<sup>[66]</sup>. More recently, evidence emerged that patients homozygous for *ANGPT2* (*Ang2* gene) rs55633437 GG genotype showed significantly longer PFS ( $P < 0.001$ ) and OS ( $P < 0.001$ ) than those with the other genotypes (GT+TT)<sup>[67]</sup>.

In the ALICE-2 study, Faloppi *et al.*<sup>[68]</sup> investigated the role of hypoxia-inducible factor 1-alpha (*HIF-1 $\alpha$* ) SNPs, confirming the results of the ALICE-1 study<sup>[65]</sup>. In multivariate analysis, rs12434438 of *HIF-1 $\alpha$* , rs2010963 of *VEGF-A* and rs4604006 of *VEGF-C* were confirmed as independent factors and may help to identify patients who are more likely to respond to sorafenib<sup>[68]</sup>. The prospective INNOVATE study is ongoing to validate the role of *VEGF*, *eNOS*, *Ang-2* and *HIF-1 $\alpha$*  SNPs in relation to clinical outcome in advanced HCC patients treated with sorafenib (NCT02786342)<sup>[69]</sup>.

**Gene amplification, gene mutations and RNA expression:** A relation between *VEGF-A* gene amplification and response to sorafenib was observed in a study performed on a mouse model of HCC<sup>[70]</sup>.

The authors found that HCC patients with tumor *VEGF-A* amplification showed markedly better survival than those with non-amplified tumors, highlighting that *VEGFA* amplification is a potential biomarker of response to *VEGF-A*-blocking drugs in HCC<sup>[70]</sup>. Arao *et al.*<sup>[71]</sup> observed that FGF3/FGF4 amplification and multiple lung metastases were frequently observed in responders to sorafenib, although the sample size was relatively small.

Sakai *et al.*<sup>[62]</sup> used targeted DNA and RNA sequencing in FFPE specimens from fine-needle biopsy to identify candidate biomarkers of response to sorafenib in 46 HCC patients. A significant difference was observed in the number of oncogene mutations between progressing and non-progressing patients ( $P = 0.045$ ), suggesting that tumor mutational burden may be predictive of sorafenib effectiveness. Tumor gene expression of NRG1, TGF $\alpha$ , and PECAM1 would also seem to be a marker of treatment response and PFS.

### MicroRNAs

MicroRNAs (miRNAs) affect drug response directly or indirectly by regulating the expression of genes involved in drug transportation, metabolism, and downstream signaling pathways. The deregulation of various miRNAs has been reported in *in vitro*, *in vivo* and population studies<sup>[72,73]</sup>, confirming its correlation with response to sorafenib. Some authors have evaluated the predictive role of miRNA expression in HCC tissue<sup>[74,75]</sup>, while others have studied circulating miRNA levels prior to sorafenib treatment<sup>[76]</sup>. To date, the most interesting tissue miRNAs are miR-425-3p<sup>[74]</sup> and miR-224<sup>[75]</sup>. High levels of miR-425-3p have been associated with longer TTP and PFS (HR = 0.4, 95%CI: 0.2-0.7,  $P = 0.0008$  and HR = 0.5, 95%CI: 0.3-0.9,  $P = 0.007$ , respectively), and elevated miR-224 expression have been correlated with increased PFS and OS (HR = 0.28, 95%CI: 0.09-0.92,  $P = 0.029$  and HR = 0.0.24, 95%CI: 0.07-0.79,  $P = 0.012$ , respectively). Circulating miRNAs have also been studied in the serum of HCC patients to predict early response to sorafenib treatment, with miR-181a-5p and miR-339-5p associated with partial response and disease progression<sup>[77]</sup>, miR-423-5p with stable disease or partial response<sup>[78]</sup> and miR-10b-3p with shorter survival<sup>[79]</sup>.

Another potentially interesting circulating miRNA is miR-221, which was studied by Fornari *et al.*<sup>[76]</sup> in both animal models and in a patient population. Patients with radiologic disease progression after 2-mo treatment had higher pretreatment miR-221 levels than responders ( $P = 0.007$ ).

Larger confirmatory studies are needed before miRNAs can be considered valid biomarkers for clinical practice.

### Tissue biomarker expression

A number of studies have analyzed tissue biomarkers

that may be very specific to the disease of interest<sup>[80]</sup>. In a phase II study of sorafenib in advanced HCC, Abou-Alfa *et al.*<sup>[81]</sup> showed that patients whose tumors expressed higher baseline phospho-ERK levels had a longer TTP. However, other studies reported conflicting results<sup>[82,83]</sup>. With regard to the expression of angiogenic markers in tumor tissue, it has been observed that high platelet-derived growth factor receptor beta expression is correlated with poor OS but not with PFS in HCC patients receiving sorafenib. High expression of the proto-oncogene c-Met may predict the therapeutic effectiveness of sorafenib in HCC patients, but no differences in terms of outcome have been seen with respect to VEGFR-2 expression<sup>[84]</sup>. Hagiwara *et al.*<sup>[85]</sup> studied another interesting tissue biomarker, phospho-c-Jun, reporting a significantly higher expression ( $P < 0.001$ ) in non-responding compared to responding patients treated with sorafenib.

## CONCLUSION

After 10 years of research into sorafenib, there are still no validated prognostic or predictive markers of response to sorafenib in hepatocellular carcinoma. Furthermore, the main results obtained to date come from 2 important randomized trials and from different subanalyses and pooled analyses rather than from normal clinical practice. The fact of there being only one drug for the treatment of these patients has certainly done nothing to stimulate research into identifying and validating predictors of response and prognosis. However, given the recent publication of a positive phase III trial<sup>[86]</sup> and the ongoing NCT01658878 immunotherapy study, the race is now on to see who will be the first to identify a prognostic and predictive factor for sorafenib and/or new drugs in this setting. In conclusion, the use of metabolomic profiling and whole genome analysis to examine the association between patient outcome and response to sorafenib could become alternative approaches to the search for new biomarkers in HCC.

## ACKNOWLEDGMENTS

The authors thank Gráinne Tierney and Cristiano Verna for editorial assistance.

## REFERENCES

- Llovet JM, Ricci S, Mazzaferro V, Hilgard P, Gane E, Blanc JF, de Oliveira AC, Santoro A, Raoul JL, Forner A, Schwartz M, Porta C, Zeuzem S, Bolondi L, Greten TF, Galle PR, Seitz JF, Borbath I, Häussinger D, Giannaris T, Shan M, Moscovici M, Voliotis D, Bruix J; SHARP Investigators Study Group. Sorafenib in advanced hepatocellular carcinoma. *N Engl J Med* 2008; **359**: 378-390 [PMID: 18650514 DOI: 10.1056/NEJMoa0708857]
- Cervello M, Bachvarov D, Lampiasi N, Cusimano A, Azzolina A, McCubrey JA, Montalto G. Molecular mechanisms of sorafenib action in liver cancer cells. *Cell Cycle* 2012; **11**: 2843-2855 [PMID: 22801548 DOI: 10.4161/cc.21193]
- Tai WT, Cheng AL, Shiau CW, Huang HP, Huang JW, Chen PJ, Chen KF. Signal transducer and activator of transcription 3 is a major kinase-independent target of sorafenib in hepatocellular carcinoma. *J Hepatol* 2011; **55**: 1041-1048 [PMID: 21354226 DOI: 10.1016/j.jhep.2011.01.047]
- Llovet JM, Peña CE, Lathia CD, Shan M, Meinhardt G, Bruix J; SHARP Investigators Study Group. Plasma biomarkers as predictors of outcome in patients with advanced hepatocellular carcinoma. *Clin Cancer Res* 2012; **18**: 2290-2300 [PMID: 22374331 DOI: 10.1158/1078-0432.CCR-11-2175]
- Forner A, Llovet JM, Bruix J. Hepatocellular carcinoma. *Lancet* 2012; **379**: 1245-1255 [PMID: 22353262 DOI: 10.1016/S0140-6736(11)61347-0]
- Bruix J, Cheng AL, Meinhardt G, Nakajima K, De Sanctis Y, Llovet J. Prognostic factors and predictors of sorafenib benefit in patients with hepatocellular carcinoma: Analysis of two phase III studies. *J Hepatol* 2017; **67**: 999-1008 [PMID: 28687477 DOI: 10.1016/j.jhep.2017.06.026]
- Lin ZZ, Hsu C, Hu FC, Shao YY, Chang DY, Yang CH, Hong RL, Hsu CH, Cheng AL. Factors impacting prognosis prediction in BCLC stage C and Child-Pugh class A hepatocellular carcinoma patients in prospective clinical trials of systemic therapy. *Oncologist* 2012; **17**: 970-977 [PMID: 22673633 DOI: 10.1634/theoncologist.2011-0411]
- Shao YY, Lin ZZ, Hsu C, Shen YC, Hsu CH, Cheng AL. Early alpha-fetoprotein response predicts treatment efficacy of antiangiogenic systemic therapy in patients with advanced hepatocellular carcinoma. *Cancer* 2010; **116**: 4590-4596 [PMID: 20572033 DOI: 10.1002/ncr.25257]
- Yau T, Yao TJ, Chan P, Wong H, Pang R, Fan ST, Poon RT. The significance of early alpha-fetoprotein level changes in predicting clinical and survival benefits in advanced hepatocellular carcinoma patients receiving sorafenib. *Oncologist* 2011; **16**: 1270-1279 [PMID: 21885876 DOI: 10.1634/theoncologist.2011-0105]
- Personeni N, Bozzarelli S, Pressiani T, Rimassa L, Tronconi MC, Sclafani F, Carnaghi C, Pedicini V, Giordano L, Santoro A. Usefulness of alpha-fetoprotein response in patients treated with sorafenib for advanced hepatocellular carcinoma. *J Hepatol* 2012; **57**: 101-107 [PMID: 22414760 DOI: 10.1016/j.jhep.2012.02.016]
- Nakazawa T, Hidaka H, Takada J, Okuwaki Y, Tanaka Y, Watanabe M, Shibuya A, Minamino T, Kokubu S, Koizumi W. Early increase in  $\alpha$ -fetoprotein for predicting unfavorable clinical outcomes in patients with advanced hepatocellular carcinoma treated with sorafenib. *Eur J Gastroenterol Hepatol* 2013; **25**: 683-689 [PMID: 23395995 DOI: 10.1097/MEG.0b013e32835d913b]
- Vincenzi B, Santini D, Russo A, Addeo R, Giuliani F, Montella L, Rizzo S, Venditti O, Frezza AM, Caraglia M, Colucci G, Del Prete S, Tonini G. Early skin toxicity as a predictive factor for tumor control in hepatocellular carcinoma patients treated with sorafenib. *Oncologist* 2010; **15**: 85-92 [PMID: 20051477 DOI: 10.1634/theoncologist.2009-0143]
- Reig M, Torres F, Rodriguez-Lope C, Forner A, Llovet JM, Rimola J, Darnell A, Ríos J, Ayuso C, Bruix J. Early dermatologic adverse events predict better outcome in HCC patients treated with sorafenib. *J Hepatol* 2014; **61**: 318-324 [PMID: 24703956 DOI: 10.1016/j.jhep.2014.03.030]
- Wang P, Tan G, Zhu M, Li W, Zhai B, Sun X. Hand-foot skin reaction is a beneficial indicator of sorafenib therapy for patients with hepatocellular carcinoma: a systemic review and meta-analysis. *Expert Rev Gastroenterol Hepatol* 2018; **12**: 1-8 [PMID: 28847184 DOI: 10.1080/17474124.2017.1373018]
- Scartozzi M, Galizia E, Chiarrini S, Giampieri R, Berardi R, Pierantoni C, Cascinu S. Arterial hypertension correlates with clinical outcome in colorectal cancer patients treated with first-line bevacizumab. *Ann Oncol* 2009; **20**: 227-230 [PMID: 18842611 DOI: 10.1093/annonc/mdn637]
- Casadei Gardini A, Scarpi E, Marisi G, Foschi FG, Donati G, Giampalma E, Faloppi L, Scartozzi M, Silvestris N, Bisulli M, Corbelli J, Gardini A, La Barba G, Veneroni L, Tamperi S, Cascinu



- S, Frassinetti GL. Early onset of hypertension and serum electrolyte changes as potential predictive factors of activity in advanced HCC patients treated with sorafenib: results from a retrospective analysis of the HCC-AVR group. *Oncotarget* 2016; **7**: 15243-15251 [PMID: 26893366 DOI: 10.18632/oncotarget.7444]
- 17 **Zhong BY**, Ni CF, Chen L, Zhu HD, Teng GJ. Early Sorafenib-related Biomarkers for Combination Treatment with Transarterial Chemoembolization and Sorafenib in Patients with Hepatocellular Carcinoma. *Radiology* 2017; **284**: 583-592 [PMID: 28263701 DOI: 10.1148/radiol.2017161975]
  - 18 **Howell J**, Pinato DJ, Ramaswami R, Bettinger D, Arizumi T, Ferrari C, Yen C, Gibbin A, Burlone ME, Guaschino G, Sellers L, Black J, Pirisi M, Kudo M, Thimme R, Park JW, Sharma R. On-target sorafenib toxicity predicts improved survival in hepatocellular carcinoma: a multi-centre, prospective study. *Aliment Pharmacol Ther* 2017; **45**: 1146-1155 [PMID: 28252185 DOI: 10.1111/apt.13977]
  - 19 **Shin SY**, Lee YJ. Correlation of skin toxicity and hypertension with clinical benefit in advanced hepatocellular carcinoma patients treated with sorafenib. *Int J Clin Pharmacol Ther* 2013; **51**: 837-846 [PMID: 24075093 DOI: 10.5414/CP201907]
  - 20 **Otsuka T**, Eguchi Y, Kawazoe S, Yanagita K, Ario K, Kitahara K, Kawasoe H, Kato H, Mizuta T; Saga Liver Cancer Study Group. Skin toxicities and survival in advanced hepatocellular carcinoma patients treated with sorafenib. *Hepatol Res* 2012; **42**: 879-886 [PMID: 22469363 DOI: 10.1111/j.1872-034X.2012.00991.x]
  - 21 **Bettinger D**, Schultheiss M, Knüppel E, Thimme R, Blum HE, Spangenberg HC. Diarrhea predicts a positive response to sorafenib in patients with advanced hepatocellular carcinoma. *Hepatology* 2012; **56**: 789-790 [PMID: 22307848 DOI: 10.1002/hep.25637]
  - 22 **Koschny R**, Gotthardt D, Koehler C, Jaeger D, Stremmel W, Ganten TM. Diarrhea is a positive outcome predictor for sorafenib treatment of advanced hepatocellular carcinoma. *Oncology* 2013; **84**: 6-13 [PMID: 23075905 DOI: 10.1159/000342425]
  - 23 **Di Costanzo GG**, de Stefano G, Tortora R, Farella N, Addario L, Lampasi F, Lanza AG, Cordone G, Imperato M, Caporaso N. Sorafenib off-target effects predict outcomes in patients treated for hepatocellular carcinoma. *Future Oncol* 2015; **11**: 943-951 [PMID: 25760975 DOI: 10.2217/fon.14.291]
  - 24 **Di Costanzo GG**, Casadei Gardini A, Marisi G, Foschi FG, Scartozzi M, Granata R, Faloppi L, Cascinu S, Silvestris N, Brunetti O, Palmieri VO, Ercolani G, Tortora R. Validation of a Simple Scoring System to Predict Sorafenib Effectiveness in Patients with Hepatocellular Carcinoma. *Target Oncol* 2017; **12**: 795-803 [PMID: 28770532 DOI: 10.1007/s11523-017-0522-5]
  - 25 **Abdel-Rahman O**, Lamarca A. Development of sorafenib-related side effects in patients diagnosed with advanced hepatocellular carcinoma treated with sorafenib: a systematic-review and meta-analysis of the impact on survival. *Expert Rev Gastroenterol Hepatol* 2017; **11**: 75-83 [PMID: 27882800 DOI: 10.1080/17474124.2017.1264874]
  - 26 **Cheng AL**, Guan Z, Chen Z, Tsao CJ, Qin S, Kim JS, Yang TS, Tak WY, Pan H, Yu S, Xu J, Fang F, Zou J, Lentini G, Voliotis D, Kang YK. Efficacy and safety of sorafenib in patients with advanced hepatocellular carcinoma according to baseline status: subset analyses of the phase III Sorafenib Asia-Pacific trial. *Eur J Cancer* 2012; **48**: 1452-1465 [PMID: 22240282 DOI: 10.1016/j.ejca.2011.12.006]
  - 27 **Hollebecque A**, Cattan S, Romano O, Sergent G, Mourad A, Louvet A, Dharancy S, Boleslawski E, Truant S, Pruvot FR, Hebbbar M, Ernst O, Mathurin P. Safety and efficacy of sorafenib in hepatocellular carcinoma: the impact of the Child-Pugh score. *Aliment Pharmacol Ther* 2011; **34**: 1193-1201 [PMID: 21958438 DOI: 10.1111/j.1365-2036.2011.04860.x]
  - 28 **Pressiani T**, Boni C, Rimassa L, Labianca R, Fagioli S, Salvagni S, Ferrari D, Cortesi E, Porta C, Mucciarini C, Latini L, Carnaghi C, Banzi M, Fanello S, De Giorgio M, Lutman FR, Torzilli G, Tommasini MA, Ceriani R, Covini G, Tronconi MC, Giordano L, Locopo N, Naimo S, Santoro A. Sorafenib in patients with Child-Pugh class A and B advanced hepatocellular carcinoma: a prospective feasibility analysis. *Ann Oncol* 2013; **24**: 406-411 [PMID: 23041587 DOI: 10.1093/annonc/mds343]
  - 29 **Marrero JA**, Kudo M, Venook AP, Ye SL, Bronowicki JP, Chen XP, Dagher L, Furuse J, Geschwind JH, de Guevara LL, Papandreou C, Takayama T, Sanyal AJ, Yoon SK, Nakajima K, Lehr R, Heldner S, Lencioni R. Observational registry of sorafenib use in clinical practice across Child-Pugh subgroups: The GIDEON study. *J Hepatol* 2016; **65**: 1140-1147 [PMID: 27469901 DOI: 10.1016/j.jhep.2016.07.020]
  - 30 **Lee S**, Kang JH, Kim DY, Ahn SH, Park JY, Kim BK, Kim SU, Han KH. Prognostic factors of sorafenib therapy in hepatocellular carcinoma patients with failure of transarterial chemoembolization. *Hepatol Int* 2017; **11**: 292-299 [PMID: 28324324 DOI: 10.1007/s12072-017-9792-3]
  - 31 **Peng S**, Zhao Y, Xu F, Jia C, Xu Y, Dai C. An updated meta-analysis of randomized controlled trials assessing the effect of sorafenib in advanced hepatocellular carcinoma. *PLoS One* 2014; **9**: e112530 [PMID: 25460347 DOI: 10.1371/journal.pone.0112530]
  - 32 **Iavarone M**, Cabibbo G, Piscaglia F, Zavaglia C, Grieco A, Villa E, Cammà C, Colombo M; SOFIA (SORafenib Italian Assessment) study group. Field-practice study of sorafenib therapy for hepatocellular carcinoma: a prospective multicenter study in Italy. *Hepatology* 2011; **54**: 2055-2063 [PMID: 21898496 DOI: 10.1002/hep.24644]
  - 33 **Casadei Gardini A**, Frassinetti GL, Foschi FG, Ercolani G, Ulivi P. Sorafenib and Regorafenib in HBV- or HCV-positive hepatocellular carcinoma patients: Analysis of RESORCE and SHARP trials. *Dig Liver Dis* 2017; **49**: 943-944 [PMID: 28546063 DOI: 10.1016/j.dld.2017.04.022]
  - 34 **Donadon V**, Balbi M, Mas MD, Casarin P, Zanette G. Metformin and reduced risk of hepatocellular carcinoma in diabetic patients with chronic liver disease. *Liver Int* 2010; **30**: 750-758 [PMID: 20331505 DOI: 10.1111/j.1478-3231.2010.02223.x]
  - 35 **Casadei Gardini A**, Marisi G, Scarpi E, Scartozzi M, Faloppi L, Silvestris N, Masi G, Vivaldi C, Brunetti O, Tamberi S, Foschi FG, Tamburini E, Tenti E, Ricca Rosellini S, Ulivi P, Cascinu S, Nanni O, Frassinetti GL. Effects of metformin on clinical outcome in diabetic patients with advanced HCC receiving sorafenib. *Expert Opin Pharmacother* 2015; **16**: 2719-2725 [PMID: 26513009 DOI: 10.1517/14656566.2015.1102887]
  - 36 **Casadei Gardini A**, Faloppi L, De Matteis S, Foschi FG, Silvestris N, Tovoli F, Palmieri V, Marisi G, Brunetti O, Vespasiani-Gentilucci U, Perrone G, Valgiusti M, Granato AM, Ercolani G, Negrini G, Tamburini E, Aprile G, Passardi A, Santini D, Cascinu S, Frassinetti GL, Scartozzi M. Metformin and insulin impact on clinical outcome in patients with advanced hepatocellular carcinoma receiving sorafenib: Validation study and biological rationale. *Eur J Cancer* 2017; **86**: 106-114 [PMID: 28985579 DOI: 10.1016/j.ejca.2017.09.003]
  - 37 **De Matteis S**, Granato AM, Napolitano R, Molinari C, Valgiusti M, Santini D, Foschi FG, Ercolani G, Vespasiani Gentilucci U, Faloppi L, Scartozzi M, Frassinetti GL, Casadei Gardini A. Interplay Between SIRT-3, Metabolism and Its Tumor Suppressor Role in Hepatocellular Carcinoma. *Dig Dis Sci* 2017; **62**: 1872-1880 [PMID: 28527050 DOI: 10.1007/s10620-017-4615-x]
  - 38 **Di Costanzo GG**, Tortora R, Morisco F, Addario L, Guarino M, Cordone G, Falco L, Caporaso N. Impact of Diabetes on Outcomes of Sorafenib Therapy for Hepatocellular Carcinoma. *Target Oncol* 2017; **12**: 61-67 [PMID: 27503006 DOI: 10.1007/s11523-016-0454-5]
  - 39 **Ferrucci PF**, Gandini S, Battaglia A, Alfieri S, Di Giacomo AM, Giannarelli D, Cappellini GC, De Galitiis F, Marchetti P, Amato G, Lazzeri A, Pala L, Cocorocchio E, Martinoli C. Baseline neutrophil-to-lymphocyte ratio is associated with outcome of ipilimumab-treated metastatic melanoma patients. *Br J Cancer* 2015; **112**: 1904-1910 [PMID: 26010413 DOI: 10.1038/bjc.2015.180]
  - 40 **Hu B**, Yang XR, Xu Y, Sun YF, Sun C, Guo W, Zhang X, Wang



- WM, Qiu SJ, Zhou J, Fan J. Systemic immune-inflammation index predicts prognosis of patients after curative resection for hepatocellular carcinoma. *Clin Cancer Res* 2014; **20**: 6212-6222 [PMID: 25271081 DOI: 10.1158/1078-0432.CCR-14-0442]
- 41 **Casadei Gardini A**, Scarpi E, Faloppi L, Scartozzi M, Silvestris N, Santini D, de Stefano G, Marisi G, Negri FV, Foschi FG, Valgiusti M, Ercolani G, Frassinetti GL. Immune inflammation indicators and implication for immune modulation strategies in advanced hepatocellular carcinoma patients receiving sorafenib. *Oncotarget* 2016; **7**: 67142-67149 [PMID: 27613839 DOI: 10.18632/oncotarget.11565]
- 42 **Howell J**, Pinato DJ, Ramaswami R, Arizumi T, Ferrari C, Gibbin A, Burlone ME, Guaschino G, Toniutto P, Black J, Sellers L, Kudo M, Pirisi M, Sharma R. Integration of the cancer-related inflammatory response as a stratifying biomarker of survival in hepatocellular carcinoma treated with sorafenib. *Oncotarget* 2017; **8**: 36161-36170 [PMID: 28212535 DOI: 10.18632/oncotarget.15322]
- 43 **da Fonseca LG**, Barroso-Sousa R, Bento Ada S, Blanco BP, Valente GL, Pffiffer TE, Hoff PM, Sabbaga J. Pre-treatment neutrophil-to-lymphocyte ratio affects survival in patients with advanced hepatocellular carcinoma treated with sorafenib. *Med Oncol* 2014; **31**: 264 [PMID: 25273866 DOI: 10.1007/s12032-014-0264-5]
- 44 **Lué A**, Serrano MT, Bustamante FJ, Iñarrairaegui M, Arenas JI, Testillano M, Lorente S, Gil C, de la Torre M, Gomez A, Sangro B. Neutrophil-to-lymphocyte ratio predicts survival in European patients with hepatocellular carcinoma administered sorafenib. *Oncotarget* 2017; **8**: 103077-103086 [PMID: 29262546 DOI: 10.18632/oncotarget.21528]
- 45 **Therasse P**, Arbuck SG, Eisenhauer EA, Wanders J, Kaplan RS, Rubinstein L, Verweij J, Van Glabbeke M, van Oosterom AT, Christian MC, Gwyther SG. New guidelines to evaluate the response to treatment in solid tumors. European Organization for Research and Treatment of Cancer, National Cancer Institute of the United States, National Cancer Institute of Canada. *J Natl Cancer Inst* 2000; **92**: 205-216 [PMID: 10655437]
- 46 **Eisenhauer EA**, Therasse P, Bogaerts J, Schwartz LH, Sargent D, Ford R, Dancy J, Arbuck S, Gwyther S, Mooney M, Rubinstein L, Shankar L, Dodd L, Kaplan R, Lacombe D, Verweij J. New response evaluation criteria in solid tumours: revised RECIST guideline (version 1.1). *Eur J Cancer* 2009; **45**: 228-247 [PMID: 19097774 DOI: 10.1016/j.ejca.2008.10.026]
- 47 **Ronot M**, Bouattour M, Wassermann J, Bruno O, Dreyer C, Larroque B, Castera L, Vilgrain V, Belghiti J, Raymond E, Faivre S. Alternative Response Criteria (Choi, European association for the study of the liver, and modified Response Evaluation Criteria in Solid Tumors [RECIST]) Versus RECIST 1.1 in patients with advanced hepatocellular carcinoma treated with sorafenib. *Oncologist* 2014; **19**: 394-402 [PMID: 24652387 DOI: 10.1634/theoncologist.2013-0114]
- 48 **Sugimoto K**, Moriyasu F, Saito K, Rognin N, Kamiyama N, Furuichi Y, Imai Y. Hepatocellular carcinoma treated with sorafenib: early detection of treatment response and major adverse events by contrast-enhanced US. *Liver Int* 2013; **33**: 605-615 [PMID: 23305331 DOI: 10.1111/liv.12098]
- 49 **Zocco MA**, Garcovich M, Lupascu A, Di Stasio E, Roccarina D, Annicchiarico BE, Riccardi L, Ainora ME, Ponziani F, Caracciolo G, Rapaccini GL, Landolfi R, Siciliano M, Pompili M, Gasbarrini A. Early prediction of response to sorafenib in patients with advanced hepatocellular carcinoma: the role of dynamic contrast enhanced ultrasound. *J Hepatol* 2013; **59**: 1014-1021 [PMID: 23811306 DOI: 10.1016/j.jhep.2013.06.011]
- 50 **Zhou C**, Liu J, Li Y, Liu L, Zhang X, Ma CY, Hua SC, Yang M, Yuan Q. microRNA-1274a, a modulator of sorafenib induced a disintegrin and metalloproteinase 9 (ADAM9) down-regulation in hepatocellular carcinoma. *FEBS Lett* 2011; **585**: 1828-1834 [PMID: 21530512 DOI: 10.1016/j.febslet.2011.04.040]
- 51 **Hsu CY**, Shen YC, Yu CW, Hsu C, Hu FC, Hsu CH, Chen BB, Wei SY, Cheng AL, Shih TT. Dynamic contrast-enhanced magnetic resonance imaging biomarkers predict survival and response in hepatocellular carcinoma patients treated with sorafenib and metronomic tegafur/uracil. *J Hepatol* 2011; **55**: 858-865 [PMID: 21338641 DOI: 10.1016/j.jhep.2011.01.032]
- 52 **Lee JH**, Park JY, Kim DY, Ahn SH, Han KH, Seo HJ, Lee JD, Choi HJ. Prognostic value of 18F-FDG PET for hepatocellular carcinoma patients treated with sorafenib. *Liver Int* 2011; **31**: 1144-1149 [PMID: 21745288 DOI: 10.1111/j.1478-3231.2011.02541.x]
- 53 **Mulé S**, Thieffn G, Costentin C, Durot C, Rahmouni A, Luciani A, Hoeffel C. Advanced Hepatocellular Carcinoma: Pretreatment Contrast-enhanced CT Texture Parameters as Predictive Biomarkers of Survival in Patients Treated with Sorafenib. *Radiology* 2018; **288**: 445-455 [PMID: 29584597 DOI: 10.1148/radiol.2018171320]
- 54 **Su WW**, Lee KT, Yeh YT, Soon MS, Wang CL, Yu ML, Wang SN. Association of circulating insulin-like growth factor 1 with hepatocellular carcinoma: one cross-sectional correlation study. *J Clin Lab Anal* 2010; **24**: 195-200 [PMID: 20486202 DOI: 10.1002/jcla.20320]
- 55 **Shao YY**, Huang CC, Lin SD, Hsu CH, Cheng AL. Serum insulin-like growth factor-1 levels predict outcomes of patients with advanced hepatocellular carcinoma receiving antiangiogenic therapy. *Clin Cancer Res* 2012; **18**: 3992-3997 [PMID: 22623732 DOI: 10.1158/1078-0432.CCR-11-2853]
- 56 **Kim HY**, Lee DH, Lee JH, Cho YY, Cho EJ, Yu SJ, Kim YJ, Yoon JH. Novel biomarker-based model for the prediction of sorafenib response and overall survival in advanced hepatocellular carcinoma: a prospective cohort study. *BMC Cancer* 2018; **18**: 307 [PMID: 29558905 DOI: 10.1186/s12885-018-4211-2]
- 57 **Miyahara K**, Nouse K, Tomoda T, Kobayashi S, Hagihara H, Kuwaki K, Toshimori J, Onishi H, Ikeda F, Miyake Y, Nakamura S, Shiraha H, Takaki A, Yamamoto K. Predicting the treatment effect of sorafenib using serum angiogenesis markers in patients with hepatocellular carcinoma. *J Gastroenterol Hepatol* 2011; **26**: 1604-1611 [PMID: 22011296 DOI: 10.1111/j.1440-1746.2011.06887.x]
- 58 **Tsuchiya K**, Asahina Y, Matsuda S, Muraoka M, Nakata T, Suzuki Y, Tamaki N, Yasui Y, Suzuki S, Hosokawa T, Nishimura T, Ueda K, Kuzuya T, Nakanishi H, Itakura J, Takahashi Y, Kurosaki M, Enomoto N, Izumi N. Changes in plasma vascular endothelial growth factor at 8 weeks after sorafenib administration as predictors of survival for advanced hepatocellular carcinoma. *Cancer* 2014; **120**: 229-237 [PMID: 24122122 DOI: 10.1002/cncr.28384]
- 59 **Hu B**, Cheng SY. Angiopoietin-2: development of inhibitors for cancer therapy. *Curr Oncol Rep* 2009; **11**: 111-116 [PMID: 19216842]
- 60 **Hayashi T**, Yamashita T, Terashima T, Suda T, Okada H, Asahina Y, Hayashi T, Hara Y, Nio K, Sunagozaka H, Takatori H, Arai K, Sakai Y, Yamashita T, Mizukoshi E, Honda M, Kaneko S. Serum cytokine profiles predict survival benefits in patients with advanced hepatocellular carcinoma treated with sorafenib: a retrospective cohort study. *BMC Cancer* 2017; **17**: 870 [PMID: 29258450 DOI: 10.1186/s12885-017-3889-x]
- 61 **Villanueva A**, Minguez B, Forner A, Reig M, Llovet JM. Hepatocellular carcinoma: novel molecular approaches for diagnosis, prognosis, and therapy. *Annu Rev Med* 2010; **61**: 317-328 [PMID: 20059340 DOI: 10.1146/annurev.med.080608.100623]
- 62 **Sakai K**, Takeda H, Nishijima N, Orito E, Joko K, Uchida Y, Izumi N, Nishio K, Osaki Y. Targeted DNA and RNA sequencing of fine-needle biopsy FFPE specimens in patients with unresectable hepatocellular carcinoma treated with sorafenib. *Oncotarget* 2015; **6**: 21636-21644 [PMID: 26046304 DOI: 10.18632/oncotarget.4270]
- 63 **Kan Z**, Zheng H, Liu X, Li S, Barber TD, Gong Z, Gao H, Hao K, Willard MD, Xu J, Hauptschein R, Rejto PA, Fernandez J, Wang G, Zhang Q, Wang B, Chen R, Wang J, Lee NP, Zhou W, Lin Z, Peng Z, Yi K, Chen S, Li L, Fan X, Yang J, Ye R, Ju J, Wang K, Estrella H, Deng S, Wei P, Qiu M, Wulur IH, Liu J, Ehsani ME, Zhang

- C, Loboda A, Sung WK, Aggarwal A, Poon RT, Fan ST, Wang J, Hardwick J, Reinhard C, Dai H, Li Y, Luk JM, Mao M. Whole-genome sequencing identifies recurrent mutations in hepatocellular carcinoma. *Genome Res* 2013; **23**: 1422-1433 [PMID: 23788652 DOI: 10.1101/gr.154492.113]
- 64 **Cleary SP**, Jeck WR, Zhao X, Chen K, Selitsky SR, Savich GL, Tan TX, Wu MC, Getz G, Lawrence MS, Parker JS, Li J, Powers S, Kim H, Fischer S, Guindi M, Ghanekar A, Chiang DY. Identification of driver genes in hepatocellular carcinoma by exome sequencing. *Hepatology* 2013; **58**: 1693-1702 [PMID: 23728943 DOI: 10.1002/hep.26540]
- 65 **Scartozzi M**, Faloppi L, Svegliati Baroni G, Loretelli C, Piscaglia F, Iavarone M, Toniutto P, Fava G, De Minicis S, Mandolesi A, Bianconi M, Giampieri R, Granito A, Facchetti F, Bitetto D, Marinelli S, Venerandi L, Vavassori S, Gemino S, D'Errico A, Colombo M, Bolondi L, Bearzi I, Benedetti A, Cascinu S. VEGF and VEGFR genotyping in the prediction of clinical outcome for HCC patients receiving sorafenib: the ALICE-1 study. *Int J Cancer* 2014; **135**: 1247-1256 [PMID: 24510746 DOI: 10.1002/ijc.28772]
- 66 **Casadei Gardini A**, Marisi G, Faloppi L, Scarpi E, Foschi FG, Iavarone M, Lauletta G, Corbelli J, Valgiusti M, Facchetti F, Della Corte C, Neri LM, Tamberi S, Cascinu S, Scartozzi M, Amadori D, Nanni O, Tenti E, Ulivi P, Frassinetti GL. eNOS polymorphisms and clinical outcome in advanced HCC patients receiving sorafenib: final results of the ePHAS study. *Oncotarget* 2016; **7**: 27988-27999 [PMID: 27058899 DOI: 10.18632/oncotarget.8569]
- 67 **Marisi G**, Scartozzi M, Faloppi L, Foschi GF, Lauletta G, Valgiusti M, Aprile G, Fornaro L, Iavarone M, Ercolani G, Frassinetti GL, Amadori D, Casadei Gardini A. Ang-2 polymorphisms in relation to outcome in advanced HCC patients receiving sorafenib. *J Clin Oncol* 2017; **35** Suppl 15: e15666 [DOI: 10.1200/JCO.2017.35.15\_suppl.e15666]
- 68 **Faloppi L**, Casadei Gardini A, Masi G, Silvestris N, Loretelli C, Ulivi P, Vivaldi C, Bianconi M, Giampieri R, Bittoni A, Andrikou K, Del Prete M, Scartozzi M, Cascinu S. Angiogenesis polymorphisms profile in the prediction of clinical outcome of advanced HCC patients receiving sorafenib: Combined analysis of VEGF and HIF-1 $\alpha$ . Final results of the ALICE-2 study. *J Clin Oncol* 2016; **34** Suppl 4: 280 [DOI: 10.1200/jco.2016.34.4\_suppl.280]
- 69 **Casadei Gardini A**, Faloppi L, Aprile G, Brunetti O, Caparello C, Corbelli J, Chessa L, Bruno D, Ercolani G, Leonetti A, de Stefano G, Farella N, Foschi FG, Lanzi A, Dadduzio V, Marisi G, Masi G, Negri FV, Pagan F, Santini D, Scarpi E, Silletta M, Silvestris N, Tamburini E, Tassinari D, Vivaldi C, Gentilucci UV, Zagonel V, Calvetti L, Cascinu S, Frassinetti GL, Scartozzi M. Multicenter Prospective Study of Angiogenesis Polymorphism Validation in HCC Patients Treated with Sorafenib. An INNOVATE Study Protocol. *Tumori* 2017; : tj5000704 Epub ahead of print [PMID: 29739298 DOI: 10.5301/tj.5000704]
- 70 **Horwitz E**, Stein I, Andreozzi M, Nemeth J, Shoham A, Pappo O, Schweitzer N, Tornillo L, Kanarek N, Quagliata L, Zreik F, Porat RM, Finkelstein R, Reuter H, Koschny R, Ganten T, Mogler C, Shibolet O, Hess J, Breuhahn K, Grunewald M, Schirmacher P, Vogel A, Terracciano L, Angel P, Ben-Neriah Y, Pikarsky E. Human and mouse VEGFA-amplified hepatocellular carcinomas are highly sensitive to sorafenib treatment. *Cancer Discov* 2014; **4**: 730-743 [PMID: 24687604 DOI: 10.1158/2159-8290.CD-13-0782]
- 71 **Arao T**, Ueshima K, Matsumoto K, Nagai T, Kimura H, Hagiwara S, Sakurai T, Haji S, Kanazawa A, Hidaka H, Iso Y, Kubota K, Shimada M, Utsunomiya T, Hirooka M, Hiasa Y, Toyoki Y, Hakamada K, Yasui K, Kumada T, Toyoda H, Sato S, Hisai H, Kuzuya T, Tsuchiya K, Izumi N, Arii S, Nishio K, Kudo M. FGF3/FGF4 amplification and multiple lung metastases in responders to sorafenib in hepatocellular carcinoma. *Hepatology* 2013; **57**: 1407-1415 [PMID: 22890726 DOI: 10.1002/hep.25956]
- 72 **He C**, Dong X, Zhai B, Jiang X, Dong D, Li B, Jiang H, Xu S, Sun X. MiR-21 mediates sorafenib resistance of hepatocellular carcinoma cells by inhibiting autophagy via the PTEN/Akt pathway. *Oncotarget* 2015; **6**: 28867-28881 [PMID: 26311740 DOI: 10.18632/oncotarget.4814]
- 73 **Xu Y**, Huang J, Ma L, Shan J, Shen J, Yang Z, Liu L, Luo Y, Yao C, Qian C. MicroRNA-122 confers sorafenib resistance to hepatocellular carcinoma cells by targeting IGF-1R to regulate RAS/RAF/ERK signaling pathways. *Cancer Lett* 2016; **371**: 171-181 [PMID: 26655273 DOI: 10.1016/j.canlet.2015.11.034]
- 74 **Vaira V**, Roncalli M, Carnaghi C, Favarsani A, Maggioni M, Augello C, Rimassa L, Pressiani T, Spagnuolo G, Di Tommaso L, Fagioli S, Rota Caremoli E, Barberis M, Labianca R, Santoro A, Bosari S. MicroRNA-425-3p predicts response to sorafenib therapy in patients with hepatocellular carcinoma. *Liver Int* 2015; **35**: 1077-1086 [PMID: 25040368 DOI: 10.1111/liv.12636]
- 75 **Gyöngyösi B**, Végh É, Járny B, Székely E, Fassan M, Bodoky G, Schaff Z, Kiss A. Pretreatment MicroRNA Level and Outcome in Sorafenib-treated Hepatocellular Carcinoma. *J Histochem Cytochem* 2014; **62**: 547-555 [PMID: 24804874 DOI: 10.1369/0022155414537277]
- 76 **Fornari F**, Pollutri D, Patrizi C, La Bella T, Marinelli S, Casadei Gardini A, Marisi G, Baron Toaldo M, Baglioni M, Salvatore V, Callegari E, Baldassarre M, Galassi M, Giovannini C, Cescon M, Ravaioli M, Negrini M, Bolondi L, Gramantieri L. In Hepatocellular Carcinoma miR-221 Modulates Sorafenib Resistance through Inhibition of Caspase-3-Mediated Apoptosis. *Clin Cancer Res* 2017; **23**: 3953-3965 [PMID: 28096271 DOI: 10.1158/1078-0432.CCR-16-1464]
- 77 **Nishida N**, Arizumi T, Hagiwara S, Ida H, Sakurai T, Kudo M. MicroRNAs for the Prediction of Early Response to Sorafenib Treatment in Human Hepatocellular Carcinoma. *Liver Cancer* 2017; **6**: 113-125 [PMID: 28275578 DOI: 10.1159/000449475]
- 78 **Stiuso P**, Potenza N, Lombardi A, Ferrandino I, Monaco A, Zappavigna S, Vanacore D, Mosca N, Castiello F, Porto S, Addeo R, Prete SD, De Vita F, Russo A, Caraglia M. MicroRNA-423-5p Promotes Autophagy in Cancer Cells and Is Increased in Serum From Hepatocarcinoma Patients Treated With Sorafenib. *Mol Ther Nucleic Acids* 2015; **4**: e233 [PMID: 25782064 DOI: 10.1038/mtna.2015.8]
- 79 **Yoon EL**, Yeon JE, Ko E, Lee HJ, Je JH, Yoo YJ, Kang SH, Suh SJ, Kim JH, Seo YS, Yim HJ, Byun KS. An Explorative Analysis for the Role of Serum miR-10b-3p Levels in Predicting Response to Sorafenib in Patients with Advanced Hepatocellular Carcinoma. *J Korean Med Sci* 2017; **32**: 212-220 [PMID: 28049231 DOI: 10.3346/jkms.2017.32.2.212]
- 80 **Carvajal-Hausdorf DE**, Schalper KA, Neumeister VM, Rimm DL. Quantitative measurement of cancer tissue biomarkers in the lab and in the clinic. *Lab Invest* 2015; **95**: 385-396 [PMID: 25502176 DOI: 10.1038/labinvest.2014.157]
- 81 **Abou-Alfa GK**, Schwartz L, Ricci S, Amadori D, Santoro A, Figuer A, De Greve J, Douillard JY, Lathia C, Schwartz B, Taylor I, Moscovici M, Saltz LB. Phase II study of sorafenib in patients with advanced hepatocellular carcinoma. *J Clin Oncol* 2006; **24**: 4293-4300 [PMID: 16908937 DOI: 10.1200/JCO.2005.01.3441]
- 82 **Personeni N**, Rimassa L, Pressiani T, Destro A, Ligorio C, Tronconi MC, Bozzarelli S, Carnaghi C, Di Tommaso L, Giordano L, Roncalli M, Santoro A. Molecular determinants of outcome in sorafenib-treated patients with hepatocellular carcinoma. *J Cancer Res Clin Oncol* 2013; **139**: 1179-1187 [PMID: 23568548 DOI: 10.1007/s00432-013-1429-x]
- 83 **Chen D**, Zhao P, Li SQ, Xiao WK, Yin XY, Peng BG, Liang LJ. Prognostic impact of pERK in advanced hepatocellular carcinoma patients treated with sorafenib. *Eur J Surg Oncol* 2013; **39**: 974-980 [PMID: 23845703 DOI: 10.1016/j.ejso.2013.06.018]
- 84 **Chu JS**, Ge FJ, Zhang B, Wang Y, Silvestris N, Liu LJ, Zhao CH, Lin L, Brunetti AE, Fu YL, Wang J, Paradiso A, Xu JM. Expression and prognostic value of VEGFR-2, PDGFR- $\beta$ , and c-Met in advanced hepatocellular carcinoma. *J Exp Clin Cancer Res* 2013; **32**: 16 [PMID: 23552472 DOI: 10.1186/1756-9966-32-16]
- 85 **Hagiwara S**, Kudo M, Nagai T, Inoue T, Ueshima K, Nishida N, Watanabe T, Sakurai T. Activation of JNK and high expression level of CD133 predict a poor response to sorafenib

in hepatocellular carcinoma. *Br J Cancer* 2012; **106**: 1997-2003 [PMID: 22596232 DOI: 10.1038/bjc.2012.145]

- 86 **Kudo M**, Finn RS, Qin S, Han KH, Ikeda K, Piscaglia F, Baron A, Park JW, Han G, Jassem J, Blanc JF, Vogel A, Komov D, Evans TRJ, Lopez C, Dutcus C, Guo M, Saito K, Kraljevic S,

Tamai T, Ren M, Cheng AL. Lenvatinib versus sorafenib in first-line treatment of patients with unresectable hepatocellular carcinoma: a randomised phase 3 non-inferiority trial. *Lancet* 2018; **391**: 1163-1173 [PMID: 29433850 DOI: 10.1016/S0140-6736(18)30207-1]

**P- Reviewer:** Lin ZY, Sun XY, Yao DF **S- Editor:** Gong ZM  
**L- Editor:** A **E- Editor:** Huang Y





Published by **Baishideng Publishing Group Inc**  
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA  
Telephone: +1-925-223-8242  
Fax: +1-925-223-8243  
E-mail: [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)  
Help Desk: <http://www.f6publishing.com/helpdesk>  
<http://www.wjgnet.com>



ISSN 1007-9327

