



University of Dundee

Nitrogen Fertilisers Shape the Composition and Predicted Functions of the Microbiota of Field-Grown Tomato Plants

Caradonia, Federica; Ronga, Domenico ; Catellani, Marcello ; Azevedo, Cleber Vinícius Giaretta ; Alegria Terrazas, Rodrigo; Robertson-Albertyn, Senga; Francia, Enrico; Bulgarelli, Davide

DOI:

[10.1094/PBIOMES-06-19-0028-R](https://doi.org/10.1094/PBIOMES-06-19-0028-R)

Publication date:

2019

Document Version

Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Caradonia, F., Ronga, D., Catellani, M., Azevedo, C. V. G., Alegria Terrazas, R., Robertson-Albertyn, S., ... Bulgarelli, D. (2019). Nitrogen Fertilisers Shape the Composition and Predicted Functions of the Microbiota of Field-Grown Tomato Plants. *Phytobiomes Journal*. <https://doi.org/10.1094/PBIOMES-06-19-0028-R>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

1 **NITROGEN FERTILISERS SHAPE THE COMPOSITION AND PREDICTED FUNCTIONS**
2 **OF THE MICROBIOTA OF FIELD-GROWN TOMATO PLANTS**

3 Federica Caradonia^{1,3*}, Domenico Ronga^{1,#}, Marcello Catellani^{1,‡}, Cleber Vinícius Giaretta
4 Azevedo^{1,2}, Rodrigo Alegria Terrazas³, Senga Robertson-Albertyn³, Enrico Francia^{1*} and
5 Davide Bulgarelli ^{3*}

6 ¹Department of Life Sciences, Centre BIOGEST-SITEIA, University of Modena and Reggio
7 Emilia, Italy; ²Faculty of Agricultural and Veterinary Sciences, São Paulo State University,
8 Jaboticabal, Brasil; ³Plant Sciences, School of Life Sciences, University of Dundee, Dundee,
9 United Kingdom.

10 Present addresses:

11 #CREA, Centre for Animal Production and Aquaculture (CREA-ZA), Lodi, Italy;

12 ‡ENEA, Department for Sustainability, C.R. Trisaia, Rotondella (MT), Italy.

13

14 * Correspondence:

15 Federica Caradonia

16 federica.caradonia@unimore.it

17 Enrico Francia

18 enrico.francia@unimore.it

19 Davide Bulgarelli

20 d.bulgarelli@dundee.ac.uk

21

22 ABSTRACT

23 The microbial communities thriving at the root-soil interface have the potential to improve
24 plant growth and sustainable crop production. Yet, how agricultural practices, such as the
25 application of either mineral or organic nitrogen fertilisers, impact on the composition and
26 functions of these communities remains to be fully elucidated. By deploying a two-pronged
27 16S rRNA gene sequencing and predictive metagenomics approach we demonstrated that
28 the bacterial microbiota of field-grown tomato (*Solanum lycopersicum*) plants is the product
29 of a selective process that progressively differentiates between rhizosphere and root
30 microhabitats. This process initiates as early as plants are in a nursery stage and it is then
31 more marked at late developmental stages, in particular at harvest. This selection acts on
32 both the bacterial relative abundances and phylogenetic assignments, with a bias for the
33 enrichment of members of the phylum Actinobacteria in the root compartment. Digestate-
34 based and mineral-based nitrogen fertilisers trigger a distinct bacterial enrichment in both
35 rhizosphere and root microhabitats. This compositional diversification mirrors a predicted
36 functional diversification of the root-inhabiting communities, manifested predominantly by
37 the differential enrichment of genes associated to ABC transporters and the two-component
38 system. Together, our data suggest that the microbiota thriving at the tomato root-soil
39 interface is modulated by and in responses to the type of nitrogen fertiliser applied to the
40 field.

41

42 Additional keywords: *Solanum lycopersicum*, Rhizosphere, Root, Microbiota, Nitrogen,
43 Fertilisers, Digestate.

44 INTRODUCTION

45 Limiting the negative impact of agricultural practices on the environment while preserving
46 sustainable crop yield is one of the key challenges facing agriculture in the years to come.

47 As an essential element for plant nutrition, nitrogen represents a paradigmatic example of
48 such a challenge. Moreover, due to the combined effect of elevated solubility and little
49 retention in soils, the lack of this element is and will be one of the major yield-limiting factors
50 worldwide (Tilman et al., 2011). At the same time, the application of synthetic nitrogen
51 fertilisers is, in many agricultural systems, a low efficiency approach which has been linked
52 with the degradation of natural resources (Elser and Bennett, 2011). One of the strategies
53 adopted to limit the economic and environmental footprint of crop production while
54 maintaining sustainable yield is the “recycling” of, mineral-rich, biodegradable products of
55 the livestock and agricultural sectors.

56 One example of this approach is the application of the digestate, a by-product of the
57 anaerobic digestion of organic waste for the production of biogas (Möller and Müller, 2012)
58 as renewable soil amendment for crop production. The digestate is a mixture of partially-
59 degraded organic matter, microbial biomass and inorganic compounds (Albuquerque et al.,
60 2012). We recently demonstrated how the digestate can be efficiently used as innovative
61 fertiliser and plant growing media (Ronga et al., 2018b; Ronga et al., 2018a; Ronga et al.,
62 2019), yet the impact of digestate applications on the agroecosystem remains to be fully
63 elucidated.

64 For instance, the digestate can be a source of phytoavailable nitrogen, in particular
65 ammonium, capable of impacting on organic matter mineralisation and emission of carbon
66 dioxide from the soil profile (Grigatti et al., 2011). Therefore, it is legitimate to hypothesize
67 that such treatments impact on the composition of the microbial communities thriving at the
68 root-soil interface, collectively referred to as the rhizosphere and root microbiota, which play

69 a critical role in mobilisation of mineral elements for plant uptake (Alegria Terrazas et al.,
70 2016). Congruently, several studies indicate that the application of biogas by-product
71 enhances soil microbial activity (Möller, 2015) and the availability of phytohormones (Scaglia
72 et al., 2015). However, the intertwined relationship among microbiota composition, soil
73 characteristics and amendments as well as host plant species-specificity (Bulgarelli et al.,
74 2013) makes it difficult to infer first principles.

75 This is particularly true for field-grown crops such as tomato (*Solanum lycopersicum L.*), one
76 of the most cultivated horticultural crops globally with plantations occupying an area of 4.8
77 million of hectares with a production of 182 million tonnes in 2017 ([FAO 2017](#)). Notably, this
78 species is also an excellent experimental model for basic science investigations: tomato was
79 one of the first crops whose genome was sequenced (Consortium, 2012) and provided a
80 superb platform to test the significance of genome editing for evolutionary studies and plant
81 breeding (Zsögön et al., 2018). Perhaps not surprisingly, tomato is gaining momentum as
82 an experimental system to study host-microbiota interactions in crop plants. Recent
83 investigations revealed novel insights into the assembly cues of the microbiota associated
84 to this plant (Bergna et al., 2018; Toju et al., 2019) and the contribution of microbes thriving
85 at the tomato root-soil interface to pathogen protection (Chialva et al., 2018; Kwak et al.,
86 2018). However, the composition and functional potential of the tomato microbiota and their
87 interdependency from nitrogen fertilisers remain to be elucidated.

88 Here we report the metagenomics characterisation of the microbiota thriving at the root-soil
89 interface of field-grown tomato plants. We hypothesize that nitrogen treatments shape and
90 modulate the contribution of the tomato microbiota for crop yield. To test this hypothesis, we
91 focused on processing tomato exposed to different nitrogen fertilisers, either digestate-
92 based or containing a mineral fraction. By using a 16S rRNA amplicon sequencing survey
93 we deciphered how the microhabitat (i.e., either rhizosphere or root) sculpts the tomato

94 microbiota which, in turn, is fine-tuned by the type of fertiliser applied. Finally, by using a
95 predictive metagenomics survey, we inferred the functional diversification imposed by the
96 nature of the fertilisers on the root microbiota.

97

98

99 MATERIALS AND METHODS

100 Field site

101 A field trial was established in a tomato farm near the city of Ravenna (44°25'40.8"N
102 12°05'53.3"E), Emilia Romagna Region, Italy, during the 2017 growing season. During the
103 period from transplant to harvest, the minimum and the maximum average temperatures
104 recorded were 17.1°C and 32.8°C, respectively, and the rainfall was 101.7 mm. The soil had
105 a silty loam texture (14% clay, 51% silt, 35% sand), a pH 8.3 (in H₂O), 1.1 g kg⁻¹ total N
106 (Kjeldahl method), 7 mg kg⁻¹ available P (Olsen method), 129 mg kg⁻¹ exchangeable K
107 (Ammonium acetate), and 9 g kg⁻¹ organic matter (Walkey-Black method). A schematic
108 illustration of the field trial is depicted in Supplementary Figure S1.

109 Plant material

110 We used the tomato cultivar 'Fokker', a processing-type genotype with blocky fruit, late fruit
111 ripening and suitable for tomato puree, for the experimentation. Seedlings were provided by
112 Bronte Soc. Coop. Agr. A.R.L. (Mira, Italy). Processing tomato seedlings were transplanted
113 at the end of May when they were 6-week old corresponding to plants at the fourth true leaf
114 stage. Plant density was 3 plants m⁻². Plants were transplanted into single row, with a
115 spacing of 0.22 m between plants in each row and 1.50 m between rows.

116 Experimental design

117 We established a randomized complete design with three replicates and seven treatments:
118 pelleted digestate (hereafter PE), liquid digestate (LD), slow-acting liquid digestate (SRLD),
119 organo-mineral fertiliser based on digestate (SC), synthetic fertiliser (MF), slow-acting
120 synthetic fertiliser (SRMF), and no fertilization treatment (NT). The composition of the
121 treatments is summarised in Table 1. For each treatment, we applied a total amount of
122 nitrogen in the ratio 150 N kg ha⁻¹ on the basis of soil analysis, crop rotation and crop
123 nutrients required. Nitrogen was supplied at transplanting time with the exception of the
124 synthetic fertiliser treatment. For this latter treatment, the amount of total Nitrogen was

125 equally divided and applied in 3 times (transplanting, full flowering and fruit ripening) using
126 ammonium nitrate in the first treatment and calcium nitrate in the second and in the third
127 ones. During the trial, 600 m³ ha⁻¹ of irrigation water was distributed by drip irrigation to each
128 treatment. The other soil and crop management practices were performed according to the
129 production rules of Emilia Romagna Region, Italy. Briefly, weeds control was performed with
130 a single treatment (on 11th June) using products based on metribuzin and propaquizafop.
131 Sulphur and Copper were used to control phytopatogenic fungi while imidacloprid,
132 abamectin and spinosad were used as insecticide.

133 **Yield traits**

134 At harvest we determined the marketable yield (t ha⁻¹), as a weight of fully ripe fruits, and
135 the solid soluble content (°Brix t ha⁻¹) as a proxy for fruit quality. The °Brix parameter was
136 determined using the digital refractometer HI 96814 (Hanna, Italy), while the °Brix t ha⁻¹ was
137 calculated by multiplying the hectare marketable yield by °Brix and dividing the result by
138 100.

139 **Root, Rhizosphere and Bulk soil Sampling and DNA Extraction**

140 At transplanting time (May 2017), 5 root specimens per treatment were collected. Upon
141 uprooting, soil particles loosely bound to roots were dislodged by hand shaking and root
142 segments of ~ 6 cm were placed in sterile 50 mL tubes. The samples were stored in a
143 portable cooler (~ 4°C), transported to the laboratory and immediately processed. Root
144 specimens were incubated in 30 mL of PBS (Phosphate buffered saline) and placed on a
145 shaker for 20 minutes in order to separate the soil tightly adhering to plant material, which
146 we operationally defined as “rhizosphere”, from the roots. The first tubes were centrifuged
147 for 20 minutes at 4,000 x g and the rhizosphere pelleted was collected in liquid nitrogen and
148 stored at -80°C. The roots were moved to a new sterile tube containing 30 mL PBS and
149 sonicated by Ultrasonics Sonomatic Cleaner (Langford Ultrasonics, Birmingham, UK) for 10
150 minutes (intervals of 30 seconds pulse and 30 seconds pause) at 150 W, as previously

151 reported (Schlaeppli et al., 2014) to enrich for endophytic microorganisms. Roots were then
152 washed in the same new buffer and dried on sterile filter paper. After few minutes, the roots
153 were moved to 50 mL tubes and frozen in liquid nitrogen for storage at -80°C. Three
154 independent soil samples were harvested from unplanted soil in different points of the field,
155 frozen in liquid nitrogen and stored at -80°C. At harvest time (September 2017) the whole
156 plants were harvested, 5 roots per treatment and 3 bulk soil samples were collected,
157 prepared and stored like the previous samples. Frozen root samples were pulverized in a
158 sterile mortar using liquid nitrogen prior DNA preparation. DNA was extracted from all the
159 specimens (i.e., bulk soil, rhizosphere and pulverized roots) using the FastDNA® SPIN Kit
160 for Soil (MP Biomedicals, Solon, USA) following the instruction manual provided by
161 manufacturer. DNA samples were diluted using 50 µL DES water and quantified using the
162 Nanodrop 1000 Spectrophotometer (Thermo Scientific, Wilmington, United States).

163 16S rRNA Gene Sequencing

164 The sequencing library was generated using primers specific (515F 5'-
165 GTGCCAGCMGCCGCGGTAA-3' and 806R 5'-GGACTACHVGGGTWTCTAAT-3') for
166 hypervariable V4 region of the 16S rRNA gene. The reverse primers included a 12-mer
167 unique "barcode" sequences (Caporaso et al., 2012) to facilitate the multiplexing of the
168 samples into a unique sequencing run. Individual PCR reactions were performed as
169 previously reported (Robertson-Albertyn et al., 2017), with the exception of the concentration
170 of the Bovine Serum Albumin, added at 10 µg/reaction, and the addition of a Peptide Nucleic
171 Acid (PNA) blocker (PNA Bio, Newbury Park, United States) at a concentration of 0.5
172 µM/reaction to inhibit plastidial amplification. For each barcoded primers, three technical
173 replicates and a no-template control (NTC) were organised and processed starting from a
174 unique master mix. Five microliters of amplified samples and cognate NTCs were inspected
175 on a 1% (w/v) agarose gel. Two independent sets of triplicated amplicons, displaying the
176 expected amplicon size and lacking detectable contaminations, were combined in a

177 barcode-wise manner and purified using the Agencourt AMPure XP kit (Beckman Coulter,
178 Brea, United States) with a ratio of 0.7 mL AMPure XP beads per 1 mL of sample. Purified
179 DNA samples were quantified using Picogreen (Thermo Fisher, United Kingdom) and
180 combined in an equimolar ratio into an amplicon pool. This latter material was used for the
181 preparation of a MiSeq run at the Genome Technology facilities of the James Hutton Institute
182 (Invergowrie, UK) as previously reported (Robertson-Albertyn et al., 2017).

183

184 OTU Table Generation and pre-processing

185 We used QIIME, version 1.9.0 (Caporaso et al., 2010) to process the sequencing output of
186 the MiSeq machine. Briefly, the command `join_paired_ends.py` was used to decompress
187 and merged (minimum overlap 5bp) forward and reverse read FASTQ files. Next, we
188 removed *in silico* low-quality sequencing reads and sequencing reads without the barcode
189 information. Then, the reads were assigned to individual samples. In these analyses, the
190 command `split_libraries_fastq.py` was used imposing a minimum PHRED score of 20. The
191 resulted high-quality reads were assembled into an Operational Taxonomic Unit (OTU) table
192 at 97% sequence identity. We used a 'closed reference' approach for OTU-picking using the
193 command `pick_closed_reference_otus.py`. We imposed the Green Genes database version
194 13_5 (DeSantis et al., 2006) as a reference database to identify microbial OTUs and prune
195 for chimeric sequences. We used SortMeRNA algorithm for OTU -picking and taxonomy
196 assignment. Finally, OTUs whose representative sequences were classified as either
197 chloroplast or mitochondria, as well as OTUs accruing only one sequencing read over the
198 entire dataset (i.e., singletons), were depleted *in silico* using the function
199 `filter_otus_from_otu_table.py`.

200

201 [Data visualisation and statistical analyses](#)

202 Agronomic traits were analysed by Analysis of Variance (ANOVA) using GenStat 17th (VSN
203 International, Hemel Hempstead, UK). Means were compared using Bonferroni's test at the
204 5% level.

205 The OTU table produced in QIIME was analysed in R using a custom script developed from
206 Phyloseq package (McMurdie and Holmes, 2013).

207 Initially, the data were filtered removing the samples with less than 1,000 reads and the
208 OTUs with less than 10 reads in at least 5% of the samples. For alpha-diversity calculation,
209 sequencing reads were rarefied at an even sequencing deep of 18,467 reads per sample
210 retaining 2,439 unique OTUs. The number of Observed OTUs and Chao1 index were used
211 as richness estimators, while the Shannon index was used for evaluating the evenness.
212 Upon inspecting distribution of the data using a Shapiro-Wilk test, the means of rhizosphere
213 and root samples at harvest time were compared using a non-parametric Wilcoxon rank sum
214 test. Next, we performed a non-parametric Kruskal–Wallis test independently on
215 rhizosphere and root samples to identify significant effect of the individual treatments on the
216 ecological indices.

217 For beta-diversity calculation, the original counts (i.e., not rarefied) were transformed to
218 relative abundances and we imposed an abundance threshold to target PCR-reproducible
219 OTUs. The differences among microbial communities of the samples were computed using
220 Bray-Curtis index and weighted Unifrac index, with this latter index including phylogenetic
221 information in the analysis (Lozupone and Knight, 2005). A Principal Coordinates Analysis
222 (PCoA) was generated to visualize similarities and dissimilarities of microhabitats and
223 treatments. In order to assess the effects of microhabitats and the treatments on the
224 bacterial community composition, a Permutational Multivariate Analysis Of Variance
225 (PERMANOVA) on distance matrices was implemented using the function Adonis in p a

226 two-pronged approach. First, we assessed the effect of nursery/harvest stage on
227 microhabitat composition. Next, we used the same test to assess the impact of the treatment
228 on rhizosphere and root specimens at harvest stage. In the two approaches, the computed
229 R^2 therefore reflects the proportion of variance explained by the given factor in the group of
230 samples tested.

231 Finally, original counts were used to perform a differential analysis to identify individual
232 bacteria differentially enriched in the tested samples using DESeq2 (Love et al., 2014).

233 The phylogenetic tree was constructed using the representative sequences of the OTUs
234 significantly enriched in rhizosphere and root specimens and annotated with iTOL (Letunic
235 and Bork, 2006).

236 [Functional predictions](#)

237 Tax4Fun (Asshauer et al., 2015) package in R was used as a predictive tool to obtain a
238 functional profile based on 16S rRNA gene data. Metabolic capabilities are calculated by
239 linking the amplicon data phylogenetic and abundance profile to a set of pre-computed
240 metabolic reference profiles, based on the KEGG Ortholog (KO) database (Kanehisa et al.,
241 2008). The input for this analysis was an OTU table obtained with the representative
242 sequences of the OTU table previously generated (see above), reclassified using
243 SILVA_115 taxonomic database (Quast et al., 2013). Similar to a previously reported
244 operational protocol (Kavamura et al., 2018), we focused our analysis in prokaryotic
245 functional categories related to amino acid metabolism, carbohydrate metabolism, cell
246 motility, energy metabolism, membrane transport, metabolism of terpenoids and polyketides
247 and two-component system, trimming the rest of predicted functions from the Tax4fun
248 output. A statistical comparison between two groups using a Welch's t-test (Bluman, 2009)
249 filtered at a p-value < 0.01 with Storey's correction for false discovery rate (Storey and

250 Tibshirani, 2003) was performed in STAMP, Statistical Analysis of Metagenomic Profiles
251 (Parks et al., 2014).

252 [Data and scripts availability](#)

253 The 16S rRNA gene sequences presented in this study are available at the European
254 Nucleotide archive under the study accession number PRJEB32219. The scripts to
255 reproduce the statistical analysis and figures are available at [https://github.com/BulgarelliD-
256 Lab/Tomato_nitrogen](https://github.com/BulgarelliD-Lab/Tomato_nitrogen). Data frames required for scripts reproducibility are included in
257 Supplementary Database 1.

258 **RESULTS**

259 [Fertiliser treatment impacts on yield and quality of processing tomato](#)

260 At harvest time the two most important parameters such as marketable yield and fruit quality
261 were measured to evaluate the effect of 7 different fertiliser performances on processing
262 tomato (Figure 1). The fertiliser treatments had a significant effect on fresh biomass of fruits
263 (ANOVA, Bonferroni's test, $P < 0.001$). Pelleted digestate registered the best performance
264 followed by synthetic fertiliser and slow acting liquid digestate. In addition, the different
265 fertilisers influenced significantly also the quality of processing tomato (Figure 1) (ANOVA,
266 Bonferroni's test, $P < 0.001$).

267 [The assembly dynamics of the bacterial microbiota of field-grown processing tomato](#)

268 To gain insights into the relationships between yield traits and microbiota composition in
269 field-grown processing tomato plants, we generated 5,546,303 high quality 16S rRNA gene
270 sequences for the 86 samples generated in this study.

271 Upon *in silico* depletion of OTUs classified as Mitochondria and Chloroplast we reduced the
272 number of analysable sequences to 4,645,503 with a retaining proportion of 83.7% of the
273 original sequences (mean per samples = 54,017.48 reads; max = 111,213 reads; min = 272
274 reads). The data were further filtered removing the samples with less than 1,000 reads as

275 well as the OTUs with less than 10 reads in 5% of samples. This allowed us to retain 2,515
276 unique OTUs accounting for 4,308,580 high quality reads and 85 samples.

277 Then, we computed alpha-diversity calculations on a dataset rarefied at 18,467 reads per
278 sample and alpha-diversity was investigated considering two microhabitats (root and
279 rhizosphere) and the seven fertiliser treatments. OTUs richness was assessed by Chao1
280 index and Observed OTUs while the OTUs evenness was assessed by Shannon index. This
281 analysis revealed a significant effect of the microhabitat on the characteristics of the
282 microbiota thriving at the tomato root-soil interface: regardless of the treatment, the root
283 microhabitat emerged as less diverse and even compared to the rhizosphere one (Wilcoxon
284 rank sum, $p < 0.01$, Figure 2). This observation suggests that root microhabitat represents a
285 gated community compared to the surrounding soil environment. Conversely, the treatment
286 impacted only the number of OTUs observed in the rhizosphere compartment (Kruskal-
287 Wallis non parametric analysis of variance followed by Dunn's post-hoc test $p < 0.05$. Figure
288 2).

289 Congruently, beta-diversity analysis computed on the non-rarefied dataset using both
290 weighted Unifrac and Bray-Curtis indicated a microhabitat-dependent microbiota
291 diversification. In particular, the weighted Unifrac matrix visualised using a Principal
292 Coordinates Analysis revealed such a microhabitat effect on samples processed at harvest
293 time along the axis accounting for the major variation. Interestingly, younger nursery
294 samples displayed a similar degree of diversification, although their communities were
295 separated from the harvest samples on the axis accounting for the second source of
296 variation (Figure 3). These data were supported by a PERMANOVA which attributed a R^2
297 of 30% to the microhabitat, a R^2 of 28% to the 'Nursery/Harvest effect' and a R^2 of 2% to
298 their interactions (Adonis test, 5,000 permutations, $p < 0.01$). The analysis conducted on
299 rhizosphere and root samples at harvest stage revealed that, congruently with the observed

300 diversification along the axis accounting for the major variation, the microhabitat remained
301 the major driver of the tomato communities (R^2 47%, Adonis test, 5,000 permutations, p
302 <0.01) while the individual fertiliser treatments impacted these plant-associated microbial
303 assemblages to a lesser, but significant, extent (R^2 13%, Adonis test, 5,000 permutations, p
304 <0.01). This suggest that, rather than on richness per se, the fertiliser treatment impacts on
305 the abundances and phylogenetic assignments of members of the tomato microbiota.
306 Remarkably, the Bray-Curtis matrix produced a congruent results, although the temporal
307 effect (i.e., nursery vs. harvest time) explained slightly more variation ($\sim 29\%$;
308 Supplementary Figure S2) than microhabitat diversification manifested along the second
309 axis of variation ($\sim 26\%$; Supplementary Figure S2). Crucially, also in this case the observed
310 diversification was supported by a PERMANOVA which attributed a R^2 of 23% to the
311 microhabitat, a R^2 of 29% to the 'Nursery/Harvest effect' and a R^2 of 3% to their interactions
312 (Adonis test, 5,000 permutations, $p <0.01$).

313 [Differential bacterial enrichments define microhabitat and treatment “signatures” on the](#) 314 [field grown tomato microbiota](#)

315 To gain insights into individual members of the tomato microbiota responsible for the
316 observed diversification we implemented a series of pair-wise comparisons among
317 microhabitats and treatments at harvest stage. We took a two-pronged approach. First, we
318 identified bacteria underpinning the microhabitat effect i.e., the selective enrichment of
319 bacteria in the roots and the rhizosphere microhabitats amended with no fertiliser. Next, we
320 assessed the effect of the fertiliser treatment on roots and rhizosphere bacterial composition
321 by comparison with bacteria enriched in untreated samples.

322 This allowed us to identify 170 bacterial OTUs whose abundance was significantly enriched
323 in and differentiated between rhizosphere specimens and unplanted soil samples (Wald test,
324 $p <0.01$, FDR corrected; Supplementary database 1). Similarly, we identified 374 bacterial
325 OTUs whose abundance was significantly enriched in and differentiated between root

326 specimens and unplanted soil samples (Wald test, $p < 0.01$, FDR corrected; Supplementary
327 database 1). Of these differentially enriched bacteria, 96 OTUs represented a set of tomato-
328 competent OTUs capable of colonising both the rhizosphere and root environments. When
329 we then looked into the taxonomic affiliations of this tomato-competent microbiota, we
330 discovered that it is dominated by members of Actinobacteria, Bacteroidetes, Alpha-, Beta-
331 , Gamma- and Deltaproteobacteria as well as Verrucomicrobia (Figure 4). Strikingly, the
332 taxonomic investigation revealed a bias for Actinobacteria in the root compartment, possibly
333 reflecting an adaptive advantage of members of this phylum in colonising the endophytic
334 environment.

335 Interestingly, each fertiliser treatment had a distinct impact on these tomato-enriched
336 microbiota. The pelleted digestate (PE) and the slow-acting synthetic fertiliser (SRMF)
337 yielded the highest number of uniquely enriched OTUs regardless of the microhabitat
338 investigated, albeit with a distinct pattern: the SRMF had a more pronounced effect on the
339 rhizosphere communities while the PE impacted more on the bacteria thriving in association
340 with root tissues. (Wald test, $p < 0.01$, FDR corrected; Figure 5; Supplementary database 1).

341 Interestingly, when we inspected the taxonomic composition of the bacteria differentially
342 impacted by the fertiliser treatment we observed an increase of the number of OTUs
343 belonging to phylum of Actinobacteria. In particular, PE had 12 OTUs out of 80 and 14 OTUs
344 out of 105, in root and rhizosphere, respectively, belonging to phylum Actinobacteria. While,
345 MF had 15 OTUs out of 38 and 22 OTUs out of 49 in root and rhizosphere, respectively,
346 belonging to phylum Actinobacteria (Supplementary database S1). Within this phylum we
347 observed the presence of OTUs classified as *Streptomyces spp.*, *Agromyces sp.*,
348 *Microbispora sp.* and *Actinoplanes spp.*

349 Together these data suggested that the enrichment of specific bacteria underpins the
350 observed microhabitat effect whose magnitude is fine-tuned by the applied fertiliser.

351 Organic- and synthetic-based fertiliser trigger different metabolic capacities in the tomato 352 root microbiota

353 To investigate the ecological significance of the observed differential recruitments among
354 fertiliser treatment we employed a predictive metagenomics approach. Briefly, we inferred
355 *in silico* the functions encoded by the tomato microbiota at harvest stage (Materials and
356 Methods) and we grouped the samples in digestate-based (i.e., PE, LD and SRLD; hereafter
357 'organic') and treatments containing at least a synthetic component (i.e., SC, MF and SRMF;
358 hereafter 'mineral'). We observed that the functions putatively encoded by the communities
359 exposed to either organic or mineral fertilisers can discriminate between treatments in both
360 microhabitats (PERMANOVA: Rhizosphere samples $R^2 = 14\%$, p value <0.01 , 5,000
361 permutations; Root samples $R^2 = 16\%$, p value <0.01 , 5,000 permutations). Congruently,
362 we identified a set of 14 functions differentially enriched between root communities exposed
363 to either group of treatments (Welch t-test, p <0.01 , FDR corrected; Figure 6). Interestingly,
364 we observed a striking dichotomy between the two groups of treatments: communities
365 exposed to mineral fertilisers are predicted to enrich for genes implicated in the ABC
366 transporter machinery while bacteria exposed to the organic treatments are predicted to
367 enrich for genes implicated in the two-component system. These two set of genes are
368 dominant in communities exposed to both treatments and are also associated to additional
369 distinct enrichment patterns, most notably including nitrogen metabolism (organic
370 communities) and tetracycline biosynthesis (mineral communities).

371 These results suggest that, within tomato roots, the observed taxonomic diversification
372 underpins a functional specialisation of the microbiota which, in turn, may impact on plant
373 growth development and health.

374 DISCUSSION

375 This study revealed that all nitrogen treatments led to an increase of tomato production in
376 comparison with the no fertilization treatment (fold change between 0.8 and 1.73) confirming

377 that, in the tested conditions, nitrogen limits the yield potential of processing tomato crops
378 as observed in previous studies (Ronga et al., 2015; Ronga et al., 2017). Yet, despite the
379 same amount of nitrogen was applied in each treatment (i.e., 150 kg ha⁻¹), all the treatments
380 were statistically different from each other. A prediction of this observation is that, under the
381 tested conditions, the nature of the fertilisers, rather than the amount of nitrogen per se,
382 affect the yield and the fruit quality of tomato plants. These observations and the putative
383 contribution to fertiliser use efficiency of the microbial communities thriving at the root-soil
384 interface (Alegria Terrazas et al., 2016), motivated us to investigate relationships between
385 yield traits and the composition of the tomato rhizosphere and root microbiota under field
386 conditions.

387 [The tomato rhizosphere and root microbiota are gated communities](#)

388 First, we characterised the rhizosphere and root microbiota of processing tomato with no
389 treatment. Both alpha and beta diversity discriminated between the communities of
390 seedlings and adult plants. Despite these differences, which could be attributed to both
391 abiotic, e.g., time of residence in soil (Dombrowski et al., 2017), and biotic factors, e.g.,
392 developmental-conditioned rhizodeposits (Chaparro et al., 2014), it is striking to note how
393 tomato plants displayed a rhizosphere and root compartmentalisation regardless of the
394 developmental stage. This is congruent with the observation that in rice, the assembly and
395 structural diversification of the microbiota is a rapid process which reaches a steady-state
396 level within a few weeks from germination (Edwards et al., 2015). Closer inspection of the
397 rhizosphere and root profiles at harvest stage indicates that these plant-associated
398 communities are phylogenetically related to those of unplanted soil, suggesting that the
399 initial tomato microbiota is further modulated by the growing conditions.

400 Despite this apparent relatedness, the selective enrichment of individual bacterial members
401 of the microbiota discriminates between rhizosphere and root communities for mature plants
402 from unplanted soil profiles (Figure 4). These enrichments displayed a bias for members of

403 the phyla Actinobacteria, Bacteroidetes, Proteobacteria (including the classes Alpha-, Beta-
404 , Delta- and Gammaproteobacteria) as well as Verrucomicrobia. Members of these taxa
405 have routinely been reported in studies focussing on plant-competent bacteria under both
406 laboratory and field conditions (Bulgarelli et al., 2013; Walters et al., 2018), suggesting that
407 the experimental approach followed in this study can be considered representative for field-
408 grown processing tomato.

409 However, we noticed a differential selective pressure on the bacteria thriving either in the
410 rhizosphere or in the root tissue: this latter environment produced more distinct profiles, i.e.
411 more differentially enriched bacteria compared to unplanted soil, than the ones retrieved
412 from the soil surrounding the roots. This indicates that the diversification of the tomato-
413 inhabiting microbial communities from the surrounding soil biota initiates in the rhizosphere
414 and progresses through the root tissue, where it produces a more pronounced microbiota
415 diversification compared to unplanted specimens. This observation is reminiscent of the
416 recruitment patterns of other crops such as barley (Bulgarelli et al., 2015) but it is in striking
417 contrast with studies conducted with both model (Bulgarelli et al., 2012) and field-grown
418 (Rathore et al., 2017) Brassicaceae, whose 'rhizosphere effect' appears negligible.

419 We further noticed that the "root effect" on the microbiota was exerted also at phylogenetic
420 level with a bias for the enrichment Actinobacteria. This observation is in apparent contrast
421 with results gathered from the recent seed-to-seed characterisation of the tomato microbiota
422 which revealed that, albeit averaging 8% of the sequencing reads across microhabitats,
423 members of this phylum did not significantly discriminate root from rhizosphere specimens
424 (Bergna et al., 2018). However, it is worth mentioning that these two studies differed in terms
425 of both soil type and plant genotype used.

426 Together, our results suggest that both species- and soil-specific traits govern the assembly
427 of the tomato microbiota in field-grown crops.

428 [Nitrogen source impacts on the structural and functional composition of the tomato](#)
429 [microbiota](#)

430 Next, we investigated the impact of the type of nitrogen fertiliser on the tomato microbiota
431 and we demonstrated that each treatment produced “distinct signatures”, represented by
432 specific selective enrichment, on both the rhizosphere and root communities. Despite
433 microhabitat-associated variation, the effect of the application of pelleted digestate (PE)
434 resulted in the most distinct microbial profile in the root compartment and the second largest
435 number of specifically enriched OTUs in the rhizosphere. Of note, the slow-acting mineral
436 fertiliser (SRMF) follow a “complementary” pattern: its application yielded the greatest and
437 the second greatest number of differentially enriched OTUs compared to untreated samples
438 in the rhizosphere and root profiles, respectively. Remarkably, these two treatments had a
439 discernible effect also on crop yield, with the PE treatment producing the best performance
440 among the various fertilisers. Our data are congruent with studies conducted on wheat which
441 observed a structural diversification of the soil and plant-associated communities exposed
442 to either mineral or organic fertilisers (Kavamura et al., 2018). Yet, the numerical shift in
443 terms of OTUs differentially enriched per se cannot explain the potential impact of these
444 communities on crop yield: owing to the fact that the SMRF treatment, which is associated
445 to a significant reduction in yield traits (compared to PE) is capable of triggering a
446 comparable OTU enrichment.

447 We therefore focused our attention on the taxonomical composition of the rhizosphere and
448 root communities. In particular, we noticed that the proliferation of Actinobacteria in the root
449 compartment was retained in the various treatments. The enriched Actinobacteria included
450 *Streptomyces spp.*, *Agromyces sp.*, *Microbispora sp.* and *Actinoplanes spp.*
451 (Supplementary Database 1). *Streptomyces spp.* are well-known bacteria able to produce a
452 wide diversity of bioactive compounds able to promote plant growth and health (de Jesus
453 Sousa and Olivares, 2016). On the other hand, members of the genus *Streptomyces* are

454 responsible of economically relevant plant diseases, most notably common scab of potato
455 caused by *S. scabies* (Loria et al., 2006).

456 Thus, the taxonomic diversification triggered by both microhabitat and treatment may
457 underpin a functional diversification of the microbiota at the cross-road of mutualism and
458 inter-species competition.

459 This functional diversification of the root communities is manifested by the differential
460 enrichments of ABC transporter genes (mineral) and the two-component system (organic).
461 Although predictive metagenomics is inherently limited by fact that the individual
462 phylogenetic marker used (i.e., the 16S rRNA gene) may fail to recapitulate the genetic
463 diversity existing among strains of the same phylogenetic lineage (Karasov et al., 2018) ,
464 ABC transporters have previously been identified as genes underpinning rhizosphere
465 competence in the microbiota of wheat and cucumber (Ofek-Lalzar et al., 2014). Likewise,
466 the two-component system is required for the rhizosphere colonisation of the biocontrol
467 agent *Pseudomonas fluorescens* WCS365 (De Weert et al., 2006). These observations
468 indirectly support the results gathered from our predictive metagenomics approach. Owing
469 the role played by these classes of genes in uptake of organic compounds (e.g., root
470 exudates, cellular secretion) and stimulus-response mechanisms (e.g., chemotaxis)
471 respectively, it is tempting to hypothesize that the different source of nitrogen define a
472 different metabolic status in and in the vicinity of tomato roots which, in turn, requires a
473 prompt adaptation of the root-inhabiting communities.

474 For instance, experimental data indicate that the abundance of phytoavailable nitrogen, i.e.,
475 the scenario of mineral fertiliser treatments, tends to repress the proliferation and activity of
476 members of the microbiota (Ramirez et al., 2012; Terrazas et al., 2019), and this in turn may
477 be reflected in the metabolism of secondary compounds (terpenoid and polyketide
478 metabolism) and membrane transport (ABC transporters).

479 A “true” comparative metagenomics investigation, whereby the individual communities are
480 subjected to shot-gun sequencing, will be ultimately necessary to test these hypotheses.

481 We further hypothesize that this adaptation is modulated by mineral nitrogen availability, as
482 manifested by the differential enrichment of functions associated to nitrogen metabolism *per*
483 *se* and aminoacids. This observation is congruent with results gathered from monocots
484 wheat (Kavamura et al., 2018) and rice (Zhang et al., 2019) and suggests a cross-species
485 pattern whereby plant’s adaptation to nitrogen forms and availability is mediated, at least in
486 part, by the associated microbiota.

487 Finally, it is interesting to note how the production of antibiotics, namely tetracycline, is also
488 among the functions differentially enriched between fertilisers. It is becoming increasingly
489 clear how plant-associated bacteria can act as a reservoir of antimicrobial genes (Cernava
490 et al., 2019) which can be deployed during inter-organismal competition in the plant
491 microbiota. This hypothesis could be tested by leveraging on indexed- and genome-
492 annotated bacterial collection for the tomato microbiota, similar to the approach pursued
493 with bacteria isolated from other plant species (Levy et al., 2018).

494 Our investigation suggests that the bacterial microbiota of field-grown processing tomato is
495 the product of a selective process that progressively differentiates between rhizosphere and
496 root microhabitats. This process initiates as early as plants are in a nursery stage and it is
497 then more marked when plants reached the harvest stage. This selection *a*) acts both on
498 the relative abundances and phylogenetic assignments of members of the tomato
499 microbiota, *b*) is modulated, at least in part, by the nitrogen fertiliser provided which, in turn,
500 *c*) triggers different microbial metabolic specialisations within tomato roots.

501 It is important to mention that the nitrogen fertiliser may also represent a microbial inoculant
502 *per se*, in particular in the case of organic-based amendments. For instance, a comparative
503 study of 29 different full-scale anaerobic digestion installations revealed that Firmicutes,

504 followed by Bacteroidetes and Proteobacteria, dominated the resulting microbial
505 communities (De Vrieze et al., 2015). Considering the plant-associated profiles observed in
506 this study, in particular the enrichment of Actinobacteria in the root communities, it is
507 legitimate to hypothesize that the input digestate bacteria may act as in inoculum for a part
508 of the tomato microbiota, which is further fine-tuned by the exposure to soil microbes. Future
509 studies, integrating the microbial profiling of the input fertiliser treatment, will be required to
510 accurately elucidate microbial dynamics associated with the application synthetic (i.e., germ-
511 free) and organic fertilisers.

512 [Towards a lab-in-the-field approach to harness the potential of plant microbiota for climate- 513 smart agriculture](#)

514 Our experiments represent an example of how cultivation-independent approaches can be
515 efficiently deployed to investigate the plant microbiota under field conditions. Although this
516 type of investigation is not novel *per se* in tomato (Toju et al., 2019), our results revealed
517 fundamentally novel insights into plant's adaptation to nitrogen fertilisers and the implication
518 for crop yield. Similar to what has recently been postulated for tomato pathogen protection
519 (Kwak et al., 2018), our results predicts that the use of field-derived, sequencing data will
520 allow scientists to identify "signatures" of the plant microbiota that can be targeted to
521 enhance plant performance. This approach, which we define as lab-in-the-field, will be key
522 towards the rationalisation of nitrogen (and other treatments) application in agriculture and
523 we anticipate will pave the way for the effective exploitation of the plant microbiota for
524 agricultural purposes (Schlaeppli and Bulgarelli, 2015; Toju et al., 2018).

525

526 **AUTHORS CONTRIBUTION**

527 DR conceived of and designed the field experiment. DR and FC harvested the field data and
528 samples. FC, RAT, EF and DB conceived of and designed the analysis of the microbiota.
529 FC, MC, CVGA, SR-A performed the microbiota experiments. FC, RAT and DB analysed

530 the sequencing data. FC, RAT and DB wrote the initial draft of the manuscript. All the authors
531 discussed the results and commented on the manuscript.

532 **FUNDING**

533 This research was partially founded by GENBACCA project (Regione Emilia Romagna,
534 POR-FESR 2014/2020 Initiative). The generation and analysis of the sequencing data was
535 supported by a Royal Society of Edinburgh/Scottish Government Personal Research
536 Fellowship co-funded by Marie Curie Actions awarded to DB. SR-A is supported by a
537 BBSRC iCASE studentship awarded to DB (BB/M016811/1) and partnered by the James
538 Hutton Limited (Invergowrie, United Kingdom). RAT and DB are supported by the H2020
539 Innovation Action 'Circles' (European Commission, Grant agreement 818290) awarded to
540 the University of Dundee.

541 **ACKNOWLEDGEMENTS**

542 We thank Dr Guido Bezzi (CIB - Italian Biogas Association, Italy) for providing us with the
543 field, plant material and the digestate treatments, Dr. Massimo Zaghi (CAT - COOP.
544 Agroenergetica Territoriale Correggio S.C.A., Italy) for providing the pelleted digestate and
545 Dr Stefano Tagliavini (SCAM Spa, Italy) for providing the organo-mineral fertiliser.

546 We thank Malcolm Macaulay, Jenny Morris and Dr Pete Hedley (The James Hutton Institute,
547 Invergowrie, UK) for their support in preparing the sequencing library and generating the
548 sequencing data.

549 We thank Prof Marco Candela (University of Bologna, Italy) for the critical comments on the
550 manuscript.

551 **CONFLICTS OF INTERESTS**

552 The authors declare no conflicts of interest.

553 **LITERATURE CITED**

- 554 Alburquerque, J.A., de la Fuente, C., Ferrer-Costa, A., Carrasco, L., Cegarra, J., Abad, M.,
555 and Bernal, M.P. 2012. Assessment of the fertiliser potential of digestates from farm
556 and agroindustrial residues. *Biomass and bioenergy* 40:181-189.
- 557 Alegria Terrazas, R., Giles, C., Paterson, E., Robertson-Albertyn, S., Cesco, S., Mimmo, T.,
558 Pii, Y., and Bulgarelli, D. 2016. Plant-Microbiota Interactions as a Driver of the Mineral
559 Turnover in the Rhizosphere. *Adv Appl Microbiol* 95:1-67.
- 560 Asshauer, K.P., Wemheuer, B., Daniel, R., and Meinicke, P. 2015. Tax4Fun: predicting
561 functional profiles from metagenomic 16S rRNA data. *Bioinformatics* 31:2882-2884.
- 562 Bergna, A., Cernava, T., Rändler, M., Grosch, R., Zachow, C., and Berg, G. 2018. Tomato
563 seeds preferably transmit plant beneficial endophytes. *Phytobiomes Journal* 2:183-
564 193.
- 565 Bluman, A.G. 2009. *Elementary statistics: A step by step approach*. McGraw-Hill Higher
566 Education New York.
- 567 Bulgarelli, D., Schlaeppli, K., Spaepen, S., van Themaat, E.V.L., and Schulze-Lefert, P.
568 2013. Structure and functions of the bacterial microbiota of plants. *Annual review of*
569 *plant biology* 64:807-838.
- 570 Bulgarelli, D., Garrido-Oter, R., Munch, P.C., Weiman, A., Droge, J., Pan, Y., McHardy, A.C.,
571 and Schulze-Lefert, P. 2015. Structure and function of the bacterial root microbiota in
572 wild and domesticated barley. *Cell host & microbe* 17:392-403.
- 573 Bulgarelli, D., Rott, M., Schlaeppli, K., van Themaat, E.V.L., Ahmadinejad, N., Assenza, F.,
574 Rauf, P., Huettel, B., Reinhardt, R., and Schmelzer, E. 2012. Revealing structure and
575 assembly cues for *Arabidopsis* root-inhabiting bacterial microbiota. *Nature* 488:91.
- 576 Caporaso, J.G., Lauber, C.L., Walters, W.A., Berg-Lyons, D., Huntley, J., Fierer, N., Owens,
577 S.M., Betley, J., Fraser, L., Bauer, M., Gormley, N., Gilbert, J.A., Smith, G., and

- 578 Knight, R. 2012. Ultra-high-throughput microbial community analysis on the Illumina
579 HiSeq and MiSeq platforms. *The ISME journal* 6:1621-1624.
- 580 Caporaso, J.G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F.D., Costello, E.K.,
581 Fierer, N., Pena, A.G., Goodrich, J.K., Gordon, J.I., Huttley, G.A., Kelley, S.T.,
582 Knights, D., Koenig, J.E., Ley, R.E., Lozupone, C.A., McDonald, D., Muegge, B.D.,
583 Pirrung, M., Reeder, J., Sevinsky, J.R., Turnbaugh, P.J., Walters, W.A., Widmann,
584 J., Yatsunenko, T., Zaneveld, J., and Knight, R. 2010. QIIME allows analysis of high-
585 throughput community sequencing data. *Nature methods* 7:335-336.
- 586 Cernava, T., Erlacher, A., Soh, J., Sensen, C.W., Grube, M., and Berg, G. 2019.
587 Enterobacteriaceae dominate the core microbiome and contribute to the resistome of
588 arugula (*Eruca sativa* Mill.). *Microbiome* 7:13.
- 589 Chaparro, J.M., Badri, D.V., and Vivanco, J.M. 2014. Rhizosphere microbiome assemblage
590 is affected by plant development. *Isme Journal* 8:790-803.
- 591 Chialva, M., Salvioli di Fossalunga, A., Daghino, S., Ghignone, S., Bagnaresi, P., Chiapello,
592 M., Novero, M., Spadaro, D., Perotto, S., and Bonfante, P. 2018. Native soils with
593 their microbiotas elicit a state of alert in tomato plants. *New Phytologist* 220:1296-
594 1308.
- 595 Consortium, T.G. 2012. The tomato genome sequence provides insights into fleshy fruit
596 evolution. *Nature* 485:635.
- 597 de Jesus Sousa, J.A., and Olivares, F.L. 2016. Plant growth promotion by streptomycetes:
598 ecophysiology, mechanisms and applications. *Chemical and Biological Technologies*
599 *in Agriculture* 3:24.
- 600 De Vrieze, J., Saunders, A.M., He, Y., Fang, J., Nielsen, P.H., Verstraete, W., and Boon, N.
601 2015. Ammonia and temperature determine potential clustering in the anaerobic
602 digestion microbiome. *Water research* 75:312-323.

- 603 De Weert, S., Dekkers, L.C., Bitter, W., Tuinman, S., Wijffes, A.H., Van Boxtel, R., and
604 Lugtenberg, B.J. 2006. The two-component colR/S system of *Pseudomonas*
605 *fluorescens* WCS365 plays a role in rhizosphere competence through maintaining
606 the structure and function of the outer membrane. *FEMS microbiology ecology*
607 58:205-213.
- 608 DeSantis, T.Z., Hugenholtz, P., Larsen, N., Rojas, M., Brodie, E.L., Keller, K., Huber, T.,
609 Dalevi, D., Hu, P., and Andersen, G.L. 2006. Greengenes, a Chimera-Checked 16S
610 rRNA Gene Database and Workbench Compatible with ARB. *Applied and*
611 *Environmental Microbiology* 72:5069-5072.
- 612 Dombrowski, N., Schlaeppli, K., Agler, M.T., Hacquard, S., Kemen, E., Garrido-Oter, R.,
613 Wunder, J., Coupland, G., and Schulze-Lefert, P. 2017. Root microbiota dynamics of
614 perennial *Arabis alpina* are dependent on soil residence time but independent of
615 flowering time. *Isme Journal* 11:43-55.
- 616 Edwards, J., Johnson, C., Santos-Medellin, C., Lurie, E., Podishetty, N.K., Bhatnagar, S.,
617 Eisen, J.A., and Sundaresan, V. 2015. Structure, variation, and assembly of the root-
618 associated microbiomes of rice. *Proceedings of the National Academy of Sciences*
619 *of the United States of America* 112:E911-920.
- 620 Elser, J., and Bennett, E. 2011. A broken biogeochemical cycle. *Nature* 478:29.
- 621 FAO 2017 <http://www.fao.org/faostat/en/#home>
- 622 Grigatti, M., Di Girolamo, G., Chincarini, R., Ciavatta, C., and Barbanti, L. 2011. Potential
623 nitrogen mineralization, plant utilization efficiency and soil CO₂ emissions following
624 the addition of anaerobic digested slurries. *biomass and bioenergy* 35:4619-4629.
- 625 Kanehisa, M., Araki, M., Goto, S., Hattori, M., Hirakawa, M., Itoh, M., Katayama, T.,
626 Kawashima, S., Okuda, S., Tokimatsu, T., and Yamanishi, Y. 2008. KEGG for linking
627 genomes to life and the environment. *Nucleic acids research* 36:D480-484.

- 628 Karasov, T.L., Almario, J., Friedemann, C., Ding, W., Giolai, M., Heavens, D., Kersten, S.,
629 Lundberg, D.S., Neumann, M., and Regalado, J. 2018. *Arabidopsis thaliana* and
630 *Pseudomonas* pathogens exhibit stable associations over evolutionary timescales.
631 *Cell host & microbe* 24:168-179. e164.
- 632 Kavamura, V.N., Hayat, R., Clark, I.M., Rossmann, M., Mendes, R., Hirsch, P.R., and
633 Mauchline, T.H. 2018. Inorganic Nitrogen Application Affects Both Taxonomical and
634 Predicted Functional Structure of Wheat Rhizosphere Bacterial Communities.
635 *Frontiers in Microbiology* 9.
- 636 Kwak, M.-J., Kong, H.G., Choi, K., Kwon, S.-K., Song, J.Y., Lee, J., Lee, P.A., Choi, S.Y.,
637 Seo, M., and Lee, H.J. 2018. Rhizosphere microbiome structure alters to enable wilt
638 resistance in tomato. *Nature biotechnology* 36:1100.
- 639 Letunic, I., and Bork, P. 2006. Interactive Tree Of Life (iTOL): an online tool for phylogenetic
640 tree display and annotation. *Bioinformatics* 23:127-128.
- 641 Levy, A., Gonzalez, I.S., Mittelviehhaus, M., Clingenpeel, S., Paredes, S.H., Miao, J.M.,
642 Wang, K.R., Devescovi, G., Stillman, K., Monteiro, F., Alvarez, B.R., Lundberg,
643 A.D.S., Lu, T.Y., Lebeis, S., Jin, Z., McDonald, M., Klein, A.P., Feltcher, M.E., Rio,
644 T.G., Grant, S.R., Doty, S.L., Ley, R.E., Zhao, B.Y., Venturi, V., Pelletier, D.A.,
645 Vorholt, J.A., Tringe, S.G., Woyke, T., and Dangl, J.L. 2018. Genomic features of
646 bacterial adaptation to plants. *Nature Genetics* 50:138-+.
- 647 Loria, R., Kers, J., and Joshi, M. 2006. Evolution of plant pathogenicity in *Streptomyces*.
648 *Annual Review of Phytopathology* 44:469-487.
- 649 Love, M.I., Huber, W., and Anders, S. 2014. Moderated estimation of fold change and
650 dispersion for RNA-seq data with DESeq2. *Genome biology* 15:550.
- 651 McMurdie, P.J., and Holmes, S. 2013. phyloseq: an R package for reproducible interactive
652 analysis and graphics of microbiome census data. *PloS one* 8:e61217.

- 653 Möller, K. 2015. Effects of anaerobic digestion on soil carbon and nitrogen turnover, N
654 emissions, and soil biological activity. A review. *Agronomy for sustainable*
655 *development* 35:1021-1041.
- 656 Möller, K., and Müller, T. 2012. Effects of anaerobic digestion on digestate nutrient
657 availability and crop growth: a review. *Engineering in Life Sciences* 12:242-257.
- 658 Ofek-Lalzar, M., Sela, N., Goldman-Voronov, M., Green, S.J., Hadar, Y., and Minz, D. 2014.
659 Niche and host-associated functional signatures of the root surface microbiome.
660 *Nature communications* 5:4950.
- 661 Parks, D.H., Tyson, G.W., Hugenholtz, P., and Beiko, R.G. 2014. STAMP: statistical
662 analysis of taxonomic and functional profiles. *Bioinformatics* 30:3123-3124.
- 663 Pulvirenti, A., Ronga, D., Zaghi, M., Tomasselli, A.R., Mannella, L., and Pecchioni, N. 2015.
664 Pelleting is a successful method to eliminate the presence of *Clostridium* spp. from
665 the digestate of biogas plants. *Biomass and Bioenergy* 81:479-482.
- 666 Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., and
667 Glockner, F.O. 2013. The SILVA ribosomal RNA gene database project: improved
668 data processing and web-based tools. *Nucleic acids research* 41:D590-596.
- 669 Ramirez, K.S., Craine, J.M., and Fierer, N. 2012. Consistent effects of nitrogen amendments
670 on soil microbial communities and processes across biomes. *Global Change Biology*
671 18:1918-1927.
- 672 Rathore, R., Dowling, D.N., Forristal, P.D., Spink, J., Cotter, P.D., Bulgarelli, D., and
673 Germaine, K.J. 2017. Crop Establishment Practices Are a Driver of the Plant
674 Microbiota in Winter Oilseed Rape (*Brassica napus*). *Frontiers in Microbiology* 8.
- 675 Robertson-Albertyn, S., Alegria Terrazas, R., Balbirnie, K., Blank, M., Janiak, A., Szarejko,
676 I., Chmielewska, B., Karcz, J., Morris, J., Hedley, P.E., George, T.S., and Bulgarelli,
677 D. 2017. Root Hair Mutations Displace the Barley Rhizosphere Microbiota. *Frontiers*
678 *in plant science* 8:1094.

- 679 Ronga, D., Loyelli, S., Zaccardelli, M., Perrone, D., Ulrici, A., Francia, E., Milc, J., and
680 Pecchioni, N. 2015. Physiological responses of processing tomato in organic and
681 conventional Mediterranean cropping systems. *Sci Hortic-Amsterdam* 190:161-172.
- 682 Ronga, D., Zaccardelli, M., Lovelli, S., Perrone, D., Francia, E., Milc, J., Ulrici, A., and
683 Pecchioni, N. 2017. Biomass production and dry matter partitioning of processing
684 tomato under organic vs conventional cropping systems in a Mediterranean
685 environment. *Sci Hortic-Amsterdam* 224:163-170.
- 686 Ronga, D., Pellati, F., Brighenti, V., Laudicella, K., Laviano, L., Fedailaine, M., Benvenuti,
687 S., Pecchioni, N., and Francia, E. 2018a. Testing the influence of digestate from
688 biogas on growth and volatile compounds of basil (*Ocimum basilicum* L.) and
689 peppermint (*Mentha x piperita* L.) in hydroponics. *Journal of Applied Research on*
690 *Medicinal and Aromatic Plants* 11:18-26.
- 691 Ronga, D., Setti, L., Salvarani, C., De Leo, R., Bedin, E., Pulvirenti, A., Milc, J., Pecchioni,
692 N., and Francia, E. 2019. Effects of solid and liquid digestate for hydroponic baby leaf
693 lettuce (*Lactuca sativa* L.) cultivation. *Sci Hortic-Amsterdam* 244:172-181.
- 694 Ronga, D., Caradonia, F., Setti, L., Hagassou, D., Giaretta Azevedo, C., Milc, J., Pedrazzi,
695 S., Allesina, G., Arru, L., and Francia, E. (2018b). Effects of innovative biofertilizers
696 on yield of processing tomato cultivated in organic cropping systems in northern Italy.
697 In XV International Symposium on Processing Tomato 1233, pp. 129-136.
- 698 Scaglia, B., Pognani, M., and Adani, F. 2015. Evaluation of hormone-like activity of the
699 dissolved organic matter fraction (DOM) of compost and digestate. *Sci Total Environ*
700 514:314-321.
- 701 Schlaeppli, K., and Bulgarelli, D. 2015. The plant microbiome at work. *Molecular plant-*
702 *microbe interactions* : *MPMI* 28:212-217.
- 703 Schlaeppli, K., Dombrowski, N., Oter, R.G., van Themaat, E.V.L., and Schulze-Lefert, P.
704 2014. Quantitative divergence of the bacterial root microbiota in *Arabidopsis thaliana*

- 705 relatives. *Proceedings of the National Academy of Sciences of the United States of*
706 *America* 111:585-592.
- 707 Storey, J.D., and Tibshirani, R. 2003. Statistical significance for genomewide studies.
708 *Proceedings of the National Academy of Sciences of the United States of America*
709 100:9440-9445.
- 710 Terrazas, R.A., Pietrangelo, L., Corral, A.M., Torres-Cortés, G., Robertson-Albertyn, S.,
711 Balbirnie-Cumming, K., Morris, J., Hedley, P.E., Barret, M., and Paterson, E. 2019.
712 Nitrogen Availability Modulates the Host Control of the Barley Rhizosphere
713 Microbiota. *bioRxiv:605204*.
- 714 Tilman, D., Balzer, C., Hill, J., and Befort, B.L. 2011. Global food demand and the
715 sustainable intensification of agriculture. *Proceedings of the National Academy of*
716 *Sciences of the United States of America* 108:20260-20264.
- 717 Toju, H., Okayasu, K., and Notaguchi, M. 2019. Leaf-associated microbiomes of grafted
718 tomato plants. *Scientific reports* 9:1787.
- 719 Toju, H., Peay, K.G., Yamamichi, M., Narisawa, K., Hiruma, K., Naito, K., Fukuda, S., Ushio,
720 M., Nakaoka, S., and Onoda, Y. 2018. Core microbiomes for sustainable
721 agroecosystems. *Nat. Plants* 4:247-257.
- 722 Walters, W.A., Jin, Z., Youngblut, N., Wallace, J.G., Sutter, J., Zhang, W., González-Peña,
723 A., Peiffer, J., Koren, O., Shi, Q., Knight, R., Glavina del Rio, T., Tringe, S.G., Buckler,
724 E.S., Dangl, J.L., and Ley, R.E. 2018. Large-scale replicated field study of maize
725 rhizosphere identifies heritable microbes. *Proceedings of the National Academy of*
726 *Sciences*.
- 727 Zhang, J., Liu, Y.-X., Zhang, N., Hu, B., Jin, T., Xu, H., Qin, Y., Yan, P., Zhang, X., and Guo,
728 X. 2019. NRT1. 1B is associated with root microbiota composition and nitrogen use
729 in field-grown rice. *Nature biotechnology:1*.

730 Zsögön, A., Čermák, T., Naves, E.R., Notini, M.M., Edel, K.H., Weinl, S., Freschi, L., Voytas,
731 D.F., Kudla, J., and Peres, L.E.P. 2018. De novo domestication of wild tomato using
732 genome editing. *Nature biotechnology*.

733

734

735 **TABLES**736 **Table 1. Composition and information on fertilisers used in this study.**737 TOC = Total organic carbon; N = Nitrogen; P = Phosphorus; K = Potassium; H₂O = water
738 content

Treatment	TOC%	(N)%	(P)%	(K)%	H ₂ O%	Additional information
Synthetic fertiliser (MF)		41				Ammonium nitrate (N 26%) and calcium nitrate (N 15%)
Pelleted digestate (PE)	39.70	1.50	2.50	2.00	7.80	(Pulvirenti et al., 2015)
Slow acting liquid digestate (SRLD)	3.74	0.34		0.95		Liquid digestate plus the nitrogen stabilizer Vizura® (BASF, 2 L ha ⁻¹),
Liquid digestate (LD)	3.74	0.34		0.95		EC 1.07 dS m ⁻¹ and pH 8.3
Organo-mineral fertiliser (SC)	10.50	10.00	5.00	15.00	7.00	Produced by SCAM Spa (Modena, Italy), based on solid digestate for the organic fraction
Slow acting synthetic fertiliser (SRMF)		15.00	15.40	15.00		NPK Original Gold® (Compo Expert)

739

740

741 **FIGURE LEGENDS**742 **Figure 1. Effect of the nitrogen treatments on tomato yield traits**

743 Mean and standard deviation of **(A)** marketable yield and **(B)** Brix of tomato plants exposed
744 to the following treatments: LD (Liquid Digestate), SRLD (Slow acting Liquid Digestate), PE
745 (Pelleted Digestate); SC (Organo-mineral fertiliser); MF (Mineral Fertiliser); SRMF (Slow
746 acting Mineral Fertiliser). Different letters denote statistically significant differences between
747 treatments by Analysis of Variance (ANOVA). Means were compared using Bonferroni's test
748 at the 5% level ($P < 0.001$).

749 **Figure 2. The tomato root microbiota is a gated community**

750 Average **(A)** number of observed OTUs, **(B)** Chao 1 index and **(C)** Shannon index computed
751 on the indicated rhizosphere and root specimens. Abbreviations LD (Liquid Digestate),
752 SRLD (Slow acting Liquid Digestate), PE (Pelleted Digestate); SC (Organo-mineral
753 fertiliser); MF (Mineral Fertiliser); SRMF (Slow acting Mineral Fertiliser). Asterisks denote
754 statistically significant differences between microhabitat by non-parametric Wilcoxon rank
755 sum test ($P < 0.01$). Different blue letters within individual microhabitats denote statistically
756 significant differences between treatment means by Kruskal-Wallis non parametric analysis
757 of variance followed by Dunn's post-hoc test ($P < 0.05$); ns, no significant differences
758 observed.

759 **Figure 3. The tomato rhizosphere and root microbiota host compositionally different
760 communities.**

761 PCoA calculated using a weighted UniFrac matrix calculated on the OTUs clustered at 97%
762 identity among the indicated microhabitat and treatments. Abbreviations: LD (Liquid
763 Digestate), SRLD (Slow acting Liquid Digestate), PE (Pelleted Digestate); SC (Organo-
764 mineral fertiliser); MF (Mineral Fertiliser); SRMF (Slow acting Mineral Fertiliser).

765 [Figure 4. The enrichment of Actinobacteria is a distinctive feature of the tomato root](#)
766 [microbiota.](#)

767 Phylogenetic relationships of the OTUs enriched in rhizosphere and root compartment.
768 Individual external nodes represent one of the OTUs enriched in either (or both) rhizosphere
769 or root samples in no treatment conditions (Wald test, P value < 0.01, FDR corrected) whose
770 colour reflects their taxonomic affiliation at Phylum level. A black bar in the outer rings
771 depicts whether that given OTU was identified in the rhizosphere- or root-enriched sub-
772 communities, respectively. Phylogenetic tree constructed using OTUs 16S rRNA gene
773 representative sequences.

774
775 [Figure 5. Nitrogen fertiliser modulates bacterial enrichment in the tomato rhizosphere and](#)
776 [root compartments.](#)

777 Number of OTUs significantly enriched (Wald test, P value < 0.01, FDR corrected) in the
778 indicated treatment versus untreated controls in **(A)** rhizosphere and **(B)** roots. In each
779 panel, blue bars denote the total number of enriched OTUs for a given treatment, the black
780 bars denote the magnitude of the enrichment in either the individual treatment or among two
781 or more treatments highlighted by the interconnected dots underneath the panels.
782 Abbreviations: LD (Liquid Digestate), SRLD (Slow acting Liquid Digestate), PE (Pelleted
783 Digestate); SC (Organo-mineral fertiliser); MF (Mineral Fertiliser); SRMF (Slow acting
784 Mineral Fertiliser).

785 [Figure 6. Digestate- and mineral-based fertilisers trigger a functional diversification of the](#)
786 [tomato root microbiota.](#)

787 Prokaryotic functions discriminating between Digestate-based (indicated as 'R_organic':
788 Liquid Digestate; Slow acting Liquid Digestate and Pelleted Digestate) and mineral-based
789 fertilisers (indicated as 'R_mineral': Organo-mineral; Mineral Fertiliser and Slow acting

790 Mineral Fertiliser) retrieved from Tax4Fun functional profiles (Welch's t-test FDR corrected,
791 $p < 0.01$).




Figure 1. Mean and standard deviation of (A) marketable yield and (B) Brix of tomato plants exposed to the following treatments: LD (Liquid Digestate), SRLD (Slow acting Liquid Digestate), PE (Pelleted Digestate); SC (Organo-mineral fertiliser); MF (Mineral Fertiliser); SRMF (Slow acting Mineral Fertiliser). Different letters denote statistically significant differences between treatments by Analysis of Variance (ANOVA). Means were compared using Bonferroni's test at the 5% level ($P < 0.001$).

159x155mm (150 x 150 DPI)




Figure 2. Average (A) number of observed OTUs, (B) Chao 1 index and (C) Shannon index computed on the indicated rhizosphere and root specimens. Abbreviations LD (Liquid Digestate), SRLD (Slow acting Liquid Digestate), PE (Pelleted Digestate), SC (Organo-mineral fertiliser); MF (Mineral Fertiliser); SRMF (Slow acting Mineral Fertiliser). Asterisks denote statistically significant differences between microhabitat by non-parametric Wilcoxon rank sum test ($P < 0.01$). Different blue letters within individual microhabitats denote statistically significant differences between treatment means by Kruskal-Wallis non parametric analysis of variance followed by Dunn's post-hoc test ($P < 0.05$); ns, no significant differences observed.

139x228mm (300 x 300 DPI)




Figure 3. PCoA calculated using a weighted UniFrac matrix calculated on the OTUs clustered at 97% identity among the indicated microhabitat and treatments. Abbreviations: LD (Liquid Digestate), SRLD (Slow acting Liquid Digestate), PE (Pelleted Digestate); SC (Organo-mineral fertiliser); MF (Mineral Fertiliser); SRMF (Slow acting Mineral Fertiliser).

166x193mm (300 x 300 DPI)




Figure 4. The enrichment of Actinobacteria is a distinctive feature of the tomato root microbiota. Phylogenetic relationships of the OTUs enriched in rhizosphere and root compartment. Individual external nodes represent one of the OTUs enriched in either (or both) rhizosphere or root samples in no treatment conditions (Wald test, P value < 0.01, FDR corrected) whose colour reflects their taxonomic affiliation at Phylum level. A black bar in the outer rings depicts whether that given OTUs was identified in the rhizosphere- or root-enriched sub-communities, respectively. Phylogenetic tree constructed using OTUs 16S rRNA gene representative sequences.

163x191mm (300 x 300 DPI)




Figure 5. Nitrogen fertiliser modulates bacterial enrichment in the tomato rhizosphere and root compartments. Number of OTUs significantly enriched (Wald test, P value < 0.01, FDR corrected) in the indicated treatment versus untreated controls in (A) rhizosphere and (B) roots. In each panel, blue bars denote the total number of enriched OTUs for a given treatment, the black bars denote the magnitude of the enrichment in either the individual treatment or among two or more treatments highlighted by the interconnected dots underneath the panels. Abbreviations LD (Liquid Digestate), SRLD (Slow acting Liquid Digestate), PE (Pelleted Digestate); SC (Organo-mineral fertiliser); MF (Mineral Fertiliser); SRMF (Slow acting Mineral Fertiliser).





Figure 6. Prokaryotic functions discriminating between Digestate-based (indicated as 'R_organic': Liquid Digestate; Slow acting Liquid Digestate and Pelleted Digestate) and mineral-based fertilisers (indicated as 'R_mineral': Organo-mineral; Mineral Fertiliser and Slow acting Mineral Fertiliser) retrieved from Tax4Fun functional profiles (Welch's t-test FDR corrected, $p < 0.01$).

91x253mm (300 x 300 DPI)



Supplementary Figure S1. Images of the field at (A) plant's transplant and details of (B) an individual tomato plant and (C) the plantation at harvest stage.

131x87mm (300 x 300 DPI)



Supplementary Figure S2. PCoA calculated using a Bray-Curtis matrix calculated on the OTUs clustered at 97% identity among the indicated microhabitat and treatments. Abbreviations: LD (Liquid Digestate), SRLD (Slow acting Liquid Digestate), PE (Pelleted Digestate); SC (Organo-mineral fertiliser); MF (Mineral Fertiliser); SRMF (Slow acting Mineral Fertiliser).

224x189mm (300 x 300 DPI)