

Accepted Manuscript

Conjugated linoleic acid production and probiotic assessment of *Lactobacillus plantarum* isolated from Pico cheese

Susana C. Ribeiro, Catherine Stanton, Bo Yang, R. Paul Ross, Célia C.G. Silva



PII: S0023-6438(17)30959-3

DOI: [10.1016/j.lwt.2017.12.065](https://doi.org/10.1016/j.lwt.2017.12.065)

Reference: YFSTL 6754

To appear in: *LWT - Food Science and Technology*

Received Date: 12 July 2017

Revised Date: 14 December 2017

Accepted Date: 26 December 2017

Please cite this article as: Ribeiro, S.C., Stanton, C., Yang, B., Ross, R.P., Silva, Cé.C.G., Conjugated linoleic acid production and probiotic assessment of *Lactobacillus plantarum* isolated from Pico cheese, *LWT - Food Science and Technology* (2018), doi: 10.1016/j.lwt.2017.12.065.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1 Conjugated linoleic acid production and probiotic assessment of

2 *Lactobacillus plantarum* isolated from Pico cheese

3

4

5 Susana C. Ribeiro¹, Catherine Stanton², Bo Yang^{2,3}, R. Paul Ross⁴, Célia C.G. Silva^{1*}

6 ¹ Instituto de Investigação e Tecnologias Agrárias e do Ambiente dos Açores (IITAA),

7 Universidade dos Açores, Angra do Heroísmo, Açores, Portugal

8 ² Teagasc Food Research Centre, Moorepark, Fermoy, Co. Cork, Ireland

9 ³ State Key Laboratory of Food Science and Technology, School of Food Science and
10 Technology, Jiangnan University, Wuxi, Jiangsu, China

11 ⁴ College of Science, Engineering and Food Science, University College Cork, Ireland

12

13

14

15

16 *Corresponding author:

17 Célia Costa Silva

18 Instituto de Investigação e Tecnologias Agrárias e do Ambiente dos Açores (IITAA)

19 Universidade dos Açores

20 Rua Capitão João D'Avila

21 9700-042 Angra do Heroísmo

22 Portugal

23 Tel:+351295402200

24 Email: celia.cg.silva@uac.pt

25

26

27 **Abstract**

28 Lactic acid bacteria isolated from a traditional Azorean cheese were screened for their
29 ability to convert free linoleic acid to conjugated linoleic acid (CLA). Two strains of
30 *Lactobacillus plantarum* were recognized as potential CLA producers. GC analysis
31 identified *cis*-9, *trans*-11 C18:2 as the predominant isomer (10-14 µg/mL), followed by
32 *trans*-9, *trans*-11 C18:2 (4-6 µg/mL). The CLA producing strains demonstrated strong
33 biofilm capacity, high cell surface hydrophobicity and good auto-aggregation ability.
34 These strains were capable of surviving in the presence of bile salts (0.3%) and
35 pancreatin (0.1%), but only the highest CLA producer (L3C1E8) was able to resist low
36 pH (2.5). Moreover, the CLA-producers showed good adhesion capacity to intestinal
37 human cells (Caco-2 and HT-29) and were able to prevent colonization of *Escherichia*
38 *coli*. Of the two strains, *Lactobacillus plantarum* L3C1E8 revealed superior probiotic
39 properties and great potential for producing food products enriched in the two CLA
40 isomers, *cis*-9, *trans*-11 C18:2 (60%) and *trans*-9, *trans*-11 C18:2 (25%).

41

42

43

44

45

46

47 **Keywords:** Functional food, conjugated linoleic acid, CLA, lactic acid bacteria,

48 *Lactobacillus plantarum*, probiotics.

49

50

51 1. Introduction

52 The 'functional foods' concept originated in Japan, but owing to the positive health
53 benefits of such foods, consumer demand has spread globally. These foods are fortified
54 with biologically active compounds that may impart beneficial effects on the body, as
55 well as decrease the risk of certain diseases (Annunziata & Vecchio, 2011; Bigliardi &
56 Galati, 2013). Conjugated linoleic acid (CLA) refers to a heterogeneous group of
57 positional and geometric (*cis* or *trans*) isomers of linoleic acid (LA) with conjugated
58 double bonds at multiple carbon positions (Pandit, Anand, Kalscheur, & Hassan, 2012).
59 CLA isomers are considered to be beneficial functional lipids due to their biological
60 activities and health promoting properties, such as anti-cancer, anti-atherogenic, anti-
61 obesity and anti-inflammatory (Chinnadurai, Kanwal, Tyagi, Stanton, & Ross, 2013;
62 Coakley et al., 2006; Hennessy, Ross, Devery, & Stanton, 2011; Kobaa & Yanagita,
63 2014; Shen et al., 2013; Sluijs, Plantinga, de Roos, Mennen, & Bots, 2010). Dietary
64 CLA can be found primarily in the meat and milk of ruminants as a result of bacterial
65 biohydrogenation of lipids in the rumen (Lin & Lee, 1997; Fuke & Nornberg, 2017),
66 with the *cis*-9, *trans*-11 C18:2 isomer being the most prevalent (Chin, Liu, Storkson,
67 Ha, & Pariza, 1992). Other isomers present in smaller quantities include *trans*-7, *cis*-9
68 C18:2, *cis*-11, *trans*-13 C18:2, *cis*-8, *trans*-10 C18:2, and *trans*-10, *cis*-12 C18:2.
69 Nonetheless, the low concentrations of CLA found in these food products (meat, milk,
70 and dairy products) are lower than the level required to obtain health benefits (Gauillier
71 et al., 2007). Consequently, increasing the concentration of CLA in food products has
72 been the target of several studies in recent years with a view to developing functional
73 food products (Fuke & Nornberg, 2017; Ozer, Kilic & Kilic, 2016; Shingfield, Bonnet,
74 & Scollan, 2013).

75 Lactic acid bacteria (LAB), especially *Lactobacillus*, may produce CLA by
76 isomerization of linoleic acid (LA) (Alonso, Cuesta, & Gilliland, 2003; Chung et al.,
77 2008; Coakley et al., 2003; Jiang, Björck, & Fondén, 1998; Kishino et al., 2003; Ogawa
78 et al., 2005; Zeng, Lin, & Gong, 2009). In this regard, the production of these bioactive
79 fatty acid metabolites may be considered a probiotic trait. Incorporation of such bacteria
80 into foods offers a viable solution for increasing CLA content. Therefore, the
81 identification of LAB cultures capable of producing CLA from a LA source is a
82 worthwhile pursuit for the food industry, particularly in relation to fermented dairy
83 products. (Ozer, Kilic & Kilic, 2016; Vieira et al., 2017).). In addition, CLA production
84 in humans can be performed by the gut microbiota (Raimondi et al., 2016), as CLA
85 production by a probiotic has been observed in the murine gut where it was linked to
86 suppression of colitis (Bassaganya-Riera et al., 2012). Thus, the ability of CLA-
87 producing strains to exhibit probiotic characteristics such as survival in the
88 gastrointestinal (GI) tract is also of significance given that they may impart beneficial
89 effects in the gut.

90 In order to be considered a probiotic, a bacterial strain must be able to survive in the
91 extreme conditions of the GI tract (low pH in stomach, bile salts), adhere to the
92 intestinal mucosa and impart beneficial effects on the host such as antimicrobial and
93 immunomodulatory properties, amongst others (Del Piano et al., 2006; Verna & Lucak,
94 2010). In addition, biofilm production is considered an important characteristic leading
95 to successful colonization (Salas-Jara, Ilabaca, Vega, & García, 2016).

96 Pico cheese is a traditional cheese with Protected Designation of Origin (PDO) status; it
97 is produced from raw cow's milk from Pico Island in the Azores, without the addition
98 of starter-cultures. In small artisanal dairy units the raw milk is coagulated with animal
99 rennet, the curds are manually cut, placed into molds and left to ripen for approx. 20

100 days. Consequently, the microbial fermentation is carried out by the indigenous
101 microbiota derived exclusively from the raw milk of grazing cows and the production
102 environment. Therefore, Pico cheese is a fertile ground for the identification and
103 isolation of novel LAB strains. The present study was aimed to screen LAB previously
104 isolated from Pico cheese (Domingos-Lopes et al., 2017), for their ability to produce
105 CLA. The highest CLA-producing strains were further evaluated for their probiotic
106 potential, which included ability to survive to the extreme conditions of the GI tract,
107 adhesion to intestinal cells and anti-adhesion assays against the pathogenic bacterium
108 *Escherichia coli*.

109

110 **2. Materials and methods**

111 **2.1 Microorganisms**

112 The LAB strains under investigation in this study were previously isolated from a
113 traditional Azorean cheese (Pico cheese) and had been phenotypically and genetically
114 identified (Domingos-Lopes et al., 2017). One hundred and twelve LAB strains
115 belonging to the genus *Lactococcus* (3), *Lactobacillus* (21), *Leuconostoc* (4) and
116 *Enterococcus* (84) were selected from the bacterial culture collection isolated from this
117 cheese. LAB cultures were activated by successive subculturing in MRS broth (Difco
118 Laboratories, Detroit, MI) and grown at 30 °C. The strain *Escherichia coli* ATCC
119 25922 was used in the assays of bacterial adhesion to intestinal cells and was grown at
120 37 °C in Nutrient Broth under aerobic conditions (Fluka, Gillingham, England).

121

122 **2.2 Screening of LAB for CLA production**

123 LAB strains were screened for CLA production using a spectrophotometric detection
124 method according to Barrett et al. (2007). Briefly, LAB strains were incubated in MRS

125 broth containing free linoleic acid (0.5 mg/mL; Sigma-Aldrich, St Louis, MO, USA)
126 and 2% (w/v) Tween 80, at 30°C for 48 h. After incubation, 1 mL of culture was
127 centrifuged at $20,800 \times g$ for 1 min, the pellet was discarded, and the supernatant was
128 mixed with 2 mL of isopropanol by vortexing and allowed to stand for 3 min. The fatty
129 acids were extracted by vortexing the solution and allowing to stand for a further 3 min,
130 following the addition of 1.5 mL of hexane. The presence of CLA in the culture
131 supernatant was assayed by dispensing 230 μ l of the fat-soluble hexane layer into a UV-
132 transparent 96-well plate (Costar, Corning, NY) and determining the absorbance at 233
133 nm using a 96-well plate spectrophotometer (GENios Plus; Tecan, Medford, MA).
134 Measurements were obtained in duplicate.

135 A standard curve was constructed for the absorbance at 233 nm *versus* the CLA
136 concentration (mg/mL), using pure *cis*-9, *trans*-11 CLA isomer (Nu-Check Prep.,
137 Elysian, MN, USA), This method was used for screening LAB for CLA production.
138 Positive results were further confirmed by gas chromatography.

139

140 **2.3 CLA quantification by gas chromatography (GC)**

141 **2.3.1 Lipid extraction from bacterial supernatant fluids and pellets**

142 CLA production by *Lactobacillus plantarum* L2C21E8 and *Lb. plantarum* L3C1E8,
143 identified as potential CLA-producing strains from the screening in section 2.2, was
144 quantified by gas chromatography, according to the method described by Yang et al.
145 (2014) with some modifications. Prior to examination of the strains, each culture was
146 subcultured twice in MRS broth. The strains were then cultured (1%) in broth
147 containing 0.5 mg/mL free linoleic acid (Sigma-Aldrich). The stock solution consisting
148 of linoleic acid (30 mg/mL) and 2% (v/v) Tween 80, was previously filter sterilized

149 through a 0.45 μm filter (Minisart, Sigma-Aldrich) and stored in the dark at -20°C . The
150 strains were incubated aerobically at 30°C .

151 After 48 h incubation, the LAB cultures were centrifuged at $5000 \times g$ for 10 min at
152 room temperature. The fat was extracted from the culture supernatant fluid as follows:
153 An internal standard, C17:0 heptadecanoic acid (99% pure; Sigma-Aldrich), was added
154 to 5 mL of the supernatant fluid to give a final concentration of 0.75 g internal standard
155 per sample. Then, 5 mL of isopropanol was added to the supernatant fluid, and the
156 samples were vortexed for 30 s. Five milliliter of *n*-hexane was added to this mixture,
157 vortexed and centrifuged at $3260 \times g$ for 5 min. The resultant hexane layer (containing
158 lipids) was dried under a stream of nitrogen. For bacterial pellet extraction, the pellet
159 from 10 mL of bacterial culture was washed in 2 mL saline solution (0.137 mol/L NaCl,
160 7.0 mmol/L K_2HPO_4 and 2.5 mmol/L KH_2PO_4). The cells were vortexed and
161 centrifuged at $3260 \times g$ for 10 min, and the washing step repeated twice. The cells were
162 suspended in 1 mL saline solution and then the samples were extracted completely as
163 described above for the bacterial supernatant fluid. Fat was extracted from supernatant
164 and pellet, independently. The lipids were stored at -20°C prior to preparation of fatty
165 acid methyl esters for GC analysis. Samples were analyzed in triplicate.

166

167 **2.3.2 Preparation of fatty acid methyl esters**

168 The extracted lipids were analyzed by gas chromatography following methylation with
169 NaOH- BF_3 in methanol as described by Yang et al. (2014). *Tert*-butyl methyl ether
170 (MTBE, Sigma-Aldrich) (0.5 mL) was added to samples prepared above, together with
171 10 mL of NaOH (0.5 M) in methanol, and the mixture was vortexed for approx. 30 s
172 and incubated for 12 min at 90°C . Then, 10 mL of BF_3 in methanol (Sigma-Aldrich)
173 was added and incubated for 12 min at 90°C . Upon incubation, 2 mL of water saturated

174 with hexane (1 mL of hexane in 100 mL of water) and 4 mL of hexane were added to
175 the mixture and vigorously vortexed for 30 s. The upper (organic) phase was collected,
176 and again, 2 mL of water saturated hexane was added. After standing for sufficient time,
177 the top layer was collected to a clean methylation tube containing 0.5 g of anhydrous
178 sodium sulphate and left in the dark for 1h. Aliquots of the samples containing fatty
179 acid methyl esters (FAME) were stored in a vial at -20 °C for further quantification of
180 CLA content by GC.

181

182 **2.3.3 Gas chromatography analysis**

183 A gas chromatograph (3500, Varian, Harbor City, CA, USA) fitted with a flame
184 ionization detector was used. Helium served as the carrier gas. The GC conditions for
185 separation of CLA isomers were as described by Coakley et al. (2003). The CLA
186 isomers were identified by comparison with the retention time of the reference CLA
187 standard mix (Sigma-Aldrich).

188

189 **2.4 Evaluation of biofilm formation**

190 Biofilm formation by *Lb. plantarum* L2C21E8 and L3C1E8 strains was evaluated in 96-
191 well microtiter plates following the method described by Pérez et al. (2014). Briefly,
192 overnight LAB cultures from MRS broth were used as inoculums and incubated in a 96-
193 well microtiter plate without shaking at 30 °C for 24, 48 and 72 h. Then, wells were
194 washed with phosphate-buffered saline (PBS) and biofilms stained for 30 min with 200
195 μ L 0.1% (w/v) crystal violet. The dye in the cells was then remobilized with 200 μ L of
196 30% (v/v) glacial acetic acid, and the absorbance of the solution (A_{570}) was determined
197 by spectrophotometer (Fluostar Omega, BMG Labtech). Based on the absorbance, the
198 strains were classified into the following categories: no biofilm producer, weak,

199 moderate or strong biofilm producers. Two independent experiments were performed,
200 each with four replicates.

201

202 **2.5 Probiotic potential of CLA producers**

203

204 **2.5.1 Bacterial cell surface hydrophobic/hydrophilic characteristics**

205 Cell surface characteristics of *Lb. plantarum* L2C21E8 and L3C1E8 strains were
206 measured according to the method of Bellon-Fontaine et al. (1996). LAB strains were
207 grown in MRS broth and harvested by centrifugation ($4500 \times g$, 10 min), washed twice
208 with 0.85% NaCl and resuspended in the same solution. Suspensions were mixed with
209 three different solvents: chloroform, ethyl acetate and n-hexadecane (Sigma-Aldrich),
210 and the two phase systems were mixed by vortexing for 1 min. After the complete
211 separation of two phases, the absorbance was measured (A_1) at 600nm. The percentage
212 of bacterial adhesion to solvents was calculated as follow: % Adhesion = $(1-A_1/A_0) \times$
213 100, where A_0 and A_1 were the absorbance values before and after extraction with the
214 organic solvent. The experiment was performed in triplicate.

215

216 **2.5.2 Auto-aggregation**

217 Auto-aggregation determination was performed according to the protocol described by
218 Todorov & Dicks (2008). *Lb. plantarum* L2C21E8 and L3C1E8 strains were grown for
219 24h in MRS broth, centrifuged, washed and resuspended in 0.85% sterile saline
220 solution. After 60 min of incubation at room temperature, the cultures were centrifuged
221 at $300 \times g$ for 2 min. Auto-aggregation was calculated by the following equation: %
222 Auto-aggregation = $[(A_0-A_1)/A_0] \times 100$, whereas A_0 represents absorbance at time 0, and
223 A_1 absorbance after 60 min. Experiments were conducted in triplicate.

224

225 2.5.3 Resistance to low pH, bile salts and pancreatin

226 *Lb. plantarum* L2C21E8 and L3C1E8 strains were tested for resistance to low pH, bile
227 salts and pancreatin according to Argyri et al. (2013). Overnight cultures of LAB were
228 harvested by centrifugation ($10,000 \times g$ for 5 min at 4 ° C) and the pellets washed with
229 sterile phosphate-buffer saline (pH 7.3). To determine acid tolerance, the cell pellet was
230 resuspended in PBS adjusted to pH 2.5. For bile salts and pancreatin resistance, the cell
231 pellets were resuspended in PBS solution (pH 7.3), containing 0.3% (w/v) of bile salts
232 (Fluka, Buchs, Switzerland) and 0.1% (w/v) of pancreatin (Sigma-Aldrich). LAB were
233 then incubated at 37 ° C for 0, 0.5, 1, 2 and 3 h. Enumeration of viable cells were
234 performed on MRS agar (Biokar, Beauvais, France). Assays were carried out as four
235 independent experiments.

236

237 2.5.4 Adhesion assays**238 2.5.4.1 LAB adhesion capacity to intestinal human cells**

239 The method described by Argyri et al. (2013) was followed to study adhesion of *Lb.*
240 *plantarum* L2C21E8 and L3C1E8 strains to HT-29 and Caco-2 cells, with some minor
241 modifications. HT-29 cells were grown and maintained in McCoy's 5A medium
242 (Sigma-Aldrich) containing 10% (v/v) fetal bovine serum (FBS, Sigma-Aldrich) and
243 1% (v/v) gentamicin (Sigma-Aldrich) at 37 ° C in an atmosphere of 5% CO₂. Caco-2
244 cells were routinely cultured in Dulbecco's modified Eagle medium (DMEM, Sigma-
245 Aldrich) supplemented with 10% (v/v) fetal bovine serum (FBS, Sigma-Aldrich), 1%
246 (v/v) non-essential amino acids (Sigma-Aldrich), and 1% (v/v) gentamicin (Sigma-
247 Aldrich). For the experiments, cells were seeded in 24-well tissue plates at 10⁴ cells
248 density and cultured until differentiation and a confluent layer was attained. The culture

249 medium was changed every 2-3 days. Medium without antibiotic was used 24 h before
250 the experiments. The ratio of cells to bacteria was $\geq 1:100$. Overnight cultures of LAB
251 strains were centrifuged, washed twice with PBS and diluted in cell culture medium
252 without antibiotic to a concentration of approximately 10^8 - 10^{10} CFU. After co-
253 incubation for 2 h at 37 °C, wells were washed three times with PBS solution to remove
254 any non-adherent bacteria. Then, 0.1 mL of trypsin-EDTA solution (Sigma-Aldrich)
255 was added to detach cells and adhered bacteria. The cells were lysed with Triton X-100
256 (0.25%, Merck) and bacterial counts were carried out in MRS agar (Biokar). The
257 adhesion ability was expressed as the number of adhered bacteria (CFU/mL). Assays
258 were performed in triplicate and three independent experiments were carried out.

259

260 **2.5.4.2 Effects of LAB on adhesion of *E. coli* to HT-29 cells**

261 The ability of *Lb. plantarum* L2C21E8 and L3C1E8 strains to inhibit the adhesion of *E.*
262 *coli* ATCC 25922 to HT-29 cells was investigated according to the method described by
263 García-Ruiz et al. (2014). Three different assays were conducted: competition,
264 inhibition and displacement. For the competition assay, LAB strains and *E. coli* were
265 simultaneously added to HT-29 cells (1:1) and incubated for 60 min. In the inhibition
266 assay, LAB strains were previously incubated for 60 min with cells before adding *E.*
267 *coli*, and incubated for another 60 min with *E. coli*. For the displacement assay, *E. coli*
268 was added firstly to cells for 60 min before addition of LAB strains for further 60 min.
269 *E. coli* counts were performed in TBX Agar (Oxoid, Basingstoke, England). Results
270 were expressed as the percentage of inhibition of *E. coli* adhesion to cells by each LAB
271 strain. Three independent experiments were carried out in triplicate.

272

273 **2.6. Statistical analyses**

274 Analyses of CLA screening were performed in duplicate and results were expressed as
275 mean \pm standard error of the mean (SEM). A one-way analysis of variance (ANOVA)
276 was used to compare data obtained for the different strains. Two-way factorial ANOVA
277 was used to determine the effect of strain on CLA isomers production in supernatant
278 and pellet. Data on biofilm formation was also processed by ANOVA. Probiotic
279 characteristics of CLA producing strains were analyzed by factorial ANOVA, the
280 factors were time or cells (Caco-2 and HT-29) and *Lb. plantarum* strains. When a
281 significant F was observed ($P < 0.05$), differences between means were evaluated by
282 Bonferroni multiple comparison test. Percentages of microbial adhesion to solvents,
283 auto-aggregation, competition, inhibition and displacement of *E. coli* in the presence of
284 *Lb. plantarum* strains were analyzed by the Kruskal-Wallis test. The significant level
285 was set at 0.05. All statistical tests were performed using IBM SPSS Statistics, version
286 22 (IBM Corporation, New York, USA).

287

288 3. Results

289 3.1 Screening of LAB for CLA production

290 A standard curve was constructed for the absorbance at 233 nm *versus* CLA
291 concentration, using pure C18:2 *cis*-9, *trans*-11. Results demonstrated that a direct
292 relationship could be established between absorbance and CLA concentration
293 ($R^2 = 0.9959$, data not shown), up to an absorbance of 2.8. Therefore, the CLA
294 concentrations in culture supernatants with an absorbance at 233nm less than or equal to
295 2.8 could be calculated from the linear trend line of the standard curve using the
296 equation $y = 0.0575x - 0.1187$. The results of LAB screening for CLA production are
297 presented in Table 1. Two strains L2C21E8 and L3C1E8, identified as *Lb. plantarum*,
298 presented significantly ($P < 0.05$) higher CLA concentrations (or the highest percentage

299 of conversion), at 17.94 and 15.36 $\mu\text{g/mL}$ of CLA, respectively, compared to the other
300 CLA-producing strains. However, it is important to emphasize that this method does not
301 distinguish between isomers of CLA, since it is based on measurement of the
302 conjugated double bond in the fatty acid. Therefore, a more accurate analysis of the
303 CLA-isomers produced by these two strains was performed by gas chromatography.

304

305 **3.2 Gas chromatography analysis**

306 The chromatogram profiles obtained by GC for the cell supernatants of *Lb. plantarum*
307 strains L2C21E8 and L3C1E8 are presented in Fig.1. The CLA isomers detected by GC
308 were *cis*-9, *trans*-11 C18:2; *trans*-10, *cis*-12 C18:2; *cis*-9, *cis*-11 C18:2; and *trans*-9,
309 *trans*-11 C18:2. As expected, the *cis*-9, *trans*-11 CLA isomer (rumenic acid) was the
310 most abundant isomer generated and was mainly found in the cell supernatant (Fig.2).
311 The level of *cis*-9, *trans*-11 CLA isomer was significantly ($P < 0.05$) higher than *cis*-9,
312 *cis*-11 and *trans*-10, *cis*-12 isomers. C18:2 *trans*-9, *trans*-11 was the second most
313 abundant CLA isomer detected in the cell supernatant, for both strains. In contrast, *cis*-
314 9, *cis*-11 C18:2 was found at equivalent concentrations in the pellet and the supernatant.
315 Other CLA isomers, such as *trans*-10, *cis*-12, were also generated, but as minor
316 compounds. Similar profiles were obtained for the two selected strains and no
317 significant ($P > 0.05$) differences were found between them.

318

319 **3.3 Biofilm formation**

320 *Lb. plantarum* L2C21E8 and L3C1E8 strains were evaluated for their capacity to form
321 biofilms during 24, 48 and 72 h of incubation. The two *Lactobacillus* strains were able
322 to form biofilm structures on polystyrene plates after 24 h of growth. However, the
323 strains differed significantly ($P < 0.05$) since *Lb. plantarum* L3C1E8 exhibited the

324 highest biofilm-forming ability in this regard, from the beginning of incubation (Fig.3).
325 However, both strains can be considered as strong biofilm producers (increase of OD by
326 four times) after 48 and 72h of incubation.

327

328 **3.4 Probiotic properties of CLA producer strains**

329 **3.4.1 Bacterial cell surface characteristics and auto-aggregation ability**

330 The hydrophobic/hydrophilic cell surface properties of *Lb. plantarum* L2C21E8 and
331 L3C1E8 were studied using three solvents: n-hexadecane, a non-polar solvent;
332 chloroform, an acidic solvent; and ethyl acetate, a basic solvent. The results, expressed
333 as percentage of microbial adhesion (Fig.4), indicated that both strains presented a more
334 hydrophobic cell surface. Strain L3C1E8 demonstrated stronger ($P<0.05$) affinity for
335 chloroform and n-hexadecane compared to strain L2C21E8. Both strains also showed
336 more affinity ($P<0.05$) for chloroform and n-hexadecane than ethyl acetate, a basic
337 solvent and electron donor. The ability of strains to aggregate was studied and results
338 are also presented in Fig.4. Both strains showed high percentages of auto-aggregation
339 ($>70\%$).

340

341 **3.4.2 Resistance to low pH, bile salt and pancreatin**

342 The survival of the two strains of *Lb. plantarum* under low pH, bile salt and pancreatin
343 was evaluated. As shown in Table 2, both strains were tolerant to bile salts (0.3%, w/v)
344 and pancreatin (0.1%, w/v), each exhibiting no change ($P>0.05$) in viability after 3 h of
345 incubation. However, strain L3C1E8 showed significantly higher resistance ($P<0.05$) to
346 bile salts and pancreatin than L2C21E8. Moreover, strain *Lb. plantarum* L3C1E8 was
347 able to survive at pH 2.5 during 2 h of incubation, although a reduction of viable cell
348 counts was observed after 2 h (from 8.84 ± 0.23 to 2.25 ± 0.14 log CFU/mL). On the

349 contrary, strain *Lb. plantarum* L2C21E8 was susceptible to acidic conditions as no
350 viable cells were found after 0.5 h of incubation.

351

352 **3.4.3 LAB adhesion capacity to intestinal human cells**

353 The *Lb. plantarum* L2C21E8 and L3C1E8 strains were further examined for their
354 ability to adhere to Caco-2 and HT-29 cells. Efficiency of each strain's ability to adhere
355 to the different cell lines are presented in Table 2. In general, both strains displayed high
356 adhesion capacity to both cell lines and no differences ($P>0.05$) were found between
357 strains. Higher adherence ($P<0.05$) was obtained for Caco-2 cell lines, with LAB counts
358 higher than 7 log CFU/mL. Adherence to HT-29 cells was of approx. 6 log CFU/mL for
359 both strains.

360

361 **3.4.4 Effects on adhesion of *E. coli* to HT-29 cells**

362 Results of the anti-adhesion assays (competition, inhibition and displacement) of *E. coli*
363 ATCC 25922 in the presence of *Lb. plantarum* L2C21E8 and L3C1E8 are shown in
364 Fig.5. For the competition assay, when LAB strains and *E. coli* were added
365 simultaneously, both strains were able to reduce adherence of the pathogen in
366 comparison to the untreated control ($> 50\%$). In the inhibition assays, when LAB strains
367 were added before the pathogen, strain *Lb. plantarum* L3C1E8 was the most effective
368 ($P<0.05$), presenting a high degree of inhibition ($>90\%$). Additionally, both strains
369 presented high inhibition percentages ($>75\%$) in the displacement assay when *E. coli*
370 was added to the cells before the LAB strains.

371

372 **4. Discussion**

373 Production of CLA by some LAB and bifidobacteria has been reported in recent years,
374 raising the question of whether CLA production may be regarded as one of the
375 mechanisms by which these bacteria exert some of their health promoting effects
376 (Gorissen et al., 2010; Hennessy et al., 2011; Andrade et al., 2012; O'Shea, Cotter,
377 Stanton, Ross, & Hill, 2012). In this work, LAB isolated from artisanal cheese were
378 screened for their ability to produce CLA. The screening procedure lead to the selection
379 of two CLA producers belonging to *Lb. plantarum* species (strains L2C21E8 and
380 L3C1E8). Similarly, in the screening of LAB isolated from naturally fermented foods,
381 other authors also identified *Lb. plantarum* strains as the highest CLA-producers (Liu et
382 al., 2011; Yang et al., 2014). CLA values observed in the present assay were lower
383 compared to values reported by other authors, but several factors could contribute to
384 this difference, such as temperature, fermentation time, linoleic acid (LA) concentration
385 and other media components (Kuhl & De Dea Lindner, 2016; Ye et al., 2013). Some
386 authors revealed a positive correlation between CLA formation and the ability to
387 tolerate free LA, which suggests that LAB convert LA to CLA as a detoxification
388 mechanism (Adamczak, Bornscheuer, & Bednarski, 2008; Wang, Lv, Chu, Cui, & Ren,
389 2007). In addition, several studies indicated that bacteria usually produced more CLA in
390 whole milk than in MRS medium (Andrade et al., 2012). The CLA produced by the *Lb.*
391 *plantarum* strains under study was mainly found in the supernatant compared with the
392 pellets (cells). This result is in agreement with other studies showing that CLA
393 production is primarily found in the extracellular phase (Rainio, Vahvaselkä,
394 Suomalainen, & Laakso, 2002), though it can also be found in less amounts in the
395 cellular membrane as a structural lipid (Oh et al., 2003).

396 When LA is used as substrate, LAB can convert this fatty acid into 10-hydroxy-12-
397 *trans*-octadecadienoic acid and 10-hydroxy-12-*cis*-octadecadienoic acid, ending mainly

398 with C18:2 *cis*-9, *trans*-11, although other isomers, such as *trans*-10, *cis*-12 C18:2 and
399 *trans*-9, *trans*-11 C18:2, may also be produced in lesser amounts (Kishino et al., 2003;
400 Kuhl & De Dea Lindner, 2016; Ogawa et al., 2005). The high variation of *cis*-9, *trans*-
401 11 C18:2 production in each replication was reflected on the large standard error of the
402 mean (SEM) obtained (Fig. 2). Several studies also revealed a great variability in the
403 CLA isomer profile produced by different LAB strains, although, for most
404 species/strains, *cis*-9, *trans*-11 C18:2 isomer represented more than 70% of the total
405 CLA formed from LA (Kuhl & De Dea Lindner, 2016). In our study, the selected
406 strains of *Lb. plantarum* produced mainly *cis*-9, *trans*-11 C18:2 (approx. 60% of CLA
407 isomers), followed by *trans*-9, *trans*-11 C18:2 (approx. 25% of CLA isomers). In
408 addition, small amounts of other CLA isomers, such as *trans*-10, *cis*-12 (9-11%) and
409 *cis*-9, *cis*-11 (3-5%), were also detected in the supernatant. In the pellet, these ratios
410 were maintained for the predominant isomers (*cis*-9, *trans*-11 and *trans*-9, *trans*-11),
411 but the proportion of *cis*-9, *cis*-11 increased to 9-14% of total CLA, while *trans*-10, *cis*-
412 12 reduced to 5-6%. Most of the studies of CLA isomers produced by *Bifidobacterium*
413 and *Lactobacillus* strains indicated that LA was mainly converted to the *cis*-9, *trans*-11
414 CLA, followed by *trans*-10, *cis*-12 CLA and small amounts of *trans*-9, *trans*-11 CLA
415 isomers (Hennessy et al., 2012; Gorissen et al., 2010; Gorissen, Leroy, De Vuyst, De
416 Smet & Raes, 2015; Rodríguez-Alcalá, Braga, Malcata, Gomes, & Fontecha, 2011).
417 Nevertheless, some *Lb. plantarum* strains were found to produce high proportions of
418 *trans*-9, *trans*-11 C18:2 (Ogawa et al., 2005). In addition, culture conditions, such as
419 low pH and restriction of oxygen, were shown to change the proportion of individual
420 isomers and favor the formation of *trans*, *trans* isomers (Macouzet, Lee & Robert,
421 2008; Panghyová, Kačenová, Matulová & Kiss, 2009).

422 Several studies have revealed that CLAs exert various health benefits, and there is
423 increasing evidence that these effects are isomer specific (O'Shea et al., 2012). Those
424 studies demonstrated that *trans*-9, *trans*-11 C18:2 has a much higher inhibitory and
425 anti-proliferative effect on the growth of the human colon and breast cancer cells, than
426 *cis*-9, *trans*-11 CLA isomer (Coakley et al., 2006; El Roz, Bard, Huvelin, & Nazih,
427 2013). Other studies showed that *cis*-9, *trans*-11 CLA has extra beneficial effects, such
428 as anti-inflammatory and anti-atherogenic effects (Ecker, Liebisch, Patsch, & Schmitz,
429 2009; Loscher et al., 2005). Furthermore, it has been demonstrated that the mixture of
430 the two CLA isomers (*cis*-9, *trans*-11 and *trans*-9, *trans*-11 CLA) had a synergistic anti-
431 proliferation effect on a human colorectal carcinoma cell line (Zhong, Luo, Huang,
432 Deng, & Lei, 2012). Interestingly, the strains of *Lb. plantarum* tested in the present
433 work, presented the highest production of both *cis*-9, *trans*-11 and *trans*-9, *trans*-11
434 CLA isomers, exhibiting a great potential for application in health promoting food
435 products.

436 In the present study, the capacity of the two CLA-producers to form biofilms was also
437 examined. Based on the results obtained, both *Lb. plantarum* strains were able to form a
438 well-structured biofilm. Aoudia et al. (2016) demonstrated that *Lactobacillus* strains are
439 able to form biofilms in a microtiter plate biofilm assay, even under growth conditions
440 mimicking the gastrointestinal environment. Biofilm capacity of these strains can
441 prevent colonization of undesirable microorganisms by covering the epithelial receptors
442 (Martin et al., 2008; Moroni, Kheadr, Boutin, Lacroix, & Fliss, 2006).

443 The two CLA producers were further evaluated for characteristics concerning other
444 relevant probiotic features. These characteristics include auto-aggregation capacity,
445 hydrophilic/hydrophobic surface properties, survival at low pH, resistance to bile salts
446 and pancreatin, as well as, adhesion to different human cell lines. Both strains of *Lb.*

447 *plantarum* studied possess desirable probiotic characteristics, as demonstrated by the in
448 vitro studies. Firstly, we examined the hydrophobicity and auto-aggregation ability of
449 strains. Some cell wall-associated characteristics of probiotics, such as hydrophobicity
450 and auto-aggregation, can contribute to adhesion properties of bacteria to host tissues
451 (Kos et al., 2003; Naidu, Bidlack, & Clemens, 1999; Vinderola & Reinheimer, 2003).
452 Attachment of bacteria to epithelial cells depends on several factors, such as van der
453 Waals attraction, gravitational forces and surface electrostatic charges (Van Loosdrecht,
454 Norde, & Zehnder, 1990). The bacterial surface can be qualitatively assessed as either
455 polar or non-polar, by using solvents with different polarity (Ocana & Nader-Macias,
456 2002; Rosenberg, Gutnick, & Rosenberg, 1980). In our study, the *Lb. plantarum* strains
457 displayed high hydrophobicity values. These results are in agreement with the finding
458 that cell surfaces of lactobacilli are commonly of hydrophobic nature (García-Cayuela
459 et al., 2014). Bacterial aggregation is a desirable property for probiotics and plays an
460 important role in the formation of biofilms (Maria Carmen Collado, Meriluoto, &
461 Salminen, 2008). The two *Lb. plantarum* strains under study demonstrated high
462 capacities to auto-aggregate, conferring them with a survival and proliferation
463 advantage over bacteria lacking this ability (Rickard, Gilbert, High, Kolenbrander, &
464 Handley, 2003). Various lactobacilli have been described in the literature with the
465 ability to form auto-aggregates (García-Ruiz et al., 2014; Hevia et al., 2013; Lozo et al.,
466 2007).

467 Two important characteristics of potentially probiotic strains are resistance to low pH of
468 the stomach and to bile salts and pancreatin secreted into the intestine (Hyronimus, Le
469 Marrec, Hadj Sassi, & Deschamps, 2000; Vinderola & Reinheimer, 2003). In this work,
470 only the strain *Lb. plantarum* L3C1E8 exhibited ability to survive in simulated
471 gastrointestinal conditions (pH 2.5 and 0.3% w/v bile salts, 0.1% w/v pancreatin).

472 Nevertheless, strain *Lb. plantarum* L2C21E8 was also able to survive in the presence of
473 bile salts and pancreatin. With respect to acid resistance, it is reported that bacteria
474 sensitive to gastric juice may have high rates of isolation from feces (Del Piano et al.,
475 2006). Food intake in vivo has been shown to protect bacteria during gastric passage
476 (Saito et al., 2014; Silva et al., 2015). As previously reported, LAB strains were highly
477 resistant to low pH (2.5) after 3 h (survival rate around 100%), when incorporated into
478 fresh cheese (Silva et al., 2015). Therefore, acid sensitive strains can resist
479 gastrointestinal digestion with the use of a food matrix, which can offer protection.

480 *In vitro* studies on the adhesion capabilities of *Lb. plantarum* strains to human epithelial
481 cells (Caco-2 and HT-29 lines) were also conducted. This ability is the most commonly
482 encountered criteria for the selection of probiotic bacteria (Collado, Isolauri, Salminen,
483 & Sanz, 2009; Lebeer, Vanderleyden, & De Keersmaecker, 2008). Implantation in the
484 intestinal mucosa has been considered as the critical feature a strain must possess, in
485 order to influence the intestinal environment (Del Piano et al., 2006). Colonization of
486 the gut mucosa by probiotic strains can provide some beneficial health effects, such as
487 the competitive exclusion of pathogens, modulation of immunity and resistance of
488 probiotic to elimination by peristalsis (Pennacchia, Vaughan, & Villani, 2006;
489 Rinkinen, Westermarck, Salminen, & Ouwehand, 2003). Our results showed that the
490 two *Lb. plantarum* strains studied were able to adhere to Caco-2 and HT-29 cells.
491 Previous studies also reported the capability of adhesion of *Lb. plantarum* strains to
492 these two cell lines (Oguntoyinbo & Narbad, 2015; Saxami et al., 2016). The adhesion
493 capacity of probiotic strains has been identified and characterized, and can be explained
494 by the binding properties of bacterial cell-surface associated proteins with mucus and
495 intestinal cells (Sánchez, Bressollier, & Urdaci, 2008; Vélez, De Keersmaecker, &
496 Vanderleyden, 2007). Another beneficial trait is the interference of probiotic strains

497 with pathogenic adhesion. In this study, both *Lb. plantarum* strains were effective in
498 inhibiting adhesion of the pathogenic bacterium *E. coli*. The high values observed could
499 indicate competition of these strains for the same binding sites on epithelial cells and
500 the overlying mucus layer, in a strain-specific manner (Lee & Puong, 2002; Morrow,
501 Gogineni, & Malesker, 2012). In addition, displacement ability of pre-adhered *E. coli*
502 presented by the *Lb. plantarum* strains studied, could also be due to the production of
503 anti-adhesion factors (Abedi, Feizizadeh, Akbari, & Jafarian-Dehkordi, 2013).

504

505 **5. Conclusion**

506 In conclusion, two *Lb. plantarum* strains presented the ability to produce CLA isomers
507 from free LA, mainly *cis*-9, *trans*-11 CLA and *trans*-9, *trans*-11 CLA, known for
508 having important biological properties. Application of these strains in fermented food
509 products can increase the CLA intake and be beneficial for human health. In addition,
510 these strains presented high adhesion ability to colonic cells and were able to inhibit *E.*
511 *coli* adhesion. These results support the probiotic character of the two strains and their
512 potential to be used in the production of novel functional foods. However, further *in*
513 *vivo* investigations are necessary in order to confirm the role of these potential probiotic
514 strains for promoting human health.

515

516 **Acknowledgements**

517 This work is financed by National funds from FCT – ‘Fundação para a Ciência e
518 Tecnologia’, Project PTDC/ AGR-ALI/104385/2008 and in part by the Science
519 Foundation of Ireland – funded Centre for Science, Engineering and Technology, the
520 APC Microbiome Institute. PhD grant of SC Ribeiro was supported by FRCT - ‘Fundo
521 Regional para a Ciência e Tecnologia’ project M3.1.2/F/011/2011.

522

523 **References**

- 524 Abedi, D., Feizizadeh, S., Akbari, V., & Jafarian-Dehkordi, A. (2013). In vitro anti-
525 bacterial and anti-adherence effects of *Lactobacillus delbrueckii* subsp
526 *bulgaricus* on *Escherichia coli*. *Research in Pharmaceutical Sciences*, 8(4), 260-
527 268.
- 528 Adamczak, M., Bornscheuer, U. T., & Bednarski, W. (2008). Properties and
529 biotechnological methods to produce lipids containing conjugated linoleic acid.
530 *European Journal of Lipid Science and Technology*, 110(6), 491-504. doi:
531 10.1002/ejlt.200800020
- 532 Alonso, L., Cuesta, E. P., & Gilliland, S. E. (2003). Production of free conjugated
533 linoleic acid by *Lactobacillus acidophilus* and *Lactobacillus casei* of human
534 intestinal origin. *Journal of Dairy Science*, 86(6), 1941-1946. doi:
535 [http://dx.doi.org/10.3168/jds.S0022-0302\(03\)73781-3](http://dx.doi.org/10.3168/jds.S0022-0302(03)73781-3)
- 536 Annunziata, A., & Vecchio, R. (2011). Functional foods development in the European
537 market: A consumer perspective. *Journal of Functional Foods*, 3(3), 223-228.
538 doi: <http://dx.doi.org/10.1016/j.jff.2011.03.011>
- 539 Andrade, J. C., Ascensão, K., Gullón, P., Henriques, S. M. S., Pinto, J. M. S., Rocha-
540 Santos, T. A. P., Freitas, A. C., & Gomes, A. M. (2012). Production of
541 conjugated linoleic acid by food-grade bacteria: A review. *International Journal*
542 *of Dairy Technology*. 65(4) 467-481. doi: 10.1111/j.1471-0307.2012.00871.x
- 543 Aoudia, N., Rieu, A., Briandet, R., Deschamps, J., Chluba, J., Jegou, G., Guzzo, J.
544 (2016). Biofilms of *Lactobacillus plantarum* and *Lactobacillus fermentum*:
545 Effect on stress responses, antagonistic effects on pathogen growth and

- 546 immunomodulatory properties. *Food Microbiology*, 53, 51-59. doi:
547 10.1016/j.fm.2015.04.009
- 548 Argyri, A. A., Zoumpopoulou, G., Karatzas, K.-A. G., Tsakalidou, E., Nychas, G.-J. E.,
549 Panagou, E. Z., & Tassou, C. C. (2013). Selection of potential probiotic lactic
550 acid bacteria from fermented olives by in vitro tests. *Food Microbiology*, 33(2),
551 282-291. doi: <http://dx.doi.org/10.1016/j.fm.2012.10.005>
- 552 Barrett, E., Ross, R. P., Fitzgerald, G. F., & Stanton, C. (2007). Rapid screening method
553 for analyzing the conjugated linoleic acid production capabilities of bacterial
554 cultures. *Applied and Environmental Microbiology*, 73(7), 2333-2337. doi:
555 10.1128/aem.01855-06
- 556 Bassaganya-Riera, J., Viladomiu, M., Pedragosa, M., De Simone, C., Carbo, A.,
557 Shaykhutdinov, R., Jobin, C., Arthur, J. C., Corl, B. A., Vogel, H., Storr, M., &
558 Hontecillas, R. (2012). Probiotic bacteria produce conjugated linoleic acid
559 locally in the gut that targets macrophage PPAR gamma to suppress colitis. *Plos*
560 *One*, 7. doi: 10.1371/journal.pone.0031238
- 561 Bellon-Fontaine, M. N., Rault, J., & van Oss, C. J. (1996). Microbial adhesion to
562 solvents: a novel method to determine the electron-donor/electron-acceptor or
563 Lewis acid-base properties of microbial cells. *Colloids and Surfaces B:*
564 *Biointerfaces*, 7(1-2), 47-53. doi: [http://dx.doi.org/10.1016/0927-](http://dx.doi.org/10.1016/0927-7765(96)01272-6)
565 [7765\(96\)01272-6](http://dx.doi.org/10.1016/0927-7765(96)01272-6)
- 566 Bigliardi, B., & Galati, F. (2013). Innovation trends in the food industry: The case of
567 functional foods. *Trends in Food Science & Technology*, 31(2), 118-129. doi:
568 <http://dx.doi.org/10.1016/j.tifs.2013.03.006>
- 569 Chin, S. F., Liu, W., Storkson, J. M., Ha, Y. L., & Pariza, M. W. (1992). Dietary
570 sources of conjugated dienoic isomers of linoleic acid, a newly recognized class

- 571 of anticarcinogens. *Journal of Food Composition and Analysis*, 5(3), 185-197.
572 doi: [http://dx.doi.org/10.1016/0889-1575\(92\)90037-K](http://dx.doi.org/10.1016/0889-1575(92)90037-K)
- 573 Chinnadurai, K., Kanwal, H., Tyagi, A., Stanton, C., & Ross, P. (2013). High
574 conjugated linoleic acid enriched ghee (clarified butter) increases the antioxidant
575 and antiatherogenic potency in female Wistar rats. *Lipids in Health and Disease*,
576 12(1), 121. doi: 10.1186/1476-511X-12-121.
- 577 Chung, S. H., Kim, I. H., Park, H. G., Kang, H. S., Yoon, C. S., Jeong, H. Y., Kim, Y. J.
578 (2008). Synthesis of conjugated linoleic acid by human-derived *Bifidobacterium*
579 *breve* LMC 017: Utilization as a functional starter culture for milk fermentation.
580 *Journal of Agricultural and Food Chemistry*, 56(9), 3311-3316. doi:
581 10.1021/jf0730789
- 582 Coakley, M., Johnson, M. C., McGrath, E., Rahman, S., Ross, R. P., Fitzgerald, G. F.,
583 Stanton, C. (2006). Intestinal bifidobacteria that produce *trans*-9, *trans*-11
584 conjugated linoleic acid: a fatty acid with antiproliferative activity against
585 human colon SW480 and HT-29 cancer cells. *Nutrition and Cancer*, 56(1), 95-
586 102. doi: 10.1207/s15327914nc5601_13
- 587 Coakley, M., Ross, R. P., Nordgren, M., Fitzgerald, G., Devery, R., & Stanton, C.
588 (2003). Conjugated linoleic acid biosynthesis by human-derived
589 *Bifidobacterium* species. *Journal of Applied Microbiology*, 94(1), 138-145. doi:
590 10.1046/j.1365-2672.2003.01814.x
- 591 Collado, M. C., Isolauri, E., Salminen, S., & Sanz, Y. (2009). The impact of probiotic
592 on gut health. *Current Drug Metabolism*, 10(1), 68-78. doi:
593 <http://dx.doi.org/10.2174/138920009787048437>

- 594 Collado, M. C., Meriluoto, J., & Salminen, S. (2008). Adhesion and aggregation
595 properties of probiotic and pathogen strains. *European Food Research and*
596 *Technology*, 226(5), 1065-1073. doi: 10.1007/s00217-007-0632-x
- 597 Del Piano, M., Morelli, L., Strozzi, G. P., Allesina, S., Barba, M., Deidda, F., Capurso,
598 L. Probiotics: from research to consumer. *Digestive and Liver Disease*, 38,
599 S248-S255. doi: 10.1016/S1590-8658(07)60004-8
- 600 Domingos-Lopes, M. F. P., Stanton, C., Ross, P. R., Dapkevicius, M. L. E., & Silva, C.
601 C. G. (2017). Genetic diversity, safety and technological characterization of
602 lactic acid bacteria isolated from artisanal Pico cheese. *Food Microbiology*, 63,
603 178-190. doi:10.1016/j.fm.2016.11.014
- 604 Ecker, J., Liebisch, G., Patsch, W., & Schmitz, G. (2009). The conjugated linoleic acid
605 isomer *trans*-9,*trans*-11 is a dietary occurring agonist of liver X receptor α .
606 *Biochemical and Biophysical Research Communications*, 388(4), 660-666. doi:
607 <http://dx.doi.org/10.1016/j.bbrc.2009.08.048>
- 608 El Roz, A., Bard, J. M., Huvelin, J. M., & Nazih, H. (2013). The anti-proliferative and
609 pro-apoptotic effects of the *trans*9,*trans*11 conjugated linoleic acid isomer on
610 MCF-7 breast cancer cells are associated with LXR activation. *Prostaglandins,*
611 *Leukotrienes and Essential Fatty Acids (PLEFA)*, 88(4), 265-272. doi:
612 <http://dx.doi.org/10.1016/j.plefa.2012.12.006>
- 613 Fuke, G., & Nornberg, J. L. (2017). Systematic evaluation on the effectiveness of
614 conjugated linoleic acid in human health. *Critical Reviews in Food Science and*
615 *Nutrition*, 57(1), 1-7. doi:10.1080/10408398.2012.716800
- 616 García-Cayuela, T., Korany, A. M., Bustos, I., P. Gómez de Cadiñanos, L., Requena, T.,
617 Peláez, C., & Martínez-Cuesta, M. C. (2014). Adhesion abilities of dairy
618 *Lactobacillus plantarum* strains showing an aggregation phenotype. *Food*

- 619 *Research International*, 57, 44-50. doi:
620 <http://dx.doi.org/10.1016/j.foodres.2014.01.010>
- 621 García-Ruiz, A., González de Llano, D., Esteban-Fernández, A., Requena, T.,
622 Bartolomé, B., & Moreno-Arribas, M. V. (2014). Assessment of probiotic
623 properties in lactic acid bacteria isolated from wine. *Food Microbiology*, 44,
624 220-225. doi: <http://dx.doi.org/10.1016/j.fm.2014.06.015>
- 625 Gaullier, J.-M., Halse, J., Høivik, H. O., Høye, K., Syvertsen, C., Nurminiemi, M.,
626 Gudmundsen, O. (2007). Six months supplementation with conjugated linoleic
627 acid induces regional-specific fat mass decreases in overweight and obese.
628 *British Journal of Nutrition*, 97(03), 550-560. doi:
629 [doi:10.1017/S0007114507381324](http://dx.doi.org/10.1017/S0007114507381324)
- 630 Gorissen, L., Leroy, F., De Vuyst, L., De Smet, S., & Raes, K. (2015). Bacterial
631 production of conjugated linoleic and linolenic acid in foods: A technological
632 challenge. *Critical Reviews in Food Science and Nutrition*, 55(11), 1561-1574.
633 doi: [10.1080/10408398.2012.706243](http://dx.doi.org/10.1080/10408398.2012.706243)
- 634 Gorissen, L., Raes, K., Weckx, S., Dannenberger, D., Leroy, F., De Vuyst, L., & De
635 Smet, S. (2010). Production of conjugated linoleic acid and conjugated linolenic
636 acid isomers by *Bifidobacterium* species. *Applied Microbiology and
637 Biotechnology*, 87(6), 2257-2266. doi: [10.1007/s00253-010-2713-1](http://dx.doi.org/10.1007/s00253-010-2713-1)
- 638 Hennessy, A. A., Barrett, E., Paul Ross, R., Fitzgerald, G. F., Devery, R., & Stanton, C.
639 (2012). the production of conjugated α -linolenic, γ -linolenic and stearidonic
640 acids by strains of bifidobacteria and propionibacteria. *Lipids*, 47(3), 313-327.
641 doi: [10.1007/s11745-011-3636-z](http://dx.doi.org/10.1007/s11745-011-3636-z)

- 642 Hennessy, A. A., Ross, R. P., Devery, R., & Stanton, C. (2011). The health promoting
643 properties of the conjugated isomers of α -linolenic acid. *Lipids*, 46(2), 105-119.
644 doi: 10.1007/s11745-010-3501-5
- 645 Hevia, A., Martínez, N., Ladero, V., Álvarez, M. A., Margolles, A., & Sánchez, B.
646 (2013). An extracellular serine/threonine-rich protein from *Lactobacillus*
647 *plantarum* NCIMB 8826 is a novel aggregation-promoting factor with affinity to
648 mucin. *Applied and Environmental Microbiology*, 79(19), 6059-6066. doi:
649 10.1128/aem.01657-13
- 650 Hyronimus, B., Le Marrec, C., Hadj Sassi, A., & Deschamps, A. (2000). Acid and bile
651 tolerance of spore-forming lactic acid bacteria. *International Journal of Food*
652 *Microbiology*, 61(2-3), 193-197. doi: <http://dx.doi.org/10.1016/S0168->
653 1605(00)00366-4
- 654 Jiang, Björck, & Fondén. (1998). Production of conjugated linoleic acid by dairy starter
655 cultures. *Journal of Applied Microbiology*, 85(1), 95-102. doi: 10.1046/j.1365-
656 2672.1998.00481.x
- 657 Kishino, S., Ogawa, J., Ando, A., Iwashita, T., Fujita, T., Kawashima, H., & Shimizu,
658 S. (2003). Structural analysis of conjugated linoleic acid produced by
659 *Lactobacillus plantarum*, and factors affecting isomer production. *Bioscience*
660 *Biotechnology and Biochemistry*, 67(1), 179-182.
- 661 Kobaa, K., & Yanagita, T. (2014). Health benefits of conjugated linoleic acid (CLA).
662 *Obesity Research & Clinical Practice*, 8, e525-e532.
- 663 Kos, B., Šušković, J., Vuković, S., Šimpraga, M., Frece, J., & Matošić, S. (2003).
664 Adhesion and aggregation ability of probiotic strain *Lactobacillus acidophilus*
665 M92. *Journal of Applied Microbiology*, 94(6), 981-987. doi: 10.1046/j.1365-
666 2672.2003.01915.x

- 667 Kuhl, C. G., & De Dea Lindner, J. (2016). Biohydrogenation of linoleic acid by lactic
668 acid bacteria for the production of functional cultured dairy products: A review.
669 *Foods*, 5(1). doi: 10.3390/foods5010013
- 670 Lebeer, S., Vanderleyden, J., & De Keersmaecker, S. C. J. (2008). Genes and molecules
671 of lactobacilli supporting probiotic action. *Microbiology and Molecular Biology*
672 *Reviews*, 72(4), 728-+. doi: 10.1128/mmbr.00017-08
- 673 Lee, Y. K., & Puong, K. Y. (2002). Competition for adhesion between probiotics and
674 human gastrointestinal pathogens in the presence of carbohydrate. *British*
675 *Journal of Nutrition*, 88, S101-S108. doi: UNSP
676 BJN88101.sgm10.1079/bjn2002635
- 677 Lin, T. Y., & Lee, F. J. (1997). Conjugated linoleic acid as affected by food source and
678 processing. *Scientific Agriculture*, 45(9,10), 284-295.
- 679 Lin, W.-H., Yu, B., Jang, S.-H., & Tsen, H.-Y. (2007). Different probiotic properties for
680 *Lactobacillus fermentum* strains isolated from swine and poultry. *Anaerobe*,
681 13(3-4), 107-113. doi: 10.1016/j.anaerobe.2007.04.006
- 682 Loscher, C. E., Draper, E., Leavy, O., Kelleher, D., Mills, K. H. G., & Roche, H. M.
683 (2005). Conjugated linoleic acid suppresses NF- κ B activation and IL-12
684 production in dendritic cells through ERK-mediated IL-10 induction. *The*
685 *Journal of Immunology*, 175(8), 4990-4998. doi: 10.4049/jimmunol.175.8.4990
- 686 Lozo, J., Jovcic, B., Kojic, M., Dalgalarrodo, M., Chobert, J.-M., Haertlé, T., &
687 Topisirovic, L. (2007). Molecular characterization of a novel bacteriocin and an
688 unusually large aggregation factor of *Lactobacillus paracasei* subsp. *paracasei*
689 BGSJ2-8, a natural isolate from homemade cheese. *Current Microbiology*,
690 55(3), 266-271. doi: 10.1007/s00284-007-0159-1

- 691 Macouzet, M., Lee, B. H., & Robert, N. (2009). Production of conjugated linoleic acid
692 by probiotic *Lactobacillus acidophilus* La-5. *Journal of Applied Microbiology*,
693 106(6), 1886-1891. doi: 10.1111/j.1365-2672.2009.04164.x
- 694 Martin, R., Soberon, N., Vaneechoutte, M., Florez, A. B., Vazquez, F., & Suarez, J. E.
695 (2008). Characterization of indigenous vaginal lactobacilli from healthy women
696 as probiotic candidates. *International Microbiology: The official journal of the*
697 *Spanish Society for Microbiology*, 11(4), 261-266.
- 698 Moroni, O., Kheadr, E., Boutin, Y., Lacroix, C., & Fliss, I. (2006). Inactivation of
699 adhesion and invasion of food-borne *Listeria monocytogenes* by bacteriocin-
700 producing *Bifidobacterium* strains of human origin. *Applied and Environmental*
701 *Microbiology*, 72(11), 6894-6901. doi: 10.1128/aem.00928-06
- 702 Morrow, L. E., Gogineni, V., & Malesker, M. A. (2012). Probiotics in the intensive care
703 unit. *Nutrition in Clinical Practice*, 27(2), 235-241. doi:
704 10.1177/0884533612440290
- 705 Naidu, A. S., Bidlack, W. R., & Clemens, R. A. (1999). Probiotic spectra of lactic acid
706 bacteria (LAB). *Critical Reviews in Food Science and Nutrition*, 39(1), 13-126.
707 doi: 10.1080/10408699991279187
- 708 O'Shea, E. F., Cotter, P. D., Stanton, C., Ross, R. P., & Hill, C. (2012). Production of
709 bioactive substances by intestinal bacteria as a basis for explaining probiotic
710 mechanisms: Bacteriocins and conjugated linoleic acid. *International Journal of*
711 *Food Microbiology*, 152(3), 189-205. doi:
712 <http://dx.doi.org/10.1016/j.ijfoodmicro.2011.05.025>
- 713 Ocana, V. S., & Nader-Macias, M. E. (2002). Vaginal lactobacilli: self- and co-
714 aggregating ability. *British Journal of Biomedical Science*, 59(4), 183-190.

- 715 Ogawa, J., Kishino, S., Ando, A., Sugimoto, S., Mihara, K., & Shimizu, S. (2005).
716 Production of conjugated fatty acids by lactic acid bacteria. *Journal of*
717 *Bioscience and Bioengineering*, 100(4), 355-364. doi:
718 <http://dx.doi.org/10.1263/jbb.100.355>
- 719 Oguntoyinbo, F. A., & Narbad, A. (2015). Multifunctional properties of *Lactobacillus*
720 *plantarum* strains isolated from fermented cereal foods. *Journal of Functional*
721 *Foods*, 17, 621-631. doi: 10.1016/j.jff.2015.06.022
- 722 Oh, D.-K., Hong, G.-H., Lee, Y., Min, S., Sin, H.-S., & Cho, S. K. (2003). Production
723 of conjugated linoleic acid by isolated *Bifidobacterium* strains. *World Journal of*
724 *Microbiology and Biotechnology*, 19(9), 907-912. doi:
725 10.1023/B:WIBI.0000007313.90368.0c
- 726 Ozer, C. O., Kilic, B., & Kilic, G. B. (2016). In-vitro microbial production of
727 conjugated linoleic acid by probiotic *L. plantarum* strains: Utilization as a
728 functional starter culture in sucuk fermentation. *Meat Science*, 114, 24-31.
729 doi:10.1016/j.meatsci.2015.12.005
- 730 Pandit, A., Anand, S., Kalscheur, K., & Hassan, A. (2012). Production of conjugated
731 linoleic acid by lactic acid bacteria in milk without any additional substrate.
732 *International Journal of Dairy Technology*, 65(4), 603-608. doi: 10.1111/j.1471-
733 0307.2012.00870.x
- 734 Panghyová, E., Kačenová, D., Matulová, M., & KISS, E. (2009). Composition of
735 conjugated linoleic acid isomers formed by *Lactobacillus* and *Bifidobacterium*
736 spp. in conversion media. *Journal of Food and Nutrition Research*, 48(4), 163-
737 170.
- 738 Pennacchia, C., Vaughan, E. E., & Villani, F. (2006). Potential probiotic *Lactobacillus*
739 strains from fermented sausages: Further investigations on their probiotic

- 740 properties. *Meat Science*, 73(1), 90-101. doi:
741 <http://dx.doi.org/10.1016/j.meatsci.2005.10.019>
- 742 Pérez Ibarreche, M., Castellano, P., & Vignolo, G. (2014). Evaluation of anti-*Listeria*
743 meat borne *Lactobacillus* for biofilm formation on selected abiotic surfaces.
744 *Meat Science*, 96(1), 295-303. doi:
745 <http://dx.doi.org/10.1016/j.meatsci.2013.07.010>
- 746 Raimondi, S., Amaretti, A., Leonardi, A., Quartieri, A., Gozzoli, C., & Rossi, M.
747 (2016). Conjugated Linoleic Acid Production by Bifidobacteria: Screening,
748 Kinetic, and Composition. *Biomed Research International*.
749 doi:10.1155/2016/8654317
- 750 Rainio, A., Vahvaselkä, M., Suomalainen, T., & Laakso, S. (2002). Production of
751 conjugated linoleic acid by *Propionibacterium freudenreichii* ssp. *shermanii*.
752 *Lait*, 82(1), 91-101.
- 753 Rickard, A. H., Gilbert, P., High, N. J., Kolenbrander, P. E., & Handle y, P. S. (2003)
754 Bacterial coaggregation: an integral process in the development of multi-species
755 biofilms. *Trends in Microbiology*, 11(2), 94-100. doi: 10.1016/S0966-
756 842X(02)00034-3
- 757 Rinkinen, M., Westermarck, E., Salminen, S., & Ouwehand, A. C. (2003). Absence of
758 host specificity for in vitro adhesion of probiotic lactic acid bacteria to intestinal
759 mucus. *Veterinary Microbiology*, 97(1-2), 55-61. doi: 10.1016/s0378-
760 1135(03)00183-4
- 761 Riquelme, C., Câmara, S., Enes Dapkevicius, M. d. L. N., Vinuesa, P., da Silva, C. C.
762 G., Malcata, F. X., & Rego, O. A. (2015). Characterization of the bacterial
763 biodiversity in Pico cheese (an artisanal Azorean food). *International Journal of*

- 764 *Food Microbiology*, 192(0), 86-94. doi:
765 <http://dx.doi.org/10.1016/j.ijfoodmicro.2014.09.031>
- 766 Rodríguez-Alcalá, L. M., Braga, T., Xavier Malcata, F., Gomes, A., & Fontecha, J.
767 (2011). Quantitative and qualitative determination of CLA produced by
768 *Bifidobacterium* and lactic acid bacteria by combining spectrophotometric and
769 Ag+-HPLC techniques. *Food Chemistry*, 125(4), 1373-1378. doi:
770 <http://dx.doi.org/10.1016/j.foodchem.2010.10.008>
- 771 Rosenberg, M., Gutnick, D., & Rosenberg, E. (1980). Adherence of bacteria to
772 hydrocarbons - a simple method for measuring cell-surface hydrophobicity.
773 *FEMS Microbiology Letters*, 9(1), 29-33. doi: 10.1111/j.1574-
774 6968.1980.tb05599.x
- 775 Saito, V. S. T., dos Santos, T. F., Vinderola, C. G., Romano, C., Nicoli, J. R., Araújo, L.
776 S., Uetanabaro, A. P. T. (2014). Viability and resistance of lactobacilli isolated
777 from cocoa fermentation to simulated gastrointestinal digestive steps in soy
778 yogurt. *Journal of Food Science*, 79(2), M208-M213. doi: 10.1111/1750-
779 3841.12326
- 780 Salas-Jara, M., Ilabaca, A., Vega, M., & García, A. (2016). Biofilm forming
781 *Lactobacillus*: New challenges for the development of probiotics.
782 *Microorganisms*, 4(3), 35.
- 783 Sánchez, B., Bressollier, P., & Urdaci, M. C. (2008). Exported proteins in probiotic
784 bacteria: adhesion to intestinal surfaces, host immunomodulation and molecular
785 cross-talking with the host. *FEMS Immunology & Medical Microbiology*, 54(1),
786 1-17. <http://dx.doi.org/10.1111/j.1574-695X.2008.00454.x>
- 787 Saxami, G., Karapetsas, A., Lamprianidou, E., Kotsianidis, I., Chlichlia, A., Tassou, C.,
788 Galanis, A. (2016). Two potential probiotic *Lactobacillus* strains isolated from

- 789 olive microbiota exhibit adhesion and anti-proliferative effects in cancer cell
790 lines. *Journal of Functional Foods*, 24, 461-471. doi: 10.1016/j.jff.2016.04.036
- 791 Shen, W., Chuang, C.-C., Martinez, K., Reid, T., Brown, J. M., Xi, L., McIntosh, M.
792 (2013). Conjugated linoleic acid reduces adiposity and increases markers of
793 browning and inflammation in white adipose tissue of mice. *Journal of Lipid*
794 *Research*, 54(4), 909-922. doi: 10.1194/jlr.M030924
- 795 Shingfield, K. J., Bonnet, M., & Scollan, N. D. (2013). Recent developments in altering
796 the fatty acid composition of ruminant-derived foods. *Animal*, 7, 132-162.
797 doi:10.1017/s1751731112001681
- 798 Silva, C. C. G., Domingos-Lopes, M. F. P., Magalhães, V. A. F., Freitas, D. A. S. R.,
799 Coelho, M. C., Rosa, H. J. D., & Dapkevicius, M. L. N. E. (2015). Latin-style
800 fresh cheese enhances lactic acid bacteria survival but not *Listeria*
801 *monocytogenes* resistance under in vitro simulated gastrointestinal conditions.
802 *Journal of Dairy Science*, 98(7), 4377-4383. doi: 10.3168/jds.2015-9420
- 803 Sluijs, I., Plantinga, Y., de Roos, B., Mennen, L. I., & Bots, M. L. (2010). Dietary
804 supplementation with *cis-9,trans-11* conjugated linoleic acid and aortic stiffness
805 in overweight and obese adults. *The American Journal of Clinical Nutrition*,
806 91(1), 175-183. doi: 10.3945/ajcn.2009.28192
- 807 Todorov, S. D., & Dicks, L. M. T. (2008). Evaluation of lactic acid bacteria from kefir,
808 molasses and olive brine as possible probiotics based on physiological
809 properties. *Annals of Microbiology* 58, 661-670.
- 810 Van Loosdrecht, M. C. M., Norde, W., & Zehnder, A. J. B. (1990). Physical chemical
811 description of bacterial adhesion. *Journal of Biomaterials Applications*, 5(2), 91-
812 106. doi: 10.1177/088532829000500202

- 813 Vélez, M. P., De Keersmaecker, S. C. J., & Vanderleyden, J. (2007). Adherence factors
814 of *Lactobacillus* in the human gastrointestinal tract. *FEMS Microbiology*
815 *Letters*, 276(2), 140-148. <http://dx.doi.org/10.1111/j.1574-6968.2007.00908.x>
- 816 Verna, E. C., & Lucak, S. (2010). Use of probiotics in gastrointestinal disorders: what
817 to recommend? *Therapeutic Advances in Gastroenterology*, 3(5), 307-319. doi:
818 10.1177/1756283X10373814
- 819 Vieira, C. P., Cabral, C. C., Lima, B., Paschoalin, V. M. F., Leandro, K. C., & Conte, C.
820 A. (2017). *Lactococcus lactis* ssp *cremoris* MRS47, a potential probiotic strain
821 isolated from kefir grains, increases cis-9, trans-11-CLA and PUFA contents in
822 fermented milk. *Journal of Functional Foods*, 31, 172-178.
823 doi:10.1016/j.jff.2017.01.047
- 824 Vinderola, C. G., & Reinheimer, J. A. (2003). Lactic acid starter and probiotic bacteria:
825 a comparative “in vitro” study of probiotic characteristics and biological barrier
826 resistance. *Food Research International*, 36(9–10), 895-904. doi:
827 [http://dx.doi.org/10.1016/S0963-9969\(03\)00098-X](http://dx.doi.org/10.1016/S0963-9969(03)00098-X)
- 828 Wall, R., Ross, R. P., Fitzgerald, G. F., & Stanton, C. (2008). Microbial conjugated
829 linoleic acid production - a novel probiotic trait? *Food Science & Technology*
830 *Bulletin Functional Foods*, 4(8), 87-97.
- 831 Wang, L.-M., Lv, J.-P., Chu, Z.-Q., Cui, Y.-Y., & Ren, X.-H. (2007). Production of
832 conjugated linoleic acid by *Propionibacterium freudenreichii*. *Food Chemistry*,
833 103(2), 313-318. doi: <http://dx.doi.org/10.1016/j.foodchem.2006.07.065>
- 834 Yang, B., Chen, H., Gu, Z., Tian, F., Ross, R. P., Stanton, C., Zhang, H. (2014).
835 Synthesis of conjugated linoleic acid by the linoleate isomerase complex in
836 food-derived lactobacilli. *Journal of Applied Microbiology*, 117(2), 430-439.
837 doi: 10.1111/jam.12524

- 838 Ye, S., Yu, T., Yang, H., Li, L., Wang, H., Xiao, S., & Wang, J. (2013). Optimal culture
839 conditions for producing conjugated linoleic acid in skim-milk by co-culture of
840 different *Lactobacillus* strains. *Annals of Microbiology*, 63(2), 707-717. doi:
841 10.1007/s13213-012-0523-7
- 842 Zeng, Z., Lin, J., & Gong, D. (2009). Identification of lactic acid bacterial strains with
843 high conjugated linoleic acid-producing ability from natural sauerkraut
844 fermentations. *Journal of Food Science*, 74(4), M154-M158. doi:
845 10.1111/j.1750-3841.2009.01123.x
- 846 Zhong, X.-F., Luo, T., Huang, G.-D., Deng, Z.-Y., & Lei, L. (2012). Equimolar mixture
847 of c9,t11 and t9,t11 CLA inhibits the growth and induces apoptosis in Caco-2
848 cells. *European Journal of Lipid Science and Technology*, 114(4), 479-485. doi:
849 10.1002/ejlt.201100338
- 850

Figure captions

Fig. 1. GC chromatograms of fatty acids of cell supernatant fluid of *Lb. plantarum* L2C21E8 (A) and *Lb. plantarum* L3C1E8 (B) grown in MRS plus linoleic acid, after 48h.

Fig. 2. Isomers profiles ($\mu\text{g/mL}$) of supernatant and pellet of *Lb. plantarum* L2C21E8 (A) and *Lb. plantarum* L3C1E8 (B) in MRS broth supplemented with 0.5 mg/mL of linoleic acid after 48 h of incubation. Error bars indicate standard error of the mean (SEM) from four different experiments. CLA isomers marked with different letters differ significantly ($P < 0.05$).

Fig. 3. Biofilm formation of *Lactobacillus plantarum* L2C21E8 and L3C1E8 after incubation at 30 °C for 24, 48 and 72h. Data are expressed as mean \pm SEM. The cut-off (ODC) was defined as the mean OD value of the negative control. Based on the OD, strains were classified as non-biofilm producers ($\text{OD} \leq \text{ODC}$), weak ($\text{ODC} < \text{OD} \leq 2 \times \text{ODC}$), moderate ($2 \times \text{ODC} < \text{OD} < 4 \times \text{ODC}$) or strong biofilm producers ($4 \times \text{ODC} < \text{OD}$). Different letters within each incubation time represent significant differences ($P < 0.05$) between strains.

Fig. 4. Percentages of microbial adhesion to solvents and auto-aggregation of strains *Lb. plantarum* L2C21E8 and L3C1E8. Results are shown as the average (\pm SEM) of three independent experiments. Different letters within each solvent represent significant differences ($P < 0.05$) between strains.

Fig. 5. Competition, inhibition and displacement of *E. coli* ATCC 25922 in the presence of *Lb. plantarum* L2C21E8 and *Lb. plantarum* L3C1E8. The data is shown as the average (\pm SEM) of three independent experiments and inhibition of *E. coli* adhesion is expressed as percentage ratios. Different letters represent significant differences ($P < 0.05$) between strains.

Table 1- LAB screened for potential CLA production in MRS broth after 48 h. Values of *cis*-9, *trans*-11 CLA concentration ($\mu\text{g/mL}$) and percentage of conversion are indicated. Values of CLA concentration are means \pm SEM of two replicates.

Strains	Accession No.	cis9. trans11 CLA ($\mu\text{g/mL}$)	% Conversion
<i>Lactobacillus paracasei</i>			
L2A21R9	KM096813	3.66 \pm 0.09 ^{a,b}	0.73
L2A1K8	KM096814	2.15 \pm 0.05 ^a	0.43
L2B21R1a	KM096816	2.56 \pm 0.17 ^a	0.51
L2B21R3	KM096817	2.21 \pm 0.01 ^a	0.44
L2B1K8	KM096818	3.65 \pm 0.12 ^{a,b}	0.73
L3A21R8	KM096819	2.73 \pm 0.41 ^a	0.55
L3B1M2	KM096820	2.17 \pm 0.05 ^a	0.43
L3B21R1	KM096821	3.00 \pm 0.29 ^a	0.60
L3B21R2	KM096822	9.16 \pm 1.72 ^b	1.83
L3B21R7	KM096823	3.28 \pm 0.19 ^a	0.66
L3B1K1	KM096824	3.43 \pm 0.75 ^{a,b}	0.69
L3B21K4	KM096825	9.96 \pm 1.30 ^b	1.99
L3C21M6	KM096826	7.12 \pm 0.76 ^b	1.42
L3C1K8	KM096827	3.10 \pm 0.55 ^a	0.62
<i>Lactobacillus otakiensis</i>			
L3C1R1	KM096828	4.31 \pm 0.78 ^{a,b}	0.86
<i>Lactobacillus plantarum</i>			
L2B21R1b	KM103932	3.32 \pm 0.20 ^a	0.66
L2C21E8	KM103933	17.94 \pm 0.13 ^c	3.59
L2A21R1	KM103931	7.00 \pm 2.10 ^b	1.40
L3A21R6	KM103934	2.24 \pm 0.09 ^a	0.45
L3C1E8	KM079361	15.36 \pm 0.15 ^c	3.07
<i>Lactobacillus paraplantarum</i>			
L2B21R5	KM079360	4.29 \pm 0.24 ^{a,b}	0.86
<i>Lactococcus lactis</i>			
L3B1M7	KM079358	2.77 \pm 0.08 ^a	0.55
L3A21M1	KF193424	5.24 \pm 0.75 ^{a,b}	1.05
<i>Lactococcus garvieae</i>			
L3B1M8	KM079359	2.78 \pm 0.30 ^a	0.56
<i>Leuconostoc mesenteroides</i>			
L2A21E7	KM079353	2.06 \pm 0.00 ^a	0.00
L2B21E3	KM079354	2.52 \pm 0.07 ^a	0.50
L3A21M4	KM079355	4.21 \pm 0.20 ^{a,b}	0.84
<i>Leuconostoc citreum</i>			
L3C1E7	KM079357	2.97 \pm 0.07 ^a	0.59
<i>Enterococcus faecalis</i>			
L2B21K3	KF193420	2.42 \pm 0.22 ^a	0.48
L3A1M6	KF193421	2.16 \pm 0.07 ^a	0.43
L3A21M3	KF193425	2.44 \pm 0.27 ^a	0.49
L3A21M8	KF193426	2.20 \pm 0.07 ^a	0.44
L3A21K6	KF193422	4.66 \pm 0.43 ^{a,b}	0.93
L3A21K7	KF193423	3.38 \pm 0.52 ^{a,b}	0.68
L3B1K3	KF193427	3.37 \pm 0.32 ^{a,b}	0.67

* Values of CLA labelled with different letters are significantly different ($P < 0.05$).

Table 2- Probiotic characteristics of CLA producing strains. Resistance to low pH (2.5), bile salts (0.3%, w/v) and pancreatin (0.1%, w/v) and adhesion to Caco-2 and HT-29 cells. Results are presented as the average values \pm SEM from four independent experiments for resistance to low pH, bile salts and pancreatin assays, and three independent experiments for adhesion assays.

Probiotic characteristics	Time (h)	Bacterial counts (log CFU/mL)*	
		<i>Lb. plantarum</i> L2C21E8	<i>Lb. plantarum</i> L3C1E8
Resistance to pH 2.5	0	6.74 \pm 0.24 ^{aA}	8.84 \pm 0.23 ^{AB}
	0.5	ND ^{bA}	7.95 \pm 0.72 ^{a,bB}
	1	ND ^{bA}	5.74 \pm 0.0 ^{bB}
	2	ND ^{bA}	2.25 \pm 0.14 ^{cB}
	3	ND ^b	ND ^c
		<i>Lb. plantarum</i> L2C21E8	<i>Lb. plantarum</i> L3C1E8
Resistance to bile salts (0.3%. w/v) and pancreatin (0.1%.w/v)	0	6.04 \pm 0.31 ^A	7.39 \pm 0.26 ^B
	0.5	5.86 \pm 0.01 ^A	6.89 \pm 0.25 ^B
	1	5.98 \pm 0.03 ^A	7.38 \pm 0.19 ^B
	2	6.25 \pm 0.29 ^A	7.42 \pm 0.23 ^B
	3	5.88 \pm 0.07 ^A	7.35 \pm 0.22 ^B
		<i>Lb. plantarum</i> L2C21E8	<i>Lb. plantarum</i> L3C1E8
Adhesion to cells	Caco-2	7.36 \pm 0.07 ^a	7.66 \pm 0.21 ^a
	HT-29	5.77 \pm 0.58 ^b	5.73 \pm 0.12 ^b


* ND: Counts of bacteria below detection limits.

Different lowercase letters indicate significant differences ($P < 0.05$) within columns (among time or cells, Caco-2 and HT-29), according to Bonferroni post hoc means comparison test.

Different uppercase letters indicate significant differences ($P < 0.05$) within rows (among strains), according to Bonferroni post hoc means comparison test.

Fig.2

A



B




Fig.3




Fig.4





Fig.5



Highlights

- LAB strains isolated from cheese were screened for *in vitro* CLA production.
- Two *Lactobacillus plantarum* strains were selected as high CLA producers.
- Strains produced a high proportion (25%) of the uncommon *trans*-9, *trans*-11 CLA.
- CLA producer strains were evaluated for probiotic characteristics.
- CLA producers also inhibited of *E. coli* adhesion to human cells.