

Is cows' qualitatively assessed behaviour towards humans related to their general stress level?

Silvia Ivemeyer¹, Asja Ebinghaus¹, Christel Simantke¹, Rupert Palme², Ute Knierim¹

¹ Farm Animal Behaviour and Husbandry Section, Faculty of Organic Agricultural Sciences, Nordbahnhofstr. 1a, 37213 Witzenhausen, Germany

² Department of Biomedical Sciences, University of Veterinary Medicine, 1210 Vienna, Austria

Aim

To investigate the relation between cows' qualitatively assessed behaviour towards humans (QBA) and their general medium-term stress level (measured by faecal cortisol metabolites, FCM)

Animals and Methods

- QBA (Fig. 1) and FCM (Fig. 2) of 316 cows (≤ 200 days in milk) on 25 organic farms
- QBA: fixed list of 20 descriptors, specifically developed for this purpose (Fig. 3)
- FCM: competitive enzyme immunoassay measuring 11,17 dioxoandrostanes in ng/g fresh faeces
- Statistics:
 - analysis of QBA data by principal component analyses (PCA); PC1 reflected valence, PC2 activation (Fig. 3)
 - division of sample into 4 groups regarding combination of valence ('pos' / 'neg') and activation ('low' / 'high') (Fig. 4)
 - group comparisons: Kruskal-Wallis test and Wilcoxon post hoc tests




Fig. 3: Component plot of PCA (no rotation, eigenvalue > 1) on QBA data (n=622 cows) (Ebinghaus et al. 2017: Appl Animal Behav Sci 196:22-29)



Fig. 1: Qualitative behaviour assessment (QBA)

related to a standardized human-animal interaction



Fig. 2: Faecal cortisol metabolites (FCM)

faeces collected freshly on the same day, reflecting adrenocortical activity over several hours




Fig. 4: Sample divided into 4 groups regarding valence and activation level of cows responses towards humans using PC1 and PC2 medians as cut-points (n=316 cows)

Results

- FCM medians and variation were generally on a low level (Fig. 5)
- cows reacting 'pos_high' during human-animal interaction (n=65), 'pos_low' (n=93) and 'neg_low' (n=99) had similar

- 'neg_high' cows (n=59) had significantly lower FCM levels than all other groups




Fig. 5: Differences in stress level measured by FCM between QBA groups (n=316 cows)

Conclusions

- 'neg_high' cows presumably had higher fear levels towards humans than 'pos' cows, and expressed this more actively than 'neg_low' cows
- unexpectedly, 'neg' reactions were not related to a higher stress level
- other factors such as social rank or health status might have affected results more profoundly, requiring a more complex analysis including individual factors

Acknowledgements

Sincere thanks to the farmers for project participation. Project has been funded by the Federal Ministry of Food and Agriculture (BMEL) based on a decision of the Parliament of the Federal Republic of Germany via the Federal Office for Agriculture and Food (BLE) under the Federal Organic Farming Scheme (BÖLN) ('ORGANICDAIRYHEALTH' project) and the Hessen State Ministry of Higher Education, Research and Arts ('LOEWE' project).



Contact: Dr. Silvia Ivemeyer
ivemeyer@uni-kassel.de



Gefördert durch:
Bundesministerium für Ernährung und Landwirtschaft



UNIKASSEL
VERSITÄT
ORGANIC
AGRICULTURAL
SCIENCES