

<https://helda.helsinki.fi>

Targeting beta 1-integrin inhibits vascular leakage in endotoxemia

Hakanpaa, Laura

2018-07-10

Hakanpaa , L , Kiss , E A , Jacquemet , G , Miinalainen , I , Lerche , M , Guzman , C , Mervaala , E , Eklund , L , Ivaska , J & Saharinen , P 2018 , ' Targeting beta 1-integrin inhibits vascular leakage in endotoxemia ' , Proceedings of the National Academy of Sciences of the United States of America , vol. 115 , no. 28 , pp. E6467-E6476 . <https://doi.org/10.1073/pnas.1722317115>

<http://hdl.handle.net/10138/304366>

<https://doi.org/10.1073/pnas.1722317115>

unspecified

publishedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.



Targeting β 1-integrin inhibits vascular leakage in endotoxemia

Laura Hakanpää^a, Elina A. Kiss^a, Guillaume Jacquemet^b, Ilkka Miinalainen^c, Martina Lerche^b, Camilo Guzmán^b, Eero Mervaala^d, Lauri Eklund^e, Johanna Ivaska^{b,f}, and Pipsa Saharinen^{a,g,h,1}

^aTranslational Cancer Biology Program, Research Programs Unit, Biomedicum Helsinki, University of Helsinki, FI-00014 Helsinki, Finland; ^bTurku Centre for Biotechnology, University of Turku and Åbo Akademi University, FI-20520 Turku, Finland; ^cBiocenter Oulu, University of Oulu, FI-90014 Oulu, Finland; ^dDepartment of Pharmacology, Faculty of Medicine, University of Helsinki, FI-00014 Helsinki, Finland; ^eOulu Center for Cell-Matrix Research, Faculty of Biochemistry and Molecular Medicine, Biocenter Oulu, FI-90014 Oulu, Finland; ^fDepartment of Biochemistry, University of Turku, FI-20520 Turku, Finland; ^gWihuri Research Institute, Biomedicum Helsinki, FI-00290 Helsinki, Finland; and ^hDepartment of Biochemistry and Developmental Biology, Faculty of Medicine, University of Helsinki, FI-00014 Helsinki, Finland

Edited by Rakesh K. Jain, Harvard Medical School and Massachusetts General Hospital, Boston, MA, and approved May 29, 2018 (received for review December 21, 2017)

Loss of endothelial integrity promotes capillary leakage in numerous diseases, including sepsis, but there are no effective therapies for preserving endothelial barrier function. Angiopoietin-2 (ANGPT2) is a context-dependent regulator of vascular leakage that signals via both endothelial TEK receptor tyrosine kinase (TIE2) and integrins. Here, we show that antibodies against β 1-integrin decrease LPS-induced vascular leakage in murine endotoxemia, as either a preventative or an intervention therapy. β 1-integrin inhibiting antibodies bound to the vascular endothelium in vivo improved the integrity of endothelial cell-cell junctions and protected mice from endotoxemia-associated cardiac failure, without affecting endothelial inflammation, serum proinflammatory cytokine levels, or TIE receptor signaling. Moreover, conditional deletion of a single allele of endothelial β 1-integrin protected mice from LPS-induced vascular leakage. In endothelial monolayers, the inflammatory agents thrombin, lipopolysaccharide (LPS), and IL-1 β decreased junctional vascular endothelial (VE)-cadherin and induced actin stress fibers via β 1- and α 5-integrins and ANGPT2. Additionally, β 1-integrin inhibiting antibodies prevented inflammation-induced endothelial cell contractility and monolayer permeability. Mechanistically, the inflammatory agents stimulated ANGPT2-dependent translocation of α 5 β 1-integrin into tensin-1-positive fibrillar adhesions, which destabilized the endothelial monolayer. Thus, β 1-integrin promotes endothelial barrier disruption during inflammation, and targeting β 1-integrin signaling could serve as a novel means of blocking pathological vascular leak.

β 1-integrin | ANGPT2 | TIE2 | permeability | sepsis

Vascular stability is crucial for normal tissue homeostasis whereas decreased vascular integrity is a hallmark of serious pathological conditions, such as sepsis, acute respiratory distress syndrome (ARDS), fatal dengue fever, and malaria. Capillary leak syndromes are also a common toxicity of certain immunotherapies. A burst in the generation of proinflammatory agents, as a part of an overwhelmed host response, contributes to endothelial barrier breakdown in these diseases, promoting shock and multiorgan failure in patients (1, 2). The underlying molecular mechanisms behind pathological capillary leakage remain incompletely understood, hindering the development of targeted therapies aimed at vascular stabilization (3).

Endothelial barrier function is maintained via a dynamic coordination of endothelial cell (EC)-cell junctions, the contractile actin cytoskeleton, and EC adhesion to the underlying basement membrane in response to extracellular stimuli. Proinflammatory cytokines, microbial components such as lipopolysaccharide (LPS), thrombin, and vascular endothelial growth factor (VEGF) induce endothelial permeability by binding to their cognate receptors on ECs. These inflammatory agents activate signals that converge on common downstream pathways, including dismantling of vascular endothelial (VE)-cadherin (Cadherin 5, CDH5) from EC-EC junctions (4, 5). Small GTPases of the Rho family and myosin light chain (MLC) kinase (MLCK) also stimulate remodeling of the

cortical actin cytoskeleton of stable endothelial monolayers into contractile actin stress fibers, generating cellular tension after inflammatory stimulation (4). Furthermore, the EC-derived growth factor angiopoietin-2 (ANG2, ANGPT2), which is up-regulated in diseases associated with inflammation and vascular leakage, synergizes with multiple inflammatory agents, promoting endothelial permeability (6, 7).

Integrins, transmembrane receptors that exist as heterodimers of α - and β -subunits, couple ECs to the underlying extracellular matrix (ECM) through their ectodomains while their intracellular domains engage the actin cytoskeleton by recruiting actin-binding proteins. In focal adhesions, integrins connect to the actin cytoskeleton and cellular signaling pathways through linker and adapter proteins, such as paxillin, talin, vinculin, and α -actinin (8). Collectively, integrin-mediated EC adhesion to the vascular basement membrane is known to mediate vascular stability and angiogenesis (9, 10), but less is known about the regulation of integrin-coupled EC adhesions during vascular leakage in mature vessels.

The α 5 β 1-integrin heterodimer is a major endothelial fibronectin receptor. The α 5- and β 1-integrins and fibronectin are essential for vascular development (10–13). During embryonic development, β 1-integrin regulates vascular sprouting and vessel lumen formation (11, 12). β 1-integrin continues to play a critical

Significance

Compromised vascular integrity is associated with capillary leakage in sepsis, but effective therapies stabilizing the vasculature are lacking. Here, we show that targeting β 1-integrin in vivo with inhibitory antibodies or deletion of a single allele of endothelial β 1-integrin inhibits lipopolysaccharide (LPS)-induced vascular leakage in murine endotoxemia. The inflammatory agents IL-1 β , thrombin, and LPS induced changes in endothelial cell-extracellular matrix (ECM) adhesion via β 1-integrin, angiopoietin-2, and the adapter protein tensin-1, leading to increased endothelial cell contractility and permeability. These results indicate that β 1-integrin actively promotes vascular leakage and that targeting β 1-integrin signaling could be a novel means of achieving vascular stabilization in pathological vascular leak.

Author contributions: L.H., E.A.K., G.J., E.M., L.E., J.I., and P.S. designed research; L.H., E.A.K., G.J., I.M., M.L., C.G., L.E., and P.S. performed research; L.H., E.A.K., G.J., M.L., C.G., L.E., J.I., and P.S. analyzed data; and L.H., E.A.K., J.I., and P.S. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

This open access article is distributed under [Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 \(CC BY-NC-ND\)](https://creativecommons.org/licenses/by-nc-nd/4.0/).

¹To whom correspondence should be addressed. Email: pipsa.saharinen@helsinki.fi.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1722317115/-DCSupplemental.

Published online June 25, 2018.

role in vessel integrity during postnatal life, via stabilization of VE-cadherin in the developing EC–EC junctions of newly formed vessels (10). However, the function of $\beta 1$ -integrin in the mature vasculature and during vascular leakage has received less attention. Interestingly, $\alpha 5\beta 1$ -integrin signaling has been recently linked to chronic inflammation as it was shown to mediate harmful endothelial remodeling in a mouse atherosclerosis model (14). Furthermore, ANGPT2 was recently reported to activate $\alpha 5\beta 1$ -integrin, leading to endothelial monolayer destabilization (15). However, it is not known if $\beta 1$ -integrin-dependent EC-basement membrane adhesions contribute to acute endothelial permeability responses in vivo.

Here, we demonstrate that several inflammatory agents, including LPS, interleukin (IL)- 1β , and thrombin, disrupt the endothelial barrier function via $\beta 1$ -integrin. Mechanistically, inflammatory agents stimulated, in an ANGPT2-dependent manner, the formation of $\beta 1$ -integrin- and tensin-1-positive fibrillar adhesions that destabilized endothelial monolayer integrity. Inhibitory antibodies against $\beta 1$ -integrin decreased inflammation-induced endothelial contractility and monolayer permeability. In LPS-induced endotoxemia in mice, $\beta 1$ -integrin antibodies decreased vascular leakage, protected from cardiac failure, and stabilized vascular integrity, without affecting the up-regulation of inflammatory markers or TEK receptor tyrosine kinase (TIE2) signaling. Further supporting a role for endothelial $\beta 1$ -integrin in vascular leakage, conditional heterozygous deletion of endothelial $\beta 1$ -integrin protected adult mice from LPS-induced vascular leakage. Thus, $\beta 1$ -integrin signaling promotes vascular leakage during inflammation and may be a potential therapeutic target for vascular stabilization in disease.

Results

Inflammatory Agents Decrease Endothelial Monolayer Integrity via $\beta 1$ -Integrin. To investigate how EC adhesion to the ECM affects inflammation-induced monolayer permeability, primary human dermal blood microvascular ECs (BECs) and human umbilical vein ECs (HUVECs) were stimulated with the inflammatory agents thrombin, IL- 1β , and LPS. $\beta 1$ -integrin, which is the major integrin subunit expressed by ECs (11, 12, 15), was simultaneously inhibited with $\beta 1$ -integrin antibody (mAb13). mAb13 binds to the $\beta 1$ -domain in the headpiece of $\beta 1$ -integrin, resulting in stabilization of the $\beta 1$ -integrin in an inactive, closed conformation with decreased binding to ECM proteins (16). Consistent with this, mAb13 ($10 \mu\text{g}\cdot\text{mL}^{-1}$) inhibited EC spreading on fibronectin, and even stronger inhibition was obtained when mAb13 was combined with cilengitide, a cyclic Arg-Gly-Asp (RGD) containing pentapeptide that inhibits $\alpha\beta 3$ -, $\alpha\beta 5$ -, and $\alpha 5\beta 1$ -integrins (SI Appendix, Fig. S1 A and B). Similarly, incubation with a high concentration of mAb13 ($10 \mu\text{g}\cdot\text{mL}^{-1}$) interfered with cell junction organization (SI Appendix, Fig. S1C) whereas a low mAb13 concentration ($0.1 \mu\text{g}\cdot\text{mL}^{-1}$) preserved EC monolayer integrity and was used in subsequent studies.

Stimulation of BECs with thrombin ($0.1 \text{ U}\cdot\text{mL}^{-1}$ for 30 min) resulted in a significant decrease in VE-cadherin in EC junctions (Fig. 1A), in line with previous reports (17). Whereas treatment with mAb13 ($0.1 \mu\text{g}\cdot\text{mL}^{-1}$) had no effect on VE-cadherin in unstimulated control cells, it prevented the thrombin-induced loss of VE-cadherin (Fig. 1A and B). BEC monolayer permeability was analyzed using noninvasive electrical cell impedance monitoring. Stimulation of BECs with thrombin increased monolayer permeability, in accordance with previous publications (17), but this was prevented by mAb13 (Fig. 1C).

A balance between the activities of MLCK and MLC phosphatase (MLCP) determines the level of MLC phosphorylation and therefore the contractile status of ECs. Thrombin induces MLC phosphorylation, leading to endothelial contractility (4). In line with this, blebbistatin, a reversible inhibitor of nonmuscle myosin II, prevented thrombin-induced VE-cadherin loss from EC junctions (Fig. 1D). We next performed traction force microscopy (TFM) to measure the mechanical stress exerted by the EC monolayer. Stimulation of HUVECs with thrombin ($0.1 \text{ U}\cdot\text{mL}^{-1}$, for 10 or 20 min) increased the force exerted on the matrix threefold, indicating a disrupted cell monolayer (Fig. 1E and F). Pretreatment

of the cells with mAb13 before thrombin treatment significantly attenuated the thrombin-induced increase in cellular force (Fig. 1E and F). These results indicate that mAb13 decreased thrombin-induced EC contractility, which contributes to the dissolution of VE-cadherin from EC junctions, leading to monolayer permeability.

To confirm these results, shRNA-expressing lentiviruses were used to silence $\beta 1$ -integrin (sh $\beta 1$) in BECs. Successful $\beta 1$ -integrin silencing was confirmed by Western blotting and by quantitative PCR [96% mRNA silencing compared with scrambled shRNA (shScr)] (SI Appendix, Fig. S2). However, $\sim 10\%$ of active $\beta 1$ -integrin remained in focal adhesions (mostly located at the cell periphery overlapping with the cortical actin cytoskeleton), even 48 h after sh $\beta 1$ transduction (Fig. 2A and B). sh $\beta 1$ did not decrease VE-cadherin in confluent monolayers, but prevented the thrombin-induced loss of VE-cadherin, enabling its retention in BEC junctions, as in the untreated, shScr-transduced control cells (Fig. 2C and D and SI Appendix, Fig. S3A). The thrombin-induced decrease in EC barrier function was also less in the sh $\beta 1$ -transduced cells than in the shScr-transduced control cells (Fig. 2E). By contrast, $\beta 3$ -integrin (sh $\beta 3$) silencing decreased VE-cadherin in BECs, in line with previous studies (9, 15), and sh $\beta 3$ did not prevent the thrombin-induced loss of VE-cadherin (Fig. 2C and D and SI Appendix, Fig. S3A). However, silencing of ANGPT2 (shANGPT2), which is constitutively secreted by ECs, prevented the thrombin-induced loss of VE-cadherin, in line with previous studies (Fig. 2C and D and SI Appendix, Fig. S3A) (17).

Thrombin is a fast inflammatory mediator acting via G protein-coupled protease-activated receptors whereas IL- 1β and LPS stimulate slower EC responses via the IL-1 receptor and Toll-like receptor 4, respectively (1, 18). Despite acting through distinct cell surface receptors, thrombin, IL- 1β , and LPS activate partially overlapping downstream signaling pathways. We thus hypothesized that $\beta 1$ -integrin may play a more universal role in inflammatory monolayer destabilization and investigated whether $\beta 1$ -integrin mediated signals downstream of IL- 1β and LPS as well. Stimulation of BECs with IL- 1β ($10 \text{ ng}\cdot\text{mL}^{-1}$) decreased VE-cadherin in BEC junctions at 2 h (SI Appendix, Fig. S4); however, this did not occur in sh $\beta 1$ - or shANGPT2-transduced BECs (Fig. 2F and SI Appendix, Fig. S3B and C). LPS ($10 \mu\text{g}\cdot\text{mL}^{-1}$, 2 h)-induced actin stress fibers were significantly decreased after silencing of $\beta 1$ -integrin in BECs (SI Appendix, Fig. S5). In addition, shRNA silencing of $\alpha 5$ -integrin (sh $\alpha 5$), which pairs with $\beta 1$ -integrin to form the fibronectin receptor $\alpha 5\beta 1$, decreased stress fiber formation in LPS-stimulated BECs (SI Appendix, Fig. S5), suggesting that both $\alpha 5$ -integrin and $\beta 1$ -integrin are involved in inflammation-induced monolayer destabilization. Moreover, expression of the tight junction protein zonula occludens (ZO)-1 was decreased after IL- 1β and LPS stimulation in control BECs but was preserved in ANGPT2-, $\alpha 5$ -integrin-, or $\beta 1$ -integrin-silenced cells (Fig. 2G and SI Appendix, Fig. S3D). By contrast, silencing of $\beta 3$ -integrin had no protective effect (Fig. 2F and SI Appendix, Figs. S3C and S5A), and actin stress fibers were prominent in $\beta 3$ -integrin-silenced BECs, both in the absence and presence of LPS (SI Appendix, Fig. S5B and C).

To confirm our results in ECs of different origin, primary human pulmonary microvascular endothelial cells (HPMECs) were used. Stimulation with thrombin or IL- 1β disrupted the cortical actin rim and VE-cadherin junctions in shScr- but not sh $\beta 1$ - or shANGPT2-transduced HPMECs (SI Appendix, Fig. S6). These results indicate that several inflammatory agents that act via distinct cell surface receptors induce EC monolayer destabilization in a $\beta 1$ -integrin- and ANGPT2-dependent, but not $\beta 3$ -integrin-dependent, manner. In addition, inflammation-induced EC permeability and contractility can be prevented by $\beta 1$ -integrin inhibiting antibodies, without affecting the monolayer stability in basal conditions.

$\beta 1$ -Integrin Antibody Reduces Vascular Leakage in LPS-Induced Endotoxemia. To investigate whether antibody targeting of $\beta 1$ -integrin also improved the integrity of EC–EC junctions in vivo, we evaluated vascular leakage in a murine model of endotoxemia.




Fig. 1. β 1-integrin antibody inhibits thrombin-induced endothelial monolayer destabilization. (A) BECs were treated with a control antibody (Ctrl Ab) or a β 1-integrin inhibiting antibody (mAb13) (both 0.1 $\mu\text{g}\cdot\text{mL}^{-1}$ for 5 min), stimulated with thrombin (Thr) (0.1 U·mL⁻¹ for 30 min), and fixed and stained for VE-cadherin or filamentous actin (F-Actin). (B) Quantification of VE-cadherin staining (150 cells per treatment). Data are the mean \pm SD of n = 3 independent experiments). (C) BECs were analyzed for electrical cell impedance (represented as the cell index). Stimulation with Ab and thrombin (1 U·mL⁻¹) was initiated once the impedance reached saturating levels (set as time 0). The mean \pm SD from a representative experiment is shown (four wells per treatment, P = 0.026). (D) BECs were treated with S-Blebbistatin (20 min) and then with thrombin (0.1 U·mL⁻¹, 20 min), and fixed and stained for VE-cadherin and F-actin. (E) Representative traction force maps exerted by HUVECs plated on fibronectin-coated polyacrylamide gels. Images were taken at baseline, at 5 min after the addition of mAb13, and after 10 and 20 min of thrombin stimulation (0.1 U·mL⁻¹). (F) Quantification of the mean force (strain energy, pJ). Data are the mean \pm SD of n = 3 independent experiments. *P < 0.05, **P < 0.01, and ***P < 0.001 (Tukey's test). Representative images of maximum intensity projections of confocal z-stacks obtained using a 63 \times objective are shown. Nuclear DAPI stain. (Scale bars: 20 μm .)




Fig. 2. Silencing of ANGPT2, β 1-integrin, or α 5-integrin, but not β 3-integrin, inhibits thrombin, IL-1 β , or LPS-induced endothelial monolayer destabilization. BECs transfected with shScr or sh β 1 lentiviral vectors (A and B) were stimulated with thrombin (1 U·mL⁻¹ for 30 min) and stained for active β 1-integrin (12G10) and F-actin (A). (B) Quantification of 12G10 staining (350 cells per treatment, $n = 3$ independent experiments). BECs transfected with shScr, sh β 1, sh β 3, shANGPT2, or sh α 5, as indicated, were stimulated with thrombin (1 U·mL⁻¹ for 30 min) (C–E), IL-1 β (10 ng·mL⁻¹ for 2 h) (F and G), or LPS (2.5 μ g·mL⁻¹ for 2 h) (G) and stained for VE-cadherin (C, D, and F) or ZO-1 (G). (C and F) Quantification of VE-cadherin staining (120–300 cells per treatment, $n = 3$ independent experiments). (E) BECs transfected with shScr or sh β 1 were analyzed for electrical cell impedance (represented as the cell index). Cells were stimulated with thrombin (1 U·mL⁻¹) once the impedance reached saturating levels (set as time 0). A representative experiment is shown (four wells per treatment, mean \pm SD, $P = 0.0036$). (G) Quantification of ZO-1-positive junction length (130 to 145 cells per treatment $n = 3$ independent experiments). Data are the mean \pm SD of $n = 3$ independent experiments. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, and **** $P < 0.0001$ (Tukey's test). Representative images of maximum intensity projections of confocal z-stacks obtained using a 63 \times objective are shown. Nuclear DAPI stain. (Scale bars: 20 μ m).

A monoclonal antibody against mouse β 1-integrin (HM β 1) (19) inhibited the spreading of murine ECs on fibronectin, similarly to mAb13-mediated inhibition of human ECs (SI Appendix, Fig. S7). Control antibodies or HM β 1 was injected [both at 2.5 mg·kg⁻¹, intraperitoneally (i.p.)] into mice 24 h before LPS administration, and vascular leakage was analyzed 16 h after LPS administration and 4 min after i.v. injection of fluorospheres (diameter of 100 nm, which is roughly equivalent to a protein molecular mass of 1,000 kDa). Vascular leakage was significantly increased in the tracheal and dermal vasculature in LPS-treated mice that had received control antibodies (Fig. 3 A–C and SI Appendix, Fig. S8). However, HM β 1 decreased the LPS-induced vascular leakage significantly without affecting the baseline vascular permeability in PBS-treated mice (Fig. 3 A–C and SI Appendix, Fig. S8).

To determine whether HM β 1 inhibited the permeability of lower molecular mass substances, we used 70-kDa fluorescent dextran (the approximate molecular mass of albumin). Importantly, HM β 1, but not control antibodies, significantly decreased the leakage of 70-kDa dextran in the tracheas of LPS-treated mice, indicating that HM β 1 reduced the permeability of inflamed vessels to large as well as smaller molecular mass substances (Fig. 3 D and E).

Capillary leakage can lead to multiorgan failure in sepsis, affecting organs such as the heart. LPS administration for 16 h decreased the cardiac output, as measured by echocardiography before and after LPS administration. Pretreatment with HM β 1, but not with control antibodies, significantly improved the cardiac output, as shown by the measurement of ejection fraction (EF) and fractional shortening (FS) (Fig. 3 F and G). This suggests that the decreased vascular leakage in mice receiving HM β 1 has a physiologically significant role in protecting heart function.

β 1-Integrin Antibody Reduces Vascular Leakage After the Onset of Acute Inflammation. LPS induced a strong inflammatory cytokine response by increasing the circulating levels of IL-6, IL-1 β , TNF- α , and ANGPT2 (Fig. 4A and SI Appendix, Fig. S9). IL-6, IL-1 β , and TNF- α peaked between 1 and 3 h after LPS administration whereas ANGPT2 was increased between 3 and 16 h after LPS administration, which coincides with the decrease in *Angpt1* expression (Fig. 4A) (20, 21). The induction of proinflammatory cytokines in response to LPS was not decreased in mice receiving

the HM β 1 antibody, indicating that HM β 1 did not diminish the LPS-induced host cytokine response (Fig. 4A and SI Appendix, Fig. S9).

Next, we tested the effect of HM β 1 on vascular leakage after the onset of systemic inflammation, which is more clinically relevant than the preventative model of HM β 1 administration. In the intervention experiment, HM β 1 or control antibodies were injected into mice 2 h after administration of LPS, when the levels of IL-6, IL-1 β , and TNF- α were already increased in the circulation and in the lungs, and vascular leakage was increased (SI Appendix, Fig. S10). Importantly, vascular leakage was significantly decreased 16 h after LPS injection (i.e., 14 h after antibody administration) in mice treated with HM β 1, but not the control antibody (Fig. 4B), indicating that HM β 1 resolved the vascular leakage.

β 1-Integrin Antibody Decreases Vascular Leakage Without Affecting Endothelial Inflammation. LPS administration to mice induced a threefold increase in the expression of vascular cell adhesion molecule (VCAM)-1 in the lungs at 16 h, but this was not affected by HM β 1 treatment (Fig. 4 C–E and SI Appendix, Fig. S11 A and B). LPS also increased neutrophil numbers in the lungs, which were also not significantly altered by the HM β 1 antibody (Fig. 4 F and G and SI Appendix, Fig. S11 C and D). Furthermore, a neutralizing antibody against α 4-integrin, which pairs with β 1-integrin to form very late antigen-4 (VLA-4), which mediates leukocyte adhesion to the inflamed endothelium via VCAM-1 (22), did not reduce LPS-induced vascular leakage (SI Appendix, Fig. S12).

The ANGPT-TIE signaling system regulates vascular stability during inflammation. It was recently demonstrated that TIE1 on the endothelium of blood vessels is cleaved after LPS administration and during chronic infection in mice, releasing the soluble TIE1 (sTIE1) ectodomain into the circulation. This correlated with loss of ANGPT2 agonist activity and increased vascular leakage (21, 23). Consistent with these results, sTIE1 was detected in the serum 3 h after LPS administration (SI Appendix, Fig. S13 A and B), and TIE1 and TIE2 protein levels were decreased by 80% and 50%, respectively, in lung lysates 16 h after LPS administration (Fig. 4 H and I and SI Appendix, Fig. S13C). However, HM β 1 did not prevent the TIE receptor loss. Furthermore, HM β 1 did not prevent the LPS-induced decrease in phospho-TIE2 (Fig. 4J).




Fig. 3. β 1-integrin antibody reduces vascular leakage and improves cardiac output in LPS-induced murine endotoxemia. (A) Analysis of leakage of 100-nm fluorospheres in mice treated with β 1-integrin (HM β 1) or control antibodies (Ctrl Ab) for 24 h, followed by LPS or PBS for 16 h. Shown are representative maximum intensity projections of tile-scanned confocal z-stacks of tracheal whole mounts stained for CD31 (green), obtained using a 10 \times objective. Quantification of fluorosphere area in the trachea (B) and ear (C, corresponding micrographs in *SI Appendix*, Fig. S8), $n = 7$ to 8 mice per group. (D) Vascular leakage in mice treated as in A was measured using Texas red-conjugated 70-kDa Dextran. Shown are representative maximum intensity projections of confocal z-stacks of tracheal whole mounts stained for VE-cadherin (green), obtained using a 10 \times objective. (E) Quantification of fluorescent dextran area in the trachea, $n = 3$ to 6 mice per group. (F and G) Echocardiographic analysis of mice treated with HM β 1 or Ctrl Ab for 24 h. Heart function was recorded before and 16 h after administration of LPS or PBS. Ejection fraction (F) and fractional shortening values (G) are presented. $n = 8$ to 13 mice per group. Data are the mean \pm SD. * $P < 0.05$, ** $P < 0.01$, and *** $P < 0.001$ (Tukey's test). (Scale bars: 100 μ m.)

Altogether, HM β 1 improved endothelial barrier function independently of the inflammation-induced adhesion molecule up-regulation, neutrophil infiltration, and loss of endothelial TIE receptor signaling.

β 1-Integrin Antibody Improves the Integrity of Endothelial Cell–Cell Junctions in Endotoxemia. To investigate whether the decreased vessel leakiness induced by HM β 1 after LPS administration was due to improved EC–EC junctions, endothelial junctions in the pulmonary veins, the site of the greatest amount of inflammatory cell margination, were analyzed by transmission electron microscopy (TEM). LPS treatment decreased the actin-rich, electron-dense EC junctions in the pulmonary vasculature, in comparison with PBS-treated control mice (expressed as the ratio of the electron-dense area to the total EC–EC contact length) (Fig. 5A–C and *SI Appendix*, Fig. S14). However, treatment with HM β 1 significantly increased actin-rich junctions in the blood vessels of

LPS-challenged mice (Fig. 5A–C). Similar results were obtained when capillaries, postcapillary venules, and collecting venules in the trachea were analyzed (Fig. 5D). These results indicate that antibody targeting of β 1-integrin in endotoxemia has a vascular stabilizing effect that is mediated via improved EC junction integrity.

Interestingly, HM β 1 was found to be localized to the vascular endothelial cell layer, overlapping with CD31 staining, but less with staining for the pericyte marker desmin, 40 h after its administration (i.e., at the end point of the analysis). The results indicate that HM β 1 homes to the vasculature for extended periods, may concentrate at sites of vascular leakage, and exerts its vascular protective effects on the endothelium (*SI Appendix*, Fig. S15). Furthermore, heterozygous *Igbl¹^{W11 Δ EC};Pdgfb-iCreERT2* mice, in which endothelial β 1-integrin was deleted after birth or in adulthood, were protected from LPS-induced vascular leakage, indicating that endothelial β 1-integrin decreases vessel integrity during inflammation (Fig. 6 and *SI Appendix*, Figs. S16 and S17).




Fig. 4. β 1-integrin antibody reduces vascular leakage after the onset of systemic inflammation. (A) Quantification of circulating levels of IL-6, IL-1 β , TNF- α , and ANGPT2 in the serum and *Angpt1* mRNA in the lungs of mice challenged with HM β 1 or control antibodies for 24 h, followed by LPS or PBS for the indicated times ($n = 3$ mice per group). (B) HM β 1 or control antibodies were administered 2 h after the administration of LPS or PBS. Quantification of 100-nm fluorescent spheres in the vasculature of the trachea and ear 14 h after antibody treatment ($n = 5$ to 6 mice per group). (C–E) Mice were challenged with HM β 1 or control antibodies for 24 h, followed by LPS or PBS for 16 h. VCAM-1 (D and E) ($n = 6$ mice per group) and Ly6G/C (F and G) ($n = 3$ mice per group) staining and quantification in the lungs. Western blot analysis of VCAM-1 (C) ($n = 6$ mice per group), TIE1 (H) ($n = 3$ mice per group), TIE2 (I) ($n = 6$ mice per group), and phospho-TIE2 (J) ($n = 6$ mice per group) in mouse lung lysates. Dashed line indicates where lanes were cropped together from a single membrane. Data are the mean \pm SD. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, and **** $P < 0.0001$ (Tukey's test). (Scale bars: 100 μ m.)

Inflammatory Agents Promote α 5 β 1-Integrin Translocation into Tensin-1-Positive Fibrillar Adhesions in an ANGPT2-Dependent Manner. To understand more about the mechanism through which endothelial β 1-integrin induces permeability, we investigated the dynamics of EC–ECM adhesions after inflammatory stimulation of

BECs. In control cells, focal adhesions dominated in the cell periphery whereas thrombin stimulation resulted in rapid formation of centrally located cell adhesions (Fig. 7A and B and Movie S1). Similarly, upon stimulation with IL-1 β (Fig. 7C–E and SI Appendix, Fig. S18), thrombin (Fig. 7F and SI Appendix, Fig. S18), or LPS




Fig. 6. Heterozygous deletion of endothelial $\beta 1$ -integrin protects from vascular leakage. Vascular leakage of 100-nm fluorospheres was measured in LPS- or PBS-treated Cre-positive tamoxifen-induced *Itgb1*^{WT/ΔEC} (*Itgb1*^{WT/ΔEC}) mice or littermate controls (Ctrl) after 16 h. (A) Representative maximum intensity projections of tile-scanned confocal z-stacks of tracheal whole mounts stained for CD31 (green) obtained using a 10 \times objective. Quantification of fluorosphere area in the trachea (B) and ear skin (C). Ctrl + PBS group: $n = 15$ (6 WT, 5 *Itgb1*^{WT/ΔEC}, 4 *Pdgfb-iCreER*^{T2} mice). Ctrl + LPS group: $n = 10$ (4 WT, 5 *Itgb1*^{WT/ΔEC}, 1 *Pdgfb-iCreER*^{T2} mouse). *Itgb1*^{WT/ΔEC} + PBS group: $n = 7$. *Itgb1*^{WT/ΔEC} + LPS group: $n = 9$. Data are the mean \pm SD. * $P < 0.05$, ** $P < 0.01$, and *** $P < 0.001$ (Tukey's test). (Scale bar: 100 μ m.)

and the thrombin-induced calcium response (7, 17). We previously showed that, in addition to TIE2, ANGPT2 can signal via $\alpha 5\beta 1$ -integrin, leading to destabilization of EC monolayers where TIE2 levels are low (15). Here, we found that ANGPT2 expression in ECs is required for LPS-, IL-1 β -, and thrombin-induced monolayer destabilization, for the formation of tensin-1-positive fibrillar adhesions, and for $\alpha 5\beta 1$ -integrin recycling to these adhesions. Thus, our results provide evidence that ANGPT2 may mediate vascular leakage, in part via its ability to regulate EC-ECM adhesions. In line with this conclusion, transgenic overexpression of ANGPT2 in mice leads to aberrant and decreased EC-basement membrane matrix interactions (27, 28).

$\beta 1$ -integrin is a key regulator of blood vessel development (11, 12), and it continues to play a critical, context-dependent role in vessel sprouting and stability during postnatal life (10). However, the function of $\beta 1$ -integrin in the mature vasculature of adult mice has not been investigated. We found that heterozygous deletion of $\beta 1$ -integrin in ECs did not compromise vascular integrity but protected the mice from LPS-induced vascular leakage. In addition, short-term use of the $\beta 1$ -integrin inhibitory

antibody did not affect EC junction integrity or induce vascular leakage in adult mice in the absence of inflammatory stimuli. This indicates that the inhibitory antibody did not affect the vascular stabilizing functions of $\beta 1$ -integrin under basal conditions. In agreement with the in vivo studies, we identified an effective concentration window in vitro where the $\beta 1$ -integrin inhibitory antibody decreased the inflammation-induced formation of actin stress fibers, loss of VE-cadherin, and EC contractility and permeability, without affecting monolayer stability under noninflammatory conditions. Consistent with these results, the transient $\beta 1$ -integrin silencing, which preserved 10% of the $\beta 1$ -integrin-positive adhesions, did not affect EC junctions of confluent monolayers in basal conditions, but protected from inflammation-induced monolayer destabilization. Thus, the present study clearly indicates that, besides the known function of endothelial $\beta 1$ -integrin in maintaining vessel integrity (10), $\beta 1$ -integrin plays a key role in actively promoting vascular leak, likely via fibrillar adhesions.

Consistent with our results in acute inflammation, others have shown that $\alpha 5\beta 1$ -integrin regulates endothelial activation in chronic inflammation (14). Elegant studies using knock-in mice, where the cytoplasmic tail of the fibronectin receptor $\alpha 5$ -integrin was replaced with that of the collagen/laminin receptor $\alpha 2$ -integrin, demonstrated that switching from $\alpha 5\beta 1$ signaling to $\alpha 2\beta 1$ signaling inhibited the formation of atherosclerotic plaques in hypercholesterolemic apolipoprotein E (*ApoE*)-deficient mice (14). Inhibiting $\alpha 5\beta 1$ -integrin signaling using ATN-161, a non-RGD-based peptide inhibitor of $\alpha 5\beta 1$ -integrin, decreased atherosclerotic plaque development in *ApoE* and low density lipoprotein (LDL) receptor (*Ldlr*) deficient mice (29, 30). The present study does not identify the possible integrin alpha subunit(s) that pair(s) with $\beta 1$ -integrin to promote vascular leakage; however, we show that $\alpha 5$ -integrin was involved in mediating inflammatory responses in cultured ECs. It has been additionally reported that antibodies inhibiting $\alpha \nu \beta 5$ -integrin are able to decrease vascular leakage in a mouse sepsis model (31), but whether $\alpha \nu \beta 5$ -integrin utilizes similar mechanisms to those identified here for $\beta 1$ -integrin remains to be investigated.

Increased capillary leakage leading to hypovolemic shock and multiorgan failure is a key event in the pathobiology of several difficult-to-treat conditions, such as sepsis (2, 3, 6). In capillary leak syndromes that occur as a severe side effect of cancer therapies employing adoptive T cell transfer, an IL-6 receptor inhibitor has shown efficacy. A recent study demonstrated that improving the host vascular integrity in mice via administration of recombinant Slit2N, which enhances VE-cadherin localization in EC-EC junctions, increased the survival of mice during endotoxemia, polymicrobial sepsis, and H5N1 influenza (32). In another study, an ANGPT2 antibody that stimulated TIE2 activation was used to stabilize the vasculature in mouse sepsis models whereas conventional ANGPT2-TIE2 neutralizing antibodies were less effective (33). TIE2 activation was not increased following treatment with the $\beta 1$ -integrin antibody in our studies, indicating a TIE-independent mechanism of vascular stabilization. In our studies, $\beta 1$ -integrin antibody was able to prevent pathological vascular leakage in endotoxemia even when the inhibitory antibody was administered after the onset of systemic inflammation and vascular leakage. In the preventative mode, the $\beta 1$ -integrin antibody significantly improved the cardiac output in mice suffering from LPS-induced sepsis, suggesting that the improved vascular integrity prevented the severe drop in cardiac output. Due to the ubiquitous expression of $\beta 1$ -integrin in different cells and tissues and its versatile function, $\beta 1$ -integrin may not be a plausible drug candidate for preventing long-term vascular leakage. However, our data indicate that ANGPT2, which is increased in inflammation, can regulate EC-ECM adhesions, and therefore understanding more about this mechanism may provide a way to control $\beta 1$ -integrin signaling in vascular leakage in disease.

Materials and Methods

A detailed description of all materials and methods can be found in *SI Appendix, SI Materials and Methods*.




Fig. 7. Inflammatory agents stimulate the formation of α 5 β 1-integrin- and tensin-1-positive fibrillar adhesions in an ANGPT2-dependent manner. (*A* and *B*) BECs expressing paxillin-TagGFP2 were imaged live, every 2 min for 1 h, in the presence or absence of thrombin (1 U·mL⁻¹). Representative temporal code images at times 0 to 60 min. (*B*) Merged images of time 0 (green) and 60 min (magenta). BECs were stimulated with IL-1 β (2 h) (*C–E*) or thrombin (0.1 U·mL⁻¹, 30 min) (*F*) and stained for active β 1-integrin (12G10) (*C*) or α 5-integrin (SNAKA51) (*D* and *F*), or stimulated with LPS (2.5 μ g·mL⁻¹, 2 h) and stained for tensin-1 (TNS1) (*G*). (*E*) Quantification of the SNAKA51-positive area (160 cells per treatment, $n = 3$ independent experiments). (*H*) Quantification of the TNS1-positive area ($n = 3$ independent experiments). (*I*) VE-cadherin staining of BECs transfected with shScr or shTNS1 and stimulated with IL-1 β (2 h). (*J*) Quantification of VE-cadherin staining in *I* (220 to 250 cells per treatment, $n = 3$ independent experiments). (*K*) Quantification of the TNS1-positive area in LPS-stimulated BECs transfected with shScr or shANGPT2 (micrographs in *SI Appendix, Fig. S19*) ($n = 3$ independent experiments). (*L*) BECs transfected with shScr or sh β 1 were stimulated with recombinant human (rh)ANGPT2 for 2 h and fixed and stained for VE-cadherin ($n = 3$ independent experiments). (*M*) BECs transfected with shScr or shANGPT2 were analyzed for α 5-integrin recycling, as described in *SI Appendix, SI Materials and Methods*. Internalized α 5-integrin was quantified in cells stimulated with LPS, IL-1 β , or thrombin relative to unstimulated, shScr cells after recycling. LPS, IL-1 β , and thrombin induced the recycling of internalized α 5-integrin from vesicles to cell adhesions in shScr-transfected but not shANGPT2-transfected cells. $n = 3$ independent experiments. Data are the mean \pm SD. * $P < 0.05$, ** $P < 0.01$, and *** $P < 0.001$ (Tukey's test). Nuclear DAPI stain. (Scale bars: 20 μ m.) (*N*) Schematic model of ANGPT2- and α 5 β 1-integrin-mediated endothelial destabilization in response to inflammatory stimuli. In quiescent endothelial monolayers, the cortical actin rim and VE-cadherin-positive cell junctions maintain monolayer integrity. Inflammatory agents induce the formation of α 5 β 1-integrin- and tensin-1-positive fibrillar adhesions, in an ANGPT2-dependent manner. This stabilizes actin stress fibers, increases tension, and destabilizes endothelial cell-cell junctions via loss of junctional VE-cadherin, leading to increased permeability.

Mouse Models. All animal experiments were approved by the National Animal Experiment Board in Finland.

Analysis of Vascular Leakage in Murine Endotoxemia. Vascular leakage was analyzed as previously described (21, 23). β 1- or α 4-integrin antibodies, or control antibodies (all 2.5 mg·kg⁻¹), were administered to 2- to 4-mo-old male C57BL/6 mice by i.p. injection. Cre-positive tamoxifen-induced *Itgb1*^{WT/Fl} (*Itgb1*^{WT/ΔEC}) mice were compared with Cre-negative *Itgb1*^{WT/WT}, *Itgb1*^{WT/Fl}, or *Itgb1*^{fl/fl} littermates or Cre-positive *Itgb1*^{WT/WT} (control mice). To induce endotoxemia, LPS (O55:B5, 11 mg·kg⁻¹) was administered i.p. for 16 h. Alexa Fluor-594-labeled fluorospheres (100 nm) or Texas red-labeled dextran (70 kDa; Invitrogen, ThermoFisher Scientific) was injected via the tail vein and allowed to circulate for 4 min, followed by sequential perfusion with PBS and 1% paraformaldehyde (PFA) in PBS via the left ventricle. Tracheas and ears were immunostained, and the fluorescence was quantified as described in *SI Appendix, SI Materials and Methods*.

Fluorescence microscopy and quantification of fluorescent and TEM images are described in *SI Appendix, SI Materials and Methods*.

Statistical Tests. For pairwise comparisons of two treatment groups, Student's *t* tests (two-tailed, unequal variance) were used, and, for comparisons of multiple groups, Tukey's multiple comparison tests were used in conjunction with ANOVA post hoc pairwise testing. A *P* value of less than 0.05 was considered statistically significant. Experimental group sizes are indicated in the figure legends.

ACKNOWLEDGMENTS. We thank Kirsi Mänttari, Riina Haajanen, and Päivi Leinikka for technical assistance; Dr. Hemant Giri and Emilia Korhonen for advice; Dr. Kari Alitalo for critical comments on the manuscript; the Laboratory Animal Center at the University of Helsinki for animal care; the staff at the Biomedicum Imaging and Genome Biology Units, the Biocenter Oulu Electron Microscopy Core, and the Turku Centre for Biotechnology Cell Imaging Core facility for microscopy services; and the Functional Genomics Unit for TRC1 library clones. This work was funded by Academy of Finland Centre of Excellence Program Grants 271845 (to P.S. and J.I.) and 251314 (to L.E.), and Postdoctoral Research Grant 308663 (to E.A.K.); European Research Council (ERC) Consolidator Program Grants 773076 (to P.S.) and 615258 (to J.I.); the Cancer Society of Finland (P.S.); the Sigrid Juselius Foundation (P.S.); and Helsinki Institute for Life Sciences (HiLife) (P.S.). M.L. is supported by the University of Turku Doctoral Program (TuDMM).

1. Pober JS, Sessa WC (2007) Evolving functions of endothelial cells in inflammation. *Nat Rev Immunol* 7:803–815.
2. Goldenberg NM, Steinberg BE, Slutsky AS, Lee WL (2011) Broken barriers: A new take on sepsis pathogenesis. *Sci Transl Med* 3:88ps25.
3. Marshall JC (2014) Why have clinical trials in sepsis failed? *Trends Mol Med* 20:195–203.
4. Amado-Azevedo J, Valent ET, Van Nieuw Amerongen GP (2014) Regulation of the endothelial barrier function: A filum granum of cellular forces, Rho-GTPase signaling and microenvironment. *Cell Tissue Res* 355:557–576.
5. Trani M, Dejana E (2015) New insights in the control of vascular permeability: Vascular endothelial-cadherin and other players. *Curr Opin Hematol* 22:267–272.
6. Parikh SM, et al. (2006) Excess circulating angiopoietin-2 may contribute to pulmonary vascular leak in sepsis in humans. *PLoS Med* 3:e46.
7. Benet AV, et al. (2013) Angiopoietin-2 is critical for cytokine-induced vascular leakage. *PLoS One* 8:e70459.
8. Parsons JT, Horwitz AR, Schwartz MA (2010) Cell adhesion: Integrating cytoskeletal dynamics and cellular tension. *Nat Rev Mol Cell Biol* 11:633–643.
9. Su G, et al. (2012) Absence of integrin α β 3 enhances vascular leak in mice by inhibiting endothelial cortical actin formation. *Am J Respir Crit Care Med* 185:58–66.
10. Yamamoto H, et al. (2015) Integrin β 1 controls VE-cadherin localization and blood vessel stability. *Nat Commun* 6:6429.
11. Francis SE, et al. (2002) Central roles of α 5 β 1 integrin and fibronectin in vascular development in mouse embryos and embryoid bodies. *Arterioscler Thromb Vasc Biol* 22:927–933.
12. Zovein AC, et al. (2010) Beta1 integrin establishes endothelial cell polarity and arteriolar lumen formation via a Par3-dependent mechanism. *Dev Cell* 18:39–51.
13. Stenzel D, et al. (2011) Integrin-dependent and -independent functions of astrocytic fibronectin in retinal angiogenesis. *Development* 138:4451–4463.
14. Yun S, et al. (2016) Interaction between integrin α 5 and PDE4D regulates endothelial inflammatory signalling. *Nat Cell Biol* 18:1043–1053.
15. Hakanpaa L, et al. (2015) Endothelial destabilization by angiopoietin-2 via integrin β 1 activation. *Nat Commun* 6:5962.
16. Su Y, et al. (2016) Relating conformation to function in integrin α 5 β 1. *Proc Natl Acad Sci USA* 113:E3872–E3881.
17. Rathnakumar K, et al. (2016) Angiopoietin-2 mediates thrombin-induced monocyte adhesion and endothelial permeability. *J Thromb Haemost* 14:1655–1667.
18. Zhu W, et al. (2012) Interleukin receptor activates a MYD88-ARNO-ARF6 cascade to disrupt vascular stability. *Nature* 492:252–255.
19. Carlson TR, Hu H, Braren R, Kim YH, Wang RA (2008) Cell-autonomous requirement for beta1 integrin in endothelial cell adhesion, migration and survival during angiogenesis in mice. *Development* 135:2193–2202.
20. Mofarrahi M, et al. (2008) Regulation of angiopoietin expression by bacterial lipopolysaccharide. *Am J Physiol Lung Cell Mol Physiol* 294:L955–L963.
21. Korhonen EA, et al. (2016) Tie1 controls angiopoietin function in vascular remodeling and inflammation. *J Clin Invest* 126:3495–3510.
22. Engelhardt B, et al. (1998) The development of experimental autoimmune encephalomyelitis in the mouse requires alpha4-integrin but not alpha4beta7-integrin. *J Clin Invest* 102:2096–2105.
23. Kim M, et al. (2016) Opposing actions of angiopoietin-2 on Tie2 signaling and FOXO1 activation. *J Clin Invest* 126:3511–3525.
24. Pankov R, et al. (2000) Integrin dynamics and matrix assembly: Tensin-dependent translocation of alpha(5)beta(1) integrins promotes early fibronectin fibrillogenesis. *J Cell Biol* 148:1075–1090.
25. Georgiadou M, et al. (2017) AMPK negatively regulates tensin-dependent integrin activity. *J Cell Biol* 216:1107–1121.
26. Arjonen A, Alanko J, Veltel S, Ivaska J (2012) Distinct recycling of active and inactive β 1 integrins. *Traffic* 13:610–625.
27. Maisonpierre PC, et al. (1997) Angiopoietin-2, a natural antagonist for Tie2 that disrupts in vivo angiogenesis. *Science* 277:55–60.
28. Holopainen T, et al. (2012) Effects of angiopoietin-2-blocking antibody on endothelial cell-cell junctions and lung metastasis. *J Natl Cancer Inst* 104:461–475.
29. Yurdagul A, Jr, et al. (2014) α 5 β 1 integrin signaling mediates oxidized low-density lipoprotein-induced inflammation and early atherosclerosis. *Arterioscler Thromb Vasc Biol* 34:1362–1373.
30. Sun X, et al. (2016) Activation of integrin α 5 mediated by flow requires its translocation to membrane lipid rafts in vascular endothelial cells. *Proc Natl Acad Sci USA* 113:769–774.
31. Su G, et al. (2013) Effective treatment of mouse sepsis with an inhibitory antibody targeting integrin α β 5. *Crit Care Med* 41:546–553.
32. London NR, Li DY (2011) Robo4-dependent Slit signaling stabilizes the vasculature during pathologic angiogenesis and cytokine storm. *Curr Opin Hematol* 18:186–190.
33. Han S, et al. (2016) Amelioration of sepsis by TIE2 activation-induced vascular protection. *Sci Transl Med* 8:335ra55.