

Nematode Genome Announcement: A Draft Genome for Rice Root-Knot Nematode, *Meloidogyne graminicola*

Vishal Singh Somvanshi,¹ Madhura Tathode,² Rohit Nandan Shukla,² and Uma Rao^{1*}

¹Division of Nematology, ICAR–Indian Agricultural Research Institute, LBS Center, PUSA Campus, New Delhi 110012, India.

²Bionivid Technology Private Limited, 209, 4th Cross, Kasturi Nagar, Bangalore 560043, India.

*Email: umarao@iari.res.in.

This paper was edited by Erik J. Ragsdale.

Abstract

The rice root-knot nematode *Meloidogyne graminicola* has emerged as a devastating pest of rice in South-East Asian countries. Here we present a draft genome sequence for *M. graminicola*, assembled using data from short and long insert libraries sequenced on Illumina GAllx sequencing platform.

Key words

Meloidogyne graminicola, Rice root-knot nematode, Genomics, Draft genome, Sequencing, Illumina GAllx.

Rice is the second most important food crop in the world after corn based on the total production. In 2016, rice was cultivated in 161.1 million ha area, and the global production was 482 million metric tons (World Rice Statistics, International Rice Research Institute, Manila, Philippines, <http://ricestat.irri.org:8080/wrsv3/entrypoint.htm>). The rice root-knot nematode, *Meloidogyne graminicola*, has emerged as a devastating pest of rice in South-East Asia (Dutta et al., 2012; Mantelin et al., 2016), where it is highly damaging under upland, rainfed lowland (Prot et al., 1994) and irrigated (Netscher and Erlan, 1993) cultivation conditions. Severe *M. graminicola* infection is known to cause 100% damage to the rice nursery. Here, we report the sequencing and assembly of the genome of *M. graminicola* IARI strain. This resource would help researchers investigate and understand the unique biology of this nematode and discover new strategies for its management.

Considering the ~30Mb genome size of *M. graminicola* as predicted by Feulgen densitometry (Lapp and Triantaphyllou, 1972), we planned to generate two libraries of varying insert length with ~150× depth of data (~4.5Gb) per library using paired-end sequencing to achieve a comprehensive assembly. The *M. graminicola* population was collected from the infected rice fields from Indian Agricultural Research Institute farm, New Delhi, and multiplied from a single egg mass in pots under greenhouse conditions. Freshly hatched second stage juveniles were used for

the genomic DNA extraction using Gentra Puregene Tissue Kit (Cat No.: 158667 Qiagen, Valencia, CA, USA). The short (150–200bp) and long (300–500bp) DNA fragments were obtained by diluting 1 µg of genomic DNA in 100 µl nuclease free water (Ambion, Waltham, MA, USA) and sonication by Bioruptor (Diagenode, Seraing (Ougrée), Belgium) at 20 and 13 pulses at 30sec ON and 30sec OFF, respectively. The resulting fragmented DNA was cleaned using QIAquick columns (Qiagen, Valencia, CA, USA). The size distribution was checked by running an aliquot of the fragmented DNA sample on Agilent high sensitivity bioanalyzer (Agilent Technologies, Santa Clara, CA, USA). Subsequently, the libraries for whole genome sequencing were constructed as per the Illumina TruSeq DNA sample preparation guide (Illumina, San Diego, CA, USA). The sequencing was performed on Illumina GAllx platform at the Genotypic Technology Pvt. Ltd., Bengaluru, India.

A total of ~130 million raw reads were generated comprising of 13Gb sequence data using 100bp paired-end sequencing. Approximately 120 million High Quality (HQ) reads were obtained from the raw data by using NGS QC Tool Kit v.2.3.3 (Patel and Jain, 2012). These ~12Gb of 120 million HQ reads were better than our planned strategy expecting nine Gb. The HQ reads obtained from both short and long insert libraries were used to generate primary assembly using Platanus assembler v.1.2.4 (Kajitani et al., 2014), and the resulting contigs were further scaffolded using

Platanus Scaffolding module to generate secondary assembly. The secondary assembly was further refined by Redundans pipeline (Pryszcz and Gabaldón, 2016) to generate the final genome assembly with a minimum sequence length of 500bp. The contaminating mitochondrial and bacterial sequences were identified by NCBI servers and removed from the draft genome assembly prior to submission to the NCBI GenBank. The mitochondrial genome was assembled separately from complete HQ reads using SPAdes assembler (Bankevich et al., 2012) with coverage cutoff of 500, wherein available *M. graminicola* mitochondrial genome sequences (accession nos. HG529223, KJ139963) obtained from GenBank were provided as trusted contigs to the SPAdes assembler. This resulted in only 4 scaffolds from the assembly. The resulted scaffolds from SPAdes assembler were further merged using EMBOSS merger tool (Rice et al., 2000) to construct full length mitochondrial genome. The assembled genome was further annotated using MITOS (Bernt et al., 2013) and ARWEN (Laslett and Canbäck, 2008) servers.

The final *M. graminicola* genome sequence assembly was of 38.18Mb size, and included 4,304 scaffolds with an average scaffold length of 8.87 Kb. The minimum and maximum scaffold length was 501 bp and 145 Kb, respectively. The N50 and N90 lengths for the final assembly were 20.4 Kb and 4.2 Kb, respectively. The GC content of the assembled genome was 23.05%, and there were 1.88% N's in the assembly. Core Eukaryotic Genes Mapping Approach (CEGMA) (Parra et al., 2007) was used to assess the completeness of the *M. graminicola* genome assembly, and out of 248 core genes, 209 complete (84.27%) and 225 partial (90.73%) core eukaryotic genes (CEGs) were found to be present. Identification of protein-coding genes was carried out by using GenMark-ES tool (Borodovsky and McIninch, 1993) which predicted 10,196 protein-coding genes, as compared with 6,712 to 20,317 in other plant-parasitic nematode genomes (summarized in Kikuchi et al., 2017). Functional annotation of predicted *M. graminicola* protein-coding genes performed using OrthoMCL (Li et al., 2003) identified 5,427 proteins that shared high homology with other *Meloidogyne* spp. In addition, 245 tRNA genes were predicted. The mitochondrial genome sequence of *M. graminicola* IARI strain was 19,019bp long and contained 12 protein-coding genes, 22 tRNA and two ribosomal RNA genes. Based on the mitochondrial genome sequence, the *M. graminicola* IARI strain appears phylogenetically closer to the *M. graminicola* strain from Philippines (HG529223, 20,030bp, Besnard et al., 2014) as compared with the Chinese strain (KJ139963, 19,589bp, Sun et al., 2014).

The present assembly size deviates from that of the ~30Mb as predicted by Feulgen densitometry (Lapp and Triantaphyllou, 1972). Using sequencing technologies that produce longer reads such as PacBio or mate pair sequencing to obtain better genome assemblies, and, inbreeding the nematode strain to be used for sequencing to reduce possible heterozygosity might help in correcting the mismatch between predicted and assembled genome sizes. However, N50 value, complete and partial CEGs and other genome statistics for our *M. graminicola* assembly are comparable to the closely related and published plant-parasitic nematode genomes solved using similar sequencing platforms (Supplementary Table A1).

This draft genome sequence would be useful for the researchers working on comparative genomics of *Meloidogyne* and other tylenchid nematodes, and enable functional genomics in *M. graminicola*. We understand that the present *M. graminicola* draft genome is incomplete, and expect to improve it in the near future. The present assembly would work as a base for the further improvement of the *M. graminicola* genome sequence.

GenBank accession numbers: The Whole Genome Shotgun project has been deposited at DDBJ/ENA/GenBank under the accession NXFT00000000. The raw DNA sequence data was deposited in GenBank under BioSample no. SAMN04041660, BioProject No. PRJNA411966 and SRX1224028 (long insert library) and SRX1223928 (short insert library), respectively. The mitochondrial genome was submitted to GenBank under accession no. MG763561.

Acknowledgements

We acknowledge funding from ICAR-IARI through the IARI Outreach project IARI:ORP: NEM:09:14, and from the Department of Biotechnology, Government of India through grant no. BT/PR/5163/7/397/2012. We thank the Director, IARI and the Joint Director (Research), IARI for the facilities and support.

References

- Bankevich, A., Nurk, S., Antipov, D., Gurevich, A.A., Dvorkin, M., Kulikov, A.S., Lesin, V.M., Nikolenko, S.I., Pham, S., Prjibelski, A.D., and Pyshkin, A.V. 2012. SPAdes: A new genome assembly algorithm and its applications to single-cell sequencing. *Journal of Computational Biology* 19: 455–77.
- Bernt, M., Donath, A., Jühling, F., Externbrink, F., Florentz, C., Fritsch, G., Pütz, J., Middendorf, M., and Stadler, P.F. 2013. MITOS: Improved de novo metazoan

mitochondrial genome annotation. *Molecular Phylogenetics and Evolution* 69: 313–9.

Besnard, G., Jühling, F., Chapuis, É., Zedane, L., Lhuillier, É., MATEILLE, T., and Bellafiore, S. 2014. Fast assembly of the mitochondrial genome of a plant parasitic nematode (*Meloidogyne graminicola*) using next generation sequencing. *Comptes Rendus Biologies* 337: 295–301.

Borodovsky, M., and McIninch, J. 1993. Genmark: Parallel gene recognition for both DNA strands. *Computers and Chemistry* 17: 123–33.

Dutta, T.K., Ganguly, A.K., and Gaur, H.S. 2012. Global status of rice root-knot nematode, *Meloidogyne graminicola*. *African Journal of Microbiology Research* 32: 6016–21.

Kajitani, R., Toshimoto, K., Noguchi, H., Toyoda, A., Ogura, Y., Okuno, M., Yabana, M., Harada, M., Nagayasu, E., and Maruyama, H. 2014. Efficient *de novo* assembly of highly heterozygous genomes from whole-genome shotgun short reads. *Genome Research* 24: 1384–95.

Kikuchi, T., Eves-van den Akker, S., and Jones, J.T. 2017. Genome evolution of plant-parasitic nematodes. *Annual Review of Phytopathology* 55: 333–54.

Lapp, N.A., and Triantaphyllou, A.C. 1972. Relative DNA content and chromosomal relationships of some *Meloidogyne*, *Heterodera*, and *Meloidodera* spp. (Nematoda: Heteroderidae). *Journal of Nematology* 4: 287–91.

Laslett, D., and Canbäck, B. 2008. ARWEN, a program to detect tRNA genes in metazoan mitochondrial nucleotide sequences. *Bioinformatics* 24: 172–5.

Li, L., Stoeckert, C.J., and Roos, D.S. 2003. OrthoMCL: Identification of ortholog groups for eukaryotic genomes. *Genome Research* 13: 2178–89.

Mantelin, S., Bellafiore, S., and Kyndt, T. 2016. *Meloidogyne graminicola*: A major threat to rice agriculture. *Molecular Plant Pathology* 18: 3–15.

Netscher, C., and Erlan, X. 1993. A root-knot nematode, *Meloidogyne graminicola*, parasitic on rice in Indonesia. Afro-Asian. *Journal of Nematology* 3: 90–5.

Parra, G., Bradnam, K., and Korf, I. 2007. CEGMA: A pipeline to accurately annotate core genes in eukaryotic genomes. *Bioinformatics* 23: 1061–7.

Patel, R.K., and Jain, M. 2012. NGS QC Toolkit: A toolkit for quality control of next generation sequencing data. *PLOS ONE* 7: e30619.

Prot, J.C., Villammeva, L.M., and Gergon, E.B. 1994. The potential of increased nitrogen supply to mitigate growth and yield reductions of upland rice cultivar UPL Ril-5 caused by *Meloidogyne graminicola*. *Fundamental and Applied Nematology* 17: 445–54.

Pryszcz, L.P., and Gabaldón, T. 2016. Redundans: An assembly pipeline for highly heterozygous genomes. *Nucleic Acids Research* 44: e113.

Rice, P., Longden, I., and Bleasby, A. 2000. EMBOSS: The European molecular biology open software suite. *Trends in Genetics* 16: 276–7.

Sun, L., Zhuo, K., Lin, B., Wang, H., and Liao, J. 2014. The complete mitochondrial genome of *Meloidogyne graminicola* (Tylenchina): A unique gene arrangement and its phylogenetic implications. *PLOS ONE* 9: e98558.

Appendix

Supplementary Table A1

Supplementary Table A1. A comparison of *Meloidogyne graminicola* genome information with the published plant-parasitic nematode genomes.

Sl. no.	Nematode	Sequencing platform	Assembly approach/ assembler/ assembly-pipeline	Assembly size	No. of scaffolds	N50 value (kbp)	CEGMA score (complete/partial %) or complete%	Reference	Year
1	<i>Meloidogyne incognita</i>	Sanger, ABI3730x1 DNA analyzer	De-novo, Arachne	86.1	2,995	62.5	77/80.6	Abad et al. (2008)	2008
2	<i>Meloidogyne hapla</i>	ABI3730, megabase dequence analyzer	De-novo, Arachne v2.0.1	53.0	3,452	37.6	94.8/96.8	Opperman et al. (2008)	2008
3	<i>Bursaphelenchus xylophilus</i>	454 FLX, illumina GAL	De-novo, Newbler v2.3, Velvet v 1.0.12, Abacasll, IMAGE, iCORN	74.6	5,527	949.8	97.6/98.4	Kikuchi et al. (2011)	2011
4	<i>Meloidogyne floridensis</i>	Illumina HiSeq2000	De-novo, Velvet v1.1.04	96.7	58,696	3.7	58.1/77.4	Lunt et al. (2014)	2014
5	<i>Globodera pallida</i>	ABI 3730 Capillary DNA Analyser, 454 GS-20 and GS-FLX sequencer, Illumina GAllx	De-novo, Celera, Newbler, Abyss v1.2.7, SSPACE v1	124.6	6,873	122	74.19/80.65	Cotton et al. (2014)	2014
6	<i>Pratylenchus coffeae</i>	Roche 454	De-novo, Newbler	19.7	5,821	10	NA	Burke et al. (2015)	2015
7	<i>Globodera rostochiensis</i>	Illumina HiSeq2000	De-novo, SGA v0.9.7, Velvet v1.3.7, SSPACE, Gapfiller	95.9	4,377	89	93.55/95.56	Eves-van den Akker et al. (2016)	2016
8	<i>Meloidogyne enterolobii</i>	Illumina L30	De-novo, Platanus	162.4	46,090	9.2	81 /NA	Szitenberg et al. (2017)	2017
9	<i>Meloidogyne floridensis</i>	Illumina SJF1	De-novo, Platanus	74.9	9,134	13.2	84	Szitenberg et al. (2017)	2017
10	<i>Meloidogyne incognita</i>	Illumina W1	De-novo, Platanus	122.1	33,735	16.4	83	Szitenberg et al. (2017)	2017
11	<i>Globodera ellingtonae</i>	HiSeq, MiSeq, PacBio	De-novo assembler Allpaths-LG, PBJelly	119.1	2,248	360	92/96	Phillips et al. (2017)	2017
12	<i>Meloidogyne javanica</i>	Illumina VW4	De-novo, Platanus	142.6	34,394	14.2	90	Szitenberg et al. (2017)	2017

13	<i>Meloidogyne arenaria</i>	Illumina HiSeqA	De-novo, Platanus	163.7	46,509	10.5	91	Szitenberg et al. (2017)	2017
14	<i>Ditylenchus destructor</i>	Illumina HiSeq2500, PacBio RSII	De-novo, ALLPATHS-LG, SSPACE, pb-jelly, Gapfiller	112	1,761	570	91	Zheng et al. (2016)	2016
15	<i>Meloidogyne incognita</i>	454, Illumina HiSeq2000	De-novo, MIRA, SSPACE, GapCloser (SOAPdenovo 2)	183.5	12,091	38.6	97	Blanc-Mathieu et al. (2017)	2017
16	<i>Meloidogyne javanica</i>	454, Illumina HiSeq2000	De-novo, MIRA, SSPACE, GapCloser (SOAPdenovo 2)	256.3	31,341	10.4	96	Blanc-Mathieu et al. (2017)	2017
17	<i>Meloidogyne arenaria</i>	454, Illumina HiSeq2000	De-novo, MIRA, SSPACE, GapCloser (SOAPdenovo 2)	235.5	26,196	16.4	95	Blanc-Mathieu et al. (2017)	2017
18	<i>Meloidogyne graminicola</i>	Illumina GAIIx	De-novo, Platanus, Redundans	38.18	4,304	20.4	84.27/90.73	This study	2018

References

- Abad, P., Gouzy, J., Aury, J. M., Castagnone-Sereno, P., Danchin, E. G., Deleury, E., Perfus-Barbeoch, L., Anthouard, V., Artiguenave, F., Blok, V. C., Caillaud, M. C., Coutinho, P. M., Dasilva, C., De Luca, F., Deau, F., Esquibet, M., Flutre, T., Goldstone, J. V., Hamamouch, N., Hewezi, T., Jaillon, O., Jubin, C., Leonetti, P., Magliano, M., Maier T. R., Markov, G. V., McVeigh, P., Pesole, G., Poulain, J., Robinson-Rechavi, M., Sallet, E., Segurens, B., Steinbach, D., Tytgat, T., Ugarte, E., van Ghelder, C., Veronico, P., Baum, T. J., Blaxter, M., Bleve-Zacheo, T., Davis, E. L., Ewbank, J. J., Favery, B., Grenier, E., Henrissat, B., Jones, J. T., Laudet, V., Maule, A. G., Quesneville, H., Rosso, M. N., Schiex, T., Smant, G., Weissenbach, J., and Wincker, P. 2008. Genome sequence of the metazoan plant-parasitic nematode *Meloidogyne incognita*. *Nature Biotechnology* 26:909–915.
- Blanc-Mathieu, R., Perfus-Barbeoch, L., Aury, J. M., Da Rocha, M., Gouzy, J., Sallet, E., Martin-Jimenez, C., Bailly-Bechet, M., Castagnone-Sereno, P., Flot, J. F., Kozlowski, D. K., Cazareth, J., Couloux, A., Da Silva, C., Guy, J., Kim-Jo, Y., Rancurel, C., Schiex, T., Abad, P., Wincker, P., and Danchin, E. G. J. 2017. Hybridization and polyploidy enable genomic plasticity without sex in the most devastating plant-parasitic nematodes. *PLoS Genetics* 13:e1006777.
- Burke, M., Scholl, E. H., Bird, D. M., Schaff, J. E., Colman, S. D., Crowell, R., Diener, S., Gordon, O., Graham, S., Wang, X., Windham, E., Wright, G. M., and Opperman, C. H. 2015. The plant parasite *Pratylenchus coffeae* carries a minimal nematode genome. *Nematology* 17:621–637.
- Cotton, J. A., Lilley, C. J., Jones, L. M., Kikuchi, T., Reid, A. J., Thorpe, P., Tsai, I. J., Beasley, H., Blok, V., Cock, P. J., Eves-van den Akker, S., Holroyd, N., Hunt, M., Mantelin, S., Naghra, H., Pain, A., Palomares-Rius, J. E., Zarowiecki, M., Berriman, M., Jones, J. T., and Urwin, P. E. 2014. The genome and life-stage specific transcriptomes of *Globodera pallida* elucidate key aspects of plant parasitism by a cyst nematode. *Genome Biology* 15:R43.
- Eves-van den Akker, S., Laetsch, D. R., Thorpe, P., Lilley, C. J., Danchin, E. G., Da Rocha, M., Rancurel, C., Holroyd, N. E., Cotton, J. A., Szitenberg, A., Grenier, E., Montarry, J., Mimeo, B., Duceppe, M. O., Boyes, I., Marvin, J. M. C., Jones, L. M., Yusup, H. B., Lafond-Lapalme, J., Esquibet, M., Sabeh, M., Rott, M., Overmars, H., Finkers-Tomczak, A., Smant, G., Koutsovoulos, G., Blok, V., Mantelin, S., Cock, P. J. A., Phillips, W., Henrissat, B., Urwin, P. E., Blaxter, M., and Jones, J. T. 2016. The genome of the yellow potato cyst nematode, *Globodera rostochiensis*, reveals insights into the basis of parasitism and virulence. *Genome Biology* 17:124.
- Kikuchi, T., Cotton, J. A., Dalzell, J. J., Hasegawa, K., Kanzaki, N., McVeigh, P., Takanashi, T., Tsai, I. J., Assefa, S. A., and Cock, P. J., Otto, T. D., Hunt, M., Reid, A. J., Sanchez-Flores, A., Tsuchihara, K., Yokoi, T., Larsson, M. C., Miwa, J., Maule, A. J., Sahashi, N., Jones, J. T., and Berriman, M. 2011. Genomic insights into the origin of parasitism in the emerging plant pathogen *Bursaphelenchus xylophilus*. *PLOS Pathogens* 7:e1002219.
- Lunt, D. H., Kumar, S., Koutsovoulos, G., and Blaxter, M. L. 2014. The complex hybrid origins of the root knot nematodes revealed through comparative genomics. *PeerJ* 2:e356.
- Opperman, C. H., Bird, D. M., Williamson, V. M., Rokhsar, D. S., Burke, M., Cohn, J., Cromer, J., Diener, S., Gajan, J., Graham, S., Houfek, T. D., Liu, Q., Mitros, T., Schaff, J., Schaffer, R., Scholl, E., Sosinski, B. R., Thomas, V. P., and Windham, E. 2008. Sequence and genetic map of *Meloidogyne hapla*: A compact nematode genome for plant parasitism. *Proceedings of the National Academy of Sciences of the United States of America* 105:14802–14807.
- Phillips, W. S., Howe, D. K., Brown, A. M. V., Eves-van den Akker, S., Dettwyler, L., Peetz, A. B., Debver, D. R., and Zasada, I. A. 2017. The Draft Genome of *Globodera ellingtonae*. *Journal of Nematology* 49:127–128.
- Szitenberg, A., Salazar-Jaramillo, L., Blok V. C., Laetsch, D. R., Joseph, S., Williamson, V. M., Blaxter, M. L., and Lunt, D. H. 2017. Comparative genomics of apomictic root-knot nematodes: Hybridization, ploidy, and dynamic genome change. *Genome Biology and Evolution* 9:2844–2861.
- Zheng, J., Peng, D., Chen, L., Liu, H., Chen, F., Xu, M., Ju, S., Ruan, L., and Sun, M. 2016. The *Ditylenchus destructor* genome provides new insights into the evolution of plant parasitic nematodes. *Proceedings of the Royal Society B: Biological Sciences* 283:20160942.