

Title: Genetic predisposition to mosaic Y chromosome loss in blood is associated with genomic instability in other tissues and susceptibility to non-haematological cancers

Deborah J. Thompson¹, Giulio Genovese²⁻⁴, Jonatan Halvardson⁵, Jacob C. Ulirsch^{3,6}, Daniel J. Wright^{7,8}, Chikashi Terao⁹⁻¹², Olafur B. Davidsson¹³, Felix R. Day^{7,14}, Patrick Sulem¹³, Yunxuan Jiang¹⁵, Marcus Danielsson⁵, Hanna Davies⁵, Joe Dennis¹, Malcolm G. Dunlop¹⁶, Douglas F. Easton¹, Victoria A. Fisher¹⁷, Florian Zink¹³, Richard S. Houlston¹⁸, Martin Ingelsson¹⁹, Siddhartha Kar²⁰, Nicola D. Kerrison⁷, Ben Kinnersley¹⁸, Ragnar P. Kristjansson¹³, Philip J. Law¹⁸, Rong Li²¹, Chey Loveday¹⁸, Jonas Mattisson⁵, Steven A. McCarroll²⁻⁴, Yoshinori Murakami²², Anna Murray²³, Pawel Olszewski²⁴, Edyta Rychlicka-Buniowska^{5,24}, Robert A. Scott⁷, Unnur Thorsteinsdottir^{13,25}, Ian Tomlinson²⁶, Behrooz Torabi Moghadam⁵, Clare Turnbull^{18,27}, Nicholas J. Wareham⁷, Daniel F. Gudbjartsson^{13,28}, INTEGRAL-ILCCO[#], The Breast Cancer Association Consortium[#], CIMBA[#], The Endometrial Cancer Association Consortium[#], The Ovarian Cancer Association Consortium[#], The PRACTICAL Consortium[#], The Kidney Cancer GWAS Meta-Analysis Project[#], eQTLGen Consortium[#], BIOS Consortium[#], 23andMe Research Team[#], Yoichiro Kamatani^{9,12,29}, Eva R. Hoffmann³⁰, Steve P. Jackson^{31,32}, Kari Stefansson^{13,25}, Adam Auton¹⁵, Ken K. Ong⁷, Mitchell J. Machiela¹⁷, Po-Ru Loh^{3,33}, Jan P. Dumanski^{5,24}, Stephen J. Chanock¹⁷, Lars A. Forsberg^{5,34*} and John R. B. Perry^{7,14*}

Affiliations

1. Centre for Cancer Genetic Epidemiology, Department of Public Health and Primary Care, University of Cambridge, UK.
2. Department of Genetics, Harvard Medical School, Boston, MA, USA.
3. Program in Medical and Population Genetics, Broad Institute of MIT and Harvard, Cambridge, MA 02142, USA.
4. Stanley Center for Psychiatric Research, Broad Institute of MIT and Harvard, Cambridge, MA, USA. .
5. Department of Immunology, Genetics and Pathology and Science for Life Laboratory, Uppsala University, Uppsala, Sweden.
6. Program in Biological and Biomedical Sciences, Harvard Medical School, Boston, MA 02115, USA.
7. MRC Epidemiology Unit, School of Clinical Medicine, University of Cambridge, Cambridge, UK.
8. Open Targets Core Genetics, Wellcome Sanger Institute, Hinxton, UK.
9. Laboratory for Statistical Analysis, RIKEN Center for Integrative Medical Sciences, Kanagawa, Japan.
10. Clinical Research Center, Shizuoka General Hospital, Shizuoka, Japan.
11. The Department of Applied Genetics, The School of Pharmaceutical Sciences, University of Shizuoka, 52-1 Yada, Suruga-ku, Shizuoka Japan.
12. Laboratory for Statistical and Translational Genetics, RIKEN Center for Integrative Medical Sciences, Kanagawa, Japan.
13. deCODE Genetics–Amgen, Inc., Reykjavik, Iceland.
14. Department of Internal Medicine, Erasmus MC, University Medical Center Rotterdam, Dr Molewaterplein 40, 3015 GD, the Netherlands.
15. 23andMe, Inc., Mountain View, CA, 94041, USA.
16. Colon Cancer Genetics Group, Medical Research Council Human Genetics Unit and CRUK Cancer Research Centre, Institute of Genetics and Molecular Medicine, University of Edinburgh, Western General Hospital, Edinburgh, United Kingdom.
17. Division of Cancer Epidemiology and Genetics, National Cancer Institute, Rockville, Maryland, USA.

18. Division of Genetics & Epidemiology, The Institute of Cancer Research, London, UK.
19. Department of Public Health and Caring Sciences / Geriatrics, Uppsala University, 751 85 Uppsala, Sweden
20. Centre for Cancer Genetic Epidemiology, Department of Oncology, University of Cambridge, Cambridge, UK.
21. Department of Cell Biology, Johns Hopkins University School of Medicine, Baltimore, United States.
22. Division of Molecular Pathology, Institute of Medical Science, The University of Tokyo, Tokyo, Japan.
23. Genetics of Complex Traits, University of Exeter Medical School, University of Exeter, RILD Level 3, Royal Devon & Exeter Hospital, Barrack Road, Exeter EX2 5DW, UK.
24. Faculty of Pharmacy, Medical University of Gdansk, Gdansk, Poland.
25. Faculty of Medicine, University of Iceland, Reykjavik, Iceland.
26. Cancer Genetics and Evolution Laboratory, Institute of Cancer and Genomic Sciences, University of Birmingham, Birmingham, UK.
27. William Harvey Research Institute, Queen Mary University, London, UK.
28. School of Engineering and Natural Sciences, University of Iceland, Reykjavik, Iceland.
29. Kyoto-McGill International Collaborative School in Genomic Medicine, Kyoto University Graduate School of Medicine, Kyoto, Japan.
30. DNRF Center for Chromosome Stability, Department of Cellular and Molecular Medicine, Faculty of Health Sciences, University of Copenhagen, Copenhagen, Denmark.
31. Department of Biochemistry, University of Cambridge, Cambridge, UK.
32. Wellcome Trust and Cancer Research UK Gurdon Institute, University of Cambridge, Cambridge, UK.
33. Division of Genetics, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, MA, USA.
34. Beijer Laboratory of Genome Research, Uppsala University, Uppsala, Sweden.

Consortium membership is listed in the supplementary section

*denotes equal contribution

Correspondence to John R.B. Perry (john.perry@mrc-epid.cam.ac.uk)

Abstract

Mosaic loss of chromosome Y (LOY) in circulating white blood cells is the most common form of clonal mosaicism, yet our knowledge of the causes and consequences of this is limited. Using a newly developed approach, we estimate that 20% of the UK Biobank male population (N=205,011) has detectable LOY. We identify 156 autosomal genetic determinants of LOY, which we replicate in 757,114 men of European and Japanese ancestry. These loci highlight genes involved in cell-cycle regulation, cancer susceptibility, somatic drivers of tumour growth and cancer therapy targets. We demonstrate that genetic susceptibility to LOY is associated with non-haematological health outcomes in both men and women, supporting the hypothesis that clonal haematopoiesis is a biomarker of genome instability in other tissues. Single-cell RNA sequencing identifies dysregulated autosomal gene expression in leukocytes with LOY, providing insights into why clonal expansion of these cells may occur. Collectively, these data highlight the utility of studying clonal mosaicism to uncover fundamental mechanisms underlying cancer and other ageing-related diseases.

Introduction

Each day the human body produces billions of highly specialised blood cells, generated from a self-renewing pool of 50,000-200,000 haematopoietic stem cells (HSCs)¹. As these cells age and divide, mutation and mitotic errors create genetic diversity within the HSC pool and their progenitors. If a genetic alteration confers a selective growth advantage to one cell over the others, clonal expansion may occur. This process propels the lineage to a disproportionately high frequency, creating a genetically distinct sub-population of cells. In the literature this is commonly referred to as clonal haematopoiesis, or more broadly (not restricting to considering leukocytes), clonal mosaicism² or aberrant clonal expansion³.

Population-based studies assessing the magnitude and effect of clonal mosaicism have been largely limited by the challenges of accurately detecting the expected low cell-fraction mosaic events in leukocytes using genotype-array or sequence read data⁴. Recent advances in statistical methodology have improved sensitivity, with approaches now able to catalogue mosaic events at higher resolution across the genome^{5,6}. Detection of large structural mosaic events can vary considerably in size – from 50kb to entire chromosomes in length – and are typically present in only a small fraction of circulating leukocytes (<5%). It is well established that loss of the sex chromosomes – particularly the Y chromosome (LOY) in men – is by far the most frequently observed somatic change in leukocytes⁷⁻⁹. It remains unclear if and why absence of a Y chromosome provides a selective growth advantage in these cells – we hypothesise this could be due to (amongst other unknown mechanisms) the loss of a putative Y-linked cell-growth suppressor gene, loss of a Y-linked transcription factor influencing expression of cell-growth related autosomal genes or the reduced energy cost of cellular divisions. However, it is also possible an age-related increase of LOY could occur without direct functional involvement of the Y chromosome.

Our understanding of why some individuals, but not others, exhibit clonal mosaicism in blood is also limited. Previous studies have demonstrated robust associations with age, sex (clonal mosaicism is more frequent in males), smoking and inherited germline genetic predisposition^{2,4,10-15}. Recent epidemiological studies have challenged the view that LOY in the hematopoietic system is a phenotypically neutral event, with epidemiological associations observed with various forms of cancer^{13,16-20}, autoimmune conditions^{21,22}, age-related macular degeneration²³, cardiovascular disease²⁴, Alzheimer's disease²⁵, type 2 diabetes¹⁵, obesity¹⁵, and all-cause mortality^{15,16}. The extent to which such observations represent a causal association, reverse causality or confounding is unclear. Furthermore, if these do represent causal effects, the mechanisms underlying such effects are unknown.

Key questions are whether loss of a Y chromosome from circulating leukocytes has a direct functional effect (for example, impairs immune function) and whether LOY in leukocytes is a barometer of broader genomic instability in leukocytes and other cell types. Understanding the mechanisms that drive clonal mosaicism and identifying genes which promote proliferative advantage to cells may help answer these questions and provide important insights into mechanisms of diseases of ageing. To this end we sought to identify novel susceptibility loci for LOY, an attractive form of clonal mosaicism to study given its relative ease of detection and high prevalence in the male population. Previous genome-wide association studies (GWAS) for LOY identified 19 common susceptibility loci and highlighted its relevance as a biomarker of cell cycle efficiency and DNA

damage response (DDR) in leukocytes^{13,14}. Here, we adapt a recently described computational approach⁶ to detect LOY in over 200,000 men from the UK Biobank study. We identify 137 novel loci which we use, along with the known 19 loci¹⁴, to demonstrate a shared genetic architecture between LOY, non-haematological cancer susceptibility and reproductive ageing in women. These data, in aggregate, support the hypothesis that LOY in leukocytes is a biomarker of genomic instability in other cell types with functional consequences across diverse biological systems.

Results

Previous studies assessing LOY have used a quantitative measure derived from the average intensity log-R ratio (termed mLRR-Y) of all array-genotyped Y chromosome single-nucleotide polymorphisms (SNPs). Here, we adapted a recently developed long-range phasing approach for mosaic event detection⁶ to estimate a dichotomous classification, which uses allele-specific genotyping intensities in the pseudo-autosomal region (we term this PAR-LOY, see **Methods**). This was applied to 205,011 men from UKBB (aged 40-70) in whom we identified 41,791 (20%) with detectable LOY. Men classified as LOY had an mLRR-Y score (derived using variants outside of the PAR) 0.9 standard deviations lower on average (95% CI 0-88-0.9) than non-LOY males (mean mLRR-Y -0.046 vs 0.009), reflecting the expected lower level of intensity due to reduced Y chromosome genetic material. Consistent with previous observations of clonal mosaicism, current smokers were at a higher risk of LOY (odds ratio (OR) 1.62 [95% CI 1.57-1.66]) and there was a strong association with age; the prevalence increased from 2.5% at age 40 to 43.6% at age 70 (**Figure 1**).

The genetic architecture of mosaic Y chromosome loss

We estimated a heritability of 31.7% (95CI 29.9 to 33.6%) for LOY, distributed across all individual chromosomes in proportion to their relative sizes (**Figure S1**). To identify individual genetic variants underlying this heritability we performed a GWAS for LOY, identifying 18,146 variants with genome-wide significant associations ($P < 5 \times 10^{-8}$). We resolved these into 156 statistically independent signals (**Table S1**), which included all 19 loci previously reported¹⁴. Effect sizes for these 156 associations ranged from OR 1.03-2.02, with LOY risk allele frequencies between 0.25% and 99.8% (**Figure 2**). An analysis excluding men with any past or current cancer diagnosis demonstrated no change in beta estimates across the 156 loci (**Figure S2**) and a drop in mean χ^2 proportionate to the drop in sample size. Despite the strong correlation between smoking behaviour and LOY, we found no evidence for a modifying effect of smoking across the 156 loci combined ($P_{\text{interaction}} > 0.05$).

We directly compared the power of our PAR-LOY calls to the previously used mLRR-Y derived measures by performing an mLRR-Y based GWAS in the same current study samples (**Table S1**). Across the 156 loci we observed an average ~ 2.5 x increase in χ^2 association statistic, exemplified by the strongest associated variant (rs17758695-*BCL2*) increasing in significance from $P_{\text{mLRRY}} = 7.5 \times 10^{-65}$ to $P_{\text{PAR-LOY}} = 4.1 \times 10^{-147}$. Only 61 of the 156 loci would have reached genome-wide significance in an mLRR-Y based analysis. Across the genome the lambda GC (ratio of expected to observed median test statistic) increased from 1.15 to 1.20 (mean χ^2 from 1.28 to 1.47), with no evidence of signal inflation due to population structure (LD score regression intercept 1.01). Simulation analyses demonstrated that the power of PAR-LOY over mLRR-Y depends on both the sample size and ratio of PAR1 / non-PAR SNPs (**Methods** and **Figure S3**).

To confirm the validity of our identified signals we sought replication in three independent datasets. Firstly, we used data generated using 653,019 male research participants from the personal genetics company 23andMe, Inc. (**Table S1**). These samples differed from the discovery samples both in terms of DNA source (saliva rather than peripheral blood) and LOY measurement type (quantitative mLRR-Y rather than dichotomous PAR-LOY calls). Despite this heterogeneity, all but one of the 154 loci (2 failed QC) had directionally concordant effects (binomial sign test $P=1.4 \times 10^{-44}$), with 126 exhibiting nominally significant association ($P < 0.05$) and 88 at a more conservative threshold ($P < 0.05/156$). Secondly, we sought further confirmation from the Icelandic deCODE study ($N=8,715$) where LOY was estimated using sequence reads from whole genome sequencing (DNA extracted from blood), rather than array data. These data demonstrated an overall directional consistency of 94% across the associated loci (140/149 variants tested, binomial sign test $P=2.3 \times 10^{-31}$) and 74 nominal associations (**Table S1**). Third, we replicated our loci in a set of 95,380 Japanese ancestry men from the BBJ project, with LOY estimated using mLRR-Y in whole blood. Of the 100/156 variants which passed QC and were polymorphic in East Asians, 92 had a consistent direction of effect (binomial $P=3.2 \times 10^{-19}$). Of these, 29 reached genome-wide significance in these data alone and 73 had at least nominal association (**Table S1**).

Finally, a negative control analysis using mLRR-Y estimated in 245,349 UKBB women (**Table S1**) – reflecting experimental noise in intensity variation – did not produce any significant associations after Bonferroni correction across the 156 loci ($P_{\max}=4.3 \times 10^{-3}$). In aggregate, these data strongly suggest that our discovery analysis identifies genetic determinants of LOY that are robust to ancestry, measurement technique and DNA source.

Implicated genes, cell types and biological pathways

We used various approaches to move from genomic association to identifying potentially causal variants, functional genes, cell types and biological pathways associated with LOY (see **Methods**). First, we performed Bayesian fine-mapping (see **Methods**) to quantify the probability that any single variant at a locus was causal for LOY by disentangling the effects of linkage disequilibrium (LD) (**Figure 3, Table S2-S3**). Fine-mapping identified at least one variant with reasonable confidence (posterior probability [PP] > 10%) in 80% (101/126) of regions, including at least one very high confidence variant (PP > 75%) in 25% (31/126) of regions (**Figure 3A**). These variants were enriched in exons of protein coding genes, their promoters, their transcribed but untranslated regions, and in hematopoietic regulatory regions marked by accessible chromatin (**Figure 3B, Table S4**).

Using both fine-mapped variants and genome-wide polygenic signal (see **Methods**), we found that hematopoietic stem and progenitor cells (HSPCs) were the most strongly enriched cell-types for LOY associated variants (**Figure 3C, Figure S4, Table S5, Table S6**). Amongst the fine-mapped variants, we further subdivided this enrichment into 3 distinct temporal modes indicative of increasing regulatory capacity across haematopoiesis (**Figure 3D**). These observations suggest that many of our identified variants exert their effects directly in hematopoietic stem cells, rather than further differentiated white blood cell types. This is in stark contrast to variants associated with the production of terminal blood cell types, which are enriched at terminal blood progenitors and depleted in HSPCs²⁶.

We next used two approaches (see **methods**) to map associated genetic variants to genes via expression effects (eQTLs) in whole blood, implicating a total of 110 unique transcripts (**Table S8-S10**). This included the *HLA-A* gene, where our lead variant in this region (6:29835518_T_A) tagged

the HLA-A*02:01 allele (**Table S11**). We also identified genes harbouring a non-synonymous variant either fine-mapped (PP > 10%) or in high LD ($r^2 > 0.8$) with an index variant, highlighting 22 genes (**Table S8**).

Biological pathway analysis using two approaches (see **methods**) identified a number of associated pathways, the majority of which converged on aspects of cell cycle regulation and DNA damage response (**Table S12 and S13**).

Overlap between LOY associated variants and cancer susceptibility loci

While detectable clonal mosaicism is clearly associated with future risk of haematological cancers⁶, its relationship with other cancers is less clear. Using data curated by the Open Targets platform and gene set enrichment analysis (see **Methods**), we found that LOY-associated variants were preferentially found near genes involved in cancer susceptibility ($P=9.9 \times 10^{-7}$), somatic drivers of tumour growth ($P=7 \times 10^{-4}$) and targets of approved or in trial cancer therapies ($P=0.05$). In total, 18 of the 156 mosaic leukocyte LOY associated variants were correlated ($r^2 > 0.1$) with known susceptibility variants for one or more type of non-haematological cancer (**Table S14**), including breast, prostate, testicular, kidney, melanoma and brain. Notable examples include a loss-of-function variant in *CHEK2* (rs186430430 $r^2 \sim 1$ with frameshift variant 1100delC) which confers a ~ 2.3 fold high risk of breast cancer²⁷, and an intronic signal (rs56345976) in the telomerase reverse transcriptase (*TERT*) gene which is in modest LD ($r^2 \sim 0.12$) with variants associated with longer telomeres and with increased risks of breast, ovarian, prostate cancers and glioblastoma, but also seen to be protective in other cancers²⁸.

To systematically assess the relationship between LOY susceptibility and cancer risk, we tested a genetic risk score (using a summary statistic Mendelian Randomization style framework, see **methods**) comprised of our 156 variants on two male-specific cancers (**Figure 4, Table S15**). Genetically-predicted LOY was associated with both increased risk of prostate cancer (OR=1.68 95% CI 1.33-2.11, $P=1.9 \times 10^{-5}$) and testicular germ cell tumour (OR 2.97 (1.45-6.07) $P=0.003$). Additional publicly available GWAS data for cancers in both sexes showed (**Figure 4, Table S15**) directionally consistent associations for glioma (OR 2.36 (1.34-4.17) $P=0.004$), renal cell carcinoma (OR 2.00 (1.24-3.21) $P=0.005$), lung cancer (OR 1.28 (0.98-1.68), $P=0.07$), colorectal cancer (OR 1.18 (0.93-1.50), $P=0.16$) and chronic lymphocytic leukaemia (OR 1.27 (0.75-2.16) $P=0.37$).

Genetic predisposition to LOY is associated with health outcomes in women

Mosaic LOY in blood cells has been associated with a broad range of diseases, which if causal is likely explained by one (or both) of two mechanisms: either LOY in leukocytes has a direct physiological effect, for example through impaired immune function, and/or it acts as a barometer and readily detectable manifestation of genomic instability occurring in parallel in other tissues. Ideally, this question would be addressed by assessing clonal mosaicism in large population studies where DNA was extracted from a broad range of cell and tissue types. In the absence of such a study, we hypothesised that testing the relevance of our identified LOY associated variants in women would help inform this – any association between the two could not be explained by a direct effect of LOY, given that females are XX.

To assess this we tested a polygenic risk score comprised of our 156 lead variants for association with three female-specific cancers – breast, endometrial and ovarian (**Figure 4, Table S15**). We observed a significant association with breast cancer (OR 1.25 (1.04-1.49) $P=0.016$) and directionally consistent results in the smaller endometrial (OR 1.18 (0.94-1.48), $P=0.14$) and ovarian (OR 1.02 (0.81-1.30), $P=0.86$) studies.

We next tested the same score on a female-specific non-cancer trait also underpinned by genome instability – age at natural menopause. Previous human and animal studies have shown menopause age is substantially biologically determined by the ability of oocytes to detect, repair and respond to DNA damage^{29,30}. We found that genetically increased risk of LOY was associated with later age at menopause ($P=0.003$, **Table S16**), with the *CHEK2* locus individually reaching genome-wide significance ($P=7.9 \times 10^{-22}$). A repeated genetic risk score analysis excluding *CHEK2* retained significance ($P=0.017$).

Given this observation that genetic susceptibility to LOY in leukocytes is impacting broader biological systems in these women, it is reasonable to speculate that actual LOY in leukocytes in men similarly represents a biomarker of genome instability in other cell and tissue types.

Exploring the impact of LOY at the level of a single cell

To help understand if and why LOY may provide a growth advantage to a cell, and the potential mechanisms linking LOY to disease, we performed single cell transcriptomic analyses (scRNAseq) using the 10X Genomics Chromium Single Cell 3' platform. This was performed on peripheral blood mononuclear cells (PBMCs) collected from 19 male donors (aged 64-89), unselected for any measure of clonal mosaicism. After standard quality control steps (see **methods**), we sequenced and profiled gene expression across 86,160 single cells. Under normal conditions, blood cells express a set of genes located in the male specific region of the Y chromosome (MSY). The LOY status of individual cells could therefore be determined by the absence of expression from these genes, which we identified in 13,418 of the cells (15.6% across all cells, ranging from 7-61% within individuals).

We next tested whether any of the genes highlighted in by our LOY GWAS were differentially expressed between cells with and without the Y chromosome (**Figure 5**). This analysis highlighted *TCL1A* (mapped LOY locus rs2887399, 162bp away), where the LOY risk increasing allele is associated with higher *TCL1A* expression in blood (**Table S10**). The single cell data showed that, among the major types of leukocytes, the *TCL1A* gene was expressed only in B-lymphocytes (**Figure 5**) and LOY was detected in 11.3% of these cells, ranging from 2% to 56% within individuals. B-lymphocytes without the Y chromosome (cell N=277) had 75% higher normalized *TCL1A* expression compared to those with a Y chromosome (N=2,459, Wilcoxon test in Seurat: fold change=1.75, $P<0.0001$). We also performed an in-house resampling test to evaluate this difference and validated a substantial upregulation of *TCL1A* in LOY cells (resampling test: fold change=1.68, $P<0.0001$) (**Figure 5**). An analysis within each individual demonstrated single cells with LOY had consistently higher *TCL1A* expression, ruling out any bias by *TCL1A* genotype (**Figure S5**).

To evaluate the magnitude of the 75% overexpression of the *TCL1A* gene in LOY B-lymphocytes, we compared the expression changes of other genes proximal to our identified GWAS loci. Of the genes we prioritized at each of our GWAS loci (“consensus genes”, **Table S8**), 71 were expressed in >5% of

the B-lymphocytes and included in the comparison, but only *TCL1A* demonstrated significant fold change (**Figure 5**).

TCL1A (encoding T cell leukemia/lymphoma 1A), functions as a co-activator of the cell survival kinase AKT and is often over expressed in T cell and B cell hematological malignancies³¹. These data provide a possible explanation for the growth advantage conferred to cells missing a Y chromosome. The independent effect of *TCL1A* genotype also suggests a possible bidirectional involvement for *TCL1A*. Ultimately further experimental work will be required to fully elucidate the aetiological implications of altered *TCL1A* expression in these cells.

Discussion

This study provides several advances in our understanding of the likely underlying biology and probable consequences of mosaic Y chromosome loss in circulating leukocytes. Our newly enhanced ability to detect LOY and expanded sample size led to an 8-fold increase in the number of associated genetic determinants, which we use to make several important observations.

The origin of LOY at the level of a single cell is perhaps most readily explained by chromosome mis-segregation events during mitosis. Consistent with this, many of the identified loci harbour nearby genes involved in key mitotic processes (**Figure 6**), notably central components of condensin which affects mitotic chromosome structure (*NCAPG2*, *SMC2*)³², and the assembly, structure and function of the kinetochore (*CENPN*, *CENPU*, *PMF1*, *ZWILCH*) and spindle (*SPDL1*), which together form the main machinery of chromosome congression and segregation^{33,34}. *MAD2L1* (alongside *MAD1L1* and *MAD2L1BP*) and *ZWILCH* are core components of the mitotic spindle assembly checkpoint³⁵, which ensures that chromatids are bi-orientated at the metaphase plate and under bipolar tension before disinhibiting the anaphase-promoting complex (of which *ANAPC5* is a component) to allow progression from metaphase. Many genes governing wider cell cycle progression, including cyclins (*CCND2*, *CCND3*), regulators of cyclin (*CDKN1B*, *CDKN1C*, *CDK5RAP1*) and major checkpoint kinases (*ATM*) are also identified here, emphasising the importance of processes across the cell cycle in determining LOY. A remainder of the genes that we identify encode proteins involved in sensing and responding to DNA damage (*SETD2*, *DDB2*, *PARP1*, *ATM*, *TP53*, *CHEK2*) and apoptotic processes (*PMAIP1*, *SPOP*, *LTBR*, *SGMS1*, *TP53INP1*, *DAP*). Of note, *FANCL* – the nearest gene to one of our lead variants - is vital to DNA interstrand crosslink repair and mutations in this gene have been linked to a rare autosomal form of Fanconi Anaemia, characterised by cytogenetic instability and chromosome breakage³⁶. The Bcl-2 family, a conserved set of proteins that regulate caspase-mediated apoptosis by controlling mitochondrial release of Cytochrome-C, are also particularly well-represented (*BCL2*, *BAX*, *BCL2L1*, *BCL2L11*)³⁷. These themes are consistent with the hypothesis that, secondary to the initial mis-segregation event, clonal expansion of LOY cells requires an environment permissive to proliferation of aneuploid cells, in which normal processes to detect and terminate these cells are avoided.

A link between LOY and cancer susceptibility seems plausible conceptually, given the nature of the genes identified. Here, we find substantial overlap of LOY associated variants across known cancer susceptibility loci, somatic drivers of tumour growth and genes targeted by licensed or in-trial cancer therapeutics. A notable example is the target of PARP inhibitors *PARP1*, where the lead SNP is highly correlated with a missense variant (V762A), the minor allele for which (the alanine substitution) is

protective for LOY and has experimentally been shown to reduce PARP-1 catalytic activity by 30-40%³⁸. More broadly, we found evidence for a systematic relationship between genetic susceptibility for LOY and risk of breast, prostate, testicular and renal cell carcinomas (**Figure 4**).

Based on our observations, we propose that LOY is determined by a “common soil” of shared mechanisms that predispose to genome instability and cancer. This is perhaps most readily apparent with the observation that genetic susceptibility to LOY is associated with cancer susceptibility in women and age at natural menopause. Although in aggregate we found that LOY associated variants tended to delay menopause age, there was substantial heterogeneity in dose-response. This was exemplified by individually significant associations for LOY risk increasing alleles with both earlier (mapped genes *MON1A*, *NR6A1*, *FBXL20*) and later menopause (*PMF1*, *JMJD1C*, *USP35/GAB2*, *APOLD1/CDKN1B*, *PHF11/RCBTB1*, *HEATR3*, *TP53* and *CHEK2*). We hypothesised that the direction of effect a LOY associated risk allele has on menopause may shed light on its mechanism of effect. Genetic determinants of LOY must broadly act either by promoting chromosomal instability or facilitating the clonal expansion of aneuploid cells, both of which are potentially cancer causing mechanisms. An allele that promotes clonal mosaicism through predisposition to chromosome mis-segregation or repair/generation of DNA damage would likely be associated with an earlier menopause due to acquired DNA damage in oocytes and their subsequent elimination. This principle is most evident in mice and humans with *BRCA1/2* loss of function, where diminished double strand break repair in oocytes triggers apoptosis and depletion of the ovarian reserve^{30,39,40}. In contrast, any process that impairs DNA damage sensing or programmed cell death may also promote clonal mosaicism (via greater tolerance of damaged cells) but lead to a *later* menopause. The clearest example of this is *CHEK2*, which in mice is essential for culling oocytes bearing unrepaired DNA double-strand breaks⁴¹. Reduced activity of this gene would therefore lead to the survival of defective oocytes (hence later menopause), supported for the first time in humans by our observation that *CHEK2* loss of function is associated with later age at menopause in women and increased LOY in men. The overall trend for LOY associated loci to be associated with delayed menopause suggests that many genetic determinants may act to inhibit apoptosis and the elimination of defective leukocytes. Further experimental work in animal and cellular models should aim to investigate more thoroughly the mechanisms linking each of our putatively highlighted genes to clonal mosaicism and broader outcomes.

We also note overlap between our identified LOY associated loci and other complex traits and diseases. For example, seven of our current LOY signals are correlated with previously reported⁴² susceptibility loci for Type 2 diabetes (*TP53INP1*, *SUGP1*, *KCNQ1*, *CCND2*, *EIF2S2*, *PTH1R* and *BCL2L11*). At six of these overlapping loci, the LOY risk-increasing allele also increases the risk of Type 2 diabetes. *CCND2* encodes Cyclin D2, the major D-type cyclin expressed in pancreatic β -cells and is essential for adult β -cell growth⁴³. *TP53INP1* is a p53-inducible gene, whose product regulates p53-dependent apoptosis. Additionally, the LOY-associated genes encoding cyclins and cyclin-dependent kinases, *CCND3*, *CDKN1B* and *CDKN1C*, are also implicated in pancreatic β -cell growth and maturation. We hypothesise that the previously reported association between clonal mosaicism in blood and T2D^{15,44} may reflect a common susceptibility to cell cycle dysregulation and genome instability, which lead to both increased clonal mosaicism and reduced pancreatic β -cell mass. Future studies should aim to more systematically assess the relationships and potential mechanisms linking LOY-associated variants and these broader health outcomes.

Finally, the “common soil” hypothesis discussed above does not preclude the possibility that LOY in leukocytes also has a direct role in disease, for example through impaired immune function⁴⁵. A growing awareness of the physiological importance of chromosome Y outside of reproductive development challenges the view of this chromosome as a “genetic wasteland”⁴⁶. The male-specific region (MSY) encodes 45 distinct proteins, with roles in fundamental processes such as chromatin modification (*KDM5D*, *UTY*), gene transcription (*ZFY*) and translation (*DDX3Y*, *EIF1AY* and *RPS4Y1*). Indeed, our observation in single-cell RNA sequencing data that leukocytes with LOY have dysregulated autosomal gene expression supports the notion of a direct physiological effect.

We hope that future experimental studies may build on these observations, yielding further insights into mechanisms that may have broad relevance to a range of cancers and other ageing-related diseases.

Acknowledgements

This research has been conducted using the UK Biobank Resource under application 9905 and 19808. This work was supported by the Medical Research Council [Unit Programme number MC_UU_12015/2]. Full study-specific and individual acknowledgements can be found in the supplementary information.

Conflicts of interest

L.A.F. and J.P.D. are cofounders and shareholders in Cray Innovation AB

References

1. Lee-Six, H. *et al.* Population dynamics of normal human blood inferred from somatic mutations. *Nature* **561**, 473–478 (2018).
2. Jacobs, K. B. *et al.* Detectable clonal mosaicism and its relationship to aging and cancer. *Nat. Genet.* **44**, 651–658 (2012).
3. Forsberg, L. A., Gisselsson, D. & Dumanski, J. P. Mosaicism in health and disease - clones picking up speed. *Nat. Rev. Genet.* **18**, 128–142 (2017).
4. Zink, F. *et al.* Clonal hematopoiesis, with and without candidate driver mutations, is common in the elderly. *Blood* **130**, 742–752 (2017).
5. Vattathil, S. & Scheet, P. Extensive Hidden Genomic Mosaicism Revealed in Normal Tissue. *Am. J. Hum. Genet.* **98**, 571–578 (2016).
6. Loh, P.-R. *et al.* Insights into clonal haematopoiesis from 8,342 mosaic chromosomal alterations. *Nature* **559**, 350–355 (2018).
7. JACOBS, P. A., COURT BROWN, W. M. & DOLL, R. Distribution of human chromosome counts in relation to age. *Nature* **191**, 1178–1180 (1961).
8. JACOBS, P. A., BRUNTON, M., COURT BROWN, W. M., DOLL, R. & GOLDSTEIN, H. Change of human chromosome count distribution with age: evidence for a sex differences. *Nature* **197**, 1080–1081 (1963).
9. Forsberg, L. A. *et al.* Mosaic loss of chromosome Y in leukocytes matters. *Nat. Genet.* **51**, 4–7 (2019).
10. Laurie, C. C. *et al.* Detectable clonal mosaicism from birth to old age and its relationship to cancer. *Nat. Genet.* **44**, 642–650 (2012).
11. Machiela, M. J. *et al.* Characterization of large structural genetic mosaicism in human autosomes. *Am. J. Hum. Genet.* **96**, 487–497 (2015).
12. Dumanski, J. P. *et al.* Mutagenesis. Smoking is associated with mosaic loss of chromosome Y. *Science* **347**, 81–83 (2015).
13. Zhou, W. *et al.* Mosaic loss of chromosome Y is associated with common variation near TCL1A. *Nat. Genet.* **48**, 563–568 (2016).
14. Wright, D. J. *et al.* Genetic variants associated with mosaic Y chromosome loss highlight cell cycle genes and overlap with cancer susceptibility. *Nat. Genet.* **49**, 674–679 (2017).
15. Loftfield, E. *et al.* Predictors of mosaic chromosome Y loss and associations with mortality in the UK Biobank. *Sci. Rep.* **8**, 12316 (2018).
16. Forsberg, L. A. *et al.* Mosaic loss of chromosome Y in peripheral blood is associated with shorter survival and higher risk of cancer. *Nat. Genet.* **46**, 624–628 (2014).
17. Noveski, P. *et al.* Loss of Y Chromosome in Peripheral Blood of Colorectal and Prostate Cancer Patients. *PLoS One* **11**, e0146264 (2016).
18. Machiela, M. J. *et al.* Mosaic chromosome Y loss and testicular germ cell tumor risk. *J. Hum. Genet.* **62**, 637–640 (2017).

19. Ganster, C. *et al.* New data shed light on Y-loss-related pathogenesis in myelodysplastic syndromes. *Genes. Chromosomes Cancer* **54**, 717–724 (2015).
20. Loftfield, E. *et al.* Mosaic Y loss is moderately associated with solid tumor risk. *Cancer Res.* (2018). doi:10.1158/0008-5472.CAN-18-2566
21. Persani, L. *et al.* Increased loss of the Y chromosome in peripheral blood cells in male patients with autoimmune thyroiditis. *J. Autoimmun.* **38**, J193–6 (2012).
22. Lleo, A. *et al.* Y chromosome loss in male patients with primary biliary cirrhosis. *J. Autoimmun.* **41**, 87–91 (2013).
23. Grassmann, F. *et al.* Y chromosome mosaicism is associated with age-related macular degeneration. *Eur. J. Hum. Genet.* (2018). doi:10.1038/s41431-018-0238-8
24. Haitjema, S. *et al.* Loss of Y Chromosome in Blood Is Associated With Major Cardiovascular Events During Follow-Up in Men After Carotid Endarterectomy. *Circ. Cardiovasc. Genet.* **10**, e001544 (2017).
25. Dumanski, J. P. *et al.* Mosaic Loss of Chromosome Y in Blood Is Associated with Alzheimer Disease. *Am. J. Hum. Genet.* **98**, 1208–1219 (2016).
26. Caleb A Lareau, Jacob C Ulirsch, Erik L Bao, Leif S Ludwig, Michael H Guo, Christian Benner, Ansuman T Satpathy, Rany Salem, Joel N Hirschhorn, Hilary K Finucane, Martin J Aryee, Jason D Buenrostro, V. G. S. Interrogation of human hematopoiesis at single-cell and single-variant resolution. *bioRxiv* (<https://www.biorxiv.org/content/early/2018/01/28/255224>)
27. Schmidt, M. K. *et al.* Age- and Tumor Subtype-Specific Breast Cancer Risk Estimates for CHEK2*1100delC Carriers. *J. Clin. Oncol.* **34**, 2750–2760 (2016).
28. Wang, Z. *et al.* Imputation and subset-based association analysis across different cancer types identifies multiple independent risk loci in the TERT-CLPTM1L region on chromosome 5p15.33. *Hum. Mol. Genet.* **23**, 6616–33 (2014).
29. Day, F. R. *et al.* Large-scale genomic analyses link reproductive aging to hypothalamic signaling, breast cancer susceptibility and BRCA1-mediated DNA repair. *Nat. Genet.* **47**, 1294–303 (2015).
30. Titus, S. *et al.* Impairment of BRCA1-related DNA double-strand break repair leads to ovarian aging in mice and humans. *Sci. Transl. Med.* **5**, 172ra21 (2013).
31. Laine, J., Künstle, G., Obata, T., Sha, M. & Noguchi, M. The protooncogene TCL1 is an Akt kinase coactivator. *Mol. Cell* **6**, 395–407 (2000).
32. Hirota, T., Gerlich, D., Koch, B., Ellenberg, J. & Peters, J.-M. Distinct functions of condensin I and II in mitotic chromosome assembly. *J. Cell Sci.* **117**, 6435–6445 (2004).
33. Petry, S. Mechanisms of Mitotic Spindle Assembly. *Annu. Rev. Biochem.* **85**, 659–83 (2016).
34. Godek, K. M., Kabeche, L. & Compton, D. A. Regulation of kinetochore-microtubule attachments through homeostatic control during mitosis. *Nat. Rev. Mol. Cell Biol.* **16**, 57–64 (2015).
35. London, N. & Biggins, S. Signalling dynamics in the spindle checkpoint response. *Nat. Rev. Mol. Cell Biol.* **15**, 736–47 (2014).

36. Meetei, A. R. *et al.* A novel ubiquitin ligase is deficient in Fanconi anemia. *Nat. Genet.* **35**, 165–70 (2003).
37. Cory, S. & Adams, J. M. The Bcl2 family: regulators of the cellular life-or-death switch. *Nat. Rev. Cancer* **2**, 647–56 (2002).
38. Zaremba, T. *et al.* Poly(ADP-ribose) polymerase-1 (PARP-1) pharmacogenetics, activity and expression analysis in cancer patients and healthy volunteers. *Biochem. J.* **436**, 671–9 (2011).
39. Lin, W., Titus, S., Moy, F., Ginsburg, E. S. & Oktay, K. Ovarian Aging in Women With BRCA Germline Mutations. *J. Clin. Endocrinol. Metab.* **102**, 3839–3847 (2017).
40. Weinberg-Shukron, A. *et al.* Essential Role of BRCA2 in Ovarian Development and Function. *N. Engl. J. Med.* **379**, 1042–1049 (2018).
41. Bolcun-Filas, E., Rinaldi, V. D., White, M. E. & Schimenti, J. C. Reversal of female infertility by Chk2 ablation reveals the oocyte DNA damage checkpoint pathway. *Science* **343**, 533–536 (2014).
42. Xue, A. *et al.* Genome-wide association analyses identify 143 risk variants and putative regulatory mechanisms for type 2 diabetes. *Nat. Commun.* **9**, 2941 (2018).
43. He, L. M. *et al.* Cyclin D2 protein stability is regulated in pancreatic beta-cells. *Mol. Endocrinol.* **23**, 1865–1875 (2009).
44. Bonnefond, A. *et al.* Association between large detectable clonal mosaicism and type 2 diabetes with vascular complications. *Nat. Genet.* **45**, 1040–3 (2013).
45. Case, L. K. *et al.* The Y chromosome as a regulatory element shaping immune cell transcriptomes and susceptibility to autoimmune disease. *Genome Res.* **23**, 1474–85 (2013).
46. Maan, A. A. *et al.* The Y chromosome: a blueprint for men’s health? *Eur. J. Hum. Genet.* **25**, 1181–1188 (2017).
47. Diskin, S. J. *et al.* Adjustment of genomic waves in signal intensities from whole-genome SNP genotyping platforms. *Nucleic Acids Res.* **36**, e126 (2008).
48. Bycroft, C. *et al.* The UK Biobank resource with deep phenotyping and genomic data. *Nature* **562**, 203–209 (2018).
49. Loh, P.-R. *et al.* Reference-based phasing using the Haplotype Reference Consortium panel. *bioRxiv* 052308 (2016). doi:10.1101/052308
50. Loh, P.-R. *et al.* Efficient Bayesian mixed-model analysis increases association power in large cohorts. *Nat. Genet.* **47**, 284–290 (2015).
51. Yang, J. *et al.* Conditional and joint multiple-SNP analysis of GWAS summary statistics identifies additional variants influencing complex traits. *Nature Genetics* **44**, 369–375 (2012).
52. Lee, S. H., Wray, N. R., Goddard, M. E. & Visscher, P. M. Estimating missing heritability for disease from genome-wide association studies. *Am. J. Hum. Genet.* **88**, 294–305 (2011).
53. Day, F. R. *et al.* Causal mechanisms and balancing selection inferred from genetic associations with polycystic ovary syndrome. *Nat. Commun.* **6**, 8464 (2015).
54. 1000 Genomes Project Consortium *et al.* A global reference for human genetic variation.

- Nature* **526**, 68–74 (2015).
55. UK10K Consortium *et al.* The UK10K project identifies rare variants in health and disease. *Nature* **526**, 82–90 (2015).
 56. Gudbjartsson, D. F. *et al.* Large-scale whole-genome sequencing of the Icelandic population. *Nat Genet* **47**, 435–444 (2015).
 57. Kanai, M. *et al.* Genetic analysis of quantitative traits in the Japanese population links cell types to complex human diseases. *Nat. Genet.* **50**, 390–400 (2018).
 58. Trynka, G. *et al.* Disentangling the Effects of Colocalizing Genomic Annotations to Functionally Prioritize Non-coding Variants within Complex-Trait Loci. *Am. J. Hum. Genet.* **97**, 139–52 (2015).
 59. Finucane, H. K. *et al.* Partitioning heritability by functional annotation using genome-wide association summary statistics. *Nat. Genet.* **47**, 1228–1235 (2015).
 60. Finucane, H. K. *et al.* Heritability enrichment of specifically expressed genes identifies disease-relevant tissues and cell types. *Nat. Genet.* **50**, 621–629 (2018).
 61. Zhu, Z. *et al.* Integration of summary data from GWAS and eQTL studies predicts complex trait gene targets. *Nat. Genet.* **48**, 481–487 (2016).
 62. Vösa, U. *et al.* Unraveling the polygenic architecture of complex traits using blood eQTL meta-analysis. *bioRxiv* (2018).
 63. Gusev, A. *et al.* Integrative approaches for large-scale transcriptome-wide association studies. *Nat. Genet.* **48**, 245–252 (2016).
 64. Szklarczyk, D. *et al.* The STRING database in 2017: quality-controlled protein-protein association networks, made broadly accessible. *Nucleic Acids Res.* **45**, D362–D368 (2017).
 65. Segrè, A. V *et al.* Common inherited variation in mitochondrial genes is not enriched for associations with type 2 diabetes or related glycaemic traits. *PLoS Genet.* **6**, (2010).
 66. Benner, C. *et al.* FINEMAP: efficient variable selection using summary data from genome-wide association studies. *Bioinformatics* **32**, 1493–501 (2016).
 67. Casper, J. *et al.* The UCSC Genome Browser database: 2018 update. *Nucleic Acids Res.* **46**, D762–D769 (2018).
 68. Machiela, M. J. & Chanock, S. J. LDlink: a web-based application for exploring population-specific haplotype structure and linking correlated alleles of possible functional variants. *Bioinformatics* **31**, 3555–3557 (2015).
 69. Burgess, S., Dudbridge, F. & Thompson, S. G. Combining information on multiple instrumental variables in Mendelian randomization: comparison of allele score and summarized data methods. *Stat. Med.* **35**, 1880–1906 (2016).
 70. Bowden, J., Davey Smith, G., Haycock, P. C. & Burgess, S. Consistent Estimation in Mendelian Randomization with Some Invalid Instruments Using a Weighted Median Estimator. *Genet. Epidemiol.* **40**, 304–14 (2016).
 71. Bowden, J., Davey Smith, G. & Burgess, S. Mendelian randomization with invalid instruments: effect estimation and bias detection through Egger regression. *Int. J. Epidemiol.* **44**, 512–25

- (2015).
72. Schumacher, F. R. *et al.* Association analyses of more than 140,000 men identify 63 new prostate cancer susceptibility loci. *Nat. Genet.* **50**, 928–936 (2018).
 73. Turnbull, C. *et al.* Variants near DMRT1, TERT and ATF7IP are associated with testicular germ cell cancer. *Nat. Genet.* **42**, 604–7 (2010).
 74. Litchfield, K. *et al.* Identification of 19 new risk loci and potential regulatory mechanisms influencing susceptibility to testicular germ cell tumor. *Nat. Genet.* **49**, 1133–1140 (2017).
 75. Scelo, G. *et al.* Genome-wide association study identifies multiple risk loci for renal cell carcinoma. *Nat. Commun.* **8**, 15724 (2017).
 76. He, Y. *et al.* Exploring causality in the association between circulating 25-hydroxyvitamin D and colorectal cancer risk: a large Mendelian randomisation study. *BMC Med.* **16**, 142 (2018).
 77. May-Wilson, S. *et al.* Pro-inflammatory fatty acid profile and colorectal cancer risk: A Mendelian randomisation analysis. *Eur. J. Cancer* **84**, 228–238 (2017).
 78. McKay, J. D. *et al.* Large-scale association analysis identifies new lung cancer susceptibility loci and heterogeneity in genetic susceptibility across histological subtypes. *Nat. Genet.* **49**, 1126–1132 (2017).
 79. Melin, B. S. *et al.* Genome-wide association study of glioma subtypes identifies specific differences in genetic susceptibility to glioblastoma and non-glioblastoma tumors. *Nat. Genet.* **49**, 789–794 (2017).
 80. Atkins, I. *et al.* Transcriptome-Wide Association Study Identifies New Candidate Susceptibility Genes for Glioma. *Cancer Res.* **79**, 2065–2071 (2019).
 81. Michailidou, K. *et al.* Association analysis identifies 65 new breast cancer risk loci. *Nature* **551**, 92–94 (2017).
 82. Milne, R. L. *et al.* Identification of ten variants associated with risk of estrogen-receptor-negative breast cancer. *Nat. Genet.* **49**, 1767–1778 (2017).
 83. Phelan, C. M. *et al.* Identification of 12 new susceptibility loci for different histotypes of epithelial ovarian cancer. *Nat. Genet.* **49**, 680–691 (2017).
 84. O'Mara, T. A. *et al.* Identification of nine new susceptibility loci for endometrial cancer. *Nat. Commun.* **9**, 3166 (2018).
 85. Law, P. J. *et al.* Genome-wide association analysis implicates dysregulation of immunity genes in chronic lymphocytic leukaemia. *Nat. Commun.* **8**, 14175 (2017).
 86. Southam, L. *et al.* Whole genome sequencing and imputation in isolated populations identify genetic associations with medically-relevant complex traits. *Nat. Commun.* **8**, 15606 (2017).
 87. Han, B. *et al.* A general framework for meta-analyzing dependent studies with overlapping subjects in association mapping. *Hum. Mol. Genet.* **25**, 1857–1866 (2016).
 88. Han, B. & Eskin, E. Random-effects model aimed at discovering associations in meta-analysis of genome-wide association studies. *Am. J. Hum. Genet.* **88**, 586–598 (2011).

Tables and Figures

Figure 1 | Prevalence of mosaic Y chromosome loss by age in UK Biobank study participants




Figure 2 | Distribution of allele frequency and effect size for the 156 identified LOY loci




Figure 3 | Results from fine-mapping analyses. Panel **a** shows the posterior expected number of causal variants (*top*) as well as the best fine-mapped variant (*bottom*) in each region. Genomic enrichments for variants stratified by posterior probability are shown in panel **b**. Fine-mapped variants were enriched for accessible chromatin in hematopoiesis, as well as in exons, promoters, and UTRs of protein coding genes, but not for introns. Panel **c** shows g-chromVAR cell-type enrichments across the hematopoietic tree for LOY. HSCs, MPPs, and CMPs meet Bonferroni threshold ($\alpha = 0.05 / 18$). Developmental patterns of accessible chromatin for variants with posterior probability > 10% are shown in panel **d**, revealing that 14 variants are fully restricted to acting within HSPCs, 14 variants can also have regulatory effects in myeloid and lymphocyte progenitors, and 17 variants are capable of acting across the majority of hematopoiesis. K-means clustering (k = 4 determined by the gap statistic) was used to identify patterns of accessibility, and cell types were hierarchically clustered. HSC, hematopoietic stem cell; MPP, multi-potent progenitor; CMP, common myeloid progenitor; HSPC, hematopoietic stem and progenitor cell; M/L, myeloid and lymphoid; PP, posterior probability; AC, accessible chromatin; UTR, untranslated region; PChIC, promoter capture Hi-C; eQTL, expression quantitative trait locus; corr, ATAC/chromatin-RNA correlations




Figure 4 | Association between a genetic risk score of the 156 LOY-associated variants and cancer.






Figure 5 | Single-cell RNA sequencing results. Panel **a** shows clustering and identification of cell types using a tSNE plot generated from a pooled dataset including 86160 PBMC's isolated from peripheral blood samples collected from 19 male donors. The *TCL1A* gene was expressed in the B-lymphocytes as indicated by blue color in panel **b**. Analysis of LOY status in the B-lymphocytes identified 277 cells with LOY, plotted in red color in panel **c**. Panel **d** display the result from a resampling test performed to compare the expression of *TCL1A* in LOY B-lymphocytes with its expression in non-LOY B-lymphocytes. The grey and red curves in panel **d** represent the resampled distribution of *TCL1A* expression in non-LOY and LOY cells, respectively. The resampling test established an increased expression of *TCL1A* in B-lymphocytes with LOY (fold change=1.68, $p<0.0001$). Panel **e** display fold changes in gene expression between LOY and non-LOY B-lymphocytes for 71 selected genes from the list of genes mapping to the 156 index variants. Genes expressed in >5% of the investigated B-lymphocytes were included. The blue line at fold change 1 in panel **e** represents no differential expression and the red line shows the level of 50% overexpression in LOY cells.



1 **Figure 6 | Many LOY-associated genes converge on mechanistic and regulatory aspects of the cell**
2 **cycle.** All genes shown have been prioritized as potentially functional genes at our reported GWAS
3 loci; gene symbols may be shown more than once. Coloured indicators next to each gene symbol
4 specify the type of evidence on which it has been prioritized at its respective locus: blue, nearest
5 protein-coding gene; green, eQTL; red, contains a highly correlated non-synonymous variant. Red
6 boxes indicate each of the three known cell cycle checkpoints. Red inhibition connectors denote that
7 a target is inhibited by degradation, green by binding. Green arrows indicate a signaling cascade and
8 its effector or final physiological effect. Bidirectional dashed green arrows indicate the formation of

9 a complex between the products of the two connected genes. Excepting p53, proteins contained
10 within green boxes have not been implicated in this GWAS, but are important interactors of
11 implicated genes. CENPA-NAC, CENPA nucleosome-associated complex; APC/C, anaphase-promoting
12 complex/cyclosome; MC, mitotic checkpoint; CDK, cyclin-dependent kinase.

13



15 **Methods**

16 **Phenotype preparation in UK Biobank**

17 We adapted a recently developed statistical approach⁶ for detecting autosomal mosaic events to
18 identify male individuals with LOY based on allele-specific genotyping intensities in the
19 pseudoautosomal region (PAR) of the sex chromosomes. In contrast to previous work that has
20 quantified Y chromosome loss based on median genotyping intensity over the non-pseudoautosomal
21 region of the Y chromosome (mLRR-Y)¹²⁻¹⁵, our approach leverages the diploid nature of the PAR to
22 ascertain mosaic Y loss based on differences between maternal (X PAR) vs. paternal (Y PAR) allelic
23 intensities at heterozygous sites: mosaic Y loss causes Y PAR intensities to decrease relative to X PAR
24 intensities. This intuition can be harnessed even in population cohorts in which absolute phase
25 information (i.e., information about maternal vs. paternal inheritance of alleles) is unavailable: we
26 can overcome this obstacle by performing statistical phasing and subsequently identifying evidence
27 of an imbalance in allelic intensities between the two statistically phased haplotypes (accounting for
28 the possibility of phase switch errors)^{5,6}. In general, the signal produced by phased allelic imbalances
29 is typically much cleaner than estimates of total genotyping intensities (e.g., mLRR-Y), as the latter
30 can vary substantially across the genome due to technical artefacts⁴⁷.

31 We applied this approach to blood DNA genotyping intensity data from the full UK Biobank cohort (a
32 study described extensively elsewhere⁴⁸, analyzing 1,239 genotyped variants on PAR1 that passed
33 QC (out of 1,301 total PAR1 variants). (We ignored the much-shorter PAR2, which only contained 56
34 genotyped variants, of which 37 passed QC.) To maximize phasing accuracy, we phased the full
35 cohort including both males and females using Eagle2⁴⁹, after which we restricted our attention to
36 males. We called mosaic chromosomal alterations (mCAs) in PAR1 using a slightly modified version
37 of the pipeline we described previously⁶. Specifically, in our hidden Markov model, we increased the
38 probability of starting in a mosaic state to 0.2: we observed in a preliminary analysis that Y loss
39 events were much more common than autosomal events, so we updated this prior accordingly in
40 our final analysis to slightly improve the model (and verified that the number of Y loss calls did not
41 drastically change, so no further update was necessary). We also post-processed our PAR1 mCA calls
42 to identify likely mosaic Y loss events based on two criteria: (i) mCA spans the full PAR1 region; and
43 (ii) observed mean \log_2 R ratio (LRR) is more consistent with a mosaic loss event than a CNN-LOH or
44 gain (after taking into account the s.e.m. of LRR and an empirical prior on mCA copy numbers⁶). This
45 procedure produced 44,709 mCA calls in PAR1 (at an estimated false discovery rate of 0.05) among
46 220,924 males passing sample QC, of which 43,306 were classified as likely LOY. These calls
47 contained an average of 321 heterozygous variants on PAR1 passing QC that were usually phased
48 perfectly (no switch errors detected by the hidden Markov model in 72% of calls).

49 We estimated FDR of PAR-LOY calls using the same phase randomization procedure (similar to
50 permutation testing) that we previously employed⁶. Specifically, we computed test statistics from
51 PAR-LOY on a batch of control data sets in which we had randomized phase assignments at
52 heterozygous variants (but otherwise kept the data unchanged). The test statistics produced by this
53 procedure gave us an approximate null distribution which we then used to estimate FDR.

54 Recalled age at natural menopause (ANM) was available in 106,237 women with genetic data. We
55 included women with ANM who were 40–60 years of age in our analyses, excluding those with

56 menopause induced by hysterectomy, bilateral ovariectomy, radiation or chemotherapy and those
57 using hormone replacement therapy (HRT) before menopause.

58 **Power comparison of PAR-LOY vs mLRR-Y**

59 The efficacy of the PAR-LOY approach relative to mLRR-Y depends primarily on two factors: (i) the
60 relative number of genotyping probes in PAR1 vs. Y-nonPAR, and (ii) the size of the cohort (which
61 determines phasing accuracy in PAR1). In UK Biobank, both (i) and (ii) are heavily in favour of PAR-
62 LOY, as the UK Biobank genotyping array contained nearly twice as many PAR1 vs. Y-nonPAR variants
63 (1301 vs. 691 variants) and the cohort size is extremely large (~500K individuals). For comparison,
64 the BioBank Japan genotyping contained only one-fifth as many PAR1 vs. Y-nonPAR variants (an
65 order of magnitude lower relative coverage of PAR1).

66 To quantify the effects of factors (i) and (ii) on PAR-LOY, we subsampled the UKBB data set to
67 simulate the effects of reduced PAR1 content and reduced phasing accuracy (due to smaller sample
68 size). Specifically, we applied PAR-LOY to 20 = 4x5 data sets in which we subsampled (i) the number
69 of PAR1 variants included in analysis (down-sampling 1x, 2x, 4x, 8x) and (ii) the number of samples
70 included in analysis (down-sampling 1x, 3.5x, 10x, 35x, 100x). In each scenario, we compared the
71 quality of PAR-LOY calls to mLRR-Y by comparing association strength at the top three Y loss GWAS
72 hits (*rs17758695 (BCL2)*, *rs2887399 (TCL1A)*, *rs59633341 (TSC22D2)*). We computed relative
73 association strength by taking the mean chi-square association test statistic across the three variants
74 for PAR-LOY and dividing by the corresponding quantity for mLRR-Y.

75 We observed (**Figure S3**) that indeed the PAR-LOY approach benefited considerably from high PAR1
76 genotyping coverage in UK Biobank as well as highly accurate phasing (with diminishing returns
77 beyond a cohort size of ~100K samples, presumably due to phasing accuracy becoming near-perfect
78 across PAR1). For large cohorts with >100K samples, our results indicate that the PAR-LOY approach
79 becomes advantageous when a genotyping array contains at least ~1/3 as many PAR1 variants as Y-
80 nonPAR variants.

81 **Genetic association testing in UK Biobank**

82 We used genetic data from the “v3” release of UK biobank⁴⁸, containing the full set of HRC and
83 1000G imputed variants. In addition to the quality control metrics performed centrally by UK
84 Biobank, we defined a subset of “white European” ancestry samples using a K-means clustering
85 approach applied to the first four principle components calculated from genome-wide SNP
86 genotypes. Individuals clustered into this group who self-identified by questionnaire as being of an
87 ancestry other than white European were excluded. After application of QC criteria, a maximum of
88 205,011 male participants were available for analysis with genotype and phenotype data.
89 Association testing was performed using a linear mixed models implemented in BOLT-LMM⁵⁰ to
90 account for cryptic population structure and relatedness. Only autosomal genetic variants which
91 were common (MAF>1%), passed QC in all 106 batches and were present on both genotyping arrays
92 were included in the genetic relationship matrix (GRM). Genotyping chip, age at baseline and 10
93 genetically derived principal components were included as covariates.

94 We defined statistically independent signals (described as lead or index variants) using 1Mb
95 distanced-based clumping across all imputed variants with $P < 5 \times 10^{-8}$, an imputation quality score >

96 0.5 and MAF > 0.1%. Genome-wide significant lead variants that shared any correlation with each
97 other due to long range linkage disequilibrium ($r^2 > 0.05$) were excluded from further consideration.
98 These loci were additionally augmented using approximate conditional analyses implemented in
99 GCTA⁵¹. Here, secondary signals were only considered if they were uncorrelated ($r^2 < 0.05$) with a
100 previously identified index variant and genome-wide significant pre and post conditional analysis.

101 The total trait variance of all genotyped SNPs was calculated genome-wide and per-chromosome
102 using restricted estimate maximum likelihood (REML) implemented in BOLT-LMM⁵⁰. The
103 corresponding observed-scale estimate was transformed to the liability-scale⁵².

104 **Replication**

105 Replication was performed in three independent studies using two separate techniques.

106 Firstly, we used data generated from the customer base of 23andMe Inc, a consumer genetics
107 company. Genotyping array quality control, imputation and downstream association testing for this
108 study has been described extensively elsewhere⁵³. All individuals provided informed consent and
109 answered surveys online according to 23andMe's human subjects protocol, which was reviewed and
110 approved by Ethical & Independent Review Services, a private institutional review board
111 (<http://www.eandireview.com>). DNA extraction and genotyping were performed on saliva samples
112 by National Genetics Institute (NGI), a CLIA licensed clinical laboratory and a subsidiary of Laboratory
113 Corporation of America Mosaic LOY was estimated by calculating the mean log-R ratio (normalised
114 signal intensity) across 274 SNPs on the male-specific region of the Y chromosome that are shared
115 and perform well across genotyping platforms, using the protocol described previously¹⁴. Imputation
116 was performed using a combination of the May 2015 release of the 1000 Genomes Phase 3
117 haplotypes⁵⁴ with the UK10K imputation reference panel⁵⁵. Genetic association testing was
118 performed using linear regression in 653,019 male research participants of European ancestry, using
119 age, genetically derived principal components and genotyping platform as covariates. Results were
120 adjusted for a genomic control inflation factor of 1.129.

121 Secondly, we analyzed whole-blood genome sequences of 8,715 Icelandic males⁵⁶ (age range 41-105
122 years, mean 63 years), that had been whole-genome sequenced by Illumina method to a mean
123 depth of 37x. As an estimate of chromosome Y copy-number we used the average read depth over
124 chromosome Y, using exclusively X-degenerate regions. This was computed by samtools from bam
125 files aligned to hg38 and normalized by genome-wide sequencing coverage for the subject. A total of
126 12 outlier individuals (copy-number greater than 1.25) were excluded. Association analysis was
127 performed using a linear mixed model implemented in BOLT-LMM⁵⁰ (to account for population
128 structure instead of genetic principal components) after inverse normal transformation and
129 adjustment for age at blood draw. Effect sizes for $\log_2(\text{chrY copy-number})$ were estimated using
130 robust linear regression (rlm from R package MASS).

131 Third, we used a sample of 95,380 Japanese ancestry men from the BBJ project, a study which has
132 been described extensively elsewhere⁵⁷. The study was approved by the ethical committees in the
133 Institute of Medical Science, the University of Tokyo and RIKEN Center for Integrative Medical
134 Science. Mosaic LOY in blood was estimated using the quantitative mLRR-Y measure, using a similar
135 protocol as previous studies¹⁴. Association testing was performed using a linear mixed model
136 implemented in BOLT-LMM⁵⁰, including age, smoking, disease status (35 case/control definitions)

137 and chip array as a covariate. Replication test statistics were assessed in a sensitivity model without
138 smoking and disease status covariates to ensure consistency between models.

139 **Genomic feature enrichment**

140 We used a previously modified version of GoShifter^{26,58} to calculate the enrichment of fine-mapped
141 (PP \geq 0.10) and not fine-mapped (PP $<$ 0.10) variants with genomic annotations by locally shifting the
142 annotations and computing overlaps to approximate the null distribution. Z-scores and odds ratios
143 were calculated from 1000 permutations, and typical two-tailed p-values are calculated from the z-
144 score statistic. All annotations were obtained as described previously²⁶.

145 In order to identify which tissue types were most relevant to genes involved in LOY, we applied LD
146 score regression⁵⁹ to specifically expressed genes (“LDSC-SEG”)⁶⁰ and g-chromVAR to hematopoietic
147 accessible chromatin²⁶. For LDSC-SEG, cell-type specific analyses using GTEx and Epigenome
148 Roadmap annotations were performed using the data available on the LDSC-SEG resource page
149 (<https://github.com/bulik/ldsc/wiki/Cell-type-specific-analyses>). For g-chromVAR, hematopoietic
150 specific analyses were performed using ATAC-seq count matrices as previously processed
151 (https://github.com/caleblareau/singlecell_bloodtraits). g-chromVAR estimates were averaged
152 across 10 different random background peak sets. We note that, similar to the derivation of cell-type
153 specific features or SEGs in LDSC, g-chromVAR z-scores represent relative enrichment for specific
154 cell-types compared to other input cell-types, which allows for discrimination between closely
155 related cell types in the hematopoietic lineage.

156 **Gene expression integration**

157 We used two approaches to map associated genetic variants to genes via expression effects (eQTLs)
158 in whole blood. Firstly, Summary Mendelian Randomization (SMR) uses summary-level gene
159 expression data to map potentially functional genes to trait-associated SNPs⁶¹. We ran this approach
160 using a meta-analysis of whole blood eQTL data from 31,684 individuals⁶². Only transcripts with no
161 evidence of pleiotropic effects, as assessed by the HEIDI metric were considered⁶¹. Secondly, we
162 used the recently described Transcriptome-wide Association Study (TWAS) approach⁶³ to infer gene
163 expression association using three whole blood datasets (Young Finns Study, Netherlands Twin
164 Registry cohorts and GTEx v6). All data used is available here: <http://gusevlab.org/projects/fusion/>.
165 For all analyses significance thresholds were set to adjust for the number of tested performed.

166 **Gene set enrichment analysis**

167 Pathway analysis was performed using two distinct approaches – STRING⁶⁴ and MAGENTA⁶⁵. For
168 STRING, only the gene closest to one of the 156 lead index variants (max distance 500kb) was
169 included in the analysis. In contrast, MAGENTA performs enrichment analysis using the full genome-
170 wide summary statistic data. In this gene set enrichment analysis (GSEA) approach, each gene in the
171 genome is mapped to a single index SNP with the lowest P-value within a 300kb window. This P-
172 value, representing a gene score, is then corrected for confounding factors such as gene size, SNP
173 density and LD-related properties in a regression model. Each mapped gene is then ranked by its
174 adjusted gene score. At a given significance threshold (here the 95th percentile of all gene scores),
175 the observed number of gene scores in a given pathway, with a ranked score above the specified

176 threshold percentile, is calculated. This observed statistic is then compared to 1,000,000 randomly
177 permuted pathways of identical size. This generates an empirical GSEA P-value for each pathway.

178 We used the Open Targets Platform (<https://www.targetvalidation.org/>) to define gene sets
179 comprising genes involved in cancer susceptibility (N=249), somatic drivers of tumour growth
180 (N=394), targets of approved or in trial cancer therapies (N=458), “affected pathways” (N=216 with
181 score = 1) and finally an overall aggregated score for involvement in cancer (N=934 with score = 1).
182 The various data sources, and approach applied by Open Targets to score and prioritise target genes
183 within each of these categories is described in full at [https://docs.targetvalidation.org/getting-](https://docs.targetvalidation.org/getting-started/scoring)
184 [started/scoring](https://docs.targetvalidation.org/getting-started/scoring)). We arbitrarily defined gene set membership based on an assigned score > 0.8
185 unless otherwise specified. These pathways were tested for enrichment in downstream analyses
186 using MAGENTA.

187 **Fine-mapping**

188 Regions for fine-mapping were defined by extended 0.5 Mb in both directions from each sentinel
189 and merging when regions overlapped, resulting in 126 total regions. All variants in these regions
190 with MAF > 0.005 and INFO > 0.6 were fine-mapped. Dosage LD was estimated from the UKB
191 genotype probability files (.bgen) using 167,020 unrelated white British male individuals
192 (<http://www.nealelab.is/uk-biobank/>). Fine-mapping was then performed using v1.3 of the
193 FINEMAP software⁶⁶ with default settings allowing for up to 5 causal variants in each region. The
194 UCSC genome browser was used to view individual variants along with hosted features⁶⁷.

195 **Integration with cancer data and modelling LOY as a causal exposure**

196 The NHGRI-EBI GWAS Catalogue database was accessed and downloaded on June 25, 2018. The
197 downloaded file was curated to only include studies in which cancer is the associated disease and
198 further filtered to remove variants with association p-values greater than 5×10^{-8} . Due to a potential
199 lag between the time a new GWAS is published and included in the NHGRI-EBI GWAS Catalogue, a
200 supplementary literature search of PubMed was performed to identify additional reports of cancer
201 susceptibility studies that were not included in the GWAS Catalogue. The literature search was
202 completed on July 18, 2018. LDlink⁶⁸ was used to identify published cancer GWAS-associated genetic
203 variants which are in linkage disequilibrium (LD) ($r^2 \geq 0.1$ based on the 1000 Genomes Project
204 European Population data) with one of the 154 LOY lead SNPs. Associations with haematological
205 malignancies were excluded and additional associations were identified by manual searches.

206 The relationship between LOY-associated variants and cancer was assessed using a series of two-
207 sample summary statistic based Mendelian randomisation-style analyses. Linear regressions of the
208 cancer log odds ratios (logOR) for each available SNP on the LOY beta coefficients were carried out,
209 weighted by the inverse of the variance of the cancer logORs. This is equivalent to an inverse-
210 variance weighted meta-analysis of the variant-specific causal estimates⁶⁹. Because of evidence of
211 over-dispersion (i.e. heterogeneity in the variant-specific causal estimates), the residual standard
212 error was estimated, making this equivalent to a random-effects meta-analysis. We repeated the
213 analyses using the Weighted Median method, which is robust to up to half of the genetic variants
214 not being valid instrumental variables⁷⁰. Unbalanced horizontal pleiotropy was tested based on the
215 significance of the intercept term in MR-Egger regression⁷¹. The analyses were carried out separately
216 for each type of cancer.

217 Summary statistics for the association between the genetic variants and risk of prostate cancer were
218 obtained from the PRACTICAL/ELLIPSE consortium, based on GWAS analyses of 67,158 prostate
219 cancer cases and 48,350 controls⁷². Testicular cancer summary statistics were obtained from two
220 GWAS studies conducted at the Institute of Cancer Research comprising 4,192 testicular cancer
221 cases and 12,368 controls^{73,74}. The renal cancer analysis used summary statistics from the Kidney
222 Cancer GWAS Meta-Analysis Project of 10,784 cases of renal cell carcinoma and 20,406 controls⁷⁵.
223 Colorectal cancer summary statistics were from eight UK-based GWAS studies, totalling 22,372
224 colorectal cancer cases and 44,271 controls^{76,77}. The summary statistics for overall lung cancer were
225 from GWAS analyses of 29,266 lung cancer cases and 56,450 controls conducted by the International
226 Lung Cancer Consortium⁷⁸. The glioma summary statistics were from GWAS studies of 12,488 cases
227 and 18,169 controls^{79,80}. The breast cancer analysis was based on summary statistics from GWAS
228 analyses of 105,974 breast cancer cases and 122,977 controls conducted by the Breast Cancer
229 Association Consortium (BCAC)⁸¹, including summary statistics from analyses restricted to cases with
230 estrogen receptor positive or estrogen receptor negative breast cancer⁸². Summary statistics for the
231 ovarian cancer analysis were from GWAS studies of 25,509 ovarian cancer cases and 48,941 controls
232 conducted by the Ovarian Cancer Association Consortium (OCAC)⁸³. The endometrial cancer results
233 were from GWAS studies of 12,906 endometrial cancer cases and 108,979 controls from the
234 Endometrial Cancer Association Consortium (ECAC)⁸⁴. In addition, analyses for breast and ovarian
235 cancer risk specifically in carriers of a *BRCA1* or a *BRCA2* mutation were carried out using results
236 from GWAS studies conducted by the CIMBA consortium^{82,83}. Although our focus was on the
237 association between LOY and risk of non-haematological cancers, we also included an analysis using
238 summary statistics from GWAS studies of 4,478 chronic lymphocytic leukaemia (CLL) cases and
239 13,213 controls⁸⁵. There was some overlap in the control subjects used by the breast, ovarian,
240 endometrial and colorectal cancer studies, between the control subjects used in the prostate,
241 colorectal and testicular cancer studies, and between the control subjects used in the endometrial
242 cancer, glioma and CLL studies. All the cancer analyses were based on summary statistics from
243 studies restricted to participants with European ancestry.

244 In addition to the analyses by cancer type, we also used summary statistics from a pan-cancer meta-
245 analysis study of breast, prostate, ovarian, and endometrial cancer to look for a more general
246 association between LOY and cancer risk. The pan-cancer summary statistics were derived using a
247 three step procedure. First, the tetrachoric correlation of binary transformed Z-scores was used to
248 estimate the correlation between individual-cancer summary statistics that is attributable to control
249 sample overlap⁸⁶. Second, individual-cancer summary statistic standard errors were decoupled to
250 account for the estimated correlation (resulting from shared controls amongst the meta-analysed
251 cancer strata)⁸⁷ and third, the METASOFT software⁸⁸ was used to perform fixed effect inverse-
252 variance weighted meta-analyses for the combination of four cancers.

253 **Sample preparation for single cell gene expression study**

254 Blood samples from 19 elderly men (median age=80, range=64-89) admitted to the Geriatrics
255 Department at Uppsala University Hospital (Uppsala, Sweden) were collected in BD Vacutainer CPT
256 cell separation tubes containing sodium citrate and stored on ice. The PBMC fraction was isolated
257 from the whole blood samples by density gradient centrifugation following manufacturer
258 instructions (Becton, Dickinson and Company, Franklin Lakes). PBMCs were collected and suspended
259 in cold 1X PBS solution with 0.04% BSA. Cell concentrations were measured using an EVE cell counter

260 (NanoEnTek, Seoul) and diluted to a concentration of 10^6 cells/ml. All the prepared samples had a
261 cell viability above 90%. The local research ethics committee approved the study and all participants
262 provided their informed consent.

263 **Single cell workflow**

264 We performed single cell RNA sequencing (scRNAseq) using the 10X Chromium Single Cell 3' gene
265 expression solution (10X Genomics, Inc.) at the SNP&SEQ Technology Platform at Uppsala University
266 (Sweden). This scRNAseq technology is based on gel beads loaded with barcoded oligos mixed with
267 single cells and enzymes, before captured in droplets (GEMs). The transcripts present in individual
268 cells are barcoded with UMI's (unique molecular identifiers) and used to prepare standard
269 sequencing libraries. All transcripts from single cells get barcoded with the same index sequence
270 allowing for the transcripts from thousands of single cells to be pooled together in a single
271 sequencing run and allowing transcriptional profiling of individual cells. The barcoding and library
272 construction were performed for the 19 PBMC samples using the Chromium Single Cell 3' Reagent
273 kit (cat# 120236/37/62) according to the manufacturer protocol (CG00052 Single Cell 3' Reagent Kit
274 v2 User Guide). The entire procedure, from blood sampling to construction of GEM's was
275 accomplished within 5 hours. The generated single cell libraries were sequenced using a NovaSeq
276 6000 instrument (Illumina, Inc., San Diego) at the SNP&SEQ Technology Platform and generated a
277 median of 64900 reads per cell (range=35213-111643).

278 **Single-cell bioinformatics pipeline**

279 Sequenced reads were mapped to the human reference (GRCh37/hg19) using the software
280 Cellranger v 2.0.2 (10X Genomics, Inc.). Cellranger produces a count-matrix for each experiment
281 containing the UMI barcodes using sequence information from the 3' end of each transcript in every
282 single cell. We used the R library Seurat (v2.3.1) for further processing and implemented the
283 standard Seurat workflow⁷. Specifically, QC-steps were performed including removal of apoptotic
284 cells (i.e. cells with more than 5% mitochondrial RNA) as well as removing cells with low sequencing
285 coverage and/or a low number of expressed genes (demanding at least 350 genes expressed and
286 800 UMIs). Following QC-steps, normalization of the gene expression within each single cell was
287 performed using the function "NormalizeData" and the built in LogNormalize normalization method.
288 After this the most variable genes was identified using the function FindVariableGenes with the
289 parameters x.low.cutoff = 0.2, x.high.cutoff = 4, y.cutoff = 0.5. All expression values was following
290 this scaled using the "ScaleData" function and principal components were calculated based on the
291 top variable genes and the number of significant principal components was determined using an
292 elbow plot. Clustering of the dataset was performed using the function "FindClusters" and cell types
293 for each cluster were determined using canonical marker genes. Refined clustering was achieved by
294 reclustering within the identified cell types using the above pipeline on subsets of the data. The tSNE
295 plots were produced using the generated principal components as input.

296 **Determination of LOY in single cells**

297 The LOY status for each sequenced cell was determined under the assumption that cells with LOY
298 would not express genes located in the male specific part of chromosome Y (MSY). Hence, non-LOY
299 blood cells are normally expressing a series of genes located in the MSY (e.g EIF1AY, RPS4Y1, KDM5D
300 and ZFY). We took advantage of this information and thus scored LOY in cells without transcripts

301 from the MSY using ensembl (v93) to identify all MSY gene measured in the data. Cells with high
302 content of mitochondrial RNA (more than 5%) as well as cells with less than 800 UMIs or less than
303 350 genes expressed was removed from the analysis.

304 **Single-cell statistical analyses**

305 To compare differences in autosomal gene expression between LOY cells and non-LOY cells we first
306 performed WilcoxDETest's, implemented in the R library Seurat (v2.3.1). We also developed an in-
307 house random sampling algorithm to compare the gene expression in LOY cells with non-LOY cells
308 within specific cell types. First, we established the observed gene expression in LOY cells in the cell
309 type under investigation, by calculating the mean normalized expression values in all subjects, within
310 all LOY cells of the tested cell type. Next, we randomly selected from all subjects, a number of cells
311 from the non-LOY cells of the examined cell type, and calculated the mean normalized expression in
312 the resampled cells. To avoid biases caused by inter-individual variation, we programmed the
313 sampling algorithm to sample an equal number of non-LOY cells as observed LOY cells within each
314 subject. For example, from subjects with 100 LOY cells of a specific cell type, the same number of
315 non-LOY cells from the same cell type was sampled from the set of non-LOY cells. The resampling of
316 non-LOY cells from all subjects was repeated 50.000 times and for each iteration, the mean
317 normalized expression of the investigated gene in the resampled cells was calculated. The resampled
318 data represents a weighted expression level of the examined gene in non-LOY cells within specific
319 cell types and thus, the resampled distribution represents the normalized expression of the
320 investigated gene in non-LOY cells. The range of variation of gene expression in LOY cells was
321 estimated in a similar fashion, by resampling of a subset of the LOY cells within each subject. Exact p-
322 values were calculated by comparing the observed mean expression in LOY cells to the resampled
323 distribution of non-LOY cells. All statistical analyses were performed using R v. 3.4.4.