

Deciphering African tropical forest dynamics in the Anthropocene: how social and historical sciences can elucidate forest research and management

G. Walters^{a,b}, J. Angus Fraser^c, N. Picard^d, O. Hymas^{b,e}, J. Fairhead^f

^aUniversity of Lausanne, Faculty of Geosciences and Environment, Institute of Geography and Sustainability, Lausanne, Switzerland, corresponding author, gretchen.walters@unil.ch

^bDepartment of Anthropology, University College London, London, United Kingdom

^cLancaster Environment Center, Lancaster University, Lancaster, United Kingdom, james.angus.fraser@gmail.com

^dFood and Agriculture Organization of the United Nations, Rome, Italy, Nicolas.picard@fao.org

^eZoological Society of London, Outer Circle, Regent's Park, London, United Kingdom, olivier@hymas.eu

^fUniversity of Sussex, School of Global Studies, Department of Anthropology, Falmer, United Kingdom j.r.fairhead@sussex.ac.uk

Corresponding author at: ^aUniversity of Lausanne, Faculty of Geosciences and Environment, Institute of Geography and Sustainability, Lausanne, Switzerland, corresponding author, gretchen.walters@unil.ch

Abstract

Forests bear the historical legacies of human activities over thousands of years, including agriculture, trade, disease and resource extraction. Many of these activities may represent indices of the proposed geological epoch of the Anthropocene. Modifications to soil, topography and vegetation evidence anthropogenic influences. Yet studies of vegetation change throughout the humid tropics tend to occlude these by focussing on forest dynamics, timber, and biodiversity through permanent sample plots or forestry inventory plots. We highlight how history and social science can be combined with ecology to help better understand human signatures in forest dynamics. We (1) critically review ecological methods in the light of the environmental and social history of the Afrotropics; (2) map current plot networks for West and Central Africa in relation to the Human Footprint Index; (3) using two case-studies, demonstrate how history and social science bring new insights and inferences to plot-based studies; all leading to (4) novel forms of interdisciplinary collaboration for sustainable forest conservation, management and restoration.

Keywords

Forest ecology, Africa, historical sciences, social sciences, forest management, interdisciplinarity

This is an authors' pre-print version of this paper. The final version of record is published by in the journal *Anthropocene* and is available here:

<https://www.sciencedirect.com/science/article/pii/S2213305419300256>

or via the DOI permalink: <https://doi.org/10.1016/j.ancene.2019.100214>

The citation of the paper is:

Walters, G., J.A. Fraser, N. Picard, O. Hymas, & J. Fairhead. 2019. Deciphering African tropical forest dynamics in the Anthropocene: How social and historical sciences can elucidate forest cover change and inform forest management. *Anthropocene* 27 September 2019, 100214.

1. Progress in linking past anthropogenic activity to present-day forest structure

The ways that tropical forests are studied and conceptualized have changed greatly in recent decades. The work of geographers and anthropologists in the Amazon (e.g. Balée and Erickson 2006) and in Africa (e.g. Fairhead and Leach 1996) have helped bring about these changes, challenging researchers to consider the influences of humans on forest structure and biodiversity. In the early 2000s, ecologists believed that people (Foster *et al.* 2003) had not affected current ecosystem dynamics to a large extent; however, it is now widely accepted that past human impacts have not only shaped ecosystems, but also have become a geological force – captured in the concept of the “Anthropocene” (e.g., Ellis *et al.* 2013). While some argue for a 1950s start to the proposed geologic epoch (e.g. Zalasiewicz *et al.* 2010), others (Lewis and Maslin 2015) argue for an earlier ca 1500 start, which coincides with the beginning of global capitalism (Moore 2015). In Africa, these ‘Anthropocene’ impacts include the slave trade, colonisation, Atlantic trade, disease epidemics and lifestyles of prehistoric societies (Maddox 2006; Kay and Kaplan 2015), often leaving their mark on forest structure and biodiversity hundreds of years later.

Ecologist P.W. Richards (1952) warned that ecologists working in the African rain forest should not ignore significant modification of their study areas by recent human activity. Foresters were aware of the relationship between cultivation and pioneer forest species (Letouzey 1957); others noted the importance of understanding forest change in relation to the life span of trees (White and Oates 1999). The forests of Oban, Nigeria, for example, now construed as ‘Old Growth’, were previously inhabited (Rosevear 1979: 78) based on the evidence of trees left by farmers when the land was depopulated hundreds of years earlier. This example suggests that the lifespan of a tree, and what occurred during that period, could be important for understanding forest dynamics (Bourland *et al.* 2015). In some cases, current forest cover, with species of economic importance to the timber trade, have their origins in past African societies (Aubréville 1948).

Throughout the humid tropics, ecologists have studied forest ecosystems through permanent sample plots (PSPs) (Lopez-Gonzalez *et al.* 2011), (often networked), using common research questions, methodology or databases (Anderson-Teixeira *et al.* 2015). A few decades ago, PSP research principally comprised community ecology, species diversity, and management (Condit 1995). Increasingly, researchers use these plots to understand changes in carbon and forest response to climate change (Talbot *et al.* 2014), indicating alternative uses of these datasets. However, these are not the only tropical forest plot networks: extensive networks of forest inventory plots (FIPs) also exist. In coastal central Africa, they cover more than 11 million ha (de Wasseige *et al.* 2009). While foresters assess timber stock with FIPs, ecologists also use PSPs to study forest ecology and biodiversity. These plots potentially can answer questions beyond their ecological or forestry remit, and so can address the historical and political contexts in which the forests have grown (Robbins 2012). Both types of plots can facilitate Anthropocene studies to understand and quantify human influence on the environment in the recent past and to inform forest management.

Forest ecologists increasingly collaborate with paleo-biologists (Lovejoy and Heinz 2007) to explore the legacy of anthropogenic activities on forests from past millennia (Willis *et al.* 2004; Hayashida 2005). Collaborative research with archaeologists (Iles 2016) suggest methodological flaws can occur when plots are not interpreted in consideration of the legacies of human history. Furthermore, PSPs were largely established in forests considered ‘intact’, ‘pristine’, and ‘old growth’, or in accessible locations (e.g. research stations) (Pitman *et al.* 2011). This first bias led

ecologists to examine such plots as if they were ‘undisturbed’, neglecting anthropogenic legacies. The second bias led them to generalise from plot-based results to the wider landscape (Hecht and Saatchi 2007).

Given the transformation of the African environment in the last 500 years, such as by the Atlantic trade, new collaborations with the social-historical sciences can elucidate how human activities transformed the forest. Land-use research, for example, although originally dominated by remote sensing, increasingly benefits from historical data, historical methods (e.g. Hunter and Sluyter 2015) and ecological collaboration (Watson *et al.* 2014) to reconstruct land-use, forest change and carbon cycling. Such methods also apply to research on forest dynamics, forest management and plots, but require new collaborations with social scientists and historians to discern legacies that archaeology is less able to detect, such as political drivers shaping landscapes. Although some research drawing on plots, paleobiological methods and aerial photography does discern the legacy of recent disturbances in the present-day forest structure (e.g. Delègue *et al.* 2001; van Gemerden *et al.* 2003), this often remains unexplained.

This article highlights how elucidating forest legacies across plot networks requires new interdisciplinary collaborations amongst ecologists, foresters, anthropologists and historians and related disciplines, while discerning lessons for sustainable management and forest recovery. The paper shows how different disciplines can coordinate methods and reflections to address common questions of forest legacies in the ‘Anthropocene’. Below, we (1) assess the coverage and methods of PSPs and FIPs in tropical Africa in light of sub-Saharan, tropical African forest history; (2) using two case studies, demonstrate how history and anthropology can enrich plot-based studies and inform sustainable forest management; and (2) suggest how future research can facilitate this type of interdisciplinary collaboration.

2. Current ecological and forest inventory plot methods

Ecological studies of forest dynamics in tropical Africa mostly rely on PSPs, by monitoring individual trees over time (Picard *et al.* 2010). The size of PSPs varies from 100 m² to 500 ha (one ha is the most common size). These plots have some methodological and interpretation limitations (Sheil 1995). They were established as part of disconnected studies, so their spatial distribution is uneven and does not follow a sampling design that enables statistically significant inferences to be drawn regionally (Picard *et al.* 2010). The balance between known ‘disturbed’ and presumed ‘intact’ forests remains uncontrolled, and given that the ecological questions that drove their establishment led to more plots being established in presumed ‘intact’ locations, there is a ‘majestic forest’ bias in the data set (Phillips *et al.* 2002). Figure 1 details the current distribution of PSPs in western and central Africa rainforests overlaid with the Human Footprint (HFP) index. The HFP measures the cumulative impact of direct pressures on nature from human activities based on eight spatial variables: extent of built environments, cropland, pastureland, human population density, night-time lights, railways, roads, and navigable waterways.¹

¹ This index was developed by WCS, CIESIN and Columbia University, with a publicly-accessible data set (<http://dx.doi.org/10.7927/H4M61H5F>).




Fig. 1. Locations of Permanent Sample Plots (PSP) with the Human Footprint (HFP) index in the background. Dots = PSPs. The HFP in the background is based on the NASA index. White indicates areas with tree cover < 50%. X-axis of the inset plot = HFP while y-axis = density of distribution.

3. African forest history

African forest history comprises a complex interplay of climatic drivers and land-use changes at different timescales (McIntosh *et al.* 2015). Global models of historical land-use suggest that significant parts of Central and West Africa had increasingly reduced natural forest cover from 1,000 AD with an associated carbon loss (Kaplan *et al.* 2011). The ‘first significant use’ of landscapes in West and Central Africa drastically increased from the start of the first millennium (Ellis *et al.* 2013), but with significant differences in forest loss and gain from the 1900s to present (Aleman *et al.* 2018).

Trade between Europe and the Afrotropics historically influenced forest dynamics. Beginning in the fifteenth century, it extended along the Atlantic coast and inland, encountering trade networks and creating social upheavals in West and Central Africa, and restructuring polities and trade routes (e.g. Coquery-Vidrovitch 1985). The introduction of new crops (e.g. manioc) drove economic transformations, while slavery, warfare, and disease epidemics depopulated entire areas, which Ford (1971: 489) understood as “biological warfare on a vast scale”

In Atlantic Central Africa, peak human population density potentially occurred in the 16th century, after which the population decreased until the 19th century (Oslisly *et al.* 2013). During this period, the Atlantic trade alone potentially resulted in a loss of 11 million people from Africa (Maddox 2006). Furthermore, the worldwide Spanish influenza epidemic likely led to the death of at least 1.5 million people in Africa, in 1918-1919 (Spinage 2012: 1201–2). This uneven demographic impact left land depopulated in humid West Africa (Fairhead and Leach 1998) and Central Africa, the latter of which was accentuated by forced, colonial resettlement along roads in the 20th century (Gray 2002, Fig. 2). Evidence from a variety of sources indicates that most of the areas studied by PSPs

and FIPs in West and Central Africa were shaped by these complex factors, as described by two cases below.

4. Evidence from Western and Central Africa

4.1 Liberia

West Africa's Upper Guinea Forest region is a 'hotspot' of global biodiversity (Poorter *et al.* 2004), threatened by logging, rubber and industrial agriculture (Fairhead and Leach 1998). Liberia holds the greatest area of this forest, with 41,238 km² or 37.7% of historic forest cover remaining (Poorter *et al.* 2004:6). However, historical observations and recent research reveal that much of this forest is 'anthropogenic' or 'domesticated' due to settlement and agro-forestry dynamics which have shaped the current forest species composition (Fairhead and Leach 1998). This history is occluded by ecological studies that assume forests are 'pristine' without evaluating their history even in areas of known anthropogenic influence. Bongers *et al.* (1999), for example, represented plot species composition as primarily an effect of climatic variation, even when these species distributions are often influenced by anthropogenic processes. Cotton [*Ceiba pentandra*], Kola [*Cola nitida*], *Terminalia ivorensis*, *Terminalia superba*) are all species propagated and managed by people in this region (Fairhead and Leach 1996; Bongers *et al.* 1999).

Much of Liberia's forests are shaped by past, long-term swidden-fallow dynamics, viz., they are embedded with overgrown, old settlements, considered 'sacred' by local peoples due to the presence of certain tree species (e.g. Cotton, Kola), ancestors and spirits, and therefore often afforded protection from felling. These 'old town spots' also feature fertile 'African Dark Earths' and are frequently cultivated as cacao agro-forests (Fraser *et al.* 2016) meriting the term 'sacred agroforest'.

A recent survey of 83 localities in Gbarpolu, Bong, Lofa, Nimba counties near forestry concessions, where plots are likely used for management, found at least one sacred agroforest at 51 locations, 94 in total (Appendix 1, Fraser *et al.* 2016). At the local scale, mapping and transects within a 3km radius (ca. 2,827 ha) of a settlement found 18.6 ha of sacred agroforests, with the majority of adjacent areas covered in variously aged fallow vegetation (Diabate *pers. comm.*). In comparing transects in sacred agroforests and secondary forest, sacred agroforest increases biodiversity at the landscape level due to differing species composition in both canopy species and seedlings. This study failed to find old growth forest in this area (see Fraser *et al.* 2016: "Baema", Figure 1, Appendix 2). This pattern appears to be typical of NW Liberia with major historical disturbance also occurred in neighbouring settlements. The diaries of two African American explorers, George L. Seymour and Benjamin J. K. Anderson, who traversed the area in 1858, reported, '*it is common to see a hundred-acre farm in one cutting*' (Fairhead *et al.* 2003). And, '*Standing upon an elevation, it seemed to me that the people had attempted to cover the whole country with their rice fields...Only here and there could be seen patches of large forest trees.*' (Fairhead *et al.* 2003:190-191).

Early foresters recognised the anthropogenic landscapes that emerged from such processes over time. In the 1940's, forester Karl Meyer walked 2,300 km through Liberia's forests observing that '*abandoned villages are, in some sections, very common,*' and areas '*with no signs of occupancy during recent centuries are few and scattered*' (Mayer 1951:25). Hence, today's forest is composed of secondary forest, historically disturbed through shifting cultivation, with largely unmanaged

succession (Fairhead and Leach 1998), dotted with sacred agroforests (Fraser et al. 2016, Appendix 1).

In 2006, Liberia revised its National Forestry Law, promoting sustainable forest management. In 2009, the Community Rights Law was enacted, empowering communities to engage in sustainable forest management on their lands. That year, industrial logging companies were granted 25% of Liberia's forests, yet the implementation of the forestry laws remains difficult (Altman et al. 2012; O'Mahoney 2019).

We make the following management recommendations: First, the government largely does not recognise the history of Liberian forests. This encourages viewing forests as a resource stock rather than a cultural artefact. It is important that the linkages between extant forest peoples, such as the Loma and their ancestors who created these forests (Fraser et al 2015), are recognised in representations of forests by government, media, and in school curricula. Second, if the two above mentioned laws are not implemented, people will lose tenurial rights over the agrobiodiversity created by their ancestors. An awareness campaign should ensure that people know their tenure rights. Together these interventions - raising historical awareness and legal awareness - could empower citizens to conserve the landscapes that are also their cultural heritage.

4.2 Gabon

In Gabon, archaeology, palynology, diatoms, phytoliths, and tree population genetics have partially reconstructed forest history (Brncic *et al.* 2007; Piñeiro et al. 2017). However, history and linguistics also explain the interactions of people and forests over time (Vansina 1990), including societal change impacting vegetation structure (Walters 2012). A dominant, sub-endemic timber species to Gabon, Okoume (*Aucoumea klaineana*), colonizes slash-and-burn openings, which forester Aubréville described as "Okoume being the son of manioc" (Aubréville 1948). Once mature, large stands indicate the presence of past villages (Biraud 1959). However, disease, brought by trade and colonial rule severely affected village placement and human demography (Headrick 1990, Chamberlin 1977, Gray 2002), resulting in changes to forest composition.

Oral histories from the Parc National de Waka (PNW) area near PSPs (Balinga *et al.* 2006), (Hymas 2016) found that current stands of Okoume originated from colonial concession plantation agriculture during the late-1800s to mid-1900s. These forests are the outcome of complex historical trajectories. Inter-ethnic conflict over natural resources depopulated the area in the early 1800s; it was repopulated in the late 1800s through trade concessions, and then again depopulated in the 1920s by trade-induced disease, and further depopulated in 1960 by the government resettlement policy (Fig. 2). In the 1960s, logging companies were attracted to the Okoume stands, which resulted in a repopulation by workers. In 2003, the PNW was created to protect elephant populations (WCS, 2007), attracted to abandoned village sites, rich in planted fruit trees (Barnes *et al.* 1991).

In another case, using the FIPs of a forestry concession in the Haut-Abanga, Engone Obiang et al. (2014) diagnosed 'old, naturally-declining' populations of Okoume through a tree diameter analysis. The modal diameter of Okoume was 50-60 cm, corresponding to 60-70-year-old trees. The historical literature showed that the area was a former communication corridor that lost its role with the establishment of the modern road network. The area, populated until the 1940s, became almost empty in the 1950s due to the resettlement policy (Peyrot 2008), Fig. 2.




Figure 2. Population distribution of Gabon in 1940 prior to the resettlement policy and in 1970, afterwards (adapted from Sautter 1966), showing the Haut Abanga Concession in the north, and PNW in the center.

Okoume is Gabon’s most important timber species but other commercial species are fast-growing and light-demanding, too. For such species, current stocks are often the legacy of past disturbances, such as slash-and-burn cultivation. Hence, forest history plays a role in determining their distribution. Nonetheless, current sustainable forest management relies on the hypothesis of demography at equilibrium, where young trees continuously replace large ones. Sustainability is assessed through the recovery rate of the species at the end of the felling cycle. When the recovery rate becomes too low, forest managers often increase the minimum size of felled trees. Although this measure may have short-term success in maintaining species stock, it does not acknowledge the role of history in shaping it. As a result, these stocks may decline, with both environmental and economic consequences, bringing the concept of ‘sustainable forest management’ into question (Karsenty 2018a).

To address unsustainable forest management, in 2018, Gabon announced that all forest concessions must become certified by the Forest Stewardship Council (FSC) standard by 2022. The standard goes beyond the issue of sustainable stocks (Karsenty 2018b), but its implementation has often been problematic (Nepomuceno et al 2019). It is unlikely that “sustainable stocks” can be achieved without considering their historical origin. Forest management plans must factor in historical dimensions and when natural recovery is no longer possible, silvicultural techniques, including planting, should be proposed.

5. Interdisciplinary collaboration to understand tropical African tropical forests in the Anthropocene

Today's West and Central African forests provide an archive of the slave trade, conflicts, diseases and depopulations that left farm and village lands abandoned. Attempts to read forest composition and dynamics only through the lens of 'natural history' and 'climate change' overlooks this history and the associated meaning for those who live there, and reduces the ability to manage these legacy forests.

PSPs and FIPs in the Afrotropics provide a research lens into both 'forest ecology' and past environmental change. This requires asking a wider range of questions of these plots, collaboratively researched using ecologically and socio-historically focused inquiry. To ask such questions, new forms of research are required; however, combining different disciplines that use diverse lexicons, timeframes, epistemologies and methodologies is one of the biggest challenges of achieving interdisciplinary research (Lele and Kurian 2011). Interdisciplinary research programs must be developed with an integrated methodology combining data from archaeology and history (artefacts and texts), anthropology and political and historical ecology, with ecology to understand forest patterns. These initiatives require researchers to "share the conceptual world of their colleagues", beyond disciplinary boundaries (Darbellay 2015:167).

In achieving this, it is imperative to recognise that people shape forests. Thinking in terms of social-ecological and biocultural systems (c.f. Fischer 2018), can demonstrate society-environment interconnections, leading to discussions on how to study these impacts collaboratively. Robbins (2012) proposes that links between socio-political forces and ecological characteristics include type and direction of environmental change, drivers of change, the environment in which changes occur, the impact of cultural practices on the system (*sensu* Maraccuci 2000), and how it recovers. The final step can, as noted in the Liberia case, explicitly link historical forests to current forest usage and land claims.

In the case of PSPs, synthesising existing historical and archaeological research for each plot and linking to specific periods, can show how present-day forest structure and diversity are linked to disturbances (Fairhead, unpublished data). When studying impacts through FIPs, documenting the species most susceptible to disturbance, as in the Gabon case, is a first step. The places where these signals are strongest (e.g. mono-dominant forests), can then be explored. By providing a view on how ecological patterns and processes have reacted in the past to environmental changes, historical ecology can inform how forest structure and process may respond to future change (Safford *et al.* 2012).

Such collaborations will also inform forest management still rooted in equilibria paradigms despite historical evidence (Morin-Rivat *et al.* 2017). Some forest managers are aware that the management of some species is not sustainable because the stock is a legacy of past human perturbations (Morin-Rivat *et al.* 2016). However, a historical perspective on the current structure of commercial species would offer a stronger basis for sustainable management. In both case-studies, forest history and species responses are different, and each country has different sustainable forest management strategies. However, neither country's policies acknowledge the impact that history has on timber stocks, nor address if and how to maintain and manage these

stocks. Furthermore, neither recognises forests as historical and cultural products of people living there today.

In this paper, we reviewed limitations of plot-based research in light of Afrotropical history, which heavily influenced the history of these forests, compared the usage of PSPs and FIPs, and provided cases demonstrating how interdisciplinary research collaborations can enrich plot-based studies. We propose future work focus on using plot networks to research new questions, in collaboration with historians and social scientists. The lack of historical perspectives on forests will limit addressing sustainability (Roberts *et al.* 2018, this issue). However, new collaborations will not only help deepen conclusions from forest ecology, but also influence study design and management options.

Interdisciplinary collaboration is one way to explore how the forests have been shaped during the Anthropocene. This new view suggests that some tree species require new forms of management and some forests deserve new recognition as cultural landscapes.

Acknowledgments

The authors thank Jos Barlow, Geoffroy de Saulieu, Melis Ece, Lindsey Gillson, Anne Chin, and anonymous reviewers, for providing constructive comments on this manuscript and to Simon Lewis for earlier discussions. This paper was presented and discussed at the University of Zurich 2016 workshop “The trouble with defining forests: semantics, ontology, territoriality”, organised by Muriel Côte, Ross Purves and Flurina Wartmann. This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Literature Cited

- Aleman, JC, MA Jarzyna, AC Staver. 2018. Forest Extent and Deforestation in Tropical Africa since 1900. *Nature Ecology & Evolution* **2**: 26-33.
- Altman, SL, SS Nichols, JT Woods. 2012. Leveraging high-value natural resources to restore the rule of law: The role of the Liberia Forest Initiative in Liberia’s transition to stability *in* High-Value Natural Resources and Peacebuilding, ed. P. Lujala and S. A. Rustad. London: Earthscan.
- Anderson-Teixeira KJ, Davies SJ, Bennett AC, *et al.* 2015. CTFS-ForestGEO: a worldwide network monitoring forests in an era of global change. *Glob Change Biol* **21**: 528–49.
- Aubréville A. 1948. La forêt d’okoumé, richesse permanente du Gabon. Vers l’évolution dirigée de la forêt gabonaise. *Bull Sci Dir Agric L’Élevage For Ministère Fr O-m* **2**: 64–80.
- Balée W and Erickson CL. 2006. Time, complexity, and historical ecology. In: Balée W, Erickson CL (Eds). Time and complexity in historical ecology. New York: Columbia University Press.
- Balinga MP, Sunderland TCH, Walters G, *et al.* 2006. A vegetation assessment of the Waka National Park, Gabon. Washington D.C.: The Smithsonian Institution.
- Barnes RFW, Barnes KL, Alers MPT, and Blom A. 1991. Man determines the distribution of elephants in the rain forests of northeastern Gabon. *Afr J Ecol* **29**: 54–63.
- Biraud J. 1959. Reconstitution naturelle et amélioration des peuplements d’okoumé au Gabon. *Bois For Trop* **66**: 3–28.
- Blake S, Deem SL, Mossimbo E, *et al.* 2009. Forest Elephants: Tree Planters of the Congo: Forest Elephants, Seed Dispersal, and Trees. *Biotropica* **41**: 459–68.
- Bongers F, Poorter L, Van Rompaey, R.S.A.R., and Parren MPE. 1999. Distribution of Twelve Moist Forest Canopy Tree Species in Liberia and Côte d’Ivoire: Response Curves to a Climatic Gradient. *J Veg Sci* **10**: 371–82.

- Bourland N, Cerisier F, Daïnou K, *et al.* 2015. How Tightly Linked Are *Pericopsis elata* (Fabaceae) Patches to Anthropogenic Disturbances in Southeastern Cameroon? *Forests* **6**: 293–310.
- Brcic TM, Willis KJ, Harris DJ, and Washington R. 2007. Culture or climate? The relative influences of past processes on the composition of the lowland Congo rainforest. *Philosophical Trans R Soc Lond Ser B Biol Sci*: 1–14.
- Chamberlin C. 1977. Competition and Conflict: The Development of the Bulk Export Trade in Central Gabon during the Nineteenth Century. PhD thesis: University of California, Los Angeles, USA.
- Condit R. 1995. Research in large, long-term tropical forest plots. *Trends Ecol Evol* **10**: 18–22.
- Coquery-Vidrovitch C. 1985. Afrique noire: permanences et ruptures. Paris: Payot.
- Darbellay F. 2015. « Rethinking Inter- and Transdisciplinarity: undisciplined knowledge and the emergence of a new thought style ». *Futures* **65**: 163–74.
- Delègue M-A, Fuhr M, Schwartz D, Mariotti A, Nasi R. 2001. Recent origin of a large part of the forest cover in the Gabon coastal area based on stable carbon isotope data. *Oecologia* **129**: 106–13.
- Ellis EC, Kaplan JO, Fuller DQ, Vavrus S, Klein Goldewijk K, Verburg PH. 2013. Used planet: A global history. *Proc Natl Acad Sci* **110**: 7978–85.
- Engone Obiang NL, Ngomanda A, Hymas O, Chezeaux E, Picard N. 2014. Diagnosing the demographic balance of two light-demanding tree species populations in central Africa from their diameter distribution. *For Ecol Manag* **313**: 55–62.
- Fairhead J, Geysbeek T, Holsoe SE, and Leach M. 2003. African-American Exploration in West Africa: Four Nineteenth-Century Diaries. Bloomington: Indiana University Press.
- Fairhead J and Leach M. 1996. Misreading the African Landscape: Society and Ecology in a Forest Savanna Mosaic. Cambridge: Cambridge University Press.
- Fairhead J and Leach M. 1998. Reframing Deforestation: Global Analyses and Local Realities: Studies in West Africa. London: Routledge.
- Fischer, AP, 2018. Forest landscapes as social-ecological systems and implications for management. *Landscape and Urban Planning* **177**, 138–147
- Ford J. 1971. The role of the trypanosomiasis in African ecology; a study of the tsetse fly problem. Oxford: Clarendon Press.
- Foster D, Swanson F, Aber J, Burke I, Brokaw N, Tilman D, Knapp A. 2003. The Importance of Land-Use Legacies to Ecology and Conservation. *Bioscience* **53**: 77–88.
- Fraser JA, Diabaté M, Narmah W, Beavogui P, Guilavogui K, de Foresta H, Junqueira AB. 2016. Cultural valuation and biodiversity conservation in the Upper Guinea forest, West Africa. *Ecol Soc* **21**: 36.
- Fraser, JA, Frausin V, Jarvis A, 2015. An intergenerational transmission of sustainability? Ancestral habitus and food production in a traditional agro-ecosystem of the Upper Guinea Forest, West Africa. *Global Environmental Change* **31**, 226–238.
- Gemerden BS van, Oloff H, Parren MPE, and Bongers F. 2003. The pristine rain forest? Remnants of historical human impacts on current tree species composition and diversity. *J Biogeogr* **30**: 1381–90.
- Gourlet-Fleury S, Rossi V, Rejou-Mechain M, *et al.* 2011. Environmental filtering of dense-wooded species controls above-ground biomass stored in African moist forests: Environmental filtering and tree biomass. *J Ecol* **99**: 981–90.
- Gray CJ. 2002. Colonial rule and crisis in Equatorial Africa: southern Gabon ca. 1850-1940. Rochester, USA: University of Rochester Press.
- Hayashida FM. 2005. Archaeology, ecological history, and conservation. *Annu Rev Anthropol* **34**: 43–65.
- Headrick R. 1990. Studying the population of French Equatorial Africa. In: Fetter BD (Ed). *Demography from scanty evidence: Central Africa in the colonial era*. Boulder, USA: Lynne Reiner.
- Hecht SB and Saatchi SS. 2007. Globalization and Forest Resurgence: Changes in Forest Cover in El Salvador. *Bioscience* **57**: 663–72.
- Hunter R and Sluyter A. 2015. Sixteenth-century soil carbon sequestration rates based on Mexican land-grant documents. *The Holocene* **25**: 880–5.
- Hymas O. 2016. L'Okoumé, fils du manioc: Post-logging in remote rural forest areas of Gabon and its long-term impacts on development and the environment. PhD Thesis: University College London, London.

- Iles L. 2016. The Role of Metallurgy in Transforming Global Forests. *J Archaeol Method Theory* **23**: 1219–41.
- Kaplan JO, KM Krumhardt, EC Ellis, WF Ruddiman, C Lemmen, KK Goldewijk et al. 2011. Holocene carbon emissions as a result of anthropogenic land cover change. *The Holocene* **21**(5): 775–791
- Karsenty A. 2018a. Is sustainable logging possible in Africa’s dense forest? *Bois et Forêts des Tropiques* **336**: 3–5.
- Karsenty A. 2018b. Forêts : l’institutionnalisation légale de la certification FSC au Gabon. WillAgri [Blog] 4 October 2018. <http://www.willagri.com/2018/10/04/forets-linstitutionnalisation-legale-de-la-certification-fsc-au-gabon/> Accessed 24 June 2019.
- Kay, AU, and JO Kaplan. 2015. Human Subsistence and Land Use in Sub-Saharan Africa, 1000BC to AD1500: A Review, Quantification, and Classification. *Anthropocene* **9**: 14–32.
- Lele, S, Kurien A, 2011. Interdisciplinary analysis of the environment: insights from tropical forest research. *Environmental Conservation* **38**, 211–233
- Letouzey R. 1957. La forêt à *Lophira alata* de la zone littorale camerounaise. *Bois For Trop* **53**: 9–20.
- Lewis SL and Maslin MA. 2015. Defining the Anthropocene. *Nature* **519**: 171–80.
- Lewis SL, Sonké B, Sunderland T, et al. 2013. Above-ground biomass and structure of 260 African tropical forests. *Philos Trans R Soc B Biol Sci* **368**: 20120295–20120295.
- Lopez-Gonzalez G, Lewis SL, Burkitt M, Phillips OL. 2011. ForestPlots.net: a web application and research tool to manage and analyse tropical forest plot data: ForestPlots.net. *J Veg Sci* **22**: 610–3.
- Lovejoy TE and Heinz HJ. 2007. Paleoeecology and the path ahead. *Front Ecol Environ* **5**: 456.
- Maddox G. 2006. Sub-Saharan Africa: an environmental history. Santa Barbara, California: ABC-CLIO.
- Maniatis D, Malhi Y, Saint André L, Mollicone D, Barbier N, Saatchi S, Henry M, Tellier L, Schwartzenberg M, White L. 2011. Evaluating the Potential of Commercial Forest Inventory Data to Report on Forest Carbon Stock and Forest Carbon Stock Changes for REDD+ under the UNFCCC. *Int J For Res* **2011**: 1–13.
- Marcucci, DJ, 2000. Landscape history as a planning tool. *Landscape and Urban Planning* **49**, 67–81
- Mayer KB. 1951. Forest Resources of Liberia. Forest Service United State Department of Agriculture.
- McIntosh RJ, J Maley, R Vernet. 2015. African Palaeoclimate and Human Response: A Special Issue of the African Archaeological Review. *African Archaeological Review*, **32**: 167–177.
- Moore JW. 2015. Capitalism in the web of life: ecology and the accumulation of capital. New York: Verso.
- Morin-Rivat J, Biwole A, Gorel A-P, Vleminckx J, Gillet J-F, Bourland N, Hardy OJ, Smith AL, Dai Nou K, Dedry L, Doucet J-L. 2016. High spatial resolution of late-Holocene human activities in the moist forests of central Africa using soil charcoal and charred botanical remains. *The Holocene* **26**: 1954–67.
- Morin-Rivat J, Fayolle A, Favier C, Bremond L, Gourlet-Fleury S, Bayol N, Lejeune P, Beeckman H, Doucet J-L. 2017. Present-day central African forest is a legacy of the 19th century human history. *eLife* **6**.
- Nepomuceno, Í, H Affonso, JA Fraser, M Torres. 2019. Counter-Conducts and the Green Grab: Forest Peoples’ Resistance to Industrial Resource Extraction in the Saracá-Taquera National Forest, Brazilian Amazonia. *Global Environmental Change* **56**: 124–33.
- O’Mahony, J. 2019. Liberia’s new land rights law hailed as victory, but critics say it’s not enough. *Mongabay*, 22 March. <https://news.mongabay.com/2019/03/liberias-new-land-rights-law-hailed-as-victory-but-critics-say-its-not-enough/>. Accessed 24 June 2019.
- Oslisly R, White L, Bentaleb I, Favier C, Fontugne M, Gillet J-F, Sebag D.. 2013. Climatic and cultural changes in the west Congo Basin forests over the past 5000 years. *Philos Trans R Soc B Biol Sci* **368**: 20120304–20120304.
- Peyrot B. 2008. Incidences écologiques, anthropiques et paléocologiques sur l’évolution des forêts du Gabon. *Cah O-m* **61**: 111–44.
- Phillips OL, Malhi Y, Vinceti B, et al. 2002. Changes in growth of tropical forests: evaluating potential biases. *Ecol Appl* **12**: 576–87.
- Picard N, Magnussen S, Banak LN, Namkossereena S, Yalibanda Y. 2010. Permanent sample plots for natural tropical forests: A rationale with special emphasis on Central Africa. *Environ Monit Assess* **164**: 279–95.

- Piñeiro, R, G Dauby, E Kaymak, OJ Hardy. 2017. Pleistocene Population Expansions of Shade-Tolerant Trees Indicate Fragmentation of the African Rainforest during the Ice Ages. *Proceedings of the Royal Society B: Biological Sciences* **284** : 20171800.
- Pitman NCA, Widmer J, Jenkins CN, Stocks G, Seales L, Paniagua F, Bruna EM. 2011. Volume and Geographical Distribution of Ecological Research in the Andes and the Amazon, 1995-2008. *Trop Conserv Sci* **4**: 64–81.
- Poorter L, Bongers F, Kouamé FN, Hawthorne WD. 2004. West African Forests: an ecological atlas of woody plant species. Cambridge, Massachusetts, USA: CAB International.
- Réjou-Méchain M, Fayolle A, Nasi R, Gourlet-Fleury S, Doucet J-L, Gally M, Hubert D, Pasquier A, Billand A. 2011. Detecting large-scale diversity patterns in tropical trees: Can we trust commercial forest inventories? *For Ecol Manag* **261**: 187–94.
- Richards PW. 1952. The tropical rain forest: an ecological study. London: Cambridge University Press.
- Robbins, P, 2012. Political ecology: a critical introduction, 2nd ed, Critical introductions to geography. J. Wiley & Sons, Chichester, West Sussex ; Malden, MA
- Roberts P, Boivin N, and Kaplan JO. 2018. Finding the Anthropocene in tropical forests. *Anthropocene* **23**: 5–16. This issue.
- Rosevear, DR. 1979. Oban Revisited. *Niger Field* **44**: 75–81.
- Safford HD, Hayward GD, Heller NE, Wiens JA . 2012. Historical ecology, climate change, and resource management: can the past still inform the future? In: Wiens JA, Hayward GD, Safford HD, Giffen CM (Eds). Historical environmental variation in conservation and natural resource management. Wiley & Sons.
- Sautter G. 1966. De l'Atlantique au fleuve Congo, une géographie du sous-peuplement. République du Congo, République du Gabon. Paris: Editions du Centre National de la Recherche Scientifique.
- Sheil D. 1995. A critique of permanent plot methods and analysis with examples from Budongo Forest, Uganda. *For Ecol Manag* **77**: 11–34.
- Spinage, CA. 2012. African Ecology- benchmarks and historical perspectives. Berlin: Springer-Verlag.
- Talbot J, Lewis SL, Lopez-Gonzalez G, *et al.* 2014. Methods to estimate aboveground wood productivity from long-term forest inventory plots. *For Ecol Manag* **320**: 30–8.
- Vansina J. 1990. Paths in the Rainforest: toward a history of political tradition in Equatorial Africa. London: James Currey.
- Walters G. 2012. Changing customary fire regimes and vegetation structure in Gabon's Batéké Plateaux. *Hum Ecol* **40**: 943–55.
- Wasseige C de, Devers D, Marcken P de, Ebaà Atyi R, Nasi R, Mayaux P. (Eds). 2009. Les forêts du bassin du Congo : état des forêts 2008. Office des publications de l'Union Européene.
- Watson SJ, Luck GW, Spooner PG, Watson DM. 2014. Land-use change: incorporating the frequency, sequence, time span, and magnitude of changes into ecological research. *Front Ecol Environ* **12**: 241–9.
- White LJ and Oates JF. 1999. New data on the history of the plateau forest of Okomu, southern Nigeria: an insight into how human disturbance has shaped the African rain forest. *Glob Ecol Biogeogr* **8**: 355–61.
- Willis KJ, Gillson L, Brncic TM. 2004. How “virgin” is virgin rainforest? *Science* **304**: 402–3.
- [dataset] Wildlife Conservation Society and Center for International Earth Science Information Network, Columbia University. 2005. Last of the Wild Project, Version 2, 2005 (LWP-2): Global Human Footprint Dataset (Geographic). Palisades, NY: NASA Socioeconomic Data and Applications Center (SEDAC). <http://dx.doi.org/10.7927/H4M61H5F>. Accessed 21 July 2018.
- Zalasiewicz, J, Williams M, Steffen W, Crutzen P. 2010. The New World of the Anthropocene. *Environmental Science & Technology* **44**: 2228-2231.