



This is a post-peer-review, pre-copyedit version of an article published in PNAS. The final authenticated version is available online at: <https://doi.org/10.1073/pnas.1819345116>

1 **Title page**

2 **Classification:** Biological Sciences, Plant Biology

3 **Title:** A conserved but plant specific CDK-mediated regulation of DNA
4 replication protein A2 in the precise control of stomatal terminal division*

5 Kezhen Yang^a, Lingling Zhu^{a,b}, Hongzhe Wang^a, Min Jiang^a, Chunwang
6 Xiao^{c,d}, Xiangyang Hu^e, Steffen Vanneste^{f,g,h}, Juan Dongⁱ and Jie Le^{a,b,1}

7 ^a Key Laboratory of Plant Molecular Physiology, CAS Center for Excellence
8 in Molecular Plant Sciences, Institute of Botany, Chinese Academy of
9 Sciences, Beijing 100093, China. ^b University of Chinese Academy of
10 Sciences, Beijing 100049, China. ^c College of Life and Environmental
11 Sciences, Minzu University of China, Beijing 100081, China. ^d Hulun Lake
12 Reserve Grassland Ecology Research Station, Minzu University of China,
13 Beijing 100081, China. ^e Shanghai Key Laboratory of Bio-Energy Crops,
14 School of Life Sciences, Shanghai University, Shanghai 200444, China. ^f VIB
15 Center for Plant Systems Biology, Ghent 9052, Belgium. ^g Department of
16 Plant Biotechnology and Bioinformatics, Ghent University, Ghent 9052,
17 Belgium. ^h Laboratory of Plant Growth Analysis, Ghent University Global
18 Campus, 21985 Incheon, Republic of Korea. ⁱ Waksman Institute of
19 Microbiology, Rutgers, the State University of New Jersey, Piscataway, NJ
20 08854, USA.

21 ¹ To whom correspondence should be addressed. Email: lejie@ibcas.ac.cn

22 * Dedicated to the memory of Fred Sack, who is a pioneer in the field of
23 stomatal development.

24 **Author contributions:** K.Y. and J.L. designed the research and analyzed
25 the data. K.Y. performed most of the experiments. L.Z. and W.H. contributed
26 to the generation of some constructs. M.J. contributed to mutant screen. K.Y.,

27 J.L., J.D., X.H., S.V., and C.X. wrote the article.

28 **Abstract:**

29 The R2R3-MYB transcription factor FOUR LIPS (FLP) controls the stomatal
30 terminal division through transcriptional repression of the cell cycle genes
31 *CDKB1s*, *CDKA;1*, and *CYCA2s*. We mutagenized the weak mutant allele
32 *flp-1* seeds with ethylmethane sulfonate (EMS) and screened out a *flp-1*
33 suppressor 1 (*fsp1*) that suppressed the *flp-1* stomatal cluster phenotype.
34 *FSP1* encodes RPA2a subunit of Replication Protein A (RPA) complexes that
35 play important roles in DNA replication, recombination, and repair. Here, we
36 show that *FSP1/RPA2a* functions together with *CDKB1s* and *CYCA2s* in
37 restricting stomatal precursor proliferation, ensuring the stomatal terminal
38 division, and maintaining a normal guard cell size and DNA content.
39 Furthermore, we provide direct evidence for the existence of an evolutionally
40 conserved, but plant-specific CDK-mediated RPA regulatory pathway.
41 Serine-11 and Serine-21 at the N-terminus of RPA2a are CDK
42 phosphorylation target residues. The expression of the
43 phosphorylation-mimic variant *RPA2a^{S11,21D}* partially complemented the
44 defective cell division and DNA damage hypersensitivity in *cdkb1;1 1;2*
45 mutants. Thus, our study provides a mechanistic understanding of the
46 CDK-mediated phosphorylation of RPA in the precise control of cell cycle and
47 DNA repair in plants.

48 **Keywords:** Stomatal Development; Cell Division; Replication Protein A;
49 CDK; DNA Damage

50 **Significance:**

51 The Arabidopsis R2R3-MYB transcription factor FOUR LIPS (FLP) is the first
52 identified key transcription factor regulating stomatal development. By
53 screening and analysis of a genetic suppressor of *flp* stomatal defects, we

54 found *FSP1/RPA2a*, encoding a core subunit of Replication Protein A (RPA)
55 complexes, acts downstream of core cell cycle genes of CDKB1s in ensuring
56 the terminal division during stomatal development to produce a pair of
57 kidney-shaped guard cells to compose a functional stomatal complex. We
58 demonstrate that the CDK-mediated phosphorylation at the N-terminus of
59 RPA2a is essential for the RPA functions in cell cycle control and response to
60 DNA damage. We provide the direct evidence for the existence of an
61 evolutionally conserved, but plant-specific RPA regulatory pathway in plants.

62 \body

63 **Introduction**

64 Stomata are plant-specific epidermal structures that consist of a pair of
65 Guard Cells (GCs) surrounding a pore. The formation of stomata requires
66 successive asymmetric cell division of the precursor cells, including the
67 Meristemoid Mother Cell (MMC) and the Meristemoid (M), and one
68 symmetric division of the Guard Mother Cell (GMC) to produce two GCs (1).
69 *FOUR LIPS (FLP)* and *MYB88* encode R2R3-MYB transcription factors and
70 function in the regulation of symmetric division of the GMCs. In a weak allele
71 *flp-1*, two stomata form abnormally in direct contact. The loss of *MYB88*
72 function dramatically enhances the phenotype of *flp* mutants, leading to
73 tumor-like stomatal clusters (2, 3). The cell cycle genes *CDKB1s*, *CYCA2s*,
74 and *CDKA;1*, as transcriptional targets, are directly suppressed by FLP and
75 MYB88 (4-6).

76 Replication Protein A (RPA) is a heterotrimeric single-stranded DNA
77 (ssDNA)-binding protein complex that is required for multiple aspects of DNA
78 metabolism, including DNA replication, recombination, repair, and telomere
79 maintenance (7). The homologues of each of the three RPA subunits
80 (RPA1-3) are well conserved in eukaryotes. In humans, phosphorylation of

81 RPA2 at the N-terminal domain is required for the RPA-ssDNA interaction. In
82 mitotic cells, Serine-23 and Serine-29 at the RPA2 N-terminus are
83 phosphorylated and activated by Cdc2/CDK to promote DNA replication
84 (8-12). Upon a DNA damaging condition, RPA2 is hyperphosphorylated by
85 the PIKK-family kinases (ATM, ATR, and DNA-PK) that facilitates mitotic exit
86 and the initiation of DNA repairing (13-15). All known RPA2 homologues
87 have a conserved N-terminal phosphorylation domain, though the specific
88 residues may be not conserved in different species (11). In contrast to yeast
89 and most mammals, plants carry multiple paralogs for each of the RPA
90 subunit (16). For instance, rice has three RPA1s, three RPA2s, and one
91 RPA3 (16, 17). The model plant Arabidopsis has five RPA1s (RPA1a-e), two
92 RPA2s (RPA2a, b), and two RPA3s (RPA3a, b). Phylogenetic analysis of the
93 RPA1 sequences suggests that Arabidopsis RPA1s diverged into two
94 subgroups, the ACE-group (RPA1a, b, c) and the BD-group (RPA1b, d) (18).
95 Previously, genetic analysis confirmed that *RPA2a* plays a critical role in the
96 maintenance of epigenetic gene silencing in plants and abiotic stresses
97 (19-21).

98 In this study, with an aim of obtaining genetic suppressors of *flp-1*, we
99 identified a genetic mutation in Arabidopsis *RPA2a* that led to inhibited
100 stomatal clustering in *flp-1* and arrested GMC divisions. Our study
101 discovered the existence of an evolutionally conserved, but plant-specific
102 CDK-mediating RPA regulatory pathway. Also, by assaying the stomatal
103 development and DNA damage responses, we established the physical and
104 genetic interactions between the RPA and CDKs in the precise control of cell
105 cycle as well DNA repair in plants.

106 **Results**

107 **Isolation of *fsp1*, a Suppressor of *flp-1* in Stomatal Development.** The

108 *flp-1* mutant is featured by extra terminal divisions during stomatal
109 development, suggesting the role of FLP in restricting cell division (3). To
110 identify new genetic players in regulating the one-time terminal division, we
111 created an ethylmethane sulfonate (EMS) mutagenized M2 population of
112 *flp-1* mutants and screened for mutants with altered stomatal phenotypes.
113 *flp-1 fsp1* (*flp-1* suppressor1) was isolated for significantly reduced stomatal
114 clusters, compared to *flp-1* (Fig.1 A-C and *SI Appendix*, Table S1). In
115 addition, aberrant cells were occasionally found in the epidermis of *flp-1 fsp1*
116 double mutant (Fig. 1C). Using a GC fate marker E1728, we confirmed that
117 the aberrantly shaped, single cells in *fsp1* have the GC identity (Fig. 1D and
118 inset), resembling the Single Guard Cell (SGC) phenotype in *cdkb1;1 1;2*
119 and *cyca2s* mutants (4,5). The Stomatal Index (SI) of *fsp1* mutant was
120 reduced as well, indicating that *FSP1* promotes stomatal production (*SI*
121 *Appendix*, Table S2).

122 ***FSP1* Encodes Arabidopsis RPA2a Subunit of the RPA Complex.** Using
123 map-based cloning, we determined that mutation of *FSP1* gene was located
124 within the BAC clone T28124 on chromosome 2. Based on growth defects
125 reminiscent of *rpa2a/ror1-1* mutants, such as dwarf seedlings, narrow leaves,
126 and early flowering (*SI Appendix*, Fig. S1A) (19, 20), we amplified and
127 sequenced the open reading frames of *RPA2a* from *fsp1* and found that *fsp1*
128 possessed the same point mutation as in *ror1-2*, ag (1343) changed to aa,
129 which induces altered splicing events (Fig. 1E). The expression of *RPA2a*
130 (cDNA) fused with GFP under the control of the native promoter fully rescued
131 the stomatal defects of *fsp1* mutants (Fig. 1F and *SI Appendix*, Table S2).
132 The functionality of *RPA2a* was further confirmed by the reappearance of
133 *flp-1*-featured stomatal clusters in *flp-1 fsp1* mutants carrying
134 *RPA2a:RPA2a-GFP* (Fig. 1G and *SI Appendix*, Table S1). In addition, the
135 introgression of the *fsp1* mutation in *flp-1 myb88* double mutants largely

136 repressed the size and number of stomatal clusters (Fig. 1 *H,I* and *SI*
137 *Appendix*, Fig. S1*B*). We also revisited the T-DNA allele, *rpa2-5/ror1-3* (Fig.
138 1*E* and *SI Appendix*, Fig. S1*A*) (19, 20), which displayed a similar stomatal
139 phenotype as *fsp1*. The *flp-1* stomatal phenotype was suppressed in *flp-1*
140 *rpa2-5* double mutants (*SI Appendix*, Fig. S1 *C,D* and Table S2). Benefiting
141 from easy genotyping, *rpa2-5* was then used in most of the later experiments
142 (renamed as *rpa2a-5*).

143 ***RPA2a* Is Expressed in Specific Stomatal Lineage Stages. A**

144 complemented *ror1-1* transgenic line harboring *ROR1:gROR1-GUS-GFP* (19)
145 (here referred as *RPA2a-GFP*) was used to investigate the *RPA2a*
146 expression pattern and localization. *RPA2a-GFP* fluorescent signals were
147 observed in a subset of stomatal lineage cells and compared with those of
148 three translational reporters of *SPEECHLESS* (*SPCH*), *MUTE*, and *FLP*,
149 which have distinct and sequential expression patterns during stomatal
150 development (Fig. 1*J*). The expression of *RPA2a-GFP* overlapped with
151 *SPCH-GFP* (22) in MMCs prior to asymmetric entry divisions but not in both
152 newly-formed meristemoids (M) and Stomatal Lineage Ground Cells (SLGC)
153 after division. In late meristemoids and early GMCs (EGMCs), where
154 *MUTE-GFP* was turned on (23), diffuse *RPA2a-GFP* signals reappeared.
155 *RPA2a-GFP* persisted in late GMCs (LGMCs), but disappeared in young
156 GCs (YGCs) after the terminal symmetric cell division. The expression of
157 *RPA2* overlapped with *FLP-GFP* (3, 24) only at LGMC stage, prior to the
158 GMC division, but not in YGCs. Taken together, *RPA2a* shows a cell type-
159 and time-specific expression profile, with the preferences in the precursor
160 cells (actively dividing) prior to either asymmetric or symmetric divisions
161 during stomatal development.

162 ***RPA2a* Functions Together with *FLP* Downstream Target Genes**

163 ***CDKB1s* and *CYCA2s*.** *CDKB1;1* gene is one of the direct transcriptional

164 targets of FLP in controlling stomatal terminal symmetric divisions. In
165 agreement with this, expression of *CDKB1;1-GFP* (24) overlaps with
166 *FLP-GFP* in LGMCs and YGCs while partially with *RPA2a-GFP* in LGMCs
167 (Fig. 1J). In comparison to SGC frequencies of $42.0 \pm 2.6\%$ in *cdkb1;1 1;2*
168 and $8.0 \pm 2.1\%$ in *rpa2a-5*, the *rpa2a-5 cdkb1;1 1;2* triple mutant produced
169 dramatically more SGCs ($64.0 \pm 1.6\%$, Fig. 2A-D). GMC divisions also
170 require *CYCA2* activity (5). While SGCs in *cyca2;3* mutants was found at a
171 low frequency of $4.0 \pm 0.8\%$ (Fig. 2E), the occurrence of SGCs in *rpa2a-5*
172 *cyca2;3* double mutants was elevated to $47.0 \pm 2.0\%$ (Fig. 2F). Consistently,
173 the frequency of SGCs in *cyca2;3 2;4* ($27.6 \pm 3.4\%$) was greatly elevated to
174 $78.0 \pm 8.5\%$ in *rpa2a-5 cyca2;3 2;4* triple mutants (Fig. 2 G and H), indicating
175 interaction between RPA2a and CDKB1s/CYCA2s functions in GMC
176 divisions.

177 An ectopic and prolonged expression of RPA2a-GFP was occasionally found
178 in two daughter cells after a GMC division in *flp-1*, suggesting that RPA2a is
179 required for subsequent GMC divisions that end up with two stomata next to
180 each other (*SI Appendix*, Fig. S2 A-H). *RPA2a* transcription level slightly but
181 not significantly increased in *flp-1 myb88* (*SI Appendix*, Fig. S2I). Although
182 12 putative FLP/MYB88 binding sequences are present within the *RPA2a*
183 promoter (4-6), neither FLP nor MYB88 showed clear binding activity to
184 *RPA2a* promoter according to the results of yeast one-hybrid assays (*SI*
185 *Appendix*, Fig. S2 J and K), indicating that *RPA2a* might not be a direct
186 transcriptional target of FLP/MYB88.

187 **Combined Loss of *RPA2a* and *CDKB1s/CYCA2s* Function Induces**
188 **Abnormal Cell Enlargement and Endoreduplication.** Differing from the
189 interlocking, continuously expanding pavement cells, the mature stomatal
190 GCs once matured remain their shape and size for the life time in leaves.
191 The area of the paired GCs mostly ranged from 200 to 500 μm^2 in

192 Arabidopsis (*SI Appendix*, Fig. S3). However, about 45% and 71% stomata
193 in *rpa2a-5 cdkb1;1 1;2* and *rpa2a-5 cyca2;3 2;4* triple mutants, respectively,
194 are larger than $500 \mu\text{m}^2$ (Fig. 2 *A-H* and *SI Appendix*, Fig. S3). It has been
195 widely accepted that cell size and differentiation are closely correlated to the
196 nuclear DNA levels (25). We therefore measured the nuclear DNA levels in
197 SGCs after DAPI (4',6-diamidino-2-phenylindole) staining. The DNA content
198 in wild-type GCs was defined as 2C-DNA level. Our results showed that
199 SGCs in the *rpa2a-5*, *cdkb1;1 1;2*, and *cyca2;3 2;4* mutants all displayed a
200 similar DAPI fluorescence intensity (4C-DNA level), indicating that *RPA2a*,
201 similar to *CDKB1s* and *CYCA2s*, is required for the mitosis of GMC divisions.
202 But in the *rpa2a-5 cdkb1;1 1;2* or *rpa2a-5 cyca2;3 2;4* triple mutants, the
203 enlarged SGCs contained single giant nuclei with a mean DNA level of
204 6C-8C (Fig. 2 *I-N*). The analysis of DNA ploidy distribution in 9- and
205 14-day-old cotyledons using flow cytometry confirmed that *RPA2a* may
206 coordinately act with *CDKB1s*/*CYCA2s* to restrict the onset of endocycle (*SI*
207 *Appendix*, Fig. S4).

208 Above results indicated that the mitosis was blocked but the endoreplication
209 was inappropriately unrestricted in the *rpa2a-5 cdkb1;1 1;2* triple mutants. In
210 *rpa2a-5* single mutant, the expression levels of G1-to-S genes *CDT1*, *MCM2*,
211 *ORC3*, and *PCNA1* were significantly increased comparing those in wild type.
212 Most of these elevations as well as *CDC6a*, *CCS52A*, *ORC1a*, *CDKA;1*,
213 were additively enhanced in *rpa2a-5 cdkb1;1 1;2* triple mutants (*SI Appendix*,
214 Fig. S5A). In addition, the expression levels of genes during endoreplication
215 initiation, such as *E2Fa*, *E2Fc*, and *E2Ff*, were largely increased in the triple
216 mutants (*SI Appendix*, Fig. S5B). These results demonstrated that *RPA2a*
217 and *CDKB1s* act together to suppress the expression of endocycling genes
218 that restrict endoreduplication.

219 **RPA2a Function Is Conservatively Regulated by CDKs through Protein**

220 **Phosphorylation.** Supported by our genetic data above, we tested the
221 hypothesis that RPA2a may directly interact with CDKs in regulating cell
222 division and cell cycle. Results of Bimolecular Fluorescent Complementation
223 (BiFC) assays using the tobacco transient expression system and
224 Co-ImmunoPrecipitation (Co-IP) assay using the proteins that were extracted
225 from the infiltrated tobacco leaves suggested that RPA2a directly interacts
226 with CDKB1;1 and CDKA;1 (Fig. 3 A-D and *SI Appendix*, Fig. S6).

227 It was reported that coexpression of the RPA2 subunit with other RPA
228 subunits could enhance the solubility and function of RPA protein complexes
229 (26). When RPA2a-GST was coexpressed with RPA1b-HIS in *E. coli*, RPA1b
230 proteins failed to be pulled down by RPA2a. Only when the RPA3a was
231 present, in which RPA2a-GST-RPA3a-S coexpressed with RPA1b-HIS, the
232 whole RPA complex could be pulled down by GST beads, indicating that the
233 pre-formation of RPA2-RPA3 subcomplex is essential for the assembly of
234 RPA complex (*SI Appendix*, Fig. S7A). However, regardless the presence of
235 RPA3 and RPA1, RPA2a proteins could be phosphorylated *in vitro* by active
236 CDK-CYCLIN protein complexes (27) (Fig. 3 E,F and *SI Appendix*, Fig. S7B).
237 Additionally, a faint delayed band next to the RPA2a-GST was observed
238 when the products of CDK phosphorylation reaction were separated on
239 SDS-PAGE gels (Fig. 3G). Once alkaline phosphatases were added to the
240 products, the delayed band disappeared, indicating the phosphorylation is
241 the cause of the slower mobility of RPA2a-GST proteins in gels (*SI Appendix*,
242 Fig. S7B). Elimination of the N-terminal 32 amino acids abolished the
243 delayed band, suggesting that this region contains the primary
244 phosphorylation target sites (Fig. 3G).

245 To predict the CDK-phosphorylation target residues in RPA2a protein, the
246 sequence of the N-terminus of RPA2a protein was compared with those in
247 human, yeast, and rice RPA2 proteins. Among the 12 Serine residues within

248 the first 32 amino acids of RPA2a N-terminus, Ser-11 is the consensus CDK
249 phosphorylation sites, corresponding to Ser-23 of human RPA2 protein.
250 Ser-21 is another predicted phosphorylation target site of CDKs
251 (www.cbs.dtu.dk/services/) that maybe conserved in plants (*SI Appendix*, Fig.
252 S7 C-F). When Ser-11 and Ser-21 were substituted with
253 non-phosphorylatable Glycines, the extent of phosphorylated band was
254 reduced (Fig. 3G), indicating that these two residues are predominant
255 CDK-phosphorylation targets, but other sites in RPA2a might get
256 phosphorylated *in vitro* as well.

257 To assess the function of phosphorylation sites in the RPA2a protein *in planta*,
258 RPA2a variants were generated and introduced into *rpa2a-5* mutant to
259 evaluate their ability to rescue stomatal defects. *RPA2a-GFP*, and individual
260 variants with single mutation, *RPA2a^{S11/G}-GFP* and *RPA2a^{S21/G}-GFP*, could
261 fully rescue the stomatal production and GMC division defects of *rpa2a-5*
262 mutant (*SI Appendix*, Table S2). By contrast, further mutation of
263 *RPA2a^{S11,21/G}-GFP* or N-terminus deletion variant *RPA2a^{Δ32}-GFP* showed a
264 reduced rescuing ability (Fig. 4 A, B and *SI Appendix*, Table S2). However, a
265 phosphorylation-mimic variant *RPA2a:RPA2a^{S11,21/D}-GFP* (Ser-11 and Ser-21
266 were substituted with Aspartic acids) rescued *rpa2a-5* GMC division defects
267 (Fig. 4C and *SI Appendix*, Table S2).

268 We further examined how RPA2a phosphorylation-variants perform in *flp-1*
269 *fsp1* double mutants. While, the wild-type RPA2a-GFP fully complemented
270 the loss function of *RPA2a* and restored the *flp-1*-like stomatal clustering, the
271 expression *RPA2a:RPA2a^{Δ32}-GFP* and *RPA2a:RPA2a^{S11,21/G}-GFP* led to a
272 mild stomatal clustering phenotype (Fig. 4 D and E and *SI Appendix*, Table
273 S1). *RPA2a^{S11/G}-GFP* and *RPA2a^{S21/G}-GFP* fully complemented stomatal
274 phenotype of *flp-1 fsp1* (*SI Appendix*, Table S1). Therefore, the

275 phosphorylation status of Ser-11 and Ser-21 within the N-terminal 32
276 residues is critical for the full function of RPA2a.

277 We introduced *RPA2a-GFP*, *RPA2a:RPA2a^{S11/D}-GFP*, *RPA2a^{S21/D}-GFP*, and
278 *RPA2a^{S11,21/D}-GFP* into *cdkb1;1 1;2* double mutants to assess the genetic
279 interaction between *RPA2a* and *CDKB1s* *in vivo*. *RPA2a^{S11,21/D}-GFP* showed
280 strong effects in reducing the formation of SGCs in *cdkb1;1 1;2*, while the
281 remaining three transgenes had certain effects (Fig. 4 *F-H* and *SI Appendix*,
282 Table S3). Therefore, these results suggest that the phosphorylation status
283 of RPA2a is epistatic to *CDKB1* mutations.

284 **CDKB1 Activity Promotes RPA2a Nuclear Localization.** The strong
285 expression of complementary RPA2a-GFP was mainly found in the nucleus,
286 but also present in the cytoplasm. RPA2a-GFP signals were more diffuse in
287 *cdkb1;1 1;2* epidermal cells (Fig. 5 *A,B* and *SI Appendix*, Fig. S8 *A* and *B*).
288 Western blot analysis confirmed the abundance of nuclear RPA2a-GFP
289 proteins was decreased in *cdkb1;1 1;2* (Fig. 5C). *RPA2a^{Δ32}-GFP* and the
290 phosphorylation-deficient version *RPA2a^{S11,21/G}-GFP* displayed a
291 pronounced fluorescent signals in the cytoplasm of epidermal cells (Fig. 5 *D*,
292 *E* and *SI Appendix*, Fig. S8 *C* and *D*). By contrast, the signals of
293 *RPA2a^{S11,21/D}-GFP*, which fully complemented the stomatal defects of
294 *rpa2a-5*, displayed a high degree of colocalization with DAPI in the nucleus,
295 with a Pearson's-Rr value of 0.89 (Fig. 5*F* and *SI Appendix*, Fig. S8*E*).
296 Quantitative analysis of relative GFP fluorescent intensity in the nucleus
297 consistently supported that CDK-mediated phosphorylation of RPA2a
298 promotes the nuclear accumulation of RPA2a (*SI Appendix*, Fig. S8*K*).

299 We fused the Nuclear Localization Sequence (NLS) or Nuclear Export Signal
300 (NES) to the C-terminus of *CDKB1;1* and co-expressed with RPA2a-mCherry
301 in tobacco leaves. A Non-Nuclear Localization Sequence (NNLS, a control

302 sequence to the NLS) and Non-Nuclear Export Signal (NNES, a control
303 sequence to the NES) (28) were also fused with CDKB1;1 as controls. When
304 individually expressed, RPA2a-mCherry remained in nucleus and cytoplasm
305 (*SI Appendix*, Fig. S9A). However, RPA2a-mCherry signals predominantly
306 retained in the nucleus when co-expressed with CDKB1;1-NLS-GFP (*SI*
307 *Appendix*, Fig. S9B). By contrast, when co-expressed with
308 CDKB1;1-NES-GFP, besides the nucleus, abundant RPA2a-mCherry signals
309 were accumulated in the cytoplasm (*SI Appendix* Fig. S9C).
310 CDKB1;1-NNLS-GFP or CDKB1;1-NNES-GFP did not change the
311 subcellular localization pattern of RPA2a-mCherry (*SI Appendix* Fig. S9 D
312 and E). Thus, the physical interaction with CDKB1;1 either in the nucleus or
313 in the cytoplasm seems to affect the subcellular localization of RPA2a.

314 To further clarify whether the subcellular localization is associated with the
315 function of RPA2a, we generated NLS-fused *RPA2a:RPA2a-NLS-GFP*, and
316 NES-fused *RPA2a:RPA2a-NES-GFP* and transformed into *rpa2a-5* mutant.
317 The fluorescent signals of RPA2a-NLS-GFP were exclusively accumulated in
318 the nuclei (Fig. 5H and *SI Appendix*, Fig. S8 G and L), in which the defective
319 overall growth and GMC division of *rpa2a-5* mutants were fully recovered
320 (Fig. 5L and *SI Appendix*, Table S4). However, a transgenic line carrying
321 *RPA2a-NES-GFP* line #12, in which the diffuse GFP signals were detected in
322 both cytoplasm and nucleus (Fig. 5J and *SI Appendix*, Fig. S8 I and L), the
323 growth and stomatal defects in *rpa2a-5* were partially rescued (Fig. 5L and *SI*
324 *Appendix*, Table S4). Whereas, *RPA2a-NES-GFP* transgenic line #5, in
325 which GFP signals were mostly restricted to the cytoplasm, showing a barely
326 colocalization with DAPI in the nucleus, a representative low value of
327 Pearson's-Rr at -0.23 (Fig. 5K and *SI Appendix*, Fig. S8J and L). Neither the
328 plant growth defect nor the stomatal defect of *rpa2a-5* mutant was rescued in
329 line #5 (Fig. 5L and *SI Appendix*, Table S4). The high expression level of

330 *RPA2a-NES-GFP* in line #5 (*SI Appendix*, Fig. S10) might provide higher
331 nuclear export signals (from the NES), which drive RPA2a proteins into the
332 cytoplasm against the endogenous nuclear 'stay force', such as being
333 phosphorylated by CDKs. The control lines *RPA2a-NNLS-GFP* and
334 *RPA2a-NNES-GFP* phenocopied the protein localization and
335 complementation ability of the wild-type RPA2a-GFP (Fig. 5 G, I, L and *SI*
336 *Appendix*, Fig. S8 F, H and Table S4).

337 **The N-terminal Domain Is also Required for RPA2a's Function in**
338 **Response to DNA Damage.** It has been reported that Arabidopsis *rpa2a*
339 alleles exhibited hypersensitivity to the genotoxic agent Methyl Methane
340 Sulfonate (MMS) treatment (19-21). However, how the RPA2a protein and its
341 subdomains are related to the DNA damage responses in plants remains
342 unknown. The 5-day MMS treatment generally arrested plant growth and
343 induced seedling chlorosis. The introduction of *RPA2a-GFP* or
344 *RPA2a^{S11,21/D}-GFP* recovered *rpa2a-5* growth to the wild-type level. However,
345 the N-terminal defectives *RPA2a^{Δ32}-GFP* and phosphorylation-deficient
346 *RPA2a^{S11,21/G}-GFP* only partially complemented the chlorosis phenotype of
347 *rpa2a-5* (*SI Appendix*, Fig. S11). Furthermore, *cdkb1;1 1;2* and *rpa2a-5*
348 *cdkb1;1 1;2* displayed hypersensitivity to MMS treatment as well. The
349 phosphorylation-mimic variant *RPA2a^{S11,21/D}* alleviated the sensitivity of
350 *cdkb1;1 1;2* to MMS (Fig. 6 A,B and *SI Appendix*, Fig. S12), that is consistent
351 with its stronger capability than wild-type RPA2a in rescuing *cdkb1;1 1;2*
352 GMC divisions (*SI Appendix*, Table S3). Therefore, our finding observation is
353 consistently with the previous findings in yeast and human that the
354 phosphorylation of N-terminus is important for RPA2 function in response to
355 DNA damage.

356 **Discussion**

357 In this study, we found that a genetic mutation in Arabidopsis *RPA2a* not only
358 suppressed *flp-1* stomatal clustering phenotype but also induced the
359 formation of SGCs. *RPA2a* functions together with *CDKB1s* and *CYCA2s* in
360 ensuring the stomatal lineage cell divisions and maintaining a normal DNA
361 content and guard cell size. Furthermore, we found that Ser-11 and Ser-21 of
362 Arabidopsis *RPA2a* are consensus sites for CDK phosphorylation. Being
363 phosphorylated by CDKs at these sites is tightly associated with the *RPA2a*
364 subcellular localization and function in cell cycle control and the response to
365 DNA damage.

366 ***RPA2a* and Core Cell Cycle Genes Coordinately Control Cell Cycle**

367 **Procession.** FLP and MYB88 directly repress the transcription of several
368 core cell cycle genes, such as *CDKB1;1*, *CYCA2;3*, and *CDKA;1*, to ensure
369 that only two GCs are produced during the terminal GMC divisions (2-6). FLP,
370 as an atypical R2R3-MYB transcription factor, recognizes downstream
371 targets through a *cis*-regulatory element that overlaps with E2Fs (4, 33, 34).
372 There are 12 putative FLP/MYB88/E2F binding sequences within the *RPA2a*
373 promoter. But, neither FLP nor MYB88 displayed a clear binding activity to
374 the *RPA2a* promoter (*SI Appendix*, Fig. S2). It was proposed that rice
375 *OsRPA1* and *OsRPA2* might be the target genes of the S-phase transcription
376 factor E2Fs (35). It will be interesting to test whether FLP and MYB88
377 compete with E2Fs in regulating *RPA2a* transcription.

378 RPA stabilizes ssDNA and interacts with replication proteins at the initiation
379 site of DNA replication to facilitate the formation of replication forks (12, 36).
380 CDT1 and CDC6 are DNA replication licensing components that help the
381 loading of the MCM DNA helicases to the DNA replication origins. Similar to
382 human Cdt1 proteins, the Arabidopsis CDT1 is degraded through a
383 CDK-mediated phosphorylation. *CDT1* and *CDC6* are also regulated by E2F
384 transcription factors (37). In *rpa2a cdkb1;1 1;2* or *rpa2a cyca2;3 2;4*,

385 transcription levels of a set of G1-to-S phase genes, including *CDT1*, *CDC6*,
386 *E2F*-family genes are significantly upregulated, indicating that *RPA2a* and
387 core cell cycle genes coordinately restrict the G1-to-S transition and prevent
388 the entry of endocycle. Recently, the roles of *CYCD5;1* and *CYCD7;1* in
389 regulating GMC division and differentiation have been elegantly described
390 (38, 39). However, the expression of *CYCD5;1* and *CYCD7;1* was not altered
391 either in *rpa2a-5*, *cdkb1;1 1;2* or in *rpa2a-5 cdkb1;1 1;2* triple mutants (*SI*
392 *Appendix*, Fig. S5C).

393 **The Functions of RPA2a Are Regulated by CDKs in Plant-specific**
394 **Manner.** RPA2 proteins get phosphorylated during the G1-to-S transition
395 and M-phase, dephosphorylated at the end of M-phase (8, 29-31). In yeast, a
396 single CDK (*cdc2*) regulates DNA replication licensing, DNA replication
397 initiation, and cell mitosis by changing its activity levels from low, to
398 intermediate and high, respectively (38, 40). CDK-mediated RPA2
399 phosphorylation at Ser-23 occurs during S-phase while phosphorylation at
400 Ser-29 happens only during mitosis (12). In Arabidopsis, generally, CDKA;1
401 activity is more important for the G1-to-S transition, CDKB1s is required for
402 the G2-to-M progression (32). Here, we found that RPA2a proteins could be
403 phosphorylated by CDKA;1 *in vitro* as well, suggesting that RPA2a might be
404 regulated by both A1- and B1-type CDKs in Arabidopsis. Therefore, in
405 contrast to the single CDK in yeast, the divergent functions of RPA2a from
406 DNA replication to mitosis might be regulated by both A1-type CDKA;1 and
407 the plant specific B1-type CDKB1s (Fig. 6C).

408 **RPA2a and CDK-CYCLIN Are Involved in DNA Repair.** DNA
409 Double-Strand Breaks (DSBs) also can occur during DNA replication.
410 Arabidopsis *atm* mutant displays chromosomal fragmentation and reduced
411 fertility (41). SUPPRESSOR OF GAMMA RESPONSE 1 (SOG1) is a
412 functional analogue of mammalian tumour suppressor p53 in plants, which is

413 directly phosphorylated and activated by ATM upon DNA damage (42, 43).
414 The activated SOG1 induces an increase of *CYCB1;1* transcript (44). In
415 *rpa2a-5*, the transcription level of *CYCB1;1*, but not *CYCA2;3*, was
416 dramatically upregulated (*SI Appendix*, Fig. 12B), suggesting the existing of
417 a regulatory circuit during DNA repair.

418 **Materials and Methods**

419 Details on plant growth conditions, positional cloning, plasmid construction,
420 flow cytometric analysis, protein-protein interaction BiFC and Co-IP assays
421 are provided in *SI Appendix, Materials and Methods*.

422 **ACKNOWLEDGMENTS.** We thank Fred Sack for providing the *flp* and
423 *CDKB1;1-GFP* seeds, Dominique Bergmann for *SPCH* and *MUTE* markers,
424 Arp Schnittger for the plasmids of CDKA;1 and CDKB1;1 kinase complexes,
425 Daniela Kobbe for the plasmids for RPA complexes, Zhizhong Gong for *ror1*
426 and *ROR1-GFP* seeds, Chongyi Xu for helps in Co-IP experiments and Lu
427 Wang for editing. This work was supported by grants from the National
428 Natural Science Foundation of China program 31071198 (K.Y.), 30971652
429 and 31271463 (J.L.), 31470348 (X.H.), and a Cooperative Programme
430 between National Natural Science Foundation of China (NSFC) - Research
431 Foundation Flanders (FWO) to J.L. and S.V. The research projects in J.D.'s
432 lab are supported by the NIH R01GM109080.

433 **Figure legends:**

434 Fig. 1. *FSP1* encodes RPA2a subunit required for stomatal precursor cell
435 divisions. (A-D) DIC images of epidermis of wild-type (A), *flp-1* (B), *flp-1*
436 *fsp1*(C), and *fsp1* (D). Brackets indicate the stomatal clusters. SGCs are
437 highlighted with yellow color and normal GCs with blue color. Inset,
438 expression of the mature GC marker E1728 in a SGC. (E) Map-based
439 isolation and gene structure of *FSP1*. The mutation site in *fsp1* is the same
440 as *ror1-2*. *rpa2a-5* (*ror1-3*) is a T-DNA insertion line. (F and G)

441 *RPA2a:RPA2a-GFP* rescued *fsp1* SGC phenotype (F), while induced the
442 reappearance of *flp-1* clusters in *flp-1 fsp1* (G). (H and I) *flp-1 myb88* cluster
443 numbers and size are repressed by *fsp1* mutation. (J) *RPA2a-GFP*
444 expression profile in the stomatal lineage cells. RPA2a-GFP (Scale bars: 20
445 μm).

446 Fig. 2. RPA2a function synergistically with CYCA2s and CDKB1s in GMC
447 division. (A-G) Micrographs of epidermis of wild-type (A), *rpa2a-5* (B),
448 *cdkb1;1 1;2* (C), *rpa2a-5 cdkb1;1 1;2* (D), *cyca2;3* (E), *rpa2a-5 cyca2;3* (F),
449 *cyca2;3 2;4* (G), and *rpa2a-5 cyca2;3 2;4* (H). SGCs are highlighted with
450 yellow color and normal GCs with blue color. The numbers at the right-upper
451 corner indicate percentage of SGCs in total stomata. (I-M) DAPI fluorescence
452 micrographs. Red arrowheads indicate the nuclei. (Scale bar: 20 μm).

453 Fig. 3. Phosphorylation of RPA2a by CDK complexes. (A-D) The RPA2a
454 protein binds CDKB1;1 protein in BiFC assays (A-C) and Co-IP assays (D).
455 (Scale bars: 20 μm). (E) *in vitro* phosphorylation assays. RPA2a is
456 phosphorylated by Cak1-CDKA;1-CYCD3;1 complex in the form of
457 RPA1b-RPA2a-RPA3a complex. (F) RPA2a is also phosphorylated by
458 Cak1-CDKB1;1-CYCB1;2 complexes. The bottom box shows the loading
459 controls after Coomassie Brilliant Blue (CBB) gel staining. MBP and GST
460 proteins were used as positive and negative controls, respectively. Red
461 arrows, the RPA2a phosphorylation band in phosphorylation
462 autoradiographs. (G) Phosphorylation of RPA2a by
463 Cak1-CDKB1;1-CYCB1;2 complexes is confirmed by finding a delayed band
464 in 12.5% SDS-PAGE. The extent of phosphorylation of the RPA2a proteins
465 was reduced when Ser-11 and Ser-21 were substituted or was barely
466 detected when the N-terminus was deleted.

467 Fig. 4. Rescue stomatal phenotypes by *RPA2a* variants. (A-C) The formation

468 of SGCs in *rpa2a-5* mutant was partially rescued by *RPA2a*^{Δ32}-GFP
469 construct (A) or *RPA2a*^{S11,21/G}-GFP (B), while was mostly rescued by
470 *RPA2a*^{S11,21/D}-GFP (C). (D and E) The formation of SGCs in *flp-1 fsp1* was
471 barely affected by *RRPA2a*^{Δ32}-GFP (D) or *RPA2a*^{S11,21/G}-GFP (E). (F-H) The
472 formation of SGCs in *cdkb1;1 1;2* mutant is partially rescued by *RPA2a*-GFP
473 (G), but is greatly repressed by *RPA2a*^{S11,21/D}-GFP (H). The numbers at the
474 right-upper corner indicate percentage of SGCs in total stomata. SGCs are
475 highlighted with yellow color and normal GCs with blue color. (Scale bars: 10
476 μm).

477 Fig. 5. Function of RPA2a is associated with its phosphorylation status and
478 subcellular localization. (A) Nuclear expression of RPA2a-GFP in a
479 complemented *rpa2a-5*. (B) Diffuse RPA2a-GFP in *cdkb1;1 1;2*. (C)
480 RPA2a-GFP proteins were found in both the cytoplasm and the nucleus by
481 western blot analysis. Amount of RPA2a-GFP in the nucleus is reduced in
482 *cdkb1;1 1;2*. (D and E) Diffuse GFP signals in *RPA2a*^{Δ32}-GFP (D) and
483 *RPA2a*^{S11,21/G}-GFP (E). (F) Predominant GFP signals in the nucleus of
484 *RPA2a*^{S11,21/D}-GFP. (G-K) The RPA2a subcellular localization in
485 *RPA2a*-NNLS-GFP (G), *RPA2a*-NLS-GFP (H), *RPA2a*-NNES-GFP (I), and
486 two *RPA2a*-NES-GFP transgenic lines, line #12 (J) and line #5 (K). Insets,
487 the enlarged region highlighted by the white dashed-line boxes. (Scale bars:
488 20 μm). (L) The retarded overall growth of *rpa2a-5* was rescued by *RPA2a*
489 variants except *RPA2a*-NES-GFP. (Scale bar: 1 cm).

490 Fig. 6. Phosphorylation-mimic *RPA2a*^{S11,21/D} alleviated the MMS effects in
491 *cdkb1;1 1;2*. (A and B) Seven-day-old seedlings were transferred to liquid
492 MS (A) or liquid MS medium containing 0.01% MMS (B) for 5 days before
493 imaging. Seedlings of *rpa2a-5*, *cdkb1;1 1;2*, and *rpa2a-5 cdkb1;1 1;2*
494 displayed hypersensitivity to MMS. Phosphorylation-mimic *RPA2a*^{S11,21/D}

495 alleviated the MMS effects on *cdkb1;1 1;2*. (Scale bar: 1 cm). (C) Model for
496 RPA2 function in stomatal terminal division regulation and DNA repair
497 progression. T-bars and dashed-lines indicate negative and positive
498 regulation, respectively.

References

1. Bergmann DC & Sack FD (2007) Stomatal development. *Annu Rev Plant Biol* 58(1):163-181.
2. Yang M & Sack FD (1995) The *too many mouths* and *four lips* mutations affect stomatal production in *Arabidopsis*. *Plant Cell* 7(12):2227-2239.
3. Lai LB, *et al.* (2005) The Arabidopsis R2R3 MYB proteins FOUR LIPS and MYB88 restrict divisions late in the stomatal cell lineage. *Plant Cell* 17(10):2754-2767.
4. Xie Z, *et al.* (2010) Regulation of cell proliferation in the stomatal lineage by the Arabidopsis MYB FOUR LIPS via direct targeting of core cell cycle genes. *Plant Cell* 22(7):2306-2321.
5. Vanneste S, *et al.* (2011) Developmental regulation of CYCA2s contributes to tissue-specific proliferation in Arabidopsis. *EMBO J* 30(16):3430-3441.
6. Yang K, *et al.* (2014) Requirement for A-type cyclin-dependent kinase and cyclins for the terminal division in the stomatal lineage of Arabidopsis. *J Exp Bot* 65(9):2449-2461.
7. Wold MS (1997) Replication protein A: a heterotrimeric, single-stranded DNA-binding protein required for eukaryotic DNA metabolism. *Annu Re Biochem* 66:61-92.
8. Dutta A & Stillman B (1992) cdc2 family kinases phosphorylate a human cell DNA replication factor, RPA, and activate DNA replication. *EMBO J* 11(6):2189-2199.
9. Henricksen LA, Carter T, Dutta A, & Wold MS (1996) Phosphorylation of human replication protein A by the DNA-dependent protein kinase is involved in the modulation of DNA replication. *Nucleic Acids Res* 24(15):3107-3112.
10. Niu H, *et al.* (1997) Mapping of amino acid residues in the p34 subunit of human single-stranded DNA-binding protein phosphorylated by DNA-dependent protein kinase and Cdc2 kinase in vitro. *J Biol Chem* 272(19):12634-12641.
11. Binz SK, Lao Y, Lowry DF, & Wold MS (2003) The phosphorylation domain of the 32-kDa subunit of replication protein A (RPA) modulates RPA-DNA interactions. Evidence for an intersubunit interaction. *J Biol Chem* 278(37):35584-35591.
12. Oakley GG & Patrick SM (2010) Replication protein A: directing traffic at the intersection of replication and repair. *Frontiers in bioscience : a journal and virtual library* 15:883-900.
13. Oakley GG, *et al.* (2003) RPA phosphorylation in mitosis alters DNA binding and protein-protein interactions. *Biochemistry* 42(11):3255-3264.
14. Anantha RW, Sokolova E, & Borowiec JA (2008) RPA phosphorylation facilitates mitotic exit in response to mitotic DNA damage. *Proc Natl Acad Sci USA*

- 105(35):12903-12908.
15. Anantha RW & Borowiec JA (2009) Mitotic crisis: the unmasking of a novel role for RPA. *Cell cycle* 8(3):357-361.
 16. Aklilu BB & Culligan KM (2016) Molecular evolution and functional diversification of replication protein A1 in plants. *Front Plant Sci* 7(33):33.
 17. Ishibashi T, Kimura S, & Sakaguchi K (2006) A higher plant has three different types of RPA heterotrimeric complex. *J Biochem* 139(1):99-104.
 18. Aklilu BB, Soderquist RS, & Culligan KM (2014) Genetic analysis of the Replication Protein A large subunit family in Arabidopsis reveals unique and overlapping roles in DNA repair, meiosis and DNA replication. *Nucleic Acids Res* 42(5):3104-3118.
 19. Xia R, *et al.* (2006) ROR1/RPA2A, a putative replication protein A2, functions in epigenetic gene silencing and in regulation of meristem development in Arabidopsis. *Plant Cell* 18(1):85-103.
 20. Elmayan T, Proux F, & Vaucheret H (2005) Arabidopsis RPA2: a genetic link among transcriptional gene silencing, DNA repair, and DNA replication. *Curr Biol* 15(21):1919-1925.
 21. Kapoor A, *et al.* (2005) Mutations in a conserved replication protein suppress transcriptional gene silencing in a DNA-methylation-independent manner in Arabidopsis. *Curr Biol* 15(21):1912-1918.
 22. MacAlister CA, Ohashi-Ito K, & Bergmann DC (2007) Transcription factor control of asymmetric cell divisions that establish the stomatal lineage. *Nature* 445(7127):537-540.
 23. Pillitteri LJ, Sloan DB, Bogenschutz NL, & Torii KU (2007) Termination of asymmetric cell division and differentiation of stomata. *Nature* 445(7127):501-505.
 24. Lee E, Lucas JR, Goodrich J, & Sack FD (2014) Arabidopsis guard cell integrity involves the epigenetic stabilization of the FLP and FAMA transcription factor genes. *Plant J* 78(4):566-577.
 25. Melaragno JE, Mehrotra B, & Coleman AW (1993) Relationship between endopolyploidy and cell-size in epidermal tissue of Arabidopsis. *Plant Cell* 5(11):1661-1668.
 26. Henricksen LA, Umbricht CB, & Wold MS (1994) Recombinant replication protein A: expression, complex formation, and functional characterization. *J Biol Chem* 269(15):11121-11132.
 27. Harashima H & Schnittger A (2012) Robust reconstitution of active cell-cycle control complexes from co-expressed proteins in bacteria. *Plant Methods* 8:9.
 28. Wang R, *et al.* (2016) The Brassinosteroid-Activated BRI1 Receptor Kinase Is Switched off by Dephosphorylation Mediated by Cytoplasm-Localized PP2A B Subunits. *Mol Plant* 9(1):148-157.
 29. Din S, Brill SJ, Fairman MP, & Stillman B (1990) Cell-cycle-regulated phosphorylation of DNA replication factor A from human and yeast cells. *Genes Dev* 4(6):968-977.
 30. Fang F & Newport JW (1993) Distinct roles of cdk2 and cdc2 in RPA phosphorylation during the cell cycle. *J Cell Sci* 106 (Pt 3)(3):983-994.

31. Binz SK, Sheehan AM, & Wold MS (2004) Replication protein A phosphorylation and the cellular response to DNA damage. *DNA Rep* 3(8-9):1015-1024.
32. Boudolf V, *et al.* (2004) B1-type cyclin-dependent kinases are essential for the formation of stomatal complexes in *Arabidopsis thaliana*. *Plant Cell* 16(4):945-955.
33. Magyar Z, *et al.* (2012) Arabidopsis E2FA stimulates proliferation and endocycle separately through RBR-bound and RBR-free complexes. *EMBO J* 31(6):1480-1493.
34. Boudolf V, *et al.* (2004) The plant-specific cyclin-dependent kinase CDKB1;1 and transcription factor E2Fa-DPa control the balance of mitotically dividing and endoreduplicating cells in Arabidopsis. *Plant Cell* 16(10):2683-2692.
35. Marwedel T, *et al.* (2003) Plant-specific regulation of replication protein A2 (OsRPA2) from rice during the cell cycle and in response to ultraviolet light exposure. *Planta* 217(3):457-465.
36. Dornreiter I, *et al.* (1992) Interaction of DNA polymerase alpha-primase with cellular replication protein A and SV40 T antigen. *EMBO J* 11(2):769-776.
37. Castellano MM, Boniotti MB, Caro E, Schnittger A, & Gutierrez C (2004) DNA replication licensing affects cell proliferation or endoreplication in a cell type-specific manner. *Plant Cell* 16(9):2380-2393.
38. Han SK, *et al.* (2018) MUTE directly orchestrates cell-state switch and the single symmetric division to create stomata. *Dev Cell* 45(3):303-315.e305.
39. Weimer AK, *et al.* (2018) Lineage- and stage-specific expressed CYCD7;1 coordinates the single symmetric division that creates stomatal guard cells. *Development* 145(6).
40. Porter AC (2008) Preventing DNA over-replication: a Cdk perspective. *Cell Division* 3(1):3.
41. Garcia V, *et al.* (2003) AtATM is essential for meiosis and the somatic response to DNA damage in plants. *Plant Cell* 15(1):119-132.
42. Ciccia A & Elledge SJ (2010) The DNA damage response: making it safe to play with knives. *Mol Cell* 40(2):179-204.
43. Yoshiyama KO, *et al.* (2013) ATM-mediated phosphorylation of SOG1 is essential for the DNA damage response in Arabidopsis. *EMBO Rep* 14(9):817-822.
44. Weimer AK, *et al.* (2016) The plant-specific CDKB1-CYCB1 complex mediates homologous recombination repair in Arabidopsis. *EMBO J* 35(19):2068-2086.
45. Xie Z, Li D, Wang L, Sack FD, & Grotewold E (2010) Role of the stomatal development regulators FLP/MYB88 in abiotic stress responses. *Plant J* 64(5):731-739.
46. Eschbach V & Kobbe D (2014) Different replication protein A complexes of Arabidopsis thaliana have different DNA-binding properties as a function of heterotrimer composition. *Plant Cell Physiol* 55(8):1460-1472.
47. Weimer AK, *et al.* (2012) Retinoblastoma related1 regulates asymmetric cell divisions in Arabidopsis. *Plant Cell* 24(10):4083-4095.











