

# Can a Tromino be Tiled with Unit Trominoes?

C. Rebecca Thomas, Siân K. Jones and Stephanie Perkins \*

## Abstract

Polominoes are well-known due to their use in the game Tetris, in which shapes made from four squares called tetrominoes are arranged within a game area. Polominoes can be constructed using any number of squares. In this article trominoes, which consist of three squares in an L-shape formation, are examined. We determine whether these can be used to fill larger L-shaped formations.

## 1 Introduction

A tromino (also known as a triomino) is a geometric shape formed by three squares. Trominoes can be either 'I' shaped or 'L' shaped and can be rotated in any orientation. In this article only the 'L' shaped tromino is used and throughout we use the word tromino to mean an 'L' shaped tromino. The possible orientations of the tromino are given in Figure 1.




Figure 1: A unit Tromino and its Rotations

---

**\*About the Authors:** Sian K. Jones and Stephanie Perkins work at the University of South Wales in Pontypridd, UK. Stephanie Perkins is Head of Mathematics and Sian K. Jones is a Lecturer; both conduct research into Combinatorics with interests ranging from Sudoku grids to Information Theory. Rebecca Thomas was an undergraduate studying Mathematics and completed her final year project on Trominoes.

Larger L-shapes can also be formed as shown in Figure 2 and the aim of this article is to show that such shapes can be completely filled with copies of the unit trominoes given in Figure 1. A regular L-shaped tromino can be considered to comprise three  $n \times n$  squares in the arrangement given by the dotted lines in Figure 2.





Figure 2: A Tromino of Size  $n$

The unit trominoes in Figure 1 are denoted  $L_1$  and the larger L-shaped tromino, in Figure 2, of size  $n$  is denoted by  $L_n$ . The aim of this article is to show that  $L_n$  for any integer  $n$  can be tiled completely with unit trominoes ( $L_1$ ).

## 2 Tiling $L_n$ for $n$ a multiple of 2 or 3


For the cases where  $n$  is a multiple of 2 or 3 it can be shown explicitly that a tiling exists. The smallest case  $L_2$  is given in Figure 3(a) using four copies of  $L_1$ .  $L_3$  is given in Figure 3(b) and uses five copies of  $L_1$  and one  $L_2$  (which is itself composed of four copies of  $L_1$ ).



(a) A Tiling of  $L_2$                       (b) A Tiling of  $L_3$

Figure 3: A Tiling of  $L_2$  and  $L_3$

In Figure 4(a) it is demonstrated that the same arrangement as  $L_2$  can be used to tile  $L_n$  when  $n$  is a multiple of 2 ( $n = 2k$ , for  $k$  an integer) using four copies of  $L_k$ . Similarly in Figure 4(b) the same arrangement as  $L_3$  can be used to tile  $L_n$  when  $n$  is a multiple of 3 ( $n = 3k$ , for  $k$  an integer) using five copies of  $L_k$  and one copy of  $L_{2k}$ .



(a) A General  $L_{2k}$                                       (b) A General  $L_{3k}$

Figure 4: A General  $L_{2k}$  and  $L_{3k}$

Therefore if there exists a tiling of a tromino  $L_k$ , of size  $k$ , then there exists a tiling of a tromino  $L_{2k}$ , of

size  $2k$  and there exists a tiling of a tromino  $L_{3k}$  of size  $3k$ .

Hence any  $L_n$  for  $n = 2^x 3^y m$  can be tiled using unit trominoes ( $L_1$ ) if  $L_m$  can be tiled using unit trominoes. Therefore it is sufficient to prove that  $L_m$  can be tiled by  $L_1$  when  $m$  is not a multiple of 2 or 3.

### 3 Tiling of $L_n$ , for $n \geq 5$

Consider  $L_n$  in Figure 5. First tile the squares marked x using an  $L_1$ . The remaining tromino consists of three  $n \times n$  squares with a corner square removed, called **deficient squares**. To give a tiling for a tromino for  $L_n$ , for  $n \geq 5$  and  $n$  not a multiple of 2 or 3, it suffices to show that an  $n \times n$  deficient square (for  $n \geq 5$  and not a multiple of 2 or 3) can be tiled with multiple copies of  $L_1$ .





Figure 5: A Tromino Constructed from Deficient Squares

### 4 Tiling a Deficient $n \times n$ Square

Chu and Johnsonbaugh [?] show that a deficient square of dimension  $n \times n$  can be tiled with unit trominoes if  $n \geq 5$  and 3 does not divide  $n$ . They consider both  $n$  odd and  $n$  even, and with the removed square located anywhere within the  $n \times n$  square. However only the case of  $n$  odd and the corner square removed is of interest in this article and is summarized in this section.


Chu and Johnsonbaugh [?] give an explicit tiling of a deficient  $5 \times 5$  square, an example of which is given in Figure 6(a). They also give a  $2 \times 3$  configuration (as shown in Figure 6(b)), which is used to show that a  $2i \times 3j$ , or  $3i \times 2j$  board can be tiled exactly using unit trominoes (using an  $i \times j$  arrangement of  $2 \times 3$  resp.  $3 \times 2$  configurations).



(a) A  $5 \times 5$  Deficient Board Tiled Using Trominoes      (b) A  $2 \times 3$  Configuration

Figure 6:  $5 \times 5$  and  $2 \times 3$  Configurations

Using the  $5 \times 5$  deficient square and the  $2 \times 3$  and  $3 \times 2$  configurations, the  $7 \times 7$  deficient square can now be tiled (Figure 7(a)).



(a) A  $7 \times 7$  deficient square      (b) A  $n \times n$  deficient square for  $n$  odd

Figure 7: A  $7 \times 7$  and  $n \times n$  (for  $n$  odd) deficient square

A  $n \times n$  deficient square for  $n \geq 11$  and  $n$  odd is given in Figure 7(b) and comprises of four shapes. This is a rearranged version of Figure 9 given in [?] using the same shapes but with a corner square removed. Consider each of these shapes in turn:

- $(n - 7) \times 6$  (and hence also  $6 \times (n - 7)$ ) rectangle can be tiled using the configurations given in Figure

6(b).

- $7 \times 7$  deficient square is given in Figure 7(a).
- $(n - 6) \times (n - 6)$  deficient square, the first case when  $(n - 6) \times (n - 6) = 5 \times 5$  is given in Figure 6(a) and when  $(n - 6) \times (n - 6) = 7 \times 7$  is given in Figure 7(a). Therefore since an  $11 \times 11$  deficient square can be created when  $(n - 6) \times (n - 6) = 5 \times 5$  and  $13 \times 13$  is created using  $(n - 6) \times (n - 6) = 7 \times 7$  then by an inductive argument all remaining cases for  $n = 5 + 6k, 7 + 6k$ , i.e.  $n$  odd and not a multiple of three can be generated.

## 5 A tiling of $L_n$

In Section 2 it was demonstrated that  $L_n$  can be tiled using unit trominoes ( $L_1$ ) for  $n$  a multiple of 2 or 3. The results of Chu and Johnsonbaugh [?] demonstrate that an  $n \times n$  deficient square can be tiled using unit trominoes for  $n \geq 5$ ,  $n$  odd and where 3 does not divide  $n$ . Since the deficient square can be chosen to be a corner square then three such squares plus one copy of  $L_1$  can be arranged as shown in Figure 5 to construct  $L_n$  where  $n \geq 5$ ,  $n$  odd and where 3 does not divide  $n$ . Hence it has been shown that there exists a tiling of  $L_n$ , for any integer  $n$  using unit trominoes.

## References

- [1] I. Chu and R. Johnsonbaugh, *Tiling deficient boards with trominoes*, Mathematics Magazine **59** (1986), no. 1, 34–40.